

Weight-at-length of the invasive lionfish *Pterois volitans* (Actinopterygii, Scorpaenidae) in the Central Mexican Caribbean, and a review of allometric growth parameters across the invasion range

Running title: Length-Weight relationships of lionfish

Abstract

Lionfish (*Pterois volitans/miles*) are an invasive species in the North-Western Atlantic and the Caribbean. To better manage the invasion, inform lionfish removal programs, and estimate biomass available for harvest, we must be able to accurately estimate their total biomass; estimating individual weight is frequently done from length observations. This work compares length--weight relationships of the invasive lionfish through the invasion range and reports the length--weight relationship for lionfish in the Central Mexican Caribbean. A review of 13 length--weight relationships reported in eight peer-reviewed studies and FishBase is provided. These parameters were used to identify spatial variation in weight-at-length. For a given length, parameters from the Caribbean yielded lower weights than those from the Gulf of Mexico and Atlantic, indicating that weight-at-length is spatially variable. This highlights the importance of using site-specific parameters to estimate biomass from length observations. This study also reports a new pair of length-weight parameters ($a = 3.2056 \times 10^{-6}$; $b = 3.235$) for organisms sampled in the Central Mexican Caribbean. Findings from this work can aid managers and decision makers to better select length-weight parameters when these are not available for their region of interest.

Resumen

El pez león (*Pterois volitans/miles*) es una especie invasora en el Atlántico Noroeste y el Mar Caribe. Para tener un mejor manejo de la invasión, informar programas de remoción y estimar la biomasa disponible para aprovechamiento, debemos ser capaces de estimar la biomasa disponible con precisión, frecuentemente a partir de observaciones de tallas. Este trabajo compara la relación longitud--peso del pez león a través del rango de invasión, y reporta un nuevo par de parámetros de crecimiento alométrico para la zona central del Caribe Mexicano. El trabajo presenta revisión de 13 relaciones longitud--peso reportadas en ocho estudios publicados en revistas arbitradas y FishBase. Para una misma talla, los parámetros del Caribe indican un mejor peso que los parámetros del Golfo de México y el Atlántico, indicando una variabilidad espacial en la relación longitud peso. Esta variabilidad resalta la importancia de utilizar parámetros específicos a un sitio al estimar biomasa a partir de observaciones de tallas. Este estudio también reporta una nueva relación longitud--peso ($a = 3.2056 \times 10^{-6}$; $b = 3.235$) para organismos muestreados en la costa central del Caribe Mexicano. Los resultados de este trabajo pueden ayudar a tomadores de decisiones en la selección de relaciones longitud-peso cuando éstas no están disponibles para su zona de interés.

Key words

biometry, invasive species, spatial variability, allometric growth, Mexico

37 Introduction

38 At least 84% of the marine eco-regions have reported the presence of an invasive species (Molnar et
39 al., 2008). These represent a major threat to local biodiversity and the economic activities that
40 depend on it, like tourism or fisheries (Bax et al., 2003). Invasive species may also threaten native
41 species through competition (DAVIS, 2003) or predation. By 2005, the economic cost of invasive
42 species to the United States was estimated at \$120 billion per year and nearly 42% of species that
43 have been included in the Endangered or Threatened species list have been labeled as such due to
44 presence of invasive species (Pimentel, Zuniga & Morrison, 2005). This highlights the importance
45 of understanding, managing, and preventing ecological invasions.

46 Lionfish (*Pterois volitans/miles* complex) are an invasive species in the North-Western Atlantic and
47 the Caribbean, likely introduced through liberation of aquarium-kept organisms (Betancur-R et al.,
48 2011). They are the first marine vertebrates to establish in North Atlantic (Schofield, 2009, 2010)
49 and Caribbean coasts (Sabido-Itza et al., 2016). Lionfish have been widely reported in coral reefs
50 (Aguilar-Perera & Tuz-Sulub, 2010), but also in other habitats such as estuaries (Jud et al., 2011),
51 mangroves (Barbour et al., 2010), areas with hard-bottoms (Muñoz, Currin & Whitfield, 2011), and
52 mesophotic reefs (Andradi-Brown et al., 2017). Due to its threat to local biodiversity, the speed of
53 their spread, and its difficulty of management, their presence in these waters has been labeled as a
54 major marine invasion (Hixon et al., 2016).

55 A significant amount of research has been done to describe lionfish feeding ecology in North
56 Carolina (Muñoz, Currin & Whitfield, 2011), the Bahamas (Morris & Akins, 2009; Cote et al.,
57 2013), Northern Gulf of Mexico (Dahl & Patterson, 2014), Mexican Caribbean (Valdez-Moreno et
58 al., 2012; Villaseñor-Derbez & Herrera-Pérez, 2014), Belize (Hackerott et al., 2017), and Costa
59 Rica (Sandel et al., 2015). Their feeding behavior and high consumption rates can reduce
60 recruitment (Albins & Hixon, 2008) and population sizes (Green et al., 2012) of native reef-fish
61 species, and further the endangerment of critically endangered reef fish (Rocha et al., 2015).
62 (However, see Hackerott et al. (2017) for a case where there was no evidence that lionfish affected
63 the density, richness, or composition of prey fishes). Major efforts have also been made to
64 understand the possible impacts of the invasion by keeping track of its range through time
65 (Schofield, 2009, 2010) and predicting invasion ranges under climate change scenarios (Grieve,
66 Curchitser & Rykaczewski, 2016). By combining information from these disciplines, researchers
67 have been able to predict the trophic impacts of lionfish (Arias-Gonzalez et al., 2011), which can
68 then be translated into ecosystem-level and economic impacts.

69 Seeking to reduce lionfish densities, governments and non-profit organizations have promoted
70 removal programs and incentivized its consumption (Chin, Aiken & Buddo, 2016). In some cases,
71 these have shown to significantly reduce -but not quite eliminate- lionfish abundances at local
72 scales (Sandel et al., 2015, Chin, Aiken & Buddo (2016), de Leon et al. (2013)). The rapid recovery
73 rates exhibited by lionfish (Barbour et al., 2011) and the persistent populations in mesophotic coral
74 ecosystems (Andradi-Brown et al., 2017) -which can contribute with recruitment to shallow-water
75 populations- make of complete eradication through fishing effort an unlikely solution. However,
76 further incentivizing its consumption might create a demand big enough to promote and sustain a
77 stable fishery (Chin, Aiken & Buddo, 2016), which can reduce local abundances and control -not
78 eradicate- the invasion while providing alternative livelihoods.

79 The feasibility of lionfish removal programs has been extensively evaluated through field
80 observations (Sandel et al., 2015, Chin, Aiken & Buddo (2016), de Leon et al. (2013); Usseglio et
81 al., 2017) and empirical modeling (Barbour et al., 2011; Morris, Shertzer & Rice, 2011; Johnston &
82 Purkis, 2015). The latter measure changes in biomass or density (Barbour et al., 2011; Johnston &
83 Purkis, 2015) in response to increased mortality (*i.e.* lionfish removal). In this case, biomass

represents the sum of all fish's individual weight. Total Weight (TW) can be estimated from Total Length (TL) observations using the allometric growth equation ($TW = aTL^b$). Parameters a and b for this equation exist for North Carolina (Barbour et al., 2011), Northern (Fogg et al., 2013) and Southern Gulf of Mexico (Aguilar-Perera & Quijano-Puerto, 2016), the Southern Mexican Caribbean (Sabido-Itza et al., 2016), Little Cayman (Edwards, Frazer & Jacoby, 2014), Jamaica (Chin, Aiken & Buddo, 2016), Bonaire (de Leon et al., 2013) and Costa Rica (Sandel et al., 2015), but remain unavailable for the central Mexican Caribbean. The weight-at-length of a species can vary across regions as a response to biotic (*e.g.* local food availability) and abiotic (*e.g.* water temperature) conditions (Johnson & Swenarton, 2016). Thus, when using biomass-informed models or estimating biomass from length observations, it is important to use site-specific parameters to obtain an accurate estimate. This is especially important when research involves identifying the total biomass available for harvest by fishers (Chin, Aiken & Buddo, 2016) or the efficacy of lionfish removals (Barbour et al., 2011; Morris, Shertzer & Rice, 2011; Johnston & Purkis, 2015).

Here, I provide the first allometric growth parameters for the invasive lionfish in the central Mexican Caribbean region. At the same time, I highlight the importance of using site-specific parameters by estimating biomass with parameters from other regions across the invasion range and comparing them to observed biomass. I also provide other 13 standardized parameters from eight studies through the invasion range, making them readily available for future research. Finally, I discuss the way in which allometric parameters are reported, and call for standardization to facilitate their use.

Materials and Methods

Area of study

The study took place off the coasts of Playa del Carmen, in the Mexican Caribbean (**Fig. 1**). The region represents the northernmost section of the Mesoamerican Barrier Reef System (Ruiz-Zarate & Arias-Gonzalez, 2004). Coral reefs and mangroves are locally important habitats that represent important sources of income in terms of extractive (*e.g.* recreational fishing) and non-extractive (*e.g.* SCUBA diving) activities related to tourism, the main source of income to the local economy (Murray, 2007).

The reef profile has been described by Arias-Gonzalez (1998), indicating that the reef lagoon extends about 500 m from the coast, until the reef crest is reached. The reef becomes deeper, leading to the reef front often found at 700 m from the coastline and extends for an additional 300 m. At approximately 1000 m away from shore and 30 - 40 m depth, the reef leads to a drop-off. Along a perpendicular profile to the coast, bands of reef are interrupted by sand patches at 8 - 12 m deep and 16-18 m deep. Along the coast, these reefs have been reported to be under significant anthropogenic pressure, likely causing a shift in structure and function (Bozec et al., 2008).




Figure 1 - Map of the study area. The black inset on the left (Yucatan Peninsula) indicates the location where study sites are distributed. On the right, circular markers indicate sampling sites and the black romboid indicates location of Puerto Aventuras, Mexico.

Fish sampling

A total of 33 SCUBA immersions were performed in 10 sampling sites along the coast in 2010 (Fig. 1, Table I). Sampling locations included wall and carpet reefs at depths between 5.7 m and 38.1 m. All observed organisms ($n = 109$) were collected using hand nets and numbered collection bottles. The use of hand nets prevented any weight loss due to bleeding and allowed a better representation of small sizes, often ignored due to gear selectivity when spearing. Organisms were euthanized and frozen within 30 minutes of completing the dive and stored for posterior analysis. Total Length (TL; mm) and Total Weight (TW; gr) were recorded for all organisms.

132

133 Table I - Coordinates, minimum, maximum and mean depth (m), and number of samples for each location.

Location	Lat.	Long.	Min. Depth	Max. Depth	Mean Depth	n
Canones	20.477	-87.233	15.0	31.2	21.6	11
Castillo	20.496	-87.220	12.5	30.5	27.5	18
Cuevitas	20.478	-87.244	7.4	12.8	11.2	4
Islas	20.490	-87.228	14.0	19.4	16.7	10
Paamul	20.513	-87.192	9.9	22.7	15.5	31
Paraiso	20.484	-87.226	9.4	38.1	17.7	16
Pared	20.502	-87.212	12.1	21.0	16.3	12
Pedregal	20.507	-87.204	14.4	14.9	14.7	3
Santos	20.493	-87.222	5.7	26.6	16.2	2
Tzimin-Ha	20.393	-87.307	21.2	24.6	22.9	2
Summary			5.7	38.1	18.6	109

134 *Data analysis*

135 The weight at length relationship between the observed variables is described by the allometric
 136 growth function:

137
$$TW = aTL^b$$

138 Where TW is the Total Weight (gr), TL is the Total Length (mm), a is the ponderal index and b is
 139 the scaling exponent or allometric parameter. When $b = 3$, it is said that the organism exhibits a
 140 perfect isometric growth. The dependent and independent variables were transformed via base-10
 141 logarithms, thus the equation becomes:

142
$$\log_{10}(TW) = b \times \log_{10}(TL) + \log_{10}(a)$$

143 This can be simplified and re-written as:

144
$$Y = mX + c$$

145 Where $Y = \log_{10}(TW)$, $X = \log_{10}(TL)$, $m = b$, and $c = \log_{10}(a)$. Since $b = m$, we will only use
 146 b throughout the paper for simplicity. The coefficients (c and b) were estimated with an Ordinary
 147 Least Square Regression and heteroskedastic-robust standard error correction was performed. Both
 148 coefficients were tested against the null hypothesis of no change (*i.e.* $H_0: c = 0$ and $H_0: b = 0$).
 149 Additionally, the allometric parameter was tested against the null hypothesis of isometric growth
 150 ($H_0: b = 3$). Coefficients were tested with a two-tailed Student's t-test. The significance of the
 151 regression was corroborated with an F-test.

152 Other length-weight relationships ($n = 13$) were extracted from peer-reviewed literature. Parameters
 153 were obtained for North Carolina ($n = 1$; Barbour et al., 2011), Northern ($n = 3$; Fogg et al., 2013)
 154 and Southern Gulf of Mexico ($n = 3$; Aguilar-Perera & Quijano-Puerto, 2016), the Southern
 155 Mexican Caribbean ($n = 1$; Sabido-Itza et al., 2016), Little Cayman ($n = 1$; Edwards, Frazer &
 156 Jacoby, 2014), Jamaica ($n = 1$; Chin, Aiken & Buddo, 2016), Bonaire ($n = 1$; de Leon et al., 2013)
 157 and Costa Rica ($n = 1$; Sandel et al., 2015) and Fishbase ($n = 1$; Froese & Pauly, 2016) were also
 158 included. When available, information on sampling methods, gender differentiation, location, and

depth ranges of each study was retrieved. Whenever gender was not specified, it was assumed that the results were presented for pooled genders.

During the review process, some papers indistinctly used a to report either the ponderal index in eq. 1 or the y-intercept (c) in eq. 3, which might sometimes be overlooked. Furthermore, some studies reported their parameters as mm-to-gr conversions, but a rapid evaluation of such parameters indicated that they were estimated as cm-to-gr conversions. Here, all parameters are reported as TL(mm) to TW(gr) conversions. When required, values from other studies were transformed for consistency.

Since uncertainty around estimated relationships was not reported in some of the reviewed studies, it was not possible to test for statistical differences between relationships. Instead, the 13 length-weight relationships were used to calculate expected weight for the organisms sampled in the Central Mexican Caribbean ($n = 109$). Expected weights were divided by the observed weights to obtain a ratio. Difference in mean weight ratios across studies were tested with a one-way Analysis of Variance (ANOVA).

All hypothesis testing was performed with an *a priori* confidence level of $\alpha = 0.01$ in R version 3.4.0 (R Core Team, 2017). Data wrangling was done with the tidyverse package (Wickham, 2017). Maps were created with a mix of functions from the sp (Pebesma & Bivand, 2005), rgdal (Bivand, Keitt & Rowlingson, 2017), tmap (Tennekes, 2017a), and tmaptools (Tennekes, 2017b) packages. Heteroskedastic-robust standard errors were calculated with the sandwich (Zeileis, 2004) and lmtest (Zeileis & Hothorn, 2002) packages. Models were manipulated with the broom package (Robinson, 2017). RefManagerR (McLean, 2014) was used to keep track of citations. Raw data and code used in this work is available at github.com/jcvdav/lionfish_biometry.

Results

Organism TL ranged between 34 and 310 mm and TW between 0.3 and 397.7 gr. The smallest organism (TL = 34.00 mm) was also the lightest organism (TW = 0.30 gr). However, the largest organism (TL = 310.00 mm) was not the heaviest (TW = 303.70 gr), and the heaviest organism (TW = 397.70 gr) was 292.00 mm in total length. Kernell density plots (Fig. 2) show the distribution for TL and TW for all sampled organisms. Both measures were positively skewed, with skewness of 0.87 for TL and 2.25 for TW.

Length-weight relationship

The model adjusted to eq. 3 estimated the coefficient values at $b = 3.2347391$ and $c = -5.4940866$. Thus, TW (gr) can be calculated from TL (mm) as a linear equation: $\log_{10}(TW) = 3.2347391 \times \log_{10}(TL) - 5.4940866$, or its exponential form: $TW = 3.2056297 \times 10^{-6} \times TL^{3.2347391}$. The intercept (c) and slope (b) were significantly different from zero ($t(107) = -66.17$; $p < 0.01$ and $t(107) = 83.24$; $p < 0.01$, respectively), rejecting the null hypothesis of no change. Additionally, the allometric factor (b) was significantly different from the value of isometric growth of $b = 3$ ($t(107) = 6.04$; $p < 0.01$), indicating that lionfish present allometric growth. More information on model fit and confidence intervals for the estimated coefficients is presented in Table II. The relationship between Total Length and Total Weight is presented in Figure 3.




Figure 2 - Kernel density plots for a) Total length (mm) and b) Total weight (gr) for 109 lionfish sampled in the central Mexican Caribbean.




Figure 3 - Length-weight relationship for 109 lionfish sampled in the central Mexican Caribbean. Points indicate samples, solid line indicates curve of best fit.

Table II - Regression table for the linear model fit between log10-transformed Total Weight (dependent variable) and Total Length (independent variable). Numbers in parentheses next to coefficient estimates indicate heteroskedastic-robust standard errors.

<i>Dependent variable:</i>	
	log10(TW)
c	-5.494 (0.083) ^{***}
b	3.235 (0.039) ^{***}
95% CI for c	(-5.657--5.331)
85% CI for b	(3.159-3.311)
F Statistic	6928.67 ^{***} (df = 1; 107)
Observations	109
Adjusted R ²	0.976
Residual Std. Error	0.096 (df = 107)

Note: * p<0.1; ** p<0.05; *** p<0.0

212 *Comparison of allometric parameters*


213 From the eight peer-reviewed studies including information on growth parameters for *P. volitans*
214 and Fishbase (Froese & Pauly, 2016), 13 parameters were identified (Table III). Two studies (Fogg
215 et al., 2013; Aguilar-Perera & Quijano-Puerto, 2016) reported gender-level and pooled parameters,
216 while the rest presented pooled results. The smallest coefficient of determination was presented by
217 Chin, Aiken & Buddo (2016) with $R^2 = 0.8715$, while Sabido-Itza et al. (2016) reported the
218 highest value at $R^2 = 0.9907$. Reviewed studies presented information for organisms obtained at
219 depths between 0.5 and 57 m. Two studies (Aguilar-Perera & Quijano-Puerto, 2016; Chin, Aiken &
220 Buddo, 2016) explicitly stated that their organisms were sampled with pole spears. Five studies
221 (Barbour et al., 2011; Fogg et al., 2013; Edwards, Frazer & Jacoby, 2014; Sandel et al., 2015;
222 Sabido-Itza et al., 2016) mentioned that some of their organisms were obtained with pole spears (or
223 other type of harpoon). A single study (de Leon et al., 2013) did not specify how organisms were
224 sampled.

225 Parameters from models fit to males or females exclusively tend to have a higher steepness (*i.e.*
226 higher allometric parameter), with mean \pm standard deviation values of $b = 3.27 \pm 0.06$ and $b =$
227 3.31 ± 0.23 for males and females respectively, compared to parameters from models for pooled
228 genders with a mean \pm standard deviation value of $b = 3.09 \pm 0.22$. In the case of the ponderal
229 index (a) and its \log_{10} transformed parameter (c), values were higher for parameters for pooled
230 genders. Figure 4 shows the predicted weights for organisms within the size range of these study
231 using the 14 parameters previously described.

232 There were significant differences in expected-to-observed weight ratios estimated for each pair of
233 parameters ($F(13, 1512) = 39.28$; $p < 0.05$). From all allometric parameters reviewed, those of
234 Edwards, Frazer & Jacoby (2014) provided the lowest weight estimates, with an expected-to-
235 observed weight ratio of 0.98 ± 0.23 (mean \pm SD). On the other hand, Barbour et al. (2011) yielded
236 the highest weight estimates, with a mean (\pm SD) expected-to-observed weight ratio of 1.76 ± 0.50 .
237 Predicted-to-observed weight ratios and groups identified by Tukey's HSD ($\alpha = 0.05$) are presented
238 in Figure 5.

239

240



242

243 Figure 4 - Length-weight relationships (n = 14) for eight studies, this study, and Fishbase. Colors indicate
 244 studies from which the parameters were extracted. Solid lines indicate that the fit was performed for males
 245 and females pooled together. Dotted lines indicate that the regression was performed on females, and dashed
 246 lines indicate it was performed for males.



Figure 5 - Box and whiskers plot showing the distribution of predicted to observed biomass ratios for 14 pairs of allometric parameters. Lines indicate median values, circles indicate mean values, notches represent 95% confidence intervals around the median, lower and upper hinges correspond to the first and third quartiles, whiskers extend to the largest and lowest values within 1.5 inter-quartile range of the hinge, small points represent outliers further away than the whiskers. Like letters indicate values that do not differ significantly (Tukey's HSD; $p < 0.05$).

Discussion

This study provides the first pair of allometric growth parameters specific to the Central Mexican Caribbean, complementing other studies performed in Mexican waters in the Alacranes Reef (Aguilar-Perera & Quijano-Puerto, 2016) and Xcalak National Park (Sabido-Itza et al., 2016). By using hand nets instead of spears, we are able to sample a wider range of sizes often ignored by pole spear samples, allowing us to include smaller organisms. Estimating parameters by including smaller organisms ensures better estimation of weight for smaller sizes. This is especially important when biomass is calculated from visual census, where small organisms can be registered. Thus, this study also increases certainty in weight estimation of small organisms.

The length-weight coefficients estimated in this study were within the range identified by studies in other regions (Barbour et al., 2011; de Leon et al., 2013; Fogg et al., 2013; Edwards, Frazer & Jacoby, 2014; Sandel et al., 2015; Aguilar-Perera & Quijano-Puerto, 2016; Chin, Aiken & Buddo, 2016; Sabido-Itza et al., 2016). However, the ones presented here provide lower weight estimates for a same length. Until about $TL = 200$ mm, there are no appreciable differences between the parameters for organisms from the Mexican Caribbean and those for little Cayman (Edwards, Frazer & Jacoby, 2014) and Jamaica (Chin, Aiken & Buddo, 2016). Yet, for larger organisms ($TL >$

270 mm) parameters from Costa Rica (Sandel et al., 2015) and Bonaire (de Leon et al., 2013) provide similar estimates to those from this study. Conversely, these same studies tend to estimate higher weights ---as compared to the ones reported here--- for smaller organisms, likely due to the lack of small organisms in the samples used to estimate their parameters. Whenever possible, future works should consider the use of hand nets to obtain the samples not only for studies on weight-at-length, but also diet, behavior and life history, where length can be an important factor.

There are evident differences in weight-at-length between organisms from the Caribbean and Gulf of Mexico / North-Western Atlantic. Weight estimates with parameters from the Gulf of Mexico and North-Western Atlantic (Barbour et al., 2011; Fogg et al., 2013; Aguilar-Perera & Quijano-Puerto, 2016; Sabido-Itza et al., 2016) tend to be higher than those from the Caribbean (de Leon et al., 2013; Edwards, Frazer & Jacoby, 2014; Sandel et al., 2015; Chin, Aiken & Buddo, 2016), except for the ones from Xcalak National Park, Mexico (Sabido-Itza et al., 2016). This indicates that there are differences between lionfish across the invasion range. Similar regional variation exist for age-at-length relationships of lionfish (Fogg et al., 2015). These differences can have major implications in management, especially when estimating biomass available for harvest or predicting effects on local ecosystems, or evaluating the effectiveness of removal programs. Using site-specific values provides a more accurate estimate of fish biomass. Future research should try to use, to the extent possible, parameters calculated for their region, or use different parameters to provide upper and lower bounds in their results. At the same time, this highlights the need for more basic research that furthers our understanding of lionfish biology. To better manage the invasion, we must perform research that can describe biologically important information of lionfish throughout its invasion range (Johnson & Swenarton, 2016).

While performing the literature review, it was often unclear if parameters were presented for eq.1 or eq. 3. Sometimes, they were even mislabeled and yielded senseless results when using the suggested conversion equation. On other cases, parameters were said to be reported for mm-to-gr conversions, when they were actually reported as cm-to-gr conversions. Perhaps these minor discrepancies can be easily solved by the trained eye, but why should they exist in the first place? It is important that we report our information in a standard way, making it readily available for other researchers and managers. In this particular case, I provide my humble opinion through 5 guidelines to report allometric parameters:

1. Be explicit in the methods section. What may seem obvious to you as an author ---because you have been deeply immersed throughout the process--- may not be clear to the reader. Specify any transformation performed on the data. When using log-transformations, mention the base used to transform. Do not assume that "data were log-transformed" means $\log_{10}(X)$. These assumptions vary across disciplines and software and can be a source of confusion. For example, in biology we often assume "log-transformed" indicates the use of base 10, however in R the proper command is `log10()` and not `log()`, which uses base e .
2. Use mm and gr to measure TL and TW, respectively. While conversion is always possible, we should aim at using standard units to report these parameters. If you prefer to use cm to gr conversions, that is certainly valid, but make sure to explicitly mention units when presenting the parameters.
3. Specify the equation for which parameters are presented by including an explicit example with the parameters substituted into it, as done by some of the papers reviewed (de Leon et al., 2013; Sandel et al., 2015; Chin, Aiken & Buddo, 2016; Sabido-Itza et al., 2016). If possible, present the relationship in their exponential (eq. 1) and linear (eq. 3) forms.

4. Report standard errors and/or confidence intervals around the obtained estimates. Given that small changes in a , c , and b can result in important changes in estimated weight, it is important that we report uncertainty around each parameter and not just general model fit and coefficient significance. Reporting uncertainty around parameters allows researchers and managers to include upper and lower bounds in their predictions.
5. Make your data and code available. Even if this is not required by the journal or publisher, you can use free cloud data storage services or third-party repositories to make your research accessible to others. Resources will always be limited and budget will rarely be enough. It is important that we take advantage of open science tools that promote the advancement of knowledge and foster collaboration. Ultimately, this promotes transparency, allows replicability of research, and advances science.

This study provides a new pair of allometric growth parameters for lionfish from the central Mexican Caribbean, where they exhibit different weight-at-length. Furthermore, regional differences in length-weight relationships were identified, highlighting the importance of using site-specific parameters.

Aknowledgements

I would like to thank Nils Van Der Haar and Michael Doodey as well as Guillermo Lotz-Cador who provided help to collect samples. I would also like to (anticipatedly) thank the editor and reviewers, who significantly improved the quality of this work. This research was partially funded by the "Consejo Nacional de Ciencias y Tecnología" (CONACyT) and the Latin American Fisheries Fellowship Program.

References

- Aguilar-Perera, A. & Quijano-Puerto, L. 2016. Relations between fish length to weight, and otolith length and weight, of the lionfish *pterois volitans* in the parque nacional arrecife alacranes, southern gulf of mexico. *Rev. biol. mar. oceanogr.* 51(2):469–474. DOI: [10.4067/S0718-19572016000200025](https://doi.org/10.4067/S0718-19572016000200025).
- Aguilar-Perera, A. & Tuz-Sulub, A. 2010. Non-native, invasive red lionfish (*pterois volitans* [linnaeus, 1758]: Scorpaenidae), is first recorded in the southern gulf of mexico, off the northern yucatan peninsula, mexico. *AI*. 5(Supplement 1):S9–S12. DOI: [10.3391/ai.2010.5.S1.003](https://doi.org/10.3391/ai.2010.5.S1.003).
- Albins, M. & Hixon, M. 2008. Invasive indo-pacific lionfish *pterois volitans* reduce recruitment of atlantic coral-reef fishes. *Mar. Ecol. Prog. Ser.* 367:233–238. DOI: [10.3354/meps07620](https://doi.org/10.3354/meps07620).
- Andradi-Brown, D.A., Grey, R., Hendrix, A., Hitchner, D., Hunt, C.L., Gress, E., Madej, K., Parry, R.L., et al. 2017. Depth-dependent effects of culling-do mesophotic lionfish populations undermine current management? *R Soc Open Sci.* 4(5):170027. DOI: [10.1098/rsos.170027](https://doi.org/10.1098/rsos.170027).
- Arias-Gonzalez, J.E. 1998. Trophic models of protected and unprotected coral reef ecosystems in the south of the mexican caribbean. *J Fish Biol.* 53(sa):236–255. DOI: [10.1111/j.1095-8649.1998.tb01030.x](https://doi.org/10.1111/j.1095-8649.1998.tb01030.x).
- Arias-Gonzalez, J.E., Gonzalez-Gandara, C., Luis Cabrera, J. & Christensen, V. 2011. Predicted impact of the invasive lionfish *pterois volitans* on the food web of a caribbean coral reef. *Environ Res.* 111(7):917–925. DOI: [10.1016/j.envres.2011.07.008](https://doi.org/10.1016/j.envres.2011.07.008).

355 Barbour, A., Montgomery, M., Adamson, A., D'az-Ferguson, E. & Silliman, B. 2010. Mangrove
356 use by the invasive lionfish *pterois volitans*. *Mar. Ecol. Prog. Ser.* 401:291–294. DOI:
357 [10.3354/meps08373](https://doi.org/10.3354/meps08373).

358 Barbour, A.B., Allen, M.S., Frazer, T.K. & Sherman, K.D. 2011. Evaluating the potential efficacy
359 of invasive lionfish (*pterois volitans*) removals. *PLoS ONE*. 6(5):e19666. DOI:
360 [10.1371/journal.pone.0019666](https://doi.org/10.1371/journal.pone.0019666).

361 Bax, N., Williamson, A., Agüero, M., Gonzalez, E. & Geeves, W. 2003. Marine invasive alien
362 species: A threat to global biodiversity. *Marine Policy*. 27(4):313–323. DOI: [10.1016/S0308-
363 597X\(03\)00041-1](https://doi.org/10.1016/S0308-597X(03)00041-1).

364 Betancur-R, R., Hines, A., Acero, A., Orti, G., Wilbur, A. & Freshwater, D. 2011. Reconstructing
365 the lionfish invasion: Insights into greater caribbean biogeography. *J Biogeography*. 38:1281–1293.
366 DOI: [10.1111/j.1365-2699.2011.02496.x](https://doi.org/10.1111/j.1365-2699.2011.02496.x).

367 Bivand, R., Keitt, T. & Rowlingson, B. 2017. *Rgdal: Bindings for the geospatial data abstraction*
368 *library*. ed. (nos.). Available: <https://CRAN.R-project.org/package=rgdal>.

369 Bozec, Y., Acosta-González, G., N'úñez-Lara, E. & Arias-González, J. 2008. Impacts of coastal
370 development on ecosystem structure and function of yucatan coral reefs, mexico. In *Proceedings of*
371 *the 11th international coral reef symposium*. ed. I.C.R.S. ICRF, Ed. (nos.). Ft. Lauderdale, Florida:
372 11th International Coral Reef Symposium.

373 Chin, D.A., Aiken, K.A. & Buddo, D. 2016. Lionfish population density in discovery bay, jamaica.
374 *International Journal of Scientific & Engineering Research*. 7(12):1327–1331.

375 Cote, I., Green, S., Morris, J., Akins, J. & Steinke, D. 2013. Diet richness of invasive indo-pacific
376 lionfish revealed by dna barcoding. *Mar. Ecol. Prog. Ser.* 472:249–256. DOI: [10.3354/meps09992](https://doi.org/10.3354/meps09992).

377 Dahl, K.A. & Patterson, W.F. 2014. Habitat-specific density and diet of rapidly expanding invasive
378 red lionfish, *pterois volitans*, populations in the northern gulf of mexico. *PLoS ONE*. 9(8):e105852.
379 DOI: [10.1371/journal.pone.0105852](https://doi.org/10.1371/journal.pone.0105852).

380 DAVIS, M.A. 2003. Biotic globalization: Does competition from introduced species threaten
381 biodiversity? *Bioscience*. 53(5):481. DOI: [10.1641/0006-3568\(2003\)053\[0481:BGDCFI\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2003)053[0481:BGDCFI]2.0.CO;2).

382 de Leon, R., Vane, K., Bertuol, P., Chamberland, V.C., Simal, F., Imms, E. & Vermeij, M.J.A.
383 2013. Effectiveness of lionfish removal efforts in the southern caribbean. *Endanger Species Res.*
384 22(2):175–182. DOI: [10.3354/esr00542](https://doi.org/10.3354/esr00542).

385 Edwards, M.A., Frazer, T.K. & Jacoby, C.A. 2014. Age and growth of invasive lionfish (*pterois*
386 *spp.*) in the caribbean sea, with implications for management. *BMS*. 90(4):953–966. DOI:
387 [10.5343/bms.2014.1022](https://doi.org/10.5343/bms.2014.1022).

388 Fogg, A.Q., Evans, J.T., Ingram JR, G.W., Peterson, M.S. & Brown-Peterson, N.J. 2015.
389 Comparing age and growth patterns of invasive lionfish among three ecoregions of the northern gulf
390 of mexico. In *Proceedings of the 68 th gulf and caribbean fisheries institute*. ed. G. GCFI & C.F.
391 Institute, Eds. (nos.). Panama City: Gulf; Caribbean Fisheries Institute.

392 Fogg, A.Q., Hoffmayer, E.R., Driggers, W.B., Campbell, M.D., Pellegrin, G.J. & Stein, W. 2013.
393 Distribution and length frequency of invasive lionfish (*pterois sp.*) in the northern gulf of mexico.
394 *GCR*. 25. DOI: [10.18785/gcr.2501.08](https://doi.org/10.18785/gcr.2501.08).

395 Froese, R. & Pauly, D. 2016. Available: <http://www.fishbase.org/> [2016, December 15].

396 Green, S. J., Akins, J. L., Maljkovic, A. & Côté, I.M. 2012. Invasive lionfish drive atlantic coral
397 reef fish declines. *PLoS ONE*. 7(3):e32596. DOI: [10.1371/journal.pone.0032596](https://doi.org/10.1371/journal.pone.0032596).

398 Grieve, B., Curchitser, E. & Rykaczewski, R. 2016. Range expansion of the invasive lionfish in the
399 northwest atlantic with climate change. *Mar. Ecol. Prog. Ser.* 546:225–237. DOI:
400 [10.3354/meps11638](https://doi.org/10.3354/meps11638).

401 Hackerott, S., Valdivia, A., Cox, C.E., Silbiger, N. J. & Bruno, J.F. 2017. Invasive lionfish had no
402 measurable effect on prey fish community structure across the belizean barrier reef. *PeerJ*. 5:e3270.
403 DOI: [10.7717/peerj.3270](https://doi.org/10.7717/peerj.3270).

404 Hixon, M., Green, S., Albins, M., Akins, J. & Morris, J. 2016. Lionfish: A major marine invasion.
405 *Mar. Ecol. Prog. Ser.* 558:161–165. DOI: [10.3354/meps11909](https://doi.org/10.3354/meps11909).

406 Johnson, E. G. & Swenarton, M. K. 2016. Age, growth and population structure of invasive lionfish
407 (pterois volitans/miles) in northeast florida using a length-based, age-structured population model.
408 *PeerJ*. 4:e2730. DOI: [10.7717/peerj.2730](https://doi.org/10.7717/peerj.2730).

409 Johnston, M. & Purkis, S. 2015. A coordinated and sustained international strategy is required to
410 turn the tide on the atlantic lionfish invasion. *Mar. Ecol. Prog. Ser.* 533:219–235. DOI:
411 [10.3354/meps11399](https://doi.org/10.3354/meps11399).

412 Jud, Z., Layman, C., Lee, J. & Arrington, D. 2011. Recent invasion of a florida (USA) estuarine
413 system by lionfish Pterois volitans / p. miles . *Aquat. Biol.* 13(1):21–26. DOI: [10.3354/ab00351](https://doi.org/10.3354/ab00351).

414 McLean, M.W. 2014. *Straightforward bibliography management in r using the refmanager*
415 *package*. ed. (nos.). Available: <http://arxiv.org/abs/1403.2036>.

416 Molnar, J.L., Gamboa, R. L., Revenga, C. & Spalding, M. D. 2008. Assessing the global threat of
417 invasive species to marine biodiversity. *Frontiers in Ecology and the Environment*. 6(9):485–492.
418 DOI: [10.1890/070064](https://doi.org/10.1890/070064).

419 Morris, J. A. & Akins, J. L. 2009. Feeding ecology of invasive lionfish (pterois volitans) in the
420 bahamian archipelago. *Environ. Biol. Fishes*. 86(3):389–398. DOI: [10.1007/s10641-009-9538-8](https://doi.org/10.1007/s10641-009-9538-8).

421 Morris, J. A., Shertzer, K. W. & Rice, J. A. 2011. A stage-based matrix population model of
422 invasive lionfish with implications for control. *Biol Invasions*. 13(1):7–12. DOI: [10.1007/s10530-010-9786-8](https://doi.org/10.1007/s10530-010-9786-8).

424 Muñoz, R., Currin, C. & Whitfield, P. 2011. Diet of invasive lionfish on hard bottom reefs of the
425 southeast usa: Insights from stomach contents and stable isotopes. *Mar. Ecol. Prog. Ser.* 432:181–
426 193. DOI: [10.3354/meps09154](https://doi.org/10.3354/meps09154).

427 Murray, G. 2007. Constructing paradise: The impacts of big tourism in the mexican coastal zone.
428 *Coastal Management*. 35(2-3):339–355. DOI: [10.1080/08920750601169600](https://doi.org/10.1080/08920750601169600).

429 Pebesma, E. J. & Bivand, R. S. 2005. Classes and methods for spatial data in R. *R News*. 5(2):9–13.
430 Available: <https://CRAN.R-project.org/doc/Rnews/>.

431 Pimentel, D., Zuniga, R. & Morrison, D. 2005. Update on the environmental and economic costs
432 associated with alien-invasive species in the united states. *Ecological Economics*. 52(3):273–288.
433 DOI: [10.1016/j.ecolecon.2004.10.002](https://doi.org/10.1016/j.ecolecon.2004.10.002).

434 R Core Team. 2017. *R: A language and environment for statistical computing*. ed. (nos.). Vienna,
 435 Austria: R Foundation for Statistical Computing. Available: <https://www.R-project.org/>.

436 Robinson, D. 2017. *Broom: Convert statistical analysis objects into tidy data frames*. ed. (nos.).
 437 Available: <https://CRAN.R-project.org/package=broom>.

438 Rocha, L. A., Rocha, C. R., Baldwin, C. C., Weigt, L. A. & McField, M. 2015. Invasive lionfish
 439 preying on critically endangered reef fish. *Coral Reefs*. 34(3):803–806. DOI: [10.1007/s00338-015-](https://doi.org/10.1007/s00338-015-1293-z)
 440 [1293-z](https://doi.org/10.1007/s00338-015-1293-z).

441 Ruiz-Zarate, M. & Arias-Gonzalez, J. 2004. Spatial study of juvenile corals in the northern region
 442 of the mesoamerican barrier reef system (mbrs). *Coral Reefs*. (September, 9). DOI:
 443 [10.1007/s00338-004-0420-z](https://doi.org/10.1007/s00338-004-0420-z).

444 Sabido-Itza, M., Medina-Quej, A., De Jesus-Navarrete, A., Gomez-Poot, J. & Garcia-Rivas, M.
 445 2016. Uso de la estructura de tallas como evidencia del establecimiento poblacional del pez le?n
 446 pterois volitans (scorpaeniformes: Scorpaenidae) en el sur del caribe mexicano. *RBT*. 64(1):353.
 447 DOI: [10.15517/rbt.v64i1.18943](https://doi.org/10.15517/rbt.v64i1.18943).

448 Sandel, V., Martínez-Fernández, D., Wangpraseurt, D. & Sierra, L. 2015. Ecology and management
 449 of the invasive lionfish pterois volitans/miles complex (perciformes: Scorpaenidae) in southern
 450 costa rica. *Rev Biol Trop*. 63(1):213–221. Available:
 451 <http://www.ncbi.nlm.nih.gov/pubmed/26299126> [2017, June 27].

452 Schofield, P. 2009. Geographic extent and chronology of the invasion of non-native lionfish
 453 (Pterois volitans [linnaeus 1758] and p. miles [bennett 1828]) in the western north atlantic and
 454 caribbean sea. *AI*. 4(3):473–479. DOI: [10.3391/ai.2009.4.3.5](https://doi.org/10.3391/ai.2009.4.3.5).

455 Schofield, P. 2010. Update on geographic spread of invasive lionfishes (pteroris volitans [linnaeus,
 456 1758] and p. miles [bennett, 1828]) in the western north atlantic ocean, caribbean sea and gulf of
 457 mexico. *AI*. 5(Supplement 1):S117–S122. DOI: [10.3391/ai.2010.5.S1.024](https://doi.org/10.3391/ai.2010.5.S1.024).

458 Tennekes, M. 2017a. *Tmap: Thematic maps*. ed. (nos.). Available: [https://CRAN.R-](https://CRAN.R-project.org/package=tmap)
 459 [project.org/package=tmap](https://CRAN.R-project.org/package=tmap).

460 Tennekes, M. 2017b. *Tmaptools: Thematic map tools*. ed. (nos.). Available: [https://CRAN.R-](https://CRAN.R-project.org/package=tmaptools)
 461 [project.org/package=tmaptools](https://CRAN.R-project.org/package=tmaptools).

462 Usseglio, P., Selwyn, J. D., Downey-Wall, A .M. & Hogan, J. D. 2017. Effectiveness of removals
 463 of the invasive lionfish: How many dives are needed to deplete a reef? *PeerJ*. 5:e3043. DOI:
 464 [10.7717/peerj.3043](https://doi.org/10.7717/peerj.3043).

465 Valdez-Moreno, M., Quintal-Lizama, C., Gómez-Lozano, R. & García-Rivas, M. D. C. 2012.
 466 Monitoring an alien invasion: DNA barcoding and the identification of lionfish and their prey on
 467 coral reefs of the mexican caribbean. *PLoS ONE*. 7(6):e36636. DOI:
 468 [10.1371/journal.pone.0036636](https://doi.org/10.1371/journal.pone.0036636).

469 Villaseñor-Derbez, J. C. & Herrera-Pérez, R. 2014. Brief description of prey selectivity and
 470 ontogenetic changes in the diet of the invasive lionfish Pterois volitans (actinopterygii,
 471 scorpaenidae) in the mexican caribbean. *PANAMJAS*. 9(2):131–135.

472 Wickham, H. 2017. *Tidyverse: Easily install and load 'tidyverse' packages*. ed. (nos.). Available:
 473 <https://CRAN.R-project.org/package=tidyverse>.

- 474 Zeileis, A. 2004. Econometric computing with hc and hac covariance matrix estimators. *Journal of*
475 *Statistical Software*. 11(10):1–17. Available: <http://www.jstatsoft.org/v11/i10/>.
- 476 Zeileis, A. & Hothorn, T. 2002. Diagnostic checking in regression relationships. *R News*. 2(3):7–10.
477 Available: <https://CRAN.R-project.org/doc/Rnews/>.