

Cochrane Database of Systematic Reviews

Physiotherapy for Parkinson's disease: a comparison of techniques (Review)



Tomlinson CL, Herd CP, Clarke CE, Meek C, Patel S, Stowe R, Deane KHO, Shah L, Sackley CM, Wheatley K, Ives N. Physiotherapy for Parkinson's disease: a comparison of techniques. *Cochrane Database of Systematic Reviews* 2014, Issue 6. Art. No.: CD002815. DOI: 10.1002/14651858.CD002815.pub2.

www.cochranelibrary.com



TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
BACKGROUND	3
OBJECTIVES	3
METHODS	3
RESULTS	5
Figure 1	7
Figure 2	g
Figure 3	10
DISCUSSION	13
AUTHORS' CONCLUSIONS	15
ACKNOWLEDGEMENTS	16
REFERENCES	17
CHARACTERISTICS OF STUDIES	25
ADDITIONAL TABLES	84
APPENDICES	99
WHAT'S NEW	103
HISTORY	104
CONTRIBUTIONS OF AUTHORS	104
DECLARATIONS OF INTEREST	104
SOURCES OF SUPPORT	104
INDEX TERMS	105



[Intervention Review]

Physiotherapy for Parkinson's disease: a comparison of techniques

Claire L Tomlinson¹, Clare P Herd², Carl E Clarke², Charmaine Meek³, Smitaa Patel¹, Rebecca Stowe¹, Katherine HO Deane⁴, Laila Shah¹, Catherine M Sackley⁵, Keith Wheatley⁶, Natalie Ives¹

¹Birmingham Clinical Trials Unit, University of Birmingham, Birmingham, UK. ²School of Clinical and Experimental Medicine, College of Medical and Dental Sciences, Birmingham, UK. ³Primary Care Clinical Sciences, University of Birmingham, Birmingham, UK. ⁴Edith Cavell Building, University of East Anglia, Norwich, UK. ⁵Faculty of Medicine and Health Sciences, University of East Anglia, Norwich, UK. ⁶Cancer Research Clinical Trials Unit, School of Cancer Sciences, University of Birmingham, Birmingham, UK

Contact address: Claire L Tomlinson, Birmingham Clinical Trials Unit, University of Birmingham, Robert Aitken Institute, Edgbaston, Birmingham, B15 2TT, UK. c.l.smith.1@bham.ac.uk.

Editorial group: Cochrane Movement Disorders Group.

Publication status and date: New search for studies and content updated (no change to conclusions), published in Issue 6, 2014.

Citation: Tomlinson CL, Herd CP, Clarke CE, Meek C, Patel S, Stowe R, Deane KHO, Shah L, Sackley CM, Wheatley K, Ives N. Physiotherapy for Parkinson's disease: a comparison of techniques. *Cochrane Database of Systematic Reviews* 2014, Issue 6. Art. No.: CD002815. DOI: 10.1002/14651858.CD002815.pub2.

Copyright © 2014 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Despite medical therapies and surgical interventions for Parkinson's disease (PD), patients develop progressive disability. The role of physiotherapy is to maximise functional ability and minimise secondary complications through movement rehabilitation within a context of education and support for the whole person. The overall aim is to optimise independence, safety and wellbeing, thereby enhancing quality of life. Trials have shown that physiotherapy has short-term benefits in PD. However, which physiotherapy intervention is most effective remains unclear.

Objectives

To assess the effectiveness of one physiotherapy intervention compared with a second approach in patients with PD.

Search methods

Relevant trials were identified by electronic searches of numerous literature databases (for example MEDLINE, EMBASE) and trial registers, plus handsearching of major journals, abstract books, conference proceedings and reference lists of retrieved publications. The literature search included trials published up to the end of January 2012.

Selection criteria

Randomised controlled trials of one physiotherapy intervention versus another physiotherapy intervention in patients with PD.

Data collection and analysis

Data were abstracted independently from each paper by two authors. Trials were classified into the following intervention comparisons: general physiotherapy, exercise, treadmill training, cueing, dance and martial arts.

Main results

A total of 43 trials were identified with 1673 participants. All trials used small patient numbers (average trial size of 39 participants); the methods of randomisation and concealment of allocation were poor or not stated in most trials. Blinded assessors were used in just over half of the trials and only 10 stated that they used intention-to-treat analysis.



A wide variety of validated and customised outcome measures were used to assess the effectiveness of physiotherapy interventions. The most frequently reported physiotherapy outcomes were gait speed and timed up and go, in 19 and 15 trials respectively. Only five of the 43 trials reported data on falls (12%). The motor subscales of the Unified Parkinson's Disease Rating Scale and Parkinson's Disease Questionnaire-39 were the most commonly reported clinician-rated disability and patient-rated quality of life outcome measures, used in 22 and 13 trials respectively. The content and delivery of the physiotherapy interventions varied widely in the trials included within this review, so no quantitative meta-analysis could be performed.

Authors' conclusions

Considering the small number of participants examined, the methodological flaws in many of the studies, the possibility of publication bias, and the variety of interventions, formal comparison of the different physiotherapy techniques could not be performed. There is insufficient evidence to support or refute the effectiveness of one physiotherapy intervention over another in PD.

This review shows that a wide range of physiotherapy interventions to treat PD have been tested. There is a need for more specific trials with improved treatment strategies to underpin the most appropriate choice of physiotherapy intervention and the outcomes measured.

PLAIN LANGUAGE SUMMARY

Physiotherapy for the treatment of Parkinson's disease

In spite of various medical and surgical treatments for Parkinson's disease (PD) patients gradually develop significant physical problems. Physiotherapists aim to enable people with PD to maintain their maximum level of mobility, activity and independence through the provision of the appropriate treatment. A range of approaches to movement rehabilitation are used, which aim to enhance quality of life by maximising physical ability and minimising secondary complications over the whole course of the disease. Evidence has shown that physiotherapy has short-term benefits in PD, however which approach of physiotherapy is most effective remains unclear.

Only randomised controlled trials were included in this review. These were studies where a group of participants were given one physiotherapy intervention and were compared with another group who received a different physiotherapy intervention. The participants were assigned to a group in a random fashion to reduce the potential for bias.

A total of 43 randomised trials involving 1673 participants (average trial size of just 39 participants) were identified as suitable for this review. The trials assessed various physiotherapy interventions, so they were grouped according to the type of intervention being used (general physiotherapy, exercise, treadmill training, cueing, dance or martial arts). However, despite this grouping, the physiotherapy interventions delivered and the outcomes assessed varied so much that the results of the individual trials could not be combined.

This review highlights that a wide range of different physiotherapy techniques have been tested to treat PD. Considering the small number of participants, the wide variety of physiotherapy interventions and the outcomes assessed, there is insufficient evidence to support the use of one approach of physiotherapy intervention over another for the treatment of PD.



BACKGROUND

Parkinson's disease (PD) is a complex neurodegenerative disorder (Rubenis 2007) with wide reaching implications for patients and their families. Whilst disability can occur at all stages of the disease (Deane 2001a), PD is progressive in nature. Patients face increased difficulties with activities of daily living (ADL) (Kwakkel 2007) and mobility such as gait, transfers, balance and posture (Keus 2007). Ultimately this leads to decreased independence, inactivity and social isolation (Keus 2007), resulting in reduced quality of life (Schrag 2000).

The management of PD has traditionally centred on drug therapy with levodopa viewed as the 'gold standard' treatment (Rascol 2002). However, even with optimal medical management, patients with PD still experience a deterioration of body function, daily activities and participation (Nijkrake 2007). For this reason there has been increasing support for the inclusion of rehabilitation therapies as an adjuvant to pharmacological and neurosurgical treatment (Gage 2004; Nijkrake 2007) and a call for the move towards multidisciplinary management of this multidimensional condition (Robertson 2003; Rubenis 2007).

The physiotherapist is a member of the multidisciplinary team (Robertson 2008; Rubenis 2007) and strives to maximise functional ability and minimise secondary complications through movement rehabilitation within a context of education and support for the whole person (Deane 2001a; Plant 2000). Physiotherapy for PD focuses on transfers, posture, upper limb function, balance (and falls), gait, physical capacity and (in)activity utilising cueing strategies, cognitive movement strategies and exercise to optimise the patient's independence, safety and wellbeing, thereby enhancing quality of life (Keus 2004; Keus 2007).

Referral rates to physiotherapy for people with PD have historically been low (Mutch 1986; Yarrow 1999). However, in recent years the number of referrals has increased, with a survey by Parkinson's UK in 2008 reporting that 54% of the 13,000 members surveyed had seen a physiotherapist, compared with 27% in a survey undertaken in 1998 (PDS 2008; Yarrow 1999). This rise in referrals may be attributed to two factors. Firstly, guidelines such as those published by the National Institute for Health and Clinical Excellence (NICE) (NICE CG35 2006) recommend that physiotherapy be made available throughout all stages of the disease, raising the profile of the intervention. This has been further supported by the publication of Dutch physiotherapy guidelines (Keus 2004) (updated guidelines were due for publication in 2013), which provide specific information for physiotherapists involved in the management of PD. Secondly, there has been a substantial increase in the number of trials completed over the last decade (particularly in the last five years), offering supportive evidence for the inclusion of physiotherapy in the management of PD (Keus 2009).

A recent Cochrane review (Tomlinson 2012) assessed the effectiveness of physiotherapy intervention versus no physiotherapy intervention in patients with PD. The review provided evidence for the short-term benefit (< three months) of physiotherapy intervention in the treatment of PD. Further, it suggested that there was no difference in treatment effect between the different types of physiotherapy interventions being used, though this was based on indirect comparisons. This now needs to be confirmed by examining head-to-head trials of physiotherapy interventions. This would be of interest to both clinicians and

patients so that appropriate physiotherapy interventions which provide greater benefit can be delivered to PD patients (Tomlinson 2012).

The present Cochrane review was first published in 2001, and included only seven randomised controlled trials with a total of 142 participants (Deane 2001b). The methods of physiotherapy varied so widely across the trials that the data could not be combined. This, along with the presence of methodological flaws, small sample sizes, and the possibility of publication bias, led Deane et al to conclude that there was insufficient evidence to support or refute the efficacy of any given approach of physiotherapy over another in PD (Deane 2001b). This review updates the previous Cochrane review. It aims to compare the effectiveness of one approach of physiotherapy intervention versus another approach of physiotherapy intervention in patients with PD.

OBJECTIVES

To assess the effectiveness of one physiotherapy intervention compared with a second approach in patients with PD.

METHODS

Criteria for considering studies for this review

Types of studies

All randomised controlled trials (including the first phase of crossover trials) comparing a physiotherapy intervention with another physiotherapy intervention were considered for inclusion in the review. Only trials that implemented random methods of treatment allocation were included.

Types of participants

Participants with a diagnosis of PD (as defined by the authors of the studies).

- PD of any disease stage (i.e. early or diagnostic, maintenance, or complex phase).
- · Any duration of PD.
- All ages.
- · Any drug therapy.
- Any duration of physiotherapy treatment (although trials of less than one day of treatment were excluded).

Types of interventions

Physiotherapy interventions aim to maximise functional ability and minimise secondary complications through movement rehabilitation within a context of education and support for the whole person. Physiotherapy encompasses a wide range of techniques, so we were inclusive in our definition of physiotherapy intervention (including those not directly delivered by a physiotherapist) with trials of general physiotherapy, exercise, treadmill training, cueing, dance and martial arts being included.

Types of outcome measures

- 1. Gait outcomes such as:
- a. two- or six-minute walk test (m), measures the number of metres a person can walk in two or six minutes thereby providing a measurement of walking endurance (Kersten 2004);



b. walking speed:

- i. 10- or 20-metre walk test (s), measures the time in seconds that a person takes to walk 10 or 20 metres thereby providing a measurement of gait speed (Kersten 2004),
- ii. velocity (m/s), measures the rate of change of position, recorded in metres per second (Trew 2005);
- c. cadence (steps/min), measures the number of steps taken in a given period of time, which is then converted into the number of steps taken per minute (Trew 2005);
- d. stride length (m), measures the average distance (in metres) between two successive placements of the same foot (Whittle 1996);
- e. step length (m), measures the average distance (in metres) between successive foot to floor contact with the opposite feet (Trew 2005);
- f. Freezing of Gait Questionnaire, a validated questionnaire for the assessment of freezing of gait. The questionnaire consists of six items and scores range from 0 to 24, with higher scores corresponding to more severe freezing of gait (Giladi 2000).
- 2. Functional mobility and balance outcomes such as:
- a. timed up and go (s), measures the time taken in seconds for a person to get up from a chair, walk a certain distance (usually three metres), turn around and walk back to the chair and sit down (Podsiadlo 1991);
- b. Functional Reach Test (cm), "the maximal distance one can reach forward beyond arm's length, while maintaining a fixed base of support in the standing position" (Duncan 1990);
- c. Berg Balance Scale, a validated questionnaire designed to measure functional standing balance of the older adult. The measure consists of 14 items and score ranges from 0 to 56; with 0 to 20 = high fall risk; 21 to 40 = medium fall risk; and 41 to 56 = low fall risk (Berg 1992; Qutubuddin 2005);
- d. Activity Specific Balance Confidence. a 16-item self-report questionnaire that asks individuals to rate their confidence that they will maintain their balance in the course of daily activities. Each item is rated from 0% (no confidence) to 100% (complete confidence) (Powell 1995; Talley 2008).
- 3. Data on falls such as:
- a. number of patients falling, e.g. falls diary;
- b. Falls Efficacy Scale, a 10-item patient-reported questionnaire that measures how confident a person is at carrying out various ADL. Items are rated from 1 to 10, with higher scores correlating with lower levels of confidence, and a total score of 70 or more indicating that a person has a fear of falling (Tinetti 1990);
- c. Falls Efficacy Scale International, a 16-item questionnaire that includes the 10 original items of the standard Falls Efficacy Scale as well as six items regarding higher functioning and social activities. Each item is rated on a scale of 1 to 4, with 1 being 'not concerned at all' and 4 being 'very concerned' (maximum score out of 64) (Yardley 2005).

- 4. Clinician-rated impairment and disability measures such as:
- a. Hoehn and Yahr, a scale used to describe how symptoms of Parkinson's disease progress. Scale ranges from 0 to 5, with higher levels indicating greater disability (Hoehn 1967);
- b. Unified Parkinson's Disease Rating Scale (UPDRS), designed to assess motor impairment and disability in Parkinson's disease. Higher scores correspond to greater disability (Fahn 1987):
- i. total, score ranges from 0 to 176,
- ii. mental, score ranges from 0 to 16,
- iii. ADL, score ranges from 0 to 52,
- iv. motor, score ranges from 0 to 108;
- c. Webster Rating Scale, an assessment of severity of disease and clinical impairment against 10 items using a scale of 0 = normal to 3 = maximum impairment (bradykinesia, rigidity, posture, upper extremity swing, gait, tremor at rest, facies, seborrhoea, speech, and self care). Scores ranges from 0 to 30, with higher scores indicating greater disease severity and disability (Webster 1968);
- d. Columbia University Rating Scale, an assessment of motor impairment and ADL against 13 items, using a five-point scale for each to give a total score between 0 = normal to 65 = maximum disability (Yahr 1969).
- 5. Patient-rated quality of life such as:
- a. Parkinson's Disease Questionnaire-39 (PDQ-39), a PD specific health-related quality of life questionnaire containing 39 items divided into eight domains. Scores range from 0 to 100 with higher scores corresponding to poorer quality of life (Jenkinson 1997; Peto 1995);
- b. PDQUALIF, a PD specific health-related quality of life questionnaire containing 32 items in seven dimensions and one item of global health-related quality of life. The total score ranges from 0 to 128 with higher scores indicating poorer quality of life (Welsh 2003);
- c. PDQL, a PD specific health-related quality of life questionnaire containing 37 items grouped into four subscales. Item scores range from 1 to 5. The PDQL-Summary Index ranges from 37 to 185, with higher scores reflecting better quality of life (Deboer 1996);
- d. Short Form-36 (SF-36) or 12 (SF-12), a generic short form health survey consisting of 36 or 12 questions. The SF-36 consists of eight scaled scores assessing vitality, physical functioning, bodily pain, general health perceptions, physical role functioning, emotional role functioning, social role functioning and mental health. Scores range from 0 to 100 with higher scores corresponding to better quality of life (Ware 1992).
- 6. Adverse events e.g. fractures, pain.
- 7. Compliance e.g. participant adherence, treatment fidelity.
- 8. Economic analysis.



Search methods for identification of studies

We undertook a systematic search of the literature up to the end of January 2012 for publications or abstracts describing relevant trials. This included searching the following.

- 1. General biomedical and science electronic databases (without date limiters) including the Cochrane Movement Disorders Specialised Register, *The Cochrane Library*, MEDLINE (1966 to 2012), EMBASE (1974 to 2012), CINAHL (1982 to 2012), ISI-SCI (1981 to 2012); the rehabilitation databases AMED (1985 to 2012), REHABDATA (1995 to 2012), REHADAT (1990 to 2012), PEDro (1929 to 2012), GEROLIT (1979 to 2012); the English language databases of foreign language research and third world publications LILACS (1982 to 2012), MedCarib (17th Century to 2012) and IMEMR (1984 to 2012).
- 2. The Cochrane Central Register of Controlled Trials (CENTRAL), the CentreWatch Clinical Trials listing service, the metaRegister of Controlled Trials, ClinicalTrials.gov, RePORT, National Institute on Disability and Rehabilitation Research (NIDRR) and National Research Register (NRR).
- 3. Handsearching of general (Lancet, BMJ, JAMA) and specific journals (Movement Disorders, Neurology, Archives of Physical Medicine and Rehabilitation, Clinical Rehabilitation, Physical Therapy) from 2001 to the end of January 2012.
- 4. The reference lists of retrieved papers and review articles.
- 5. Abstract books and conference proceedings. This included The XIII International Congress on Parkinson's disease (1999), The International Congress of Parkinson's Disease and Movement Disorders (1990, 1992, 1994, 1996, 1998, 2000, 2002, 2004, 2005, 2006, 2007, 2008, 2009, 2010, 2011, 2012), World Congress on Parkinson's Disease and Related Disorders (2009) and The American Academy of Neurology 51st annual meeting (1999).
- 6. Grey literature databases (including theses): Conference Proceedings Citation Index (1982 to 2012), DISSABS (1999 to 2012), Conference Papers Index (1982 to 2012), Index to Theses (1970 to 2012), Electronic Theses Online Service (EThOS) (16th Century to 2012) and ProQuest dissertations and theses databases (1861 to 2012).

The search strategies are listed in full in Appendix 1.

Data collection and analysis

Selection of studies

From the search results, two review authors (CLT, CPH, SP or LS) independently screened the abstracts of potentially relevant studies, with the full paper being obtained if the abstract did not provide sufficient information to determine eligibility for inclusion in the review. Disagreement was resolved by referral to a third review author (RS, CM or NI). Authors of potentially eligible studies were contacted for further information if details of their trial were unclear.

Data extraction and management

Two review authors (CLT, CPH, SP or CM) independently assessed the eligible papers or abstracts for trial details and outcome data. These were validated by discussion with any discrepancies resolved by consensus. Trial details were recorded on a standard trial description form and included: trial name, trial group, authors, randomised comparison, treatment schedule (including duration, number of sessions, type of intervention), other therapy, eligibility criteria, method of randomisation, allocation concealment, blinding, accrual period, number of participants randomised, number of dropouts, duration of follow-up, outcomes reported, use of intention-to-treat analysis and publication date(s). The outcome data extracted included data on gait, functional mobility and balance, falls, clinician-rated disability scale and patient-rated quality of life, adverse events, compliance or withdrawals and health economics where available.

Authors of any eligible unpublished studies were contacted to ask if further details and the data for their trial could be provided.

Assessment of risk of bias in included studies

Two review authors (CLT, CPH, SP or CM) assessed the methodological quality of the full papers by recording the eligibility criteria (for example specified inclusion (and exclusion) criteria low risk), method of randomisation (for example used computer random number generator - low risk) and blinding (for example blinding of assessors - low risk), concealment of allocation (for example use of central randomisation service - low risk), similarity of participants in treatment groups at baseline (no difference in baseline characteristics between treatment groups as stated in trial publication - low risk), co-intervention(s) constant (for example drug therapy stable - low risk), comparable treatment arms (for example similar treatment duration and frequency - low risk), whether an intention-to-treat analysis was performed (intentionto-treat analysis and withdrawals < 10% - low risk, withdrawals > 10% - unclear risk, per protocol analysis - all unclear risk) and the number of participants lost to follow-up and missing values (withdrawals below 10% - low risk) (see the risk of bias tables under 'Characteristics of included studies').

Data synthesis

Treatment arms of the included studies were classified and divided according to the types of interventions administered:

- 1. general physiotherapy;
- 2. exercise;
- 3. treadmill training;
- 4. cueing;
- 5. dance;
- 6. martial arts.

Disparate study designs in the included trials resulted in a lack of overlap in the collated physiotherapy methods and outcome measures such that data could not be combined in a meaningful way.

RESULTS

Description of studies

A total of 78 randomised trials of physiotherapy intervention in PD patients were identified; 29 studies were excluded (see



Characteristics of excluded studies). The reasons for excluding these trials were: not randomised or not properly randomised (n=12), cross-over study with data for the different phases not adequately separated (n=4), treatment given in trial not usually used by physiotherapists (such as whole body vibration technique) (n=4), no outcome measures relevant to our review (n=2), multidisciplinary therapy rehabilitation trial (n=2), trial

duration under one day (n=1), insufficient information available for inclusion in review (n=1), unsuitable comparator arm (n=1), study was confounded (n=1) and comparison of physiotherapy delivery rather than technique (n=1). There were also six ongoing trials for which data were not yet available (see Characteristics of ongoing studies). Therefore, there were 43 trials available for inclusion in the review, compared to seven in the 2001 review (Figure 1).



Figure 1. PRISMA flow diagram.






Figure 1. (Continued)






Figure 1. (Continued)

0 studies included in quantitative synthesis (meta-analysis)

The number of participants randomised into each of the 43 trials ranged from eight to 210 participants, with 1673 participants randomised in total (giving an average trial size of 39 participants) (see Characteristics of included studies). The assessment period ranged from two weeks to 24 months. The mean age of the participants in the trials was 67 years, 62% were male, the mean Hoehn and Yahr stage was 2.4, and they had had PD for approximately seven years.

There were 14 three-arm trials. Eight trials compared two different experimental physiotherapy interventions with placebo or no intervention (Almeida 2012; Fisher 2008; Li 2012; Mak 2008; McGinley 2012; Shankar 2009; Talakkad 2011; Thaut 1996). The

placebo or no intervention arms of these trials were not included in any analysis for this review (see Tomlinson 2012). Six trials compared three different physiotherapy techniques (Chaiwanichsiri 2011; Ebersbach 2010; Juncos 2006; Reuter 2011; Schenkman 2012a; Toole 2005). There was also one four-arm trial comparing two types of dance (waltz or foxtrot and tango) and martial arts with no intervention (Hackney 2009).

Risk of bias in included studies

See the Characteristics of included studies risk of bias in included studies tables, risk of bias graph (Figure 2) and risk of bias summary (Figure 3).

Figure 2. Risk of bias graph: review authors' judgements about each risk of bias item presented as percentages across all included studies.







Figure 3. Risk of bias summary: review authors' judgements about each risk of bias item for each included study.

	Eligibility Criteria	Randomisation Method	Concealment of Allocation	Similarity at Baseline	Intention to Treat Analysis	Withdrawals Described	Cointerventions Constant	Comparable arms	Blinded Assessors
Almeida 2012	•	•	•	•	?	•	•	•	?
Braun 2011	•	?	•	?	?	•	?	•	•
Burini 2006	•	?	•	•	?	•	•	•	•
Chaiwanichsiri 2011	•	?	?	•	?	?	•	?	•
Dias 2005	•	?	?	•	?	?	?	•	?
Diehl 2011	?	?	?	•	?	?	?	•	?
Ebersbach 2010	•	•	•	•	?	•	?	?	?
Fisher 2008	•	•	•	•	?	•	•	•	•
Frazzitta 2009	•	?	?	•	?	•	?	•	?
Hackney 2007	•	•	•	•	?	•	•	•	•
Hackney 2009	•	•	•	•	?	•	•	•	•
Hackney 2010	•	•	•	•	?	•	•	•	•
Hass 2006	?	?	?	?	?	?	?	•	•
Hirsch 1996	•	•	?	•	?	•	•	•	•
Joudoux 2011	?	?	?	?	?	?	?	•	?
Juncos 2006	•	?	?	?	?	•	•	•	•
Khallaf 2011	?	?	?	?	?	?	?	?	?
Li 2012	•	?	•	?	•	•	?	•	•
Loureiro 2010	•	?	?	?	?	?	?	•	?
Mak 2008	•	•	•	•	?	•	•	•	•
McGinley 2012	•	?	?	?	•	•	?	•	•
Miyai 2000	•	?	•	•	•	•	•	•	•
Miyai 2002	•	?	•	•	?	•	•	•	•
Morris 2009	•	•	•	?	?	•	?	•	•
Palmer 1986	•	?	?	•	?	•	•	•	•
Pelosin 2010	•	•	•	•	?	•	•	•	•



Figure 3. (Continued)



Trial design

A total of 40 trials had a parallel design and three had a crossover design (Burini 2006; Miyai 2000; Shiba 1999). Most trials looked at the short-term effect of therapy by assessing the participants at baseline and immediately or shortly after the physiotherapy intervention period (which ranged from two weeks to 24 months). Of the parallel design trials 21 reported additional data at assessment points after the treatment period had finished; this may have been at only two weeks or up to 12 months after the end of the treatment period.

Sample size

Only 13 studies (30%) (Braun 2011; Hackney 2007; Hackney 2009; Hackney 2010; Hirsch 1996; Li 2012; McGinley 2012; Morris 2009; Picelli 2012; Reuter 2011; Schenkman 2012a; Smania 2010; Yang 2010) reported a sample size calculation in the trial report, which was achieved by seven studies.

Eligibility criteria

The eligibility criteria for the trials were broad and varied considerably across the trials. The level of detail provided on the eligibility criteria was also variable, with some studies providing a detailed description of the entry criteria and others just stating

'patients with Parkinson's disease'. Only four trials (Pelosin 2010; Reuter 2011; Schenkman 2012a; Yang 2010) stated that a diagnosis of PD by the United Kingdom Brain Bank Criteria (Gibb 1988) was required. It is vital that eligibility criteria are well-defined so that the trial participant population can be determined.

Randomisation method and concealment of allocation

A total of 24 trials (56%) described the randomisation method used, of which only eight trials used low risk methods (for example computer random number generators). No details of the randomisation method used were provided for the remaining 19 trials. Further, only 20 trials (47%) either stated or gave adequate information that allowed the assessment of whether an adequate concealment of treatment allocation procedure had been used. Four trials were considered to be low risk by virtue of having used a central independent randomisation service, with the other 16 considered high risk (that is concealment of treatment allocation was potentially compromised as sealed envelopes, picking a card or picking from a hat were used).

Blinding of assessors

It would be impossible to blind participants and therapists to randomised treatment allocation in trials of physiotherapy. Therefore, such trials are open label by nature, and they are



consequently liable to the possibility of both performance and attrition bias. However, blinding of assessors could be employed to try and reduce the possibility of bias; 23 (53%) of the 43 studies used blinded assessors, seven used unblinded assessors so were classed as high risk, and in the other 13 studies this information was not provided (classed as unclear risk).

Co-interventions

Information on co-interventions was provided in 24 trials (56%), with participants continuing with their standard PD medication. In 20 trials the drug therapy was kept stable (low risk) throughout the duration of the trial, whereas five trials allowed variation (unclear risk). The remaining trials did not describe drug therapy (unclear risk).

Similarity of treatment groups at baseline

A description of the baseline characteristics of the trial participants is important to determine whether the trial results are generalisable and to compare characteristics of the two arms to ensure that the randomisation methods were successful. Six trials (Diehl 2011; Joudoux 2011; Khallaf 2011; Shankar 2009; Shen 2011; Talakkad 2011) did not provide any information on the baseline characteristics of the participants entered into the trial; 31 (of the 37) trials that reported baseline data gave this information split by treatment group; 25 trials reported sufficient data that showed participants to be similar at baseline. In six trials the baseline characteristics of the withdrawn participants were not given (Hackney 2009; Mak 2008; Miyai 2002; Picelli 2012; Smania 2010; Yang 2010). Along with the six trials that did not supply baseline data, this meant that 274 (16%) of the 1693 randomised participants were not characterised.

Data analysis

Ten trials stated intention to treat as the primary method of analysis, although it was not always clear if participants who withdrew from the trial were included in the analysis. The number of participant withdrawals was classed as low risk (\leq 10% of trial participants withdrew) in five of the 10 trials. Four trials used per protocol as the method of analysis (unclear risk). In the other 29 trials the method of analysis was not described (unclear risk), of these trials seven were considered high risk in terms of the proportion of participants that withdrew (that is > 10%), and in 13 trials the number of participant withdrawals (if any) was not described (unclear risk).

Available trial information and data

A total of 13 trials were reported in abstract form; further information was requested from the authors (two were not contactable: Khallaf 2011; Shiba 1999) with four (Juncos 2006; Poliakoff 2009; Robichaud 2012; Shankar 2009) providing additional information and seven (Diehl 2011; Hass 2006; Joudoux 2011; Loureiro 2010; Shen 2011; Sigurgeirsson 2009; Talakkad 2011) being unsuccessful. A total of 30 trials were reported as full publications; further information was requested from authors for 26 trials with 13 providing additional information.

Effects of interventions

See 'Summary of results' table for included trials (Table 1).

The physiotherapy interventions were placed into one of the six categories (general physiotherapy, exercise, treadmill training, cueing, dance and martial arts) according to the type of treatment administered. However, the content and delivery of the interventions within each category were diverse and varied substantially. Further, a wide variety of validated and customised outcome measures were used to assess the effectiveness of the physiotherapy interventions. Consequently, it was inappropriate to combine the results of studies or perform any statistical analysis.

Nevertheless, results of the most regularly used outcome measures could be examined on a trial by trial basis. The most frequently reported physiotherapy outcome measures were gait speed and timed up and go. The motor subscales of the UPDRS and PDQ-39 were the most commonly reported clinician-rated disability and patient-rated quality of life outcome measures respectively. Falls data were also considered to be an important outcome in PD. Even in the case of these more widely reported outcome measures, quantitative meta-analysis could not be performed due to the wide variety of interventions employed by the included studies such that no two studies with comparable interventions assessed the same outcome measure.

Gait speed (m/s)

Gait speed was measured in 19 studies with data available from 15 studies; data were inadequately or not reported for four trials (Juncos 2006; Khallaf 2011; McGinley 2012; Shen 2011). Nine (Almeida 2012; Chaiwanichsiri 2011; Fisher 2008; Hackney 2007; Hass 2006; Li 2012; Miyai 2000; Vivas 2011; Werner 2010) of the 15 studies reported no difference between the two intervention arms. In five studies (Dias 2005 (mean difference between arms 0.34 m/s); Frazzitta 2009 (mean difference between arms 0.1 m/s); Miyai 2002 (mean difference between arms 0.16 m/s); Thaut 1996 (mean difference between arms 0.093 m/s); Yang 2010 (mean difference between arms 0.15 m/s)) the gait speed was significantly increased in the novel experimental arm compared to the comparator arm. The remaining study (Hackney 2009) was a three-arm trial. Hackney (Hackney 2009) recorded significantly greater gait speed in the tango arm compared to the waltz/foxtrot arm (mean difference between arms 0.06m/s), but not the Tai Chi arm.

Timed up and go (s)

The timed up and go test was reported in 15 studies with data available from 14 studies; data were inadequately or not reported for one study (McGinley 2012). There was no difference between the two intervention arms for 12 studies (Almeida 2012; Braun 2011; Chaiwanichsiri 2011; Hackney 2007; Hackney 2010; Li 2012; Loureiro 2010; Morris 2009; Pelosin 2010; Robichaud 2012; Sigurgeirsson 2009; Vivas 2011). In the Ebersbach (Ebersbach 2010) study, the time taken to complete the timed up and go test was significantly improved (that is reduced) with the Lee Silverman voice treatment (LSVT) BIG arm compared to the Nordic walking and home exercise arms. In the Hackney (Hackney 2009) study, the timed up and go test was significantly improved in the tango arm compared to the waltz or foxtrot and Tai Chi arms.

Falls

Outcome measures that report data on falls are important and pertinent in PD studies. However, only five trials reported data on falls (11%) (Hirsch 1996; Juncos 2006; Li 2012; McGinley 2012; Smania 2010).



Hirsch 1996 reported the effect of training on mean latency to fall (average number of seconds participants swayed before stepping or falling, touching the surrounding panels with hands, or needing assistance from the technician to keep from sitting in the harness) and the proportion of falls (number of trials resulting in falls). There were no significant differences between the combined balance and resistance arm and balance only arm for either outcome.

Juncos 2006 was published in abstract form. There was insufficient information on the falls data collected to allow a description of the manner in which falls were analysed or to give any indication of the result.

Li 2012 monitored falls using daily 'falls calendars' that were maintained by study participants. There were no differences between the Tai Chi and resistance training arms.

McGinley 2012 measured the number of fallers, the number of multiple fallers and falls rate. The number of falls during the intervention phase (eight weeks) was significantly lower in the progressive strength training arm compared to the movement strategy training arm (n = 10 versus n = 24; P = 0.006), with the frequency of falls varying markedly. However, the time to first fall did not differ significantly between groups (P = 0.4).

Smania 2010 reported the number of falls by means of a falls diary. A diary of the number of falls, their circumstances, and the consequences for the patient's health were kept for one month prior to each evaluation session. There was no significant difference between the balance training experimental arm and the general physical exercises arm.

Clinician-rated disability

UPDRS motor subscale

The motor subscale of the UPDRS was reported in 23 studies with data and information available from 17; data were inadequately or not reported for six studies (Joudoux 2011; Khallaf 2011; Reuter 2011; Shankar 2009; Shen 2011; Toole 2005). There was no difference between the two intervention arms for 14 studies (Almeida 2012; Burini 2006; Fisher 2008; Frazzitta 2009; Hackney 2007; Hackney 2009; Juncos 2006; Li 2012; McGinley 2012; Miyai 2000; Miyai 2002; Poliakoff 2009; Robichaud 2012; Schenkman 2012a). In the Ebersbach 2010 study, the UPDRS motor score was significantly improved in the LSVT BIG arm compared to the Nordic walking and home exercise arms. In the Ridgel 2009 study, UPDRS motor scores showed a significantly greater improvement in the forced exercise arm compared to the voluntary exercise arm. In the Talakkad 2011 study, partial weight supported (-20%) treadmill training had a significantly greater improvement in UPDRS motor score compared to conventional gait training (no data in publication).

Patient-rated quality of life

PDQ-39 (Summary Index)

A total of 13 studies described using data from the PDQ-39 but the data were only available from eight studies; data were inadequately or not reported in five studies (Joudoux 2011; McGinley 2012; Reuter 2011; Shankar 2009; Sigurgeirsson 2009). Seven studies (Burini 2006; Ebersbach 2010; Juncos 2006; Morris 2009; Pelosin 2010; Poliakoff 2009; Schenkman 2012a) reported no difference between the intervention arms. Only the Hackney 2009 trial

reported a significant difference, with quality of life scores significantly improved in the tango arm compared to the waltz or foxtrot and Tai Chi arms.

Adverse events

Nine trials recorded adverse event data (Chaiwanichsiri 2011; Fisher 2008; Li 2012; McGinley 2012; Picelli 2012; Poliakoff 2009; Reuter 2011; Schenkman 2012a; Yang 2010). Minor adverse events such as muscle soreness, falls and dizziness were reported, with none of these trials reporting events significant enough to cause concern over the safety of the intervention.

Compliance

Only 18 of the 43 trials discussed participant compliance, with 13 (Burini 2006; Ebersbach 2010; Hackney 2007; Hackney 2009; Hirsch 1996; Li 2012; McGinley 2012; Miyai 2000; Miyai 2002; Poliakoff 2009; Reuter 2011; Thaut 1996; Toole 2005) quantifying it in some form; however this was difficult to analyse.

Health economics

Only one trial (McGinley 2012) intended to look at health economics. Watts 2008 (See McGinley 2012) published a protocol for economic analysis alongside the McGinley trial. They proposed to evaluate cost-effectiveness using a three-way comparison of the cost per fall averted and the cost per quality adjusted life year saved across two physical therapy interventions and a control group.

DISCUSSION

Summary of main results

This review updates the previous Cochrane review published in 2001 (Deane 2001b) comparing the effectiveness of one approach of physiotherapy intervention versus a second approach of physiotherapy intervention for the treatment of PD. The review now includes 43 randomised trials and 1693 participants (compared with seven trials and 142 participants in the 2001 review). Many recent systematic reviews have focused on specific areas of physiotherapy such as exercise, cueing and treadmill training (Allen 2011; Crizzle 2006; Goodwin 2008; Lim 2005; Mehrholz 2010; Nieuwboer 2008). Physiotherapy for PD encompasses a wide range of methods and techniques ranging from standard UK National Health Service (NHS) physiotherapy to exercise regimens and martial arts (Tomlinson 2012). Therefore, it is important that all approaches of physiotherapy intervention are included. Physiotherapy interventions were placed into six categories according to the type of treatment administered. However, the content and delivery of the interventions within each category varied substantially. In view of the disparate study designs, variety of interventions, and the array of outcome measures used the results of individual studies could not be combined using quantitative meta-analysis methods.

Comparison of different physiotherapy interventions

A recent Cochrane review provided evidence on the short-term benefit of physiotherapy in the treatment of PD (Tomlinson 2012). However, it did not identify whether any specific type of physiotherapy intervention provides greater benefit. This review aimed to assess this by comparing the effectiveness of one approach of physiotherapy intervention with a second approach of physiotherapy intervention. The various



physiotherapy interventions used in the trials included in this review were categorised according to the type of treatment administered to aid comparisons (general physiotherapy, exercise, treadmill training, cueing, dance, martial arts). However, despite categorisation, the techniques employed within each category were diverse. Therefore, it was not possible or appropriate to combine the results by meta-analysis as any such analysis would be difficult to interpret. It is also difficult to summarise such large amounts of heterogeneous data using a qualitative approach. Consequently, we conclude that there is no robust trial evidence to support any one approach of physiotherapy over another in the treatment of PD.

The content and delivery of the interventions used in the trials included within this review were diverse in nature. Although attempts were made to compare trials that were 'like for like' through the creation of different categories, the interventions delivered varied substantially within these categories. In the future it may be useful to further subcategorise by the primary aim of the trial. For example, categories for the primary aim might be to improve gait or improve a specific problem such as gait freezing or balance, or improve overall PD performance or function. The variation in the therapy delivered is unsurprising. Physiotherapy is an autonomous profession. Physiotherapists use different sets of skills and work within their own scope of practice (Chartered Society of Physiotherapy), and so this variation in the interventions delivered within clinical trials may reflect the diversity of clinical practice. Over the past decade, steps have been taken to try and provide best practice consensus in the form of the Dutch KNGF guidelines for physical therapy in patients with Parkinson's disease (Keus 2004). However, this publication provides a guidance framework rather than a 'recipe' for treatment. It is therefore important that physiotherapy interventions are compared against each other within rigorous trial designs to determine which are the most effective. This will provide therapists with a menu of treatment strategies that are known to be effective, from which they can devise individualised interventions. However, given the complexity of physiotherapy interventions, it is important that such trials follow the Medical Research Council guidelines for developing and evaluating complex interventions (MRC 2008), which will help standardise interventions. In addition, many of the trials included in this review had interventions that were intensive and for long durations, which may not be feasible in main stream care. Therefore, future trials should be designed such that the interventions are transferable and cost-effective in main stream care.

Outcome measures

There was a large variety of outcome measures utilised in the different studies included in this review. The majority of the outcomes were standard physiotherapy and PD outcomes. Gait speed, timed up and go, UPDRS motor score and quality of life measured using the PDQ-39 were the most frequently reported outcome measures within their respective categories. PD is a multidimensional disease and several important outcomes were either poorly or not reported; this includes data on the number of the falls, depression and anxiety and adverse events. Further, many of the outcomes measured were not relevant to the PD patient or carer. There is a need for the use of functional gait outcomes in future trials which look at the impact on the patient during everyday living rather than in a laboratory environment. Further,

only one trial intended to look at a health economics analysis of physiotherapy intervention, therefore little is known about the cost-effectiveness and economic value of this therapy. Future trials should include these outcomes.

Gait speed

Six trials (Dias 2005; Frazzitta 2009; Hackney 2009; Miyai 2002; Thaut 1996; Yang 2010) reported significant differences between the treatment arms for gait speed. The observed differences between treatment arms ranged from 0.06 to 0.34 m/s. In all these trials, with the exception of Hackney 2009, the intervention arm that used cueing or treadmill training methods, or both, was the better treatment arm. The Hackney trial (Hackney 2009) examined dance with the tango arm outperforming the waltz or foxtrot arm. The possible relevance and benefit of these significant differences to patients with PD must be put into context in terms of what is considered a minimally clinically important change (MCIC). Data on what is considered an MCIC are lacking for PD patients, but some data have been reported in stroke patients. In one study, it was reported that an increase in speed of just 0.03 and 0.13 m/s could translate into a change from a limited household to an unlimited household walker, and from an unlimited household walker to a most-limited community walker, respectively (Perry 1995). Therefore, these differences in gait speed between arms are consistent with the findings reported by Perry (Perry 1995).

Timed up and go

The MCIC in PD patients is thought to be 11 seconds (Steffen 2008). The differences seen between treatment arms in the Ebersbach (Ebersbach 2010) and Hackney trials (Hackney 2009) were much smaller than this (the mean difference between arms ranged from 1.1 to 2 seconds).

UPDRS motor subscale

Three trials reported a significant benefit of one treatment arm in improving UPDRS motor over the other arms. The UPDRS motor score in the LSVT BIG arm in the Ebersbach (Ebersbach 2010) trial improved by 5.05 points, which was significantly different from the two other arms in which the score increased (that is got worse) by 0.53 and 1.68 points in the Nordic walking and home exercise arms, respectively. Ridgel (Ridgel 2009) reported a significant difference in UPDRS motor score between the forced exercise arm (-16.6 points) and the voluntary exercise arm (3.6 points). Additionally, Talakkad (Talakkad 2011) reported a significantly greater improvement of the UPDRS motor score in the partial weight supported (-20%) treadmill training compared to conventional gait training, although there were no data in the abstract publication. The MCIC for the UPDRS motor score has been reported in two studies. One analysed data from two independent randomised controlled trials and concluded that the MCIC was five points for the motor score (Schrag 2006). The second study performed a cross-sectional analysis on 653 PD participants and reported a MCIC of 2.3 to 2.7 points for the motor score (Shulman 2010). Considering the recommendations of both Schrag (Schrag 2006) and Shulman et al (Shulman 2010) the improvements seen in the Ebersbach (Ebersbach 2010) and Ridgel (Ridgel 2009) trials may reflect a MCIC.



Patient-rated quality of life

A study performed by Peto et al (Peto 2001) to determine the MCIC for the PDQ-39 Parkinson's questionnaire reported that a difference of 1.6 on the summary index was a meaningful change. Only the Hackney 2009 trial reported a significant difference, with quality of life scores significantly improved by 7.10 points in the tango arm, which was significantly different from the waltz or foxtrot arm that improved by 0.68 points and the Tai Chi arm which had an increased score of 1.55 points (that is got worse). This improvement in quality of life that was seen in the tango arm is therefore meaningful to patients.

Quality of the evidence

There has been an improvement in the trial methodological quality and reporting since the last Cochrane review (Deane 2001b). The use of more robust randomisation methods, blinding and intention-to-treat analyses has increased since the previous review, although it was still inadequate. Of the 43 trials, only 24 trials provided information on the randomisation method (of which eight were considered low risk) and only four used a central independent randomisation procedure to ensure concealment of treatment allocation; 23 trials used blinded assessors and only 10 stated that they used intention-to-treat analysis methods. The lack of information in many trial reports may not necessarily indicate lack of implementation within the trial, but without this information provided in the trial publications the level of bias within the individual trials is difficult to assess. The need for further improvement in the methodological quality of trials in physiotherapy for PD was noted recently (Kwakkel 2007; Tomlinson 2012). Future trials must be methodologically sound, large, randomised, and controlled with reporting following CONSORT guidelines (Boutron 2008a; Boutron 2008b).

The trials included in the review were relatively small, with the majority assessing the effect of one approach of physiotherapy intervention versus a second approach of physiotherapy intervention over a short period of time, with limited follow-up. The overall size of trials has increased (with an average of 39 participants per trial in this review compared to 20 in the previous review), but the number of small and underpowered trials remains a problem. Small trials may be subject to 'random error' (Doll 1980) and consequently may give rise to false negative or positive results. Further, it must be noted that the mean age of onset of PD in the participants in the trials was 60 years. This is relatively young as the average age of onset in the PD MED trial is 67 to 69 years (Patel 2010), therefore the results of the trials may not be relevant to the general PD population, in particular older PD patients.

It should also be noted that only 18 of the 43 trials discussed participant compliance. This is surprising as compliance can be an important determinant of the outcomes measured and the acceptability of the interventions being assessed in the trials. Therefore, it would be beneficial if the level of compliance is measured in future trials.

Another limitation is that the follow-up period in the trials included in this review was relatively short. Outcome measures were assessed in all trials at baseline and immediately or shortly after the intervention had ceased (one or two weeks, with one trial (Ebersbach 2010) assessing at 12 weeks post-intervention). PD is a long-term neurodegenerative disease, so it is important that

the long-term effect of treatment is assessed. Only half of the 43 trials followed-up participants and reported further data during the post-treatment period (but this could have been only two weeks or up to 12 months post-treatment). The previous review's recommendations were for participants to be followed-up for at least six months. Only two trials did this, reporting follow-up data at 12 months (McGinley 2012) and at both 6 and 12 months (Werner 2010) post-treatment. Long-term data will provide valuable information about the duration of any improvement following therapy.

Reporting biases

Many trials used multiple outcome measures, and in the majority of trials the primary outcome measure was not explicitly stated. Therefore, it was difficult to assess and identify if studies were free of selective outcome reporting.

In order to minimise the risk of publication bias, a comprehensive search was performed of multiple databases, including searching of unpublished and ongoing studies, without any language restrictions. Also, where necessary, authors were contacted to request additional information. However, as with any systematic review, publication bias should still be taken into consideration.

In summary, large, well-designed randomised trials with improved specific treatment strategies and a follow-up of at least 12 months that assess the impact of treatment on all aspects of a patient's PD, alongside a health economics assessment, are needed.

AUTHORS' CONCLUSIONS

Implications for practice

Considering the small number of participants, the methodological flaws in many of the studies, the wide variety of physiotherapy interventions and outcome measures used, there is insufficient evidence to support the use of one approach of physiotherapy intervention over another for the treatment of PD.

Implications for research

- The majority of the studies included in this review were small and had a short follow-up period. It is clear that larger randomised controlled trials with longer-term follow-up are required, particularly focusing on improving trial methodology and reporting. Rigorous methods of randomisation should be used and the allocation should be adequately concealed. Data should be analysed according to intention-to-treat principles and trials should be reported according to the guidelines set out in the CONSORT statement (Boutron 2008a; Boutron 2008b).
- This review highlights the variety of physiotherapy interventions being tested for the treatment of PD. There is a need for more specific trials with improved treatment strategies to underpin the most appropriate choice of physiotherapy intervention and the outcomes measured. This review also reinforces the need for the universal employment of clinically relevant, reliable and sensitive outcome measures with a predefined outcome in each trial.
- Future trials should, where appropriate, try to follow the Medical Research Council guidelines for developing and evaluating complex interventions (MRC 2008).



• Future trials should be designed such that the interventions are transferable and cost-effective in main stream care.

ACKNOWLEDGEMENTS

We recognise the contributions of all the original trialists and the individuals who performed the trials that contributed to this systematic review, and thank the participants who agreed to help improve the assessment of PD treatment by taking part in these

We acknowledge Parkinson's UK for funding and the Department of Health whose core support for BCTU made this review possible.

We would also like to thank Ana Verissimo who provided a translation for the Dias 2005 paper.



REFERENCES

References to studies included in this review

Almeida 2012 (published and unpublished data)

Almeida QJ, Bhatt H. A manipulation of visual feedback during gait training in Parkinson's disease. *Parkinsons Disease* 2012:508720.

Braun 2011 (published data only)

Braun S, Beurskens A, Kleynen M, Schols J, Wade D. Rehabilitation with mental practice has similar effects on mobility as rehabilitation with relaxation in people with Parkinson's disease: a multicentre randomised trial. *Journal of Physiotherapy* 2011;**57**(1):27-34.

Burini 2006 {published and unpublished data}

Burini D, Farabollini B, Iacucci S, Rimatori C, Riccardi G, Capecci M, et al. A randomised controlled cross-over trial of aerobic training versus Qigong in advanced Parkinson's disease. *Europa Medicophysica* 2006;**42**(3):231-8.

Chaiwanichsiri 2011 {published data only}

Chaiwanichsiri D. Treadmill and music cueing for gait training in mild to moderate Parkinson's disease. clinicaltrials.gov 2009.

* Chaiwanichsiri D, Wangno W, Kitisomprayoonkul W, Bhidayasiri R. Treadmill training with music cueing: a new approach for Parkinson's gait facilitation. *Asian Biomedicine* 2011;**5**(5):649-54.

Dias 2005 {published data only}

Dias NP, Fraga DA, Cacho EWA, Oberg TD. Training of gait with visual cues in the patient with Parkinson's disease. *Fisioterapia em Movimento* 2005;**18**(4):43-51.

Diehl 2011 {published data only}

Diehl D, Combs S, Staples B. Comparison of boxing training and traditional exercise on balance and quality of life outcomes in persons with Parkinson's disease: a pilot study. *Movement Disorders* 2011;**26**:S124.

Ebersbach 2010 {published and unpublished data}

Ebersbach G, Ebersbach A, Edler D, Kaufhold O, Kusch M, Kupsch A, et al. Comparing exercise in Parkinson's disease - The Berlin LSVT (R) BIG study. *Movement Disorders* 2010;**25**(12):1902-8.

Fisher 2008 {published data only}

Fisher BE, Wu AD, Salem GJ, Song J, Lin CH, Yip J, et al. The effect of exercise training in improving motor performance and corticomotor excitability in people with early Parkinson's disease. *Archives of Physical Medicine and Rehabilitation* 2008;**89**(7):1221-9.

Frazzitta 2009 (published data only)

Frazzitta G, Maestri R, Uccellini D, Bertotti G, Abelli P. Rehabilitation treatment of gait in patients with Parkinson's disease with freezing: A comparison between two physical therapy protocols using visual and auditory cues with or without treadmill training. *Movement Disorders* 2009;**24**(8):1139-43.

Hackney 2007 (published and unpublished data)

Hackney ME, Kantorovich S, Levin R, Earhart GM. Effects of tango on functional mobility in Parkinson's disease: A preliminary study. *Journal of Neurologic Physical Therapy* 2007;**31**(4):173-9.

Hackney 2009 (published and unpublished data)

Hackney ME, Earhart GM. Effects of Dance on Movement Control in Parkinson's Disease: A Comparison of Argentine Tango and American Ballroom. *Journal of Rehabilitation Medicine* 2009;**41**(6):475-81.

* Hackney ME, Earhart GM. Health-related quality of life and alternative forms of exercise in Parkinson disease. *Parkinsonism & Related Disorders* 2009;**15**(9):644-8.

Hackney ME, Earhart GM. Tai Chi improves balance and mobility in people with Parkinson disease. *Gait & Posture* 2008;**28**(3):456-60.

Hackney 2010 {published and unpublished data}

Hackney ME, Earhart GM. Effects of dance on gait and balance in Parkinson's disease: A comparison of partnered and nonpartnered dance movement. *Neurorehabilitation and Neural Repair* 2010;**24**(4):384-92.

Hass 2006 {published data only}

Hass CJ, Waddell DE, Wolf SL, Juncos JL, Gregor RJ. The influence of tai chi training on locomotor ability in Parkinson's disease. American Society of Biomechanics (www.asbweb.org/conferences/2006/pdfs/154.pdf). 2006:154.

Hirsch 1996 {published and unpublished data}

Hirsch MA. Activity dependent enhancement of balance following strength and balance training. PhD Thesis, Florida State University 1996. [UMI Number: 9622856]

* Hirsch MA, Toole T, Maitland CG, Rider RA. The effects of balance training and high-intensity resistance training on persons with idiopathic Parkinson's disease. *Archives of Physical Medicine and Rehabilitation* 2003;**84**(8):1109-17.

Joudoux 2011 {published data only}

Joudoux S, Santiago T, Hutin E, Bayle N, Gracies JM. Randomised controlled single-blind trial comparing two rehabilitation programs in Parkinson's disease at a moderate stage: Methodology. *Annals of Physical and Rehabilitation Medicine. Conference proceeding* 2011;**54**:e104.

Juncos 2006 {published and unpublished data}

Juncos JL. Chinese exercise modalities in Parkinson's disease. controlled-trials.com 2010.

* Juncos JL, Haber M, Waddell D, Gregor R, Wohl SL. A controlled study of aerobic exercise and Chinese exercise modalities in Parkinson disease. *Neurology* 2006;**66**(5 Suppl 2):A314.



Khallaf 2011 (published data only)

Khallaf MES, Fathy H. Influence of moderate aerobic exercises on activities of daily living and depression in patients with Parkinson's disease. *European Journal of Neurology. Conference proceeding* 2011;**18**:439.

Li 2012 {published data only}

Li F, Fitzgerald K. Postural stability in Parkinson's disease patients after Tai Chi training: A randomised controlled trial. WFN XIX World Congress on Parkinson's Disease and Related Disorders. 2012; Vol. 18:s155.

* Li F, Harmer P, Fitzgerald K, Eckstrom E, Stock R, Galver J, et al. Tai chi and postural stability in patients with Parkinson's disease. *New England Journal of Medicine* 2012;**366**(6):511-9.

Sieber B, Fuzhong L. Tai Chi and Parkinson's disease. Report Database 2008.

Loureiro 2010 (published data only)

Loureiro APC, Stori FR, Chen R, Ribas CG, Loureiro CC. Comparative study of conventional physiotherapy and virtual rehabilitation using Wii Fit for improvement of balance in Parkinson's disease patients. Movement Disorders 2010; Vol. 25 Suppl 3:S714.

Mak 2008 (published data only)

Mak MK, Hui-Chan CW. Effect of 4-week training with audiovisual cues on sit-to-stand in Parkinsonian patients. *Movement Disorders* 2005;**20 Suppl 10**:388.

* Mak MKY, Hui-Chan CWY. Cued task-specific training is better than exercise in improving sit-to-stand in patients with Parkinson's disease: A randomized controlled trial. *Movement Disorders* 2008;**23**(4):501-9.

McGinley 2012 {published and unpublished data}

* McGinley JL, Martin C, Huxham FE, Menz HB, Danoudis M, Murphy AT, et al. Feasibility, safety, and compliance in a randomized controlled trial of physical therapy for Parkinson's disease. *Parkinson's Disease* 2012:795294.

Morris ME, Menz HB, McGinley JL, Huxham FE, Murphy AT, Iansek R, et al. Falls and mobility in Parkinson's disease: protocol for a randomised controlled clinical trial. *BMC Neurology* 2011;**11**:93.

Watts JJ, McGinley JL, Huxham F, Menz HB, Iansek R, Murphy AT, et al. Cost effectiveness of preventing falls and improving mobility in people with Parkinson disease: protocol for an economic evaluation alongside a clinical trial. *BMC Geriatrics* 2008;**8**:23.

Miyai 2000 {published and unpublished data}

Miyai I, Fujimoto Y, Ueda Y, Yamamoto H, Nozaki S, Saito T, et al. Treadmill training with body weight support: Its effect on Parkinson's disease. *Archives of Physical Medicine and Rehabilitation* 2000;**81**(7):849-52.

Miyai 2002 (published and unpublished data)

Miyai I, Fujimoto Y, Yamamoto H, Ueda Y, Saito T, Nozaki S, et al. Long-term effect of body weight-supported treadmill training in Parkinson's disease: A randomized controlled trial. *Archives of Physical Medicine and Rehabilitation* 2002;**83**(10):1370-3.

Morris 2009 (published data only)

Martin CL, Morris ME, Iansek R. Carry-over effects of inpatient rehabilitation in Parkinson's disease. *Movement Disorders* 2008;**23 Suppl 1**:675.

Morris ME, Iansek R. Effects of strategy training compared to exercises for gait rehabilitation in Parkinson disease: A randomized controlled trial. *Movement Disorders* 2006;**21**:688.

* Morris ME, Iansek R, Kirkwood B. A randomized controlled trial of movement strategies compared with exercise for people with Parkinson's disease. *Movement Disorders* 2009;**24**(1):64-71.

Morris ME, Iansek R, Kirkwood B, Danoudis M, Martin CL. Carryover effects of movement strategy training in Parkinson's disease. *Movement Disorders* 2009;**24 Suppl 1**:S270.

Schenkman M. Reply: A randomized controlled trial of movement strategies compared with exercise for people with Parkinson's disease. *Movement Disorders* 2010;**25**(4):524.

Palmer 1986 {published data only}

Palmer SS, Mortimer JA, Webster DD, Bistevins R, Dickinson GL. Comparison of stretch exercises and karate training as therapy for Parkinson's disease. *Archives of Physical Medicine and Rehabilitation* 1984;**65**(10):626.

* Palmer SS, Mortimer JA, Webster DD, Bistevins R, Dickinson GL. Exercise therapy for Parkinson's disease. *Archives of Physical Medicine and Rehabilitation* 1986;**67**(10):741-5.

Pelosin 2010 {published data only}

Pelosin E, Avanzino L, Bove M, Stramesi P, Nieuwboer A, Abbruzzese G. Action observation improves freezing of gait in patients with Parkinson's disease. *Neurorehabilitation and Neural Repair* 2010;**24**(8):746-52.

Picelli 2012 (published data only)

Picelli A, Melotti C, Origano F, Waldner A, Fiaschi A, Santilli V, et al. Robot-assisted gait training in patients with Parkinson's disease: A randomized controlled trial. *Neurorehabilitation and Neural Repair* 2012;**26**(4):353-61.

Poliakoff 2009 {published and unpublished data}

Poliakoff E, Galpin A, McDonald K, Dick J, Kellett M, Hayes S, et al. The effect of gym training on multiple outcomes in Parkinson's disease: A pilot randomised waiting-list controlled trial. *Movement Disorders* 2009;**24 Suppl 1**:S374.

Reuter 2011 {published data only}

Reuter I, Mehnert S, Leone P, Kaps M, Oechsner M, Engelhardt M. Effects of a flexibility and relaxation programme, walking, and nordic walking on Parkinson's disease. *Journal of Aging Research* 2011;**2011**:232473.

Ridgel 2009 (published data only)

Ridgel AL, Vitek JL, Alberts JL. Forced, not voluntary, exercise improves motor function in Parkinson's disease patients. *Neurorehabilitation and Neural Repair* 2009;**23**(6):600-8.



Robichaud 2012 (published and unpublished data)

Chen D, Corcos DM. Motor deficits - Experimental and clinical correlates. Report Database 2009.

Corcos D, Robichaud JA, David FJ, Vaillancourt DE, Poon C, Rafferty M, et al. 24 months of exercise improves the motor symptoms in Parkinson's disease. *Neurology* 2012;**78**:S02.003.

Rafferty MR, Robichaud JA, David FJ, Poon C, Vaillancourt DE, Comella CL, et al. Long-term exercise improves and maintains physical function in people with Parkinson's disease. *Movement Disorders* 2012;**27 Suppl 1**:S135.

Robichaud JA, Corcos D. The effect of exercise on individuals with Parkinson's disease. ClinicalTrials.gov 2007.

Robichaud JA, Corcos D. The effect of exercise on individuals with Parkinson's disease. controlled-trials.gov 2009.

* Robichaud JA, David FJ, Poon C, Rafferty M, Vaillancourt DE, Comella CL, et al. Long-term progressive exercise improves bradykinesia and muscle weakness. *Movement Disorders* 2012;**27 Suppl 1**:S136.

Schenkman 2012a {published data only}

* Schenkman M, Hall DA, Baron AE, Schwartz RS, Mettler P, Kohrt WM. Exercise for people in early- or mid-stage Parkinson disease: a 16-month randomized controlled trial. *Physical Therapy* 2012;**92**(11):1395-410.

Schenkman M, Shinowara N. Exercise, physical function and Parkinson's disease. Report Database 2009.

Shinowara N. Gait and step training to prevent fall in Parkinson's disease. Report Database 2009.

Shankar 2009 {published data only}

Shankar A, Labelle N, Derwent L, Bonfield S, Eliasziw M, Hu B, et al. Treadmill-walking with music shows a synergistic improvement in gait and balance in patients with Parkinson's disease: A randomized controlled trial. *Movement Disorders* 2009;**24 Suppl 1**:S281-2.

Shen 2011 {published and unpublished data}

* Shen X, Mak MKY. Effects of 12-week strategy-focused balance training on balance performance and balance confidence in patients with Parkinson's disease. *Hong Kong Physiotherapy Journal* 2011;**29**(2):100.

Shiba 1999 {published data only}

* Shiba Y, Obuchi S, Toshima H, Yamakita H. Comparison between visual and auditory stimulation in gait training of patients with idiopathic Parkinson's disease. World Congress of Physical Therapy Conference. 1999.

Sigurgeirsson 2009 {published data only}

Sigurgeirsson AT, Bjarnadottir OH, Thorsteinsdottir MH, Ingvarsson PE. The effect of high-volume walking with visual cues on gait in Parkinson's patients. A randomised controlled trial. Parkinsonism & Related Disorders 2009; Vol. 15 Suppl 2:S192.

Smania 2010 {published data only}

Smania N, Corato E, Tinazzi M, Stanzani C, Fiaschi A, Girardi P, et al. Effect of balance training on postural instability in patients with idiopathic Parkinson's disease. *Neurorehabilitation and Neural Repair* 2010;**24**(9):826-34.

Talakkad 2011 {published data only}

Gupta A, Ganesan M, Pal P, Talakkad S. Effect of partial weightsupported treadmill gait training on balance in patients with Parkinson disease. 2011 Annual Assembly of the American Academy of Physical Medicine and Rehabilitation 2011; Vol. 3, issue 10:S163-4.

* Talakkad S, Ganesan M, Gupta A, Pal PK, Trichur R. Effect of partial weight supported treadmill gait training on cardiovascular dynamics in patients with Parkinson's disease. *Movement Disorders* 2011;**26 Suppl 2**:s173.

Thaut 1996 {published and unpublished data}

* Thaut MH, McIntosh GC, Rice RR, Miller RA, Rathbun J, Brault JM. Rhythmic auditory stimulation in gait training for Parkinson's disease patients. *Movement Disorders* 1996;**11**(2):193-200.

Toole 2005 {published data only}

Maitland CG, Toole T, Warren E, Hubmann MF. The effects of loading and unloading on gait and balance control in Parkinsonism. *Archives of Physical Medicine and Rehabilitation* 2004;**85**:E13.

* Toole T, Maitland CG, Warren E, Hubmann MF, Panton L. The effects of loading and unloading treadmill walking on balance, gait, fall risk, and daily function in Parkinsonism. *NeuroRehabilitation* 2005;**20**(4):307-22.

Vivas 2011 (published data only)

Vivas J, Arias P, Cudeiro J. Aquatic therapy versus conventional land-based therapy for Parkinson's disease: an open-label pilot study. *Archives of Physical Medicine & Rehabilitation* 2011;**92**(8):1202-10.

Werner 2010 {published and unpublished data}

Werner WG, Gentile AM. Improving gait and promoting retention in individuals with Parkinson's disease: a pilot study. *Journal of Neurology* 2010;**257**(11):1841-7.

Yang 2010 {published data only}

Yang YR, Lee YY, Cheng SJ, Wang RY. Downhill walking training in individuals with Parkinson's disease: A randomized controlled trial. *American Journal of Physical Medicine & Rehabilitation* 2010;**89**(9):706-14.

References to studies excluded from this review

Byl 2009 {published data only}

Byl N. Enhancing safe mobility in patients with Parkinson's disease: effect of dual task training during aerobic and moderate exercise. *Parkinsonism & Related Disorders* 2009;**15 Suppl 2**:S122.



Cheon 2012 (published data only)

Cheon SM, Sung HR, Chae BK, Kim JW. Effects of different exercises in the patients with Parkinson's disease. *Parkinsonism & Related Disorders* 2012;**18**:s154.

Chouza 2011 {published data only}

Chouza M, Arias P, Vinas S, Cudeiro J. Acute effects of whole-body vibration at 3, 6, and 9 hz on balance and gait in patients with Parkinson's disease. *Movement Disorders* 2011;**26**:920-1.

Cianci 2010 (published data only)

Cianci H, Robinson K, Bunting-Perry L, Sollenberger J, Nooregian J, Duda J. Are wheeled walkers with visual cues efficacious to treat freezing of gait in Parkinson's disease?. Parkinsonism & Related Disorders 2010; Vol. 16 Suppl 1:S64.

Donovan 2011 (published data only)

Donovan S, Lim C, Diaz N, Browner N, Rose P, Sudarsky LR, et al. Laserlight cues for gait freezing in Parkinson's disease: An openlabel study. *Parkinsonism & Related Disorders* 2011;**17**(4):240-5.

Farley 2005 (published data only)

Farley BG, Koshland GF. Learning Big (TM): Efficacy of a large-amplitude exercise approach for patients with Parkinson's disease-bradykinesia to balance. *Movement Disorders* 2005;**20 Suppl 10**:466.

Filippin 2010 (published data only)

Filippin NT, da Costa PHL, Mattioli R. Effects of treadmill-walking training with additional body load on quality of life in subjects with Parkinson's disease. *Revista Brasileira de Fisioterapia* 2010;**14**(4):344-50.

Frazzitta 2012 (published data only)

Frazzitta G, Bertotti G, Riboldazzi G, Turla M, Uccellini D, Boveri N, et al. A new approach to rehabilitation of Parkinsonian patients: The intensive rehabilitation treatment. *Movement Disorders* 2011;**26**:s127.

* Frazzitta G, Bertotti G, Riboldazzi G, Turla M, Uccellini D, Boveri N, et al. Effectiveness of intensive inpatient rehabilitation treatment on disease progression in Parkinsonian patients: A randomized controlled trial with 1-year follow-up. Neurorehabilitation and Neural Repair 2012;**26**(2):144-50.

Gobbi 2009 {published data only}

Gobbi L, Oliveira-Ferreira M, Aetano MJ, irani-Silva E, Arbieri F, Tella F, et al. Exercise programs improve mobility and balance in people with Parkinson's disease. Parkinsonism & Related Disorders 2009; Vol. 15 Suppl 2:S6.

* Gobbi LTB, Oliveira-Ferreira MDT, Caetano MJD, Lirani-Silva E, Barbieri FA, Stella F, et al. Exercise programs improve mobility and balance in people with Parkinson's disease. *Parkinsonism & Related Disorders* 2009;**15 Suppl 3**:S49-52.

Haase 2011 {published data only}

Haase M. The immediate effects of rhythmic arm swing and finger tapping exercises on gait of Parkinson's patients. ProQuest 2011.

Kamsma 1995 (published data only)

Kamsma YPT, Brouwer WH, Lakke JPWF. Training of compensational strategies for impaired gross motor skills in Parkinson' disease. *Physiotherapy Theory and Practice* 1995;**11**(4):209-29.

Klassen 2007 (published data only)

Klassen L, Dal Bello-Haas V, Sheppard M, Metcalfe A. Evaluating the benefits of group exercise and group exercise and education programs for individuals with Parksinson's disease (PD). *Physiotherapy* 2007;**93 Suppl 1**:S91.

Knobl 2011 {published data only}

Knobl PE. An evaluation of motor imagery and exercise interventions in Parkinson's disease. ProQuest 2011.

* Knobl PE, Almeida QJ. Rehabilitation for Parkinson's disease: Is it the sensory or attention focus that improves disease severity in sensory-attention focused interventions?. *Movement Disorders* 2011;**26 Suppl 2**:s153.

Lee 2011 {published data only}

Lee KS, Lee WH, Hwang S. Modified constraint-induced movement therapy improves fine and gross motor performance of the upper limb in Parkinson disease. *American Journal of Physical Medicine & Rehabilitation* 2011;**90**(5):380-6.

Ma 2009 {published data only}

Ma HI, Hwang WJ, Lin KC. The effects of two different auditory stimuli on functional arm movement in persons with Parkinson's disease: a dual-task paradigm. *Clinical Rehabilitation* 2009;**23**(3):229-37.

Marchese 2000 (published data only)

Abbruzzese G, Diverio M, Zucchi F, Marchese R. Comparison of two physical therapy approaches in the rehabilitation of parkinsonian patients. *Parkinsonism and Related Disorders* 1999;**5 Suppl**:S49.

* Marchese R, Diverio M, Zucchi F, Lentino C, Abbruzzese G. Comparison of two physical therapy approaches in the rehabilitation of parkinsonian patients: a comparison of two physical therapy protocols. *Movement Disorders* 2000;**15**(5):879-83.

Modugno 2010 {published data only}

Modugno N. Long-term program of social theatre for the management of motor and motor symptoms of Parkinson's disease. Movement Disorders 2010; Vol. 25 Suppl 3:S721.

* Modugno N, Iaconelli S, Fiorilli M, Lena F, Kusch I, Mirabella G. Active theater as a complementary therapy for Parkinson's disease rehabilitation: A pilot study. *The Scientific World Journal* 2010;**10**:2301-13.

Munneke 2010 {published data only}

Keus SH, Bloem BR, Nijkrake M, Lim LI, Munneke M. The ParkNet trial: Implementation of an evidence-based guideline for physical therapy in Parkinson's disease. *Movement Disorders* 2006;**21**:S122.



Keus SHJ, Nijkrake MJ, Borm GF, Kwakkel G, Roos RAC, Berendse HW, et al. The ParkinsonNet trial: design and baseline characteristics. *Movement Disorders* 2010;**25**(7):830-7.

Munneke M, Kens S, Nijkrake M, Kwakkel G, Berendse H, Roos R, et al. Efficiency of evidence-based physiotherapy for Parkinson's disease: the ParkinsonNet Trial. *Movement Disorders* 2008;**23 Suppl 1**:669.

* Munneke M, Nijkrake M, Keus SHJ, Kwakkel G, Berendse HW, Roos RAC, et al. Efficacy of community-based physiotherapy networks for patients with Parkinson's disease: a cluster-randomised trial. *Lancet Neurology* 2010;**9**(1):46-54.

Munneke M, Nijkrake MJ, Keus SHJ, Lim IIK, Borm GF, van der Wilt GJ, et al. Efficiency of physiotherapy in Parkinson's disease: The ParkNet trial. *Movement Disorders* 2007;**22 Suppl 16**:629.

Nijkrake MJ, Keus SHJ, Overeem S, Oostendorp RAB, Vlieland TPMV, Mulleners W, et al. The ParkinsonNet concept: development, implementation and initial experience. *Movement Disorders* 2010;**25**(7):823-9.

Pacchetti 2000 {published data only}

Pacchetti C, Mancini F, Aglieri R, Fundaro C, Martignoni E, Nappi G. Active music therapy in Parkinson's disease: An integrative method for motor and emotional rehabilitation. *Psychosomatic Medicine* 2000;**62**(3):386-93.

Pohl 2003 (published data only)

Pohl M, Rockstroh G, Ruckriem S, Mrass G, Mehrholz J. Immediate effects of speed-dependent treadmill training on gait parameters in early Parkinson's disease. *Archives of Physical Medicine and Rehabilitation* 2003;**84**(12):1760-6.

Rochester 2011 {published data only}

Rochester L, Baker K, Nieuwboer A, Burn D. Targeting dopasensitive and dopa-resistant gait dysfunction in Parkinson's disease: selective responses to internal and external cues. *Movement Disorders* 2011;**26**(3):430-5.

Sage 2009a {published data only}

Sage MD, Almeida QJ. Symptom and gait changes after sensory attention focused exercise vs aerobic training in Parkinson's disease. *Movement Disorders* 2009;**24**(8):1132-8.

Sage 2009b {published and unpublished data}

Sage MD, Almeida QJ. A Canadian approach to exercise rehabilitation: a systematic evaluation of strategies to reduce the symptoms of Parkinson's disease. *Movement Disorders* 2009;**24 Suppl 1**:S276-7.

Sage 2010 (published data only)

Sage MD, Almeida QJ. A positive influence of vision on motor symptoms during sensory attention focused exercise for Parkinson's disease. *Movement Disorders* 2010;**25**(1):64-9.

Stallibrass 2001 {published data only}

Stallibrass C. Controlled trial to evaluate the effects of lessons in the Alexander Technique on the management of disability

by people with Parkinson's disease. National Research Register 2001.

Tamir 2007 {published data only}

Tamir R, Dickstein R, Huberman M. Integration of motor imagery and physical practice in group treatment applied to subjects with Parkinson's disease. *Neurorehabilitation and Neural Repair* 2007;**21**(1):68-75.

Tickle-Degnen 2010 (published data only)

Tickle-Degnen L, Ellis T, Saint-Hilaire M, Thomas CA, Wagenaar RC. Efficacy of self-management rehabilitation on quality of life outcomes in Parkinson's disease. *Movement Disorders* 2009;**24 Suppl 1**:S377-8.

* Tickle-Degnen L, Ellis T, Saint-Hilaire M, Thomas CA, Wagenaar RC. Self-management rehabilitation and health-related quality of life in Parkinson's disease: a randomized controlled trial. *Movement Disorders* 2010;**25**(2):194-204.

White DK, Wagenaar RC, Ellis TD, Tickle-Degnen L. Changes in walking activity and endurance following rehabilitation for people with Parkinson disease. *Archives of Physical Medicine and Rehabilitation* 2009;**90**(1):43-50.

Wulf 2009 {published data only}

Wulf G, Landers M, Lewthwaite R, Tollner T. External focus instructions reduce postural instability in individuals with Parkinson's disease. *Physical Therapy* 2009;**89**(2):162-8.

Yen 2011 {published data only}

Yen CY, Lin KH, Hu MH, Wu RM, Lu TW, Lin CH, et al. Effects of virtual reality-augmented balance training on sensory organization and attentional demand for postural control in people with Parkinson disease: a randomized controlled trial. *Physical Therapy* 2011;**91**(6):862-78.

References to ongoing studies

Brauer 2011 (published and unpublished data)

Brauer SG, Woollacott MH, Lamont R, Clewett S, O'Sullivan J, Silburn P, et al. Single and dual task gait training in people with Parkinson's disease: a protocol for a randomised controlled trial. *BMC Neurology* 2011;**11**:90.

Protas 2005 (published data only)

Protas EJ, Mitchell K, Williams A, Qureshy H, Caroline K, Lai EC. Gait and step training to reduce falls in Parkinson's disease. *NeuroRehabilitation* 2005;**20**(3):183-90.

Schenkman 2012b {published data only}

Schenkman M. Endurance exercise in Parkinson's disease. controlled_trials.com 2012.

* Schenkman M. Study in Parkinson's disease of exercise (SPARX). ClinicalTrials.gov 2011.

Smith 2009 {published data only}

Smith S, Dibble L. High force resistance training and dopamine replacement effects on hypokinesia in Parkinson disease. ClinicalTrials.gov 2009.



Uc 2010 {published data only}

Uc EY, Newman SR. Effects of aerobic exercise in Parkinson's disease. ClinicalTrials.gov 2010.

van Nimwegen 2010 (published and unpublished data)

Speelman AD, van Nimwegan M, Smulders K, Overeem S, Borm G, Backx F, et al. Rationale and design of the Parkfit study: a randomized controlled trial to increase physical activity in patients with Parkinson's disease. Parkinsonism & Related Disorders 2010; Vol. 16 Suppl 1:S58.

Speelman AD, van Nimwegen ML, Smulders K, Overeem S, Borm GF, Backx FJG, et al. Design of a randomized controlled trial evaluating the effectiveness of a multifaceted program to increase physical activity in Parkinson patients. Movement Disorders 2010; Vol. 25 Suppl 3:S679.

van Nimwegen M, Speelman A, Smulders S, Overeem S, Borm G, Backx F, et al. Rationale and design of the ParkFit study: A randomized controlled trial to increase physical activity in patient with Parkinson's disease. *Movement Disorders* 2009;**24 Suppl 1**:S284.

* van Nimwegen M, Speelman AD, Smulders K, Overeem S, Borm GF, Backx FJG, et al. Design and baseline characteristics of the ParkFit study, a randomized controlled trial evaluating the effectiveness of a multifaceted behavioral program to increase physical activity in Parkinson patients. *BMC Neurology* 2010;**10**:70.

van Nimwegen ML, Speelman AD, Gerrits MS, Bloem BR, Munneke M. Monitoring treatment fidelity in the ParkFit Study, a multifaceted behavioral program to increase physical activity in Parkinson patients. Movement Disorders 2010; Vol. 25 Suppl 3:S678.

Additional references

Allen 2011

Allen NE, Sherrington C, Paul SS, Canning CG. Balance and falls in Parkinson's disease: a meta-analysis of the effect of exercise and motor training. *Movement Disorders* 2011;**26**(9):1605-15.

Berg 1992

Berg KO, Wooddauphinee SL, Williams JI. Measuring balance in the elderly - validation of an instrument. *Canadian Journal of Public Health-Revue Canadienne de Sante Publique* 1992;**83**:S7-S11.

Boutron 2008a

Boutron I, Moher D, Altman DG, Schulz K, Ravaud P, for the CONSORT group. Methods and processes of the CONSORT Group: example of an extension for trials assessing nonpharmacologic treatments. *Annals of Internal Medicine* 2008;**148**(4):W60-6.

Boutron 2008b

Boutron I, Moher D, Altman DG, Schulz K, Ravaud P, for the CONSORT group. Extending the CONSORT statement to randomized trials of nonpharmacologic treatment: explanation and elaboration. *Annals of Internal Medicine* 2008:**148**(4):295-309.

Chartered Society of Physiotherapy

The Chartered Society of Physiotherapy. In: The Chartered Society of Physiotherapy, editor(s). Rules of professional conduct. 2nd Edition. London: The Chartered Society of Physiotherapy, 2002.

Crizzle 2006

Crizzle AM, Newhouse IJ. Is physical exercise beneficial for persons with Parkinson's disease?. *Clinical Journal of Sports Medicine* 2006;**16**(5):422-5.

Deane 2001a

Deane K, Jones DE, Playford ED, Ben-Shlomo Y, Clarke CE. Physiotherapy versus placebo or no intervention in Parkinson's disease. *Cochrane Database of Systematic Reviews* 2001, (3):CD002817.

Deboer 1996

Deboer AGEM, Wijker W, Speelman JD, Dehaes JCJM. Quality of life in patients with Parkinson's disease: Development of a questionnaire. *Journal of Neurology, Neurosurgery, and Psychiatry* 1996;**61**(1):70-4.

Doll 1980

Doll R, Peto R. Randomized controlled trials and retrospective controls. *British Medical Journal* 1980;**280**(6206):44.

Duncan 1990

Duncan PW, Weiner DK, Chandler J, Studenski S. Functional Reach - A new clinical measure of balance. *Journals of Gerontology* 1990;**45**(6):M192-7.

Fahn 1987

Fahn S, Elton RL. UPDRS Development Committee. Unified Parkinson's Disease Rating Scale. In: Fahn S, Marsden CD, Calne DB, Goldestein M editor(s). Recent developments in Parkinson's disease. Florham Park, NJ: Macmillan Health Care Information, 1987:153-63.

Gage 2004

Gage H, Storey L. Rehabilitation for Parkinson's disease: a systematic review of available evidence. *Clinical Rehabilitation* 2004;**18**(5):463-82.

Gibb 1988

Gibb WRG, Lees AJ. The relevance of the Lewy Body to the pathogenesis of idiopathic Parkinsons disease. *Journal of Neurology, Neurosurgery, and Psychiatry* 1988;**51**(6):745-52.

Giladi 2000

Giladi N, Shabtai H, Simon ES, Biran S, Tal J, Korczyn AD. Construction of freezing of gait questionnaire for patients with Parkinsonism. *Parkinsonism & Related Disorders* 2000;**6**(3):165-70.

Goodwin 2008

Goodwin VA, Richards SH, Taylor RS, Taylor AH, Campbell JL. The effectiveness of exercise interventions for people with



Parkinson's disease: A systematic review and meta-analysis. *Movement Disorders* 2008;**23**(5):631-40.

Hoehn 1967

Hoehn MM, Yahr MD. Parkinsonism - onset, progression and mortality. *Neurology* 1967;**17**(5):427-42.

Jenkinson 1997

Jenkinson C, Fitzpatrick R, Peto V, Greenhall R, Hyman N. The Parkinson's Disease Questionnaire (PDQ-39): development and validation of a Parkinson's disease summary index score. *Age and Ageing* 1997;**26**(5):353-7.

Kersten 2004

Kersten P. Principles of physiotherapy assessment and outcome measures. In: Stokes M editor(s). Physical management in neurological rehabilitation. 2nd Edition. London: Elsevier Mosby, 2004:29-46.

Keus 2004

Keus S, Hendriks HJ, Bloem BR, Bredero-Cohen AB, De Goede CJ, van Haaren M, et al. Clinical practice guidelines for physical therapy in patients with Parkinsons disease. *Dutch Journal of Physiotherapy* 2004;**114 Suppl 3**:1-94.

Keus 2007

Keus SHJ, Bloem BR, Hendriks EJM, Bredero-Cohen AB, Munneke M. Evidence-based analysis of physical therapy in Parkinson's disease with recommendations for practice and research. *Movement Disorders* 2007;**22**(4):451-60.

Keus 2009

Keus SHJ, Munneke M, Nijkrake MJ, Kwakkel G, Bloem BR. Physical therapy in Parkinson's disease: Evolution and future challenges. *Movement Disorders* 2009;**24**(1):1-14.

Kwakkel 2007

Kwakkel G, de Goede CJT, van Wegen EEH. Impact of physical therapy for Parkinson's disease: A critical review of the literature. *Parkinsonism & Related Disorders* 2007;**13**:S478-87.

Lim 2005

Lim I, van Wegen E, de Goede C, Deutekom M, Nieuwboer A, Willems A, et al. Effects of external rhythmical cueing on gait in patients with Parkinson's disease: a systematic review. *Clinical Rehabilitation* 2005;**19**(7):695-713.

Mehrholz 2010

Mehrholz J, Friis R, Kugler J, Twork S, Storch A, Pohl M. Treadmill training for patients with Parkinson's disease. *Cochrane Database of Systematic Reviews* 2010, (1):CD007830.

MRC 2008

Medical Research Council. Developing and evaluating complex interventions: new guidance. http://www.mrc.ac.uk/Utilities/Documentrecord/index.htm?d=MRC004871 2008.

Mutch 1986

Mutch WJ, Strudwick A, Roy SK, Downie AW. Parkinson's Disease - disability, review and management. *British Medical Journal* 1986;**293**(6548):675-7.

NICE CG35 2006

National Institute for Health and Clinical Excellence. NICE clinical guideline 35. Parkinson's disease: diagnosis and management in primary and secondary care. www.nice.org.uk 2006.

Nieuwboer 2008

Nieuwboer A, Rochester L, Jones D. Cueing gait and gait-related mobility in patients with Parkinson's disease - Developing a therapeutic method based on the international classification of functioning, disability, and health. *Topics in Geriatric Rehabilitation* 2008;**24**(2):151-65.

Nijkrake 2007

Nijkrake MJ, Keus SH, Kalf JG, Sturkenboom IH, Munneke M, Kappelle AC, Bloem BR. Allied health care interventions and complementary therapies in Parkinson's disease. *Parkinsonism & Related Disorders* 2007;**13 Suppl 3**:S488-94.

Patel 2010

Patel S, Cheed V, Ives N, Clarke C, Rick C, Wheatley K, Gray R, on behalf of the PD MED Collaborators. A large randomised assessment of the relative cost-effectiveness of different classes of drugs for Parkinson's disease (PD MED Trial). *Movement Disorders* 2010;**25 Suppl 3**:S681.

PDS 2008

PDS. Life with Parkinson's today - room for improvement. Parkinson's Disease Society, London 2008.

Perry 1995

Perry J, Garrett M, Gronley JK, Mulroy SJ. Classification of walking handicap in the stroke population. *Stroke* 1995;**26**(6):982-9.

Peto 1995

Peto V, Jenkinson C, Fitzpatrick R, Greenhall R. The development and validation of a short measure of functioning and well-being for individuals with Parkinson's disease. *Quality of Life Research* 1995;**4**(3):241-8.

Peto 2001

Peto V, Jenkinson C, Fitzpatrick R. Determining minimally important differences for the PDQ-39 Parkinson's disease questionnaire. *Age and Ageing* 2001;**30**:299-302.

Plant 2000

Plant R, Jones D, Walton G, Ashburn A, Lovgreen B, Handford F, Kinnear E. Physiotherapy for people with Parkinson's disease: UK best practice short report. Institution of Rehabilitation, University of Northumbria 2000.

Podsiadlo 1991

Podsiadlo D, Richardson S. The Timed Up and Go - A test of basic functional mobility for frail elderly persons. *Journal of the American Geriatrics Society* 1991;**39**(2):142-8.

Powell 1995

Powell LE, Myers AM. The Activities-Specific Balance Confidence (Abc) Scale. *Journals of Gerontology Series A, Biological Sciences and Medical Sciences* 1995;**50**(1):M28-34.



Qutubuddin 2005

Qutubuddin AA, Pegg PO, Cifu DX, Brown R, McNamee S, Carne W. Validating the Berg Balance Scale for patients with Parkinson's disease: A key to rehabilitation evaluation. *Archives of Physical Medicine and Rehabilitation* 2005;**86**(4):789-92.

Rascol 2002

Rascol O, Payoux P, Ferreira J, Brefel-Courbon C. The management of patients with early Parkinson's disease. *Parkinsonism & Related Disorders* 2002;**9**(1):61-7.

Robertson 2003

Robertson D. Developing and delivering services. In: Playford D, Thompson AJ editor(s). Neurological Rehabilitation of Parkinson's Disease. London: Martin Dunitz, 2003.

Robertson 2008

Robertson D, Aragon A, Moore G, Whelan L. Rehabilitation and the interdisciplinary team. In: Playfer J, Hindle J editor(s). Parkinson's Disease in the Older Patient. 2nd Edition. Oxford: Radcliffe Publishing Ltd, 2008.

Rubenis 2007

Rubenis J. A rehabilitational approach to the management of Parkinson's disease. *Parkinsonism & Related Disorders* 2007;**13 Suppl 3**:S495-7.

Schrag 2000

Schrag A, Jahanshahi M, Quinn N. How does Parkinson's disease affect quality of life? A comparison with quality of life in the general population. *Movement Disorders* 2000;**15**(6):1112-8.

Schrag 2006

Schrag A, Sampaio C, Counsell N, Poewe W. Minimal clinically important change on the Unified Parkinson's Disease Rating Scale. *Movement Disorders* 2006;**21**(8):1200-7.

Shulman 2010

Shulman LM, Gruber-Baldini AL, Anderson KE, Fishman PS, Reich SG, Weiner WJ. The clinically important difference on the Unified Parkinson's Disease Rating Scale. *Archives of Neurology* 2010;**67**(1):64-70.

Steffen 2008

Steffen T, Seney M. Test-retest reliability and minimal detectable change on balance and ambulation tests, the 36-Item Short-Form Health Survey, and the Unified Parkinson Disease Rating Scale in people with parkinsonism. *Physical Therapy* 2008;**88**(6):733-46.

Talley 2008

Talley KMC, Wyman JF, Gross CR. Psychometric properties of the activities-specific balance confidence scale and the survey of activities and fear of falling in older women. *Journal of the American Geriatrics Society* 2008;**56**(2):328-33.

Tinetti 1990

Tinetti ME, Richman D, Powell L. Falls efficacy as a measure of fear of falling. *Journals of Gerontology* 1990;**45**(6):239-43.

Tomlinson 2012

Tomlinson CL, Patel S, Meek C, Clarke CE, Stowe R, Shah L, et al. Physiotherapy versus placebo or no intervention in Parkinson's disease. *Cochrane Database of Systematic Reviews* 2012, (8):CD002817.

Trew 2005

Trew M, Everett T. Human movement: an introductory text. 5th Edition. Edinburgh: Churchill Livingstone, 2005.

Ware 1992

Ware JE, Sherbourne CD. The MOS 36-Item Short-Form Health Survey (Sf-36) .1. Conceptual framework and item selection. *Medical Care* 1992;**30**(6):473-83.

Webster 1968

Webster DD. Critical analysis of disability in Parkinson's disease. *Modern Treatment* 1968;**5**(2):257-82.

Welsh 2003

Welsh M, McDermott MP, Holloway RG, Plumb S, Pfeiffer R, Hubble J, et al. Development and testing of the Parkinson's disease quality of life scale. *Movement Disorders* 2003;**18**(6):637-45.

Whittle 1996

Whittle M. Gait analysis: an introduction. 2nd Edition. Oxford: Butterworth-Heinemann, 1996.

Yahr 1969

Yahr MD, Duvoisin RC, Schear MJ, Barrett RE, Hoehn MM. Treatment of Parkinsonism with levodopa. *Archives of Neurology* 1969;**21**(4):343-54.

Yardley 2005

Yardley L, Beyer N, Hauer K, Kempen G, Piot-Ziegler C, Todd C. Development and initial validation of the Falls Efficacy Scale-International (FES-I). *Age and Ageing* 2005;**34**(6):614-9.

Yarrow 1999

Yarrow S. Members' 1998 survey of the Parkinson's Disease Society of the United Kingdom. In: Percival R, Hobson P editor(s). Parkinson's disease: Studies in psychological and social care. Leicester: BPS Books, 1999:79-92.

References to other published versions of this review

Deane 2001b

Deane K, Jones DE, Ellis-Hill C, Clarke CE, Playford ED, Ben-Shlomo Y. Physiotherapy for Parkinson's disease: a comparison of techniques. *Cochrane Database of Systematic Reviews* 2001, (1):CD002815.

^{*} Indicates the major publication for the study



CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Almeida 2012

Methods	Parallel group design
	Randomised by pulling allocation out of a hat
	Analysed on a per protocol basis
	Treated as outpatients for 9 hours over 6 weeks
	Assessed at baseline, 6 weeks and 12 weeks
	Assessors were blinded for UPDRS III evaluation
Participants	14 participants in the Overground walking group (OG), 14 in the Treadmill walking group (TM) and 14 in the control group (CL). 2 dropouts in TM group, 1 dropout in CL group.
	Participants' mean age 73.9 years (OG), 63.9 years (TM), 67.4 years (CL); male/female 12/2 (OG), 8/6 (TM), 11/3 (CL); Hoehn and Yahr stage not stated; duration of PD not stated.
	Inclusion criteria: Confirmed as having clinically typical Parkinson's disease by at least one movement disorders neurologist. Exclusion criteria: Past history of neurological conditions other than Parkinson's disease, orthopaedic or visual disturbances that severely impaired walking ability, unable to independently walk down an 8 metre GAITRite carpet for a total of 10 trials.
Interventions	OG: Walk down equally spaced transverse lines presented on a 16m carpet. The cues were white lines of tape. Participants asked to walk across the lines, turn and continue back. Spacings were set at 8% greater than the initial step length of any of the groups (70 cm). 30 minute session with mandatory 2 min break every 8 mins, additional rest allowed if necessary but a total of 24 mins walking was required to consider gait session complete.
	TM: Walk on a treadmill presented with equally distributed standardised transverse white lines. Spacings were set at 8% greater than the initial step length of any of the groups (70 cm). 30 minute session with mandatory 2 min break every 8 mins, additional rest allowed if necessary but a total of 24 mins walking was required to consider gait session complete.
	CL: Instructed to continue their usual activities.
	Participants were optimally medicated at time of all training and testing sessions and remained on stable regimen throughout trial period.
Outcomes	Step length
	UPDRS III
	Timed up and go
	Gait speed
	Cadence
	Double support time
	Step time
	Step-to-step variability, step time variability
	30 second chair stand
Notes	CL arm not included in review



Almeida 2012 (Continued)

Risk of bias

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Low risk	Inclusion and exclusion criteria stated
Randomisation Method	High risk	Allocation pulled out of hat
Concealment of Allocation	High risk	Allocation pulled out of hat
Similarity at Baseline	Low risk	Group characteristics similar at baseline
Intention to Treat Analysis	Unclear risk	Analysed on a per protocol basis
Withdrawals Described	Low risk	Withdrawals at less than 10%
Cointerventions Constant	Low risk	Participants maintained stable drug regiment throughout trial period
Comparable arms	Low risk	Same time and attention given to both arms
Blinded Assessors	Unclear risk	Assessors blind for UPDRS III evaluation only

Braun 2011

Methods	Parallel group design
	Random allocation list used for each site with block sizes of 4
	Both intention-to-treat and per protocol analysis were carried out
	Treated as outpatients for either 1 hour per week (groups) or 30 mins per week (individuals) over 6 weeks
	Assessed at baseline (week 0-1), at 7-8 weeks and at 3 months (12-13 weeks)
	Assessors were blinded
Participants	25 participants in the physiotherapy with mental practice group (PT+MP) and 22 in the physiotherapy and relaxation group (PT+R). There were 3 dropouts PT+MP group and 4 in the PT+R group prior to the post-intervention assessment, and a further 4 dropouts in the PT-MP group and 3 in the PT-R group prior to the 3 month follow-up assessment.
	Participants' mean age 70 years (PT-MP), 69 years (PT-R); Male/female ratio, 17/8 (PT-MP), 15/7 (PT-R); Mean Hoehn and Yahr not given; Mean duration of PD 5.2 years (PT-MP), 6.6 years (PT-R).
	Inclusion criteria: Clinically diagnosed adults with PD, sufficient cognitive level and communication skills to engage mental practice. Exclusion criteria: Other conditions such as stroke, rheumatic disease or dementia prior to onset of PD and sufficient to cause persistent premorbid disability.
Interventions	Participants entering the trial were already receiving physiotherapy. This pre-existing treatment was continued. The randomly allocated new treatment was incorporated into the participants existing program.
	PT-MP: In half hour sessions 10 mins were spent on mental practice, in group sessions of 1 hour, 20 mins were spent on mental practice. Therapy was recorded in pre-structured files which detailed content and duration. As soon as possible therapists encouraged unguided mental practice. Logs were given to participants to record unguided mental practice behaviour. The main goal of mental practice was to improve locomotor tasks like walking, standing up from a chair or the floor. This was achieved



Braun 2011 (Continued)

through four steps, explaining the concept, developing imagery techniques, applying mental practice and consolidating. During therapy imagery attempts and overt movements were combined. This information was embedded in the imagery attempts to make them as vivid as possible. The proportions of actual movements and imagery attempts were based on individual preferences.

PT-R: In half hour sessions 10 mins were spent on relaxation, in group sessions of 1 hour, 20 mins were spent on relaxation. Therapy was recorded in pre-structured files. As soon as possible therapists encouraged unguided relaxation. Logs were given to participants to record unguided practice. Used to control for attention and consisted of treatment according to the national Dutch guidelines with relaxation therapy being incorporated into each session. The amount of relaxation incorporated matched the amount of mental practice in the experimental group. Relaxation followed the principles of progressive muscle relaxation.

It was not stated whether drug therapy was kept constant during the trial.

Outcomes

Patient and therapist perceived effect on walking performance using visual analogue scale

Timed up and go

10 m walk test

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Low risk	Inclusion and exclusion criteria
Randomisation Method	Unclear risk	Random allocation list not clear how generated
Concealment of Allocation	Low risk	Lists kept by independent 3rd party
Similarity at Baseline	Unclear risk	Hoehn and Yahr not stated
Intention to Treat Analysis	Unclear risk	Primary analysis was ITT then per protocol was carried out
Withdrawals Described	High risk	Withdrawals > 15% prior to post-intervention assessment
Cointerventions Constant	Unclear risk	Not stated whether drug therapy was kept constant during the trial
Comparable arms	Low risk	Same time and attention given in both arms
Blinded Assessors	Low risk	Assessors were blinded

Burini 2006

Methods Cross-over design

Participants coupled consecutively with list of random numbers. Numbers correspond to a sealed envelope containing group allocation

Method of analysis not stated

Treatment delivered to groups in 45-50 minute sessions, 3 times per week up to a total of 20 sessions

Assessed at Baseline, after 7 weeks of first treatment, after 8 weeks wash-out period, after 7 weeks of second treatment



Burini 2006 (Continued)	Assessors were blinded
Participants	13 participants in the aerobic training group and 13 in the Qi-gong group. There were 2 dropouts in each group.
	Participants' mean age 65.7 years (aerobic training), 62.7 years (Qi-gong); Male/female ratio, 5/8 (aerobic training), 4/9 (Qi-gong); Mean Hoehn and Yahr 2.8 (aerobic training), 2.7 (Qi-gong); Mean duration of PD 11.2 years (aerobic training), 10.6 years (Qi-gong).
	Inclusion criteria: PD subjects, stable medication, Hoehn and Yahr stage II to III. Exclusion criteria: Severe cognitive impairment (MMSE < 24), concomitant severe neurologic cardiopulmonary or orthopaedic disorders, specific contraindication to the execution of a cardiopulmonary test or aerobic training, recent participation in any physiotherapy/rehabilitation program during 2 months prior to start of trial.
Interventions	Aerobic training: Cycle ergometer used with warm-up, endurance and cool-down phases in each 45 min session occurring 3 times per week.
	Qi-gong: Breathing exercises, stretches, neck and trunk rotation exercises and balance training in the upright position, 50 minute sessions 3 times per week.
	Drug regimen remained constant throughout trial period.
Outcomes	UPDRS III
	UPDRS II
	Brown's disability scale (BDS)
	6-minute walk test
	Borg scale for breathlessness
	Beck depression inventory (BDI)
	PDQ-39

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Low risk	Inclusion and exclusion criteria stated
Randomisation Method	Unclear risk	Not stated how random number list was generated
Concealment of Allocation	High risk	Sealed envelopes
Similarity at Baseline	Low risk	Group characteristics similar at baseline
Intention to Treat Analysis	Unclear risk	Method of analysis not stated
Withdrawals Described	High risk	Dropouts at 15%
Cointerventions Constant	Low risk	Drug regimen remained constant throughout trial period

Spirometry test

Maximum cardiopulmonary exercise test



Burini 2006 (Continued)		
Comparable arms	Low risk	Same time and attention given to both groups
Blinded Assessors	Low risk	Assessors blinded, patients instructed not to discuss treatment schedule

Methods	Parallel group design
	Randomisation method not stated
	Method of analysis not stated
	Treated as outpatients for 3 sessions per week and using home practice for 3 sessions per week each lasting 30 minutes over 4 weeks plus 4 week home program
	Assessed at baseline, immediately after treatment (4 weeks), and after home program at 8 weeks
	Assessors were blinded
Participants	10 participants in the treadmill + music + home-walking (TMH) group, 10 participants in the treadmill and home-walking (TH) group, 10 participants in the home-walking group (H). No dropouts described.
	Participants' mean age 67.1 years (TMH group), 67.9 years (TH group), 68.6 years (H group); Male/female ratio 10/0 (TMH group), 10/0 (TH group), 10/0 (H group); Mean Hoehn and Yahr score 2.3 (TMH group), 2.5 (TH group), 2.1 (H group); Mean duration of PD 3.7 years (TMH group), 7.4 years (TH group), 4.4 years (H group).
	Inclusion criteria: Male, 60-80 years old, PD as diagnosed by attending neurologists, Thai Mental State Examination > 23, stable medications without freezing, no exercise program in last 2 months, no contraindication for exercise, H&Y II-III, independent walking without use of gait aids. Exclusion criteria: Medication change during study, inability to walk on treadmill, cannot complete 80% of prescribed program.
Interventions	TMH: 10 minutes of stretching followed by 20 minutes of treadmill training with cueing (music) delivered 3 days per week and home-walking practised 3 days per week. After treadmill speed was set, the step frequency would be identified by adjustable electrical metronome. The prepared music with the same rhythmic frequency was chosen. Then the participants were training to walk synchronised with the matched music rhythm on the treadmill. This music would be recorded in MP3 for the participants to take home and listen during home practice. Home program consisted of 10 minutes of stretching and 20 minutes walking. After 4 weeks treadmill training ceased but home program was continued for a further 4 weeks.
	TH: 10 minutes of stretching followed by 20 minutes of treadmill training without cueing delivered 3 days per week and home-walking, as described above, practised 3 days per week. After 4 weeks treadmill training ceased but home program was continued for a further 4 weeks.
	H: Home-walking practised 6 days per week.
	Patients were excluded if medication was changed during trial period. Patients did not participate in training in 2 months prior to study start.
Outcomes	Timed up and go
	Walking speed
	Step length
	Cadence
	Stride length



Chaiwanichsiri 2011 (Continued)

6-minute walk test

6 metre walk time

Single leg stance

UPDRS I, II, III

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Low risk	Inclusion and exclusion criteria stated
Randomisation Method	Unclear risk	Method not stated
Concealment of Allocation	Unclear risk	Method of randomisation not stated
Similarity at Baseline	Low risk	Group characteristics similar at baseline
Intention to Treat Analysis	Unclear risk	Analysis method not stated
Withdrawals Described	Unclear risk	Withdrawals not described
Cointerventions Constant	Low risk	Patients were excluded if medication changed during trial period
Comparable arms	Unclear risk	Home group no therapist time
Blinded Assessors	Low risk	Assessors were blinded

Dias 2005

Methods	Parallel group design					
	Randomisation method not stated					
	Method of analysis not stated					
	Treatment delivered in a total of 20 hour long sessions					
	Assessed at baseline, immediately after treatment and 30 days after end of treatment					
	Not stated whether assessors were blinded					
Participants	8 participants in the physiotherapy and cardiovascular exercise with visual cues (PTCV) group and 8 in the physiotherapy only (PT) group. No dropouts described.					
	Participants mean age 61.5 years (PTCV group), 64.3 years (PT group); Male/female ratio 4/4 (PTCV group), 7/1 (PT group); Mean Hoehn and Yahr 1.6 (PTCV group), 1.7 (PT group); Mean duration of PD 7.4 years (PTCV group), 8.4 years (PT group).					
	Inclusion criteria: Hoehn and Yahr stage ≤3. Exclusion criteria: Any degree of dementia, joint deformations, arthritis, severe pain, other neurological disturbance, submitted to neurological surgery, severe associated pathologies that would impair physiotherapy.					



Dias 2005 (Continued)

Interventions PTCV: 20 sessions of physiotherapy following a protocol that included 15 mins muscular stretching,

30 mins gait training on stable ground with visual cues, 15 mins cardiovascular exercise using ergomet-

ric bicycle.

PT: Conventional physiotherapy.

It was not stated whether drug therapy was kept constant during the trial.

Outcomes UPDRS

Functional independence measurement scale

Berg balance scale

Hoehn and Yahr scale

Notes Article in Portuguese

Risk of bias

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Low risk	Inclusion and exclusion criteria stated
Randomisation Method	Unclear risk	Method of randomisation not stated
Concealment of Allocation	Unclear risk	Method of randomisation not stated
Similarity at Baseline	Low risk	Group characteristics similar at baseline
Intention to Treat Analysis	Unclear risk	Method of analysis not stated
Withdrawals Described	Unclear risk	Number of dropouts not stated
Cointerventions Constant	Unclear risk	Not stated whether drug regimen remained constant throughout trial period
Comparable arms	Low risk	Same number of sessions of active therapy in both arms
Blinded Assessors	Unclear risk	Not stated whether assessors were blinded

Diehl 2011

Methods	Parallel group design		
	Randomisation method not stated		
	Analysis method not stated		
	Treated for 24-36 sessions, lasting 90 mins each over 12 weeks		
	Assessed at baseline and immediately after treatment		
	Not stated whether assessors were blinded		
Participants	20 participants were recruited, group split not stated. Dropouts not stated.		
	Baseline characteristics not stated but it was reported that no significant differences were found between groups in pre-test demographic variables.		



Diehl 2011 (Continued)			
	Inclusion and exclusion	n criteria not stated.	
Interventions	Group box training: Stretching, lateral foot work, punching various targets, resistance exercises and aerobic training.		
	Traditional group exercises: Stretching, resistance exercises, aerobic and balance activities.		
	It was not stated whether drug therapy was kept constant during the trial.		
Outcomes	Berg balance scale		
	Activities-specific balance confidence scale		
	Functional reach test		
	Parkinson's disease quality of life scale		
Notes	Abstract only		
Risk of bias			
Bias	Authors' judgement	Support for judgement	
Eligibility Criteria	Unclear risk	Eligibility criteria not stated	
Randomisation Method	Unclear risk	Method not stated	

Eligibility Criteria	Unclear risk	Eligibility criteria not stated
Randomisation Method	Unclear risk	Method not stated
Concealment of Allocation	Unclear risk	Randomisation method not stated
Similarity at Baseline	Low risk	No significant differences were found between groups in pretest demographic variables
Intention to Treat Analysis	Unclear risk	Method of analysis not stated
Withdrawals Described	Unclear risk	Withdrawals not described
Cointerventions Constant	Unclear risk	Not stated whether drug therapy was kept constant during the trial
Comparable arms	Low risk	Same time and attention given to both groups
Blinded Assessors	Unclear risk	Not stated whether assessors were blinded

Ebersbach 2010

Methods	Parallel group design		
	Randomly allocated by drawing lots		
	Analysed on a per protocol basis		
	Treatment took place for 1 hour sessions 4 times per week for 4 weeks		
	Assessed at baseline and at 16 weeks		
	Assessors blinded for UPDRS III, not stated for other outcomes		
Participants	20 participants in LSVT BIG (BIG) group, 19 participants in Nordic walking (WALK) group and 19 participants in the Home exercise (HOME) group. 1 dropout in WALK group and 1 in HOME group.		



Ebersbach 2010 (Continued)

Participants mean age 67.1 years (LSVT BIG group), 65.5 years (WALK group), 69.3 years (HOME group); Male/female ratio 7/13 (LSVT-BIG group), 7/12 (WALK group), 8/11 (HOME group); Mean Hoehn and Yahr 2.8 (LSVT BIG group), 2.6 (WALK group), 2.5 (HOME group); Mean duration of PD 6.1 years (LSVT-BIG group), 7.8 years (WALK group), 7.4 years (HOME group).

Inclusion criteria: Fulfil diagnostic criteria for IPD, Hoehn & Yahr stage I-III outpatient treatment, stable medication 4 weeks prior to inclusion. Exclusion criteria: Dementia (MMSE<25), severe depression, disabling dyskinesias, comorbidity affecting mobility or ability to exercise.

Interventions

LSVT BIG: 50% of exercises consisted of standardized whole-body movements with maximal amplitude, repetitive multidirectional movements and stretching. 50% of exercise included goal-directed ADL according to individual needs and preferences. ADL were performed using high amplitude LSVT BIG movements. LSVT BIG was delivered one-to-one with intensive motivation and feedback. Participants were constantly encouraged to work with at least '80% of their maximal energy' on every repetition and taught to use bigger movements in routine activities to provide continuous exercise in every-day movements. Participants in all groups were encouraged to exercise regularly at home. Diaries were used to document type and duration of exercise performed in addition to supervised LSVT BIG.

WALK: Each session consisted of a standardized protocol for beginners including warming up, practicing Nordic walking and finally a cooling down. Sessions were performed in a local park in groups of 4 or 6 and constantly supervised by the therapist. Participants in all groups were encouraged to exercise regularly at home. Diaries were used to document type and duration of exercise performed in addition to supervised WALK sessions.

HOME: 1 hours instruction of domestic training with practical demonstration and training. Exercises included stretching, high-amplitude movements, as well as active workouts for muscular power and posture. Participants in all groups were encouraged to exercise regularly at home. They received a diary to document type and duration of exercise performed.

Changes in medication occurred in 6 patients from each group.

Outcomes

UPDRS III

PDQ-39

Timed up and go

Time to walk 10 m

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Low risk	Inclusion and exclusion criteria stated
Randomisation Method	High risk	Randomised by drawing lots
Concealment of Allocation	High risk	Randomised by drawing lots
Similarity at Baseline	Low risk	Group characteristics similar at baseline
Intention to Treat Analysis	Unclear risk	Analysed on a per protocol basis
Withdrawals Described	Low risk	Withdrawals <10%
Cointerventions Constant	Unclear risk	Minor changes to medication
Comparable arms	Unclear risk	Time and attention greater in LSVT BIG and WALK groups than in HOME group



Ebersbach 2010 (Continued)

Blinded Assessors Unclear risk Assessors blinded for primary outcome, not stated for secondary outcomes

Fisher 2008

Methods

Parallel group design

Subjects closed their eyes and selected a card corresponding to 1 of the 3 groups

Analysis method not stated

Treated as outpatients for 24 sessions over 8 weeks for both treatment arms, 6 sessions over 8 weeks

for control group

Assessed at baseline and immediately post-treatment

Assessors were blinded

Participants

10 participants in treadmill group, 10 participants in physiotherapy group and 10 in the control arm. No dropouts.

Participants' mean age, 64.0 years (treadmill), 61.5 years (physiotherapy), 63.1 years (control). Male/female ratio, 6/4 (treadmill), 5/5 (physiotherapy), 8/2 (control). Mean Hoehn and Yahr 1.9 in all 3 groups. Mean duration of PD 1.2 years (treadmill), 0.7 years (physiotherapy), 1.5 years (control).

Inclusion criteria: Early stage PD, diagnosis of PD within 3 years of study participation, Hoehn and Yahr stage 1 or 2, 18 years or older, medical clearance from primary care physician to participate in exercise programme, ability to walk. Exclusion criteria: Medical or physical screening examination showed a score of less than 24 on the MMSE, there were physician determined major medical problems such as cardiac dysfunction that would interfere with participation, they had musculoskeletal impairments or excessive pain in any joint that could limit participation in an exercise programme, had insufficient endurance and stamina to participate in exercise 3 times per week for a 1 hour session.

Interventions

Treadmill: Level of intensity was defined by metabolic equivalents (MET). High intensity exercise greater than 3 METs. Body weight supported (BWS) treadmill training. Goal of each session was to reach and maintain a MET > 3. Exercise progressed by decreasing BWS (initially 10% of participants' bodyweight) and physical assistance, increasing the treadmill speed and time on the treadmill, with the end goal for each participant to walk on the treadmill continuously for 45 min within the MET range.

Physiotherapy: Less than 3 METs. This group was representative of general or traditional physical therapy. Each 45 min session was individualised and consisted of activities from 6 categories 1) passive range of motion and stretching 2) active range of motion 3) balance activities 4) gait 5) resistance training 6) practice of functional activities and transitional movements.

Control: Zero intensity group. Six 1 hour education classes taken over an 8 week period.

All participants were allowed to continue their customary exercise routines and filled out a daily exercise diary.

Drug therapy was constant during the trial.

Outcomes

UPDRS (Total, I, II and III subscores)

Hoehn and Yahr

Functional assessments

Walking tests: average gait velocity, step length, stride length, cadence, double limb support time, ankle, knee, hip rotation

Sit-to-stand test



Fis	her 2008	(Continued)
-----	----------	-------------

Transcranial magnetic stimulation (subset)

All participants took their customary medications at the same time relative to each assessment

Notes Control arm not included in this review

Risk of bias

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Low risk	Inclusion and exclusion criteria stated
Randomisation Method	High risk	Subjects self-selected a card with eyes closed
Concealment of Allocation	High risk	Subjects self-selected a card with eyes closed
Similarity at Baseline	Low risk	Group characteristics similar at baseline
Intention to Treat Analysis	Unclear risk	Method of analysis not described
Withdrawals Described	Low risk	No dropouts
Cointerventions Constant	Low risk	All medication kept stable during course of study
Comparable arms	Low risk	Same time and attention given to participants in both arms
Blinded Assessors	Low risk	Assessors were blinded to group allocation

Frazzitta 2009

Methods	Parallel group design		
	Randomisation method not stated		
	Analysis method not stated		
	Treatment was delivered over 4 weeks in 28 sessions lasting 20 mins each		
	Assessed at baseline and immediately after treatment		
	Not stated whether assessors were blinded		
Participants	20 participants in treadmill with cues group, 20 in Auditory and visual cues group. No dropouts.		
	Participants mean age 71 years for both groups; Male/female ratio not stated; Mean Hoehn and Yahr 3 for both groups; Mean duration of condition 13.2 years (treadmill group), 12.9 years (cued group).		
	Inclusion criteria: Clinically probable IPD, ability to walk without assistance, visual and hearing capacity sufficient to perceive the cues, freezing of gait at the time of peak medication, stable pharmacological treatment, Hoehn & Yahr stage III, no cognitive impairment, MMSE > 26. Exclusion criteria: Neurological condition other than IPD, postural hypertension, cardiovascular disorders, musculoskeletal disorders, vestibular dysfunction limiting locomotion or balance.		
Interventions	Treadmill: Rehabilitation protocol for gait disturbance & freezing that used treadmill training associated with auditory & visual cues. Maximum tolerated walking speed -40% used for 2 day warm-up, then increased by 0.05 stride cycles/second. Visual cue of target on screen that participant had to reach with stride, auditory cue of music at matched frequency.		



Cued: Visual cue of lines spaced according to individual stride length for gait training coupled with auditory cue of music at frequency matching that used in Treadmill group.

It was not stated whether drug therapy was kept constant during the trial.

Outcomes UPDRS III

Gait speed

Freezing of gait questionnaire

Stride length

6 minute walking test

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Low risk	Inclusion and exclusion criteria stated
Randomisation Method	Unclear risk	Method not stated
Concealment of Allocation	Unclear risk	Randomisation method not stated
Similarity at Baseline	Low risk	Group characteristics similar at baseline
Intention to Treat Analysis	Unclear risk	Method of analysis not stated
Withdrawals Described	Low risk	No dropouts
Cointerventions Constant	Unclear risk	Not stated whether drug regimen remained constant throughout trial period
Comparable arms	Low risk	Same time and attention given to participants in both arms
Blinded Assessors	Unclear risk	Not stated whether assessors were blinded

Hackney 2007

Methods Parallel group design			
	Randomisation method not stated		
	Method of analysis not given		
	Treated as outpatients for 20 sessions each lasting 1 hour delivered over 13 weeks		
	Assessed 1 week prior to treatment and 1 week after treatment		
	Assessors were blinded		
Participants	9 participants in Tango group, 10 in Exercise group. No dropouts were reported.		
	Participants mean age 72.6 years (Tango), 69.6 years (exercise); Male/female ratio 6/3 (Tango), 6/4 (exercise); Hoehn and Yahr 2.3 (Tango), 2.2 (exercise); Mean duration of condition 6.2 years (Tango), 3.3 years (exercise).		



Hackney 20	07 (Continued)
------------	----------------

Inclusion criteria: Diagnosed with clinically defined IPD, clear benefit from PD medications.

Interventions

Tango: Progressive Tango dance lessons with postural stretches, balance exercises, Tango-style, footwork patterns and experimentation with timing of steps to music with and without partner.

Exercise: Structured flexibility exercise classes designed for people with PD and/or elderly individuals. Breathing and stretching exercises progressed to resistance and dexterity exercises sometimes using water bottles or yard sticks to provide resistance or leverage. Some exercises done standing or using chair for support and last 10 minutes consisted of core strengthening exercises using floor matts or modified exercises in chair.

Drug therapy was kept constant during the trial.

Outcomes

UPDRS III

Berg balance scale

Freezing of gait

Timed up and go

Velocity of walking and dual-task walking

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Low risk	Inclusion criteria stated, clear definition of disease
Randomisation Method	High risk	Numbers drawn from a hat
Concealment of Allocation	High risk	Numbers drawn from a hat
Similarity at Baseline	Low risk	Group characteristics similar at baseline
Intention to Treat Analysis	Unclear risk	Method of analysis not given
Withdrawals Described	Low risk	No dropouts reported
Cointerventions Constant	Low risk	Drug regimen remained constant throughout trial period
Comparable arms	Low risk	Same time and attention given to participants in both arms
Blinded Assessors	Low risk	Assessors were blinded

Hackney 2009

Methods Parallel group design

Randomised by selecting allocation from a hat, carried out by trial investigator

Method of analysis not given

Treatment delivered in 1 hour sessions twice weekly for a total of 20 sessions

Assessed at baseline and within 1 week of completing 20 classes



Hac	kney	2009	(Continued)
-----	------	------	-------------

Assessors were blinded

Participants

19 participants in tango group, 19 in Waltz/Foxtrot group, 17 in Tai Chi and 20 in control. 5 dropouts in Tango group, 2 dropouts in Waltz/Foxtrot, 4 dropouts in Tai Chi and 3 in control group.

Participants mean age 68.2 years (Tango group), 66.8 years (Waltz/Foxtrot group), 64.9 (Tai Chi), 66.5 (control); Male/female ratio 11/3 (Tango group), 11/6 (Waltz/Foxtrot), 11/2 (Tai Chi),12/5 (control); Hoehn & Yahr 2.1 (Tango), 2.0 (Waltz/Foxtrot), 2.0 (Tai Chi), 2.2 (control); Mean duration of condition 6.9 years (Tango), 9.2 years (Waltz/Foxtrot), 8.7 (Tai Chi), 5.9 (control).

Inclusion criteria: At least 40 years of age, could stand for at least 30 min, could walk independently 3 or more metres with or without assistive device, diagnosis of IPD using diagnostic criteria for critically defined "definite PD" based upon published standards, patients demonstrated clear benefit from L-dopa, Hoehn & Yahr I-III. Exclusion criteria: History of neurological deficit other than PD, patients had been previously screened for dementia by their neurologists and none were diagnosed with dementia.

Interventions

Waltz/Foxtrot and Progressive Tango: Experience professional ballroom dancer taught progressive Tango, Waltz/Foxtrot lessons. Instructor, equally versed in both dances, attempted to give all students equal time in leading and following dance roles. All steps done in closed practice position, participants maintain contact through upper extremities and face one another.

Tai Chi: Progressive lessons on Tai Chi's 1st and 2nd circles of the Yang short style of Cheng Manching taught by experienced instructor.

Control: No intervention.

Drug therapy was constant during the trial or patients were excluded.

Outcomes

PDQ-39

UPDRS III

Berg balance scale

Tandem stance test

Timed up and go test

One leg stance test

6 minute walk test

Gait

Notes

Control arm not included in this review

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Low risk	Inclusion and exclusion criteria stated
Randomisation Method	High risk	Allocation pulled out of hat
Concealment of Allocation	High risk	Allocation pulled out of hat
Similarity at Baseline	Low risk	Group characteristics similar at baseline
Intention to Treat Analysis	Unclear risk	Method of analysis not given
Withdrawals Described	High risk	Withdrawals at 19%



Hackney 2009 (Continued)		
Cointerventions Constant	Low risk	Patients excluded if change in medication was required
Comparable arms	Low risk	Same time and attention given to both groups
Blinded Assessors	Low risk	Assessors blinded

Hackney 2010

Methods	Parallel group design
	Randomised by selecting allocation from a hat
	Analysed on an intention-to-treat basis
	Treatment was delivered in 1 hour classes twice a week for 10 weeks
	Assessed in the week prior to start of treatment, the week following completion of 10 week treatment and 4 weeks after completion of treatment
	Assessors were blinded
Participants	19 participants in the partnered tango group, 20 in the non-partnered tango group. 7 dropouts in partnered group, 5 in non-partnered group.
	Participants mean age 69.6 years in both groups; Male/female ratio 13/6 (partnered), 15/5 (non-partnered); Mean Hoehn and Yahr 2.5 (partnered) 2.0 (non-partnered); Duration of condition 9.5 years (partnered), 7.9 years (non-partnered).
	Inclusion criteria: PD without a history of other neurological deficits, at least 40 years of age, able to stand for at least 30 minutes and walk independently for 3 or more metres with or without an assistive device, diagnosis of IPD Hoehn and Yahr stages I-III using diagnostic criteria for clinically defined 'definite PD' and demonstrated clear benefit from L-dopa. Exclusion criteria not stated.
Interventions	Both partnered and non-partnered groups began with identical warm-ups to upbeat latin music. After warm-up both classes listened to and danced to identical commercial tango music selections in the same order of presentation.
	Partnered: both sexes spent equal time leading and following dance steps, performed in a 'closed practice' position, an adaptation of the traditional ballroom frame. Participants with PD always danced with individuals without PD. These individuals included caregivers and loved ones who elected to participate in classes as well as young adult volunteers.
	Non-partnered: learned the same Argentine 'leading' and 'following' tango-based steps as the partner group but performed them without a partner. Caregivers, loved ones and volunteers participated in the nonpartner class also.
	Participants remained on a steady drug regimen throughout the study
Outcomes	Tandem stance
	One leg stance
	Timed up and go
	6 minute walk test
	Gait velocity
	Cadence



Hackney 2010 (Continued)

Stride length

Swing percentage

Double support percentage

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Low risk	Inclusion criteria stated
Randomisation Method	High risk	Randomised by selecting allocation from a hat
Concealment of Allocation	High risk	Randomised by selecting allocation from a hat
Similarity at Baseline	Low risk	Group characteristics similar at baseline
Intention to Treat Analysis	Unclear risk	Analysed on an intention-to-treat basis
Withdrawals Described	High risk	Withdrawals at 31%
Cointerventions Constant	Low risk	Participants remained on a steady drug regimen throughout the study
Comparable arms	Low risk	Same time and attention given to participants in both arms
Blinded Assessors	Low risk	Assessors were blinded

Hass 2006

Methods	Parallel group design		
	Randomisation method not stated		
	Analysis method not stated		
	Treatment delivered in 1 hour sessions twice a week for a total of 32 sessions		
	Assessment intervals not stated		
	Investigators were blinded		
Participants	23 participants were recruited into the study. Number of dropouts was not stated.		
	Baseline characteristics for groups not stated. Total mean age 67 years; Hoehn Yahr 2.2.		
	Inclusion criteria: IPD. Exclusion criteria not stated.		
Interventions	Tai Chi: Emphasised physical movements, mind/body coordination & meditation. 8 forms of Tai Chi performed in sessions.		
	Qi-gong: Emphasised prolonged intense contemplative or deep meditation in 2 postures, seated or lying supine on floor mats.		
	It was not stated whether drug therapy was kept constant during the trial.		
Outcomes	Gait initiation		



Hass 2006 (Continued)

Gait velocity

Stride length

Stance

Double limb support

Step duration

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Unclear risk	Inclusion criteria IPD only
Randomisation Method	Unclear risk	Method not stated
Concealment of Allocation	Unclear risk	Method of randomisation not stated
Similarity at Baseline	Unclear risk	Baseline characteristics not split by group
Intention to Treat Analysis	Unclear risk	Method of analysis not stated
Withdrawals Described	Unclear risk	Withdrawals not described
Cointerventions Constant	Unclear risk	Not stated whether drug therapies were kept constant throughout trial period
Comparable arms	Low risk	Same time and attention given to participants in both arms
Blinded Assessors	Low risk	Single blind study

Hirsch 1996

Methods	Parallel group design Randomised according to a computer generated random number list but concealment of allocation not stated
	Analysed on a per protocol basis Treated as outpatients for an unspecified period of time (3 days/week) for 10 weeks Assessed at baseline, immediately after treatment and 4 weeks later
	Assessors were not blinded
Participants	6 patients in novel combined balance and resistance training group and 9 patients in balance training group. No dropouts stated. Patients mean age 70.8 years (combined), 75.7 years (balance); Male/female ratio not stated; Hoehn and Yahr 1.8 (combined), 1.9 (balance); Duration of condition 5.5 years (combined), 8.3 years (balance). Inclusion criteria: Diagnosed with IPD by neurologist, not participated in any organised balance or muscle strengthening activities before being pre-tested, ambulatory, not acutely ill, able to follow simple commands, not suffering from unstable cardiovascular disease. Exclusion criteria: Uncontrolled chronic conditions that would interfere with safety and conduct of the training and testing.
Interventions	Combined: Group training in strengthening and balance exercises. Resistance exercises used Nautilus leg extension and side-lying leg-flexion machines and therabands. Balance training consisting of gen-



Hirsc	h 1996 i	(Continued)
-------	----------	-------------

tle sternal or dorsal perturbation and leaning movements designed to enhance limit of stability whilst

standing on a firm or a compliant surface.

Balance: 'standard' group balance therapy as described above.

Medications kept stable throughout trial period.

Outcomes Balance

Muscle strength (subset): knee extensors, knee flexors, ankle plantar flexors

Latency to fall

% of trials resulting on falls

Notes Randomisation violation; 1 patient who was allocated to the combined therapy group was reassigned

to the balance group after 2 weeks of training due to an inguinal hernia making it impossible for him to

carry out the strength training

Risk of bias

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Low risk	Inclusion and exclusion criteria stated
Randomisation Method	Low risk	Computer generated random number list
Concealment of Allocation	Unclear risk	Method not stated
Similarity at Baseline	Low risk	Group characteristics similar at baseline
Intention to Treat Analysis	Unclear risk	Per protocol analysis
Withdrawals Described	Low risk	13% withdrawal rate for muscle strength outcomes only
Cointerventions Constant	Low risk	Medications kept stable throughout trial period
Comparable arms	Low risk	Same time and attention given to participants in both arms
Blinded Assessors	High risk	Assessors not blinded

Joudoux 2011

Methods	Parallel group design	
	Randomisation method not stated	
	Analysis method not stated	
	Treated for 3 weekly 1 hour sessions for 8 weeks	
	Assessed at baseline, after treatment and after 3 months follow-up	
	Not stated whether assessors were blinded	
Participants	50 participants recruited, group split not stated. Dropouts not stated.	
	Patients baseline characteristics not stated. Inclusion criteria: Mild to moderate PD (Hoehn and Yahr II-III). Exclusion criteria not stated.	



Joudoux 2011 (Continued)

Interventions

Assymmetric motor training Program: Designed to enhance only the agonist activity of the 'body openers' i.e. extension/supination/abduction/external rotation - which is more reduced than their antagonist activity of flexion/protonation/adduction/internal rotation in PD - aiming at re-balancing forces around joints.

Broad program: Standard techniques of passive and active joint mobilisations, balance and gait training, relaxation techniques and respiratory techniques and respiratory work.

Not stated whether drug regimen remained constant throughout trial period.

Outcomes

UPDRS III

GMT score

Rapid alternating movements

Hand writing and spiralography

PDQ-39

Dpression (GDS-15)

Video recording of 8 activities of daily living and biomechanical evaluations

Notes

Abstract only

Risk of bias

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Unclear risk	Inclusion and exclusion criteria stated
Randomisation Method	Unclear risk	Method not stated
Concealment of Allocation	Unclear risk	Randomisation method not stated
Similarity at Baseline	Unclear risk	Group characteristics not stated
Intention to Treat Analysis	Unclear risk	Method of analysis not stated
Withdrawals Described	Unclear risk	No dropouts described
Cointerventions Constant	Unclear risk	Not stated whether drug regimen remained constant throughout trial period
Comparable arms	Low risk	Same time and attention given to participants in both arms
Blinded Assessors	Unclear risk	Not stated whether assessors were blinded

Juncos 2006

Methods Parallel group design

Method of randomisation not stated

Analysed on an intention-to-treat basis

Treated for 6 months

Assessed at baseline then monthly and after last training session



Juncos 2006 (Continued)	Assessors were blinded	i
Participants	56 participants were randomised into Aerobic exercise (AE), Tai Chi Chung (TCC) or Qi Gong (QG) groups. 16 dropouts in total. Group splits not stated.	
	Baseline characteristic	s for groups not stated. Total mean age 65 years.
	Inclusion criteria: Amb criteria not stated.	ulatory subjects with IPD, stable medication regimen, MMSE > 24/30. Exclusion
Interventions	AE: Moderate intensity	walk-run program with moderate caloric expenditure.
	TCC: Intermediate inte	nsity to that of AE, low caloric expenditure.
	QG: Meditation in stilln	ess, minimal caloric expenditure.
	Drugs stable during the	erapy.
Outcomes	UPDRS total	
	UPDRS motor	
	UPDRS ADL	
	PDQ-39	
	Clinical global impress	ion
	Walking speed	
	Falls	
Notes	Abstract and trial registration only	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Low risk	Inclusion criteria stated
Randomisation Method	Unclear risk	Method not stated
Concealment of Allocation	Unclear risk	Method of randomisation not stated
Similarity at Baseline	Unclear risk	Hoehn and Yahr not stated
Intention to Treat Analysis	Unclear risk	Intention-to-treat analysis used
Withdrawals Described	High risk	Withdrawals to 29%
Cointerventions Constant	Low risk	Drugs stable during therapy
Comparable arms	Low risk	Same time and attention given to both groups
Blinded Assessors	Low risk	Assessors were blinded

Khallaf 2011

Methods	Parallel group design
METHORS	rafallel group design



Khallaf 2011 (Continued)			
	Randomisation method not stated		
	Analysis method not stated		
	Treatment dose not stated		
	Assessment intervals not stated		
	Not stated whether assessors were blinded		
Participants	15 participants in Physiotherapy and treadmill group (PT + T) and 15 in Physiotherapy only group (PT). Dropouts not described.		
	Participant baseline characteristics not stated.		
	Inclusion/exclusion criteria not stated.		
Interventions	PT + T: Designed physiotherapy program in addition to 20 min of treadmill training.		
	PT: Designed physiotherapy program only.		
	Not stated whether drug therapies were kept constant throughout trial period.		
Outcomes	UPDRS II and III		
	Hamilton rating scale of depression		
	Walking speed		
	Walking distance		
Notes	Abstract only, no contact details found for author		

Risk of bias

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Unclear risk	Inclusion criteria not stated
Randomisation Method	Unclear risk	Method not stated
Concealment of Allocation	Unclear risk	Method of Randomisation not stated
Similarity at Baseline	Unclear risk	Participant baseline characteristics not stated
Intention to Treat Analysis	Unclear risk	Method of analysis not stated
Withdrawals Described	Unclear risk	Dropouts not described
Cointerventions Constant	Unclear risk	Not stated whether drug therapies were kept constant throughout trial period
Comparable arms	Unclear risk	No details of intervention schedules
Blinded Assessors	Unclear risk	Not stated whether assessors were blinded

Li 2012

Methods	Parallel group design



Li 2012 (Continued)

Randomised by permuted blocks, allocation delivered in sealed envelopes

Analysed on an intention-to-treat basis

Treatment delivered in 2 sessions per week each lasting an hour over 24 weeks

Assessed at baseline and post intervention (at 6 months) and at 3 months follow-up

Assessors were blinded

Participants

65 participants in Tai Chi group, 65 in resistance group, 65 in the stretching group. 9 dropouts in Tai Chi group, 6 in resistance group, 4 in stretching group.

Participants mean age 68 years (Tai Chi), 69 years (resistance), 69 years (stretching); Male/female ratio 45/20 (Tai Chi), 38/27 (resistance), 39/26 (stretching); Mean Hoehn and Yahr not given; Duration of condition 8 years (Tai Chi), 8 years (resistance), 6 years (stretching).

Inclusion criteria: Clinical diagnosis of PD, aged 40 - 85 years, Hoehn and Yahr stage I - IV, at least one score of at least 2 or more for at least one limb for the tremor, rigidity, postural stability or bradykinesia items in the motor section of the UPDRS, stable medication use, ability to stand unaided and walk with or without an assistive device, medical clearance for participation and willingness to be assigned to any of the three interventions. Exclusion criteria: Current participation in any other behavioural or pharmacologic study or instructor led exercise program, a MMSE score lower than 24, debilitating conditions or vision impairment that would impede full participation in the study, unavailability during the study period.

Interventions

Tai Chi: Six Tai Chi movements integrated into an eight-form routine. The protocol was specifically designed to tax balance and gait by having participants perform symmetric and diagonal movements, such as weight shifting, controlled displacement of the centre of mass over the base of support, ankle sways an anterior –posterior and lateral stepping. The first 10 weeks emphasized the mastery of single forms through multiple repetitions; later weeks focused on repetitions to enhance balance and increase locomotion. Natural breathing was integrated into the training routine.

Resistance: Focused on strengthening the muscles that are important for posture, balance and gait. Resistance (with weighted vests and ankle weights) was introduced at week 10. Weight vest resistance was initially set at 1% of body weight and was increased by approx. 1 to 2% of body weight, depending on each participants tolerance, every fifth week until 5% of body weight was achieved. Ankle weights started at 0.45 Kg per limb and were gradually increased to 1.36 Kg. the routine involved 8 to 10 exercises, including forward and side steps, squats, forward and side lunges, and heel and toe raises, performed in 1 to 3 sets of 10 to 15 repetitions. Progression was modified for participants with physical limitations. Natural breathing was emphasized during the training routine.

Stretching: This control condition was designed to provide a low intensity exercise program with the social interaction and enjoyment inherent in the two other interventions but without similar training benefits in lower-extremity weight bearing, strength or balance. The core activities encompassed a variety of seated and standing stretches involving the upper body (neck, upper back, shoulders, chest and arms) and lower extremities (quadriceps, hamstrings, calves and hips) with the use of gentle joint extension and flexion and trunk rotation. Abdominal breathing with an emphasis on inhaling and exhaling to maximum capacity and relaxation of major muscle were also included.

There were no significant changes reported in outside physical activity or use of anti-parkinsonian medication.

Outcomes

Two indicators of postural stability: maximum excursion and directional control

Stride length

Walking velocity

Strength of bilateral knee extensors and flexors

Functional reach test



Li 2012 (Continued)	Timed up and go UPDRS III Number of falls
Notes	122/195 participants continued with their exercise out to 3 month follow-up so this is not a clean follow up comparison Stretching arm classed as attention control and not included in this review

Risk of bias

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Low risk	Inclusion and exclusion criteria stated
Randomisation Method	Unclear risk	Permuted blocks
Concealment of Allocation	High risk	Sealed envelopes used
Similarity at Baseline	Unclear risk	Mean Hoehn and Yahr not given
Intention to Treat Analysis	Low risk	Intention-to-treat analysis used
Withdrawals Described	Low risk	Withdrawals <10%
Cointerventions Constant	Unclear risk	No significant change in medications
Comparable arms	Low risk	Same time and attention given to each group
Blinded Assessors	Low risk	Assessors blinded

Loureiro 2010

Randomisation method not stated Analysis method not stated Treatment delivered in 2 sessions per week over 6 weeks Assessment intervals not stated Not stated whether assessors were blinded Participants 6 participants in conventional physiotherapy (Physio) group, 6 in complimentary activities (CA) group. Mean age 57 (Physio group), 65 (CA group). Inclusion criteria: Hoehn and Yahr II-III. Exclusion criteria not stated. Interventions Physio: Conventional physiotherapy. CA: Wii fit in addition to conventional physiotherapy. Not stated whether the drug therapy was constant during trial.	Methods	Parallel group design	
Treatment delivered in 2 sessions per week over 6 weeks Assessment intervals not stated Not stated whether assessors were blinded Participants 6 participants in conventional physiotherapy (Physio) group, 6 in complimentary activities (CA) group. Mean age 57 (Physio group), 65 (CA group). Inclusion criteria: Hoehn and Yahr II-III. Exclusion criteria not stated. Interventions Physio: Conventional physiotherapy. CA: Wii fit in addition to conventional physiotherapy.		Randomisation method not stated	
Assessment intervals not stated Not stated whether assessors were blinded Participants 6 participants in conventional physiotherapy (Physio) group, 6 in complimentary activities (CA) group. Mean age 57 (Physio group), 65 (CA group). Inclusion criteria: Hoehn and Yahr II-III. Exclusion criteria not stated. Interventions Physio: Conventional physiotherapy. CA: Wii fit in addition to conventional physiotherapy.		Analysis method not stated	
Participants 6 participants in conventional physiotherapy (Physio) group, 6 in complimentary activities (CA) group. Mean age 57 (Physio group), 65 (CA group). Inclusion criteria: Hoehn and Yahr II-III. Exclusion criteria not stated. Physio: Conventional physiotherapy. CA: Wii fit in addition to conventional physiotherapy.		Treatment delivered in 2 sessions per week over 6 weeks	
Participants 6 participants in conventional physiotherapy (Physio) group, 6 in complimentary activities (CA) group. Mean age 57 (Physio group), 65 (CA group). Inclusion criteria: Hoehn and Yahr II-III. Exclusion criteria not stated. Physio: Conventional physiotherapy. CA: Wii fit in addition to conventional physiotherapy.		Assessment intervals not stated	
Mean age 57 (Physio group), 65 (CA group). Inclusion criteria: Hoehn and Yahr II-III. Exclusion criteria not stated. Physio: Conventional physiotherapy. CA: Wii fit in addition to conventional physiotherapy.		Not stated whether assessors were blinded	
Inclusion criteria: Hoehn and Yahr II-III. Exclusion criteria not stated. Interventions Physio: Conventional physiotherapy. CA: Wii fit in addition to conventional physiotherapy.	Participants	6 participants in conventional physiotherapy (Physio) group, 6 in complimentary activities (CA) group.	
Interventions Physio: Conventional physiotherapy. CA: Wii fit in addition to conventional physiotherapy.		Mean age 57 (Physio group), 65 (CA group).	
CA: Wii fit in addition to conventional physiotherapy.		Inclusion criteria: Hoehn and Yahr II-III. Exclusion criteria not stated.	
	Interventions	Physio: Conventional physiotherapy.	
Not stated whether the drug therapy was constant during trial.		CA: Wii fit in addition to conventional physiotherapy.	
		Not stated whether the drug therapy was constant during trial.	



Loureiro	2010	(Continued)
----------	------	-------------

Outcomes Timed up and go

Anterior functional reach

Notes Abstract only

Risk of bias

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Low risk	Inclusion criteria stated
Randomisation Method	Unclear risk	Method not stated
Concealment of Allocation	Unclear risk	Method of randomisation not stated
Similarity at Baseline	Unclear risk	Hoehn and Yahr not stated
Intention to Treat Analysis	Unclear risk	Method of analysis not stated
Withdrawals Described	Unclear risk	Dropouts not described
Cointerventions Constant	Unclear risk	Not stated whether drug therapies were kept constant throughout trial period
Comparable arms	Low risk	Same time and attention given to participants in both arms
Blinded Assessors	Unclear risk	Not stated whether assessors were blinded

Mak 2008

Methods	Parallel-group design
	Participants randomly allocated to groups by drawing lots
	Method of analysis not described
	Treated as outpatients for 4 hours (audio-visual), 6 hours (exercise) over 4 weeks
	Assessed at baseline, at 2 weeks, immediately after and 2 weeks after treatment had ended
	Assessor was blinded
Participants	21 participants in the cueing group, 21 participants in the exercise group and 18 in the control group. 2 dropouts from the cueing group, 2 from the exercise group and 4 from the control group.
	No baseline characteristics given for drop-outs. Participants' mean age 63 (cueing), 66 (exercise), 63 (control). No data given for the sex of participants. Hoehn and Yahr stage 2.8 (cueing), 2.7 (exercise) and 2.7 (control). Duration of PD 5.9 years (cueing), 6.1 years (exercise), 5.9 years (control).
	Inclusion criteria: diagnosed with PD according to Quinn, stable on anti-PD medications without dyskinesia, orthopaedic, arthritic or heart problems, aged between 50-75 years old, perform sit to stand independently, can follow instructions. No exclusion criteria stated.
Interventions	Cueing: Audio-visual cued task-specific training for 20 min three times per week. Received cued sit-to-stand training using Equitest-Balance Master. Visual cue was given on a computer screen with verbal command as auditory cue. Each task lasted 2 min, repeated once with a 30 second rests in between.



Mak 2008 (Continued)			
, ,		ventional exercise twice a week. Conventional mobility and strengthening exertensors of trunk, hips, knees and ankles followed by sit-to-stand practice.	
	Control: No treatment.		
	Drugs stable during the	erapy.	
Outcomes	Peak horizontal velocit	у	
	Peak vertical velocity		
	Movement time		
	3D Kinematics data of	sit-to-stand	
	Not stated when during the day tests took place		
Notes	Control group not inclu	uded in this review	
Risk of bias			
Bias	Authors' judgement	Support for judgement	
Eligibility Criteria	Low risk	Inclusion criteria stated	
Randomisation Method	High risk	Participants randomised by drawing lots	
Concealment of Allocation	High risk	Participants randomised by drawing lots	
Similarity at Baseline	Low risk	Group characteristics similar at baseline	
Intention to Treat Analysis	Unclear risk	Method of analysis not stated	
Withdrawals Described	High risk	Withdrawals at 13%	
Cointerventions Constant	Low risk	Medications kept stable throughout trial period	
Comparable arms	Low risk	Same time and attention given to participants in both arms	

McGinley 2012

Blinded Assessors

Methods	Parallel group design
	Method of randomisation not stated
	Analysed on an intention-to-treat basis
	Treated as outpatients for a 2 hour session once per week for 8 weeks plus home practice
	Assessed at baseline, one week after intervention, at 3 months and 12 months after intervention
	Assessors blinded.
Participants	69 participants in Movement strategy training group (MST), 70 in Progressive strength training group (PST), 71 in Life-skills control group (LS). 14 dropouts prior to post-intervention time point.

Assessors blinded

Low risk



McGinley 2012 (Continued)

Baseline characteristics not split by group. Participants mean age 67.9 years; Male/female ratio 140/70; Hoehn and Yahr mean score not stated; Duration of condition 6.7 years.

Inclusion criteria: Confirmed Idiopathic Parkinson's disease, Hoehn & Yahr stages 0-4, able to participate in an outpatient exercise program including strength training, willing to complete a falls calendar for 12 months after therapy, able to walk, willing and able to attend the therapy and assessment program. Exclusion criteria: Score less than 24 on mini-mental state examination, rating greater than 4 on the modified Hoehn & Yahr scale, on major tranquillisers, other medical conditions that could limit or prevent exercising safely at the required intensity, other prior neurological conditions affecting and dementia.

Interventions

MST: Movement strategies emphasise task specific practice of everyday functional actions such as rolling over, standing up, walking, crossing obstacles and turning. These tasks are practised with strategies such as visual or auditory cues, mental rehearsal and movement planning, conscious attention during the task and breaking the task into a sequence of smaller components. The program is tailored to movement impairments, activity limitations, and cognitive status and learning ability.

An individualised home practice session of strategies to practice within the home or community will be completed once a week. Structured falls risk education will also be provided each week. A single home visit will be conducted by a trained therapist or nurse to check compliance with the therapy program.

PST: Strengthening exercises for quadriceps, hip and trunk extensor muscles, hip abductors, calf, and ankle dorsiflexors, tailored to individual's strength and functional ability. Where possible training is performed in functional tasks such as standing up from a chair, stepping up onto a step etc, using body weight, weighted vests and Thera-band® to progress the resistance. Structured falls risk education will be provided each week. An individualised home practice session of strengthening exercises will also be completed once a week. A single home visit will be conducted by a trained therapist or nurse to check compliance with the therapy program.

LS: Equivalent duration to the MST and PST groups. Each session led by OTs, PTs, Speech pathologists or social workers and included content such as relaxation, games, or communication activities. Guided discussion. Guided discussion will also include topics such as the impact of PD on the individual and family, support and resources available and fatigue management. None of the content will relate to walking, balance or falls risk education. A home session of reflection activities and relaxation practice will also be completed once per week.

Not stated whether the drug therapy was constant during treatment.

Outcomes

Falls: no. of fallers per group, no. of multiple fallers per group, falls rate over 12 months in each group

Number of injurious falls

Walking speed

6 minute walk test

Timed up and go

UPDRS II and III

PDQ-39

EuroQol-5D

Notes

LS arm not included in this review

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Low risk	Inclusion and exclusion criteria stated



McGinley 2012 (Continued)		
Randomisation Method	Unclear risk	Method not stated
Concealment of Allocation	Unclear risk	Method of randomisation not stated
Similarity at Baseline	Unclear risk	Baseline characteristics not split into groups
Intention to Treat Analysis	Low risk	Analysed on an intention-to-treat basis
Withdrawals Described	Low risk	Withdrawals at 7% for post-intervention time point
Cointerventions Constant	Unclear risk	Not stated whether the drug therapy was constant during treatment
Comparable arms	Low risk	Same time and attention given to participants in each physio group
Blinded Assessors	Low risk	Assessors were blinded

Miyai 2000

-			
Methods	Cross-over design		
	Randomisation using the envelope method		
	Analysed on an intention-to-treat basis		
	Inpatients treated for 12 sessions, each lasting 45 minutes, over 4 weeks		
	Assessed at baseline, after first set of treatment at 4 weeks and after second set of treatment at 8 weeks		
	Assessors were not blinded		
Participants	5 participants in body-weight supported treadmill training (BWSTT) and 5 in conventional physiotherapy (PT). No dropouts described.		
	Participants mean age 66 years (BWSTT), 69.3 years (PT); Male/female ratio 2/3 (BWSTT), 3/2 (PT); Hoehn and Yahr 2.8 (BWSTT), 2.9 (PT); Duration of condition 4.9 years (BWSTT), 3.6 years (PT).		
	Inclusion criteria: Hoehn and Yahr stage 2.5 or 3 Parkinson's disease, no dementia MMSE >27. Exclusion criteria not stated.		
Interventions	BWSTT: Walking with 20% bodyweight support (BWS) for 12 minutes, then 4.5 minutes rest, followed by 10% BWS for 12 minutes, another 4.5 minutes rest and finally 12 minutes walking with 0% BWS. Walking speed started at 0.5 km/hr and was ramped up in 0.5 km/hr increments to 3.0 km/hr as tolerated.		
	PT: General conditioning, range of motion exercise, activities of daily living training and gait training.		
	Drug therapy was stable during trial period.		
Outcomes	UPDRS		
	UPDRS subscales (mental, ADL, motor and complications)		
	Overground ambulation endurance		
	Gait speed		
	No. steps taken for 10 metre walk		
Notes	Pre-crossover data only used		



Miyai 2000 (Continued)

Risk of bias

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Low risk	Inclusion criteria stated
Randomisation Method	Unclear risk	Method not stated
Concealment of Allocation	High risk	By the envelope method
Similarity at Baseline	Low risk	Group characteristics similar at baseline
Intention to Treat Analysis	Low risk	Analysed on an intention-to treat basis
Withdrawals Described	Low risk	No withdrawals
Cointerventions Constant	Low risk	Drug therapy was stable during therapy period
Comparable arms	Low risk	Same time and attention given to participants in both arms
Blinded Assessors	High risk	Assessors were not blinded

Miyai 2002

Methods	Parallel group design
	Randomisation was the envelope method
	Analysed on an intention-to-treat basis
	Treatment delivered for 12 sessions over 1 month, sessions lasted 45 minutes
	Assessed at baseline, at 1 month, 2 months, 3 months, 4 months, 5 months and 6 months
	Assessors were not blinded
Participants	11 participants in body-weight supported treadmill training (BWSTT) group and 9 in conventional physiotherapy (PT) group. 4 participants not analysed due to changes in their medication.
	Participants mean age 69.5 years (BWSTT), 69.8 years (PT); Male/female ratio 5/6 (BWSTT), 5/4 (PT); Hoehn and Yahr 2.9 (BWSTT), 2.8 (PT); Duration of condition 4.1 years (BWSTT), 4.5 years (PT). Inclusion criteria: Diagnosis of Parkinson's disease based on the presence of rest tremor, bradykinesia, rigidity, positive response to l-dopa and no evidence of vascular lesions on MRI. Hoehn and Yahr stage 2.5 or 3. Exclusion criteria: Dementia, MMSE <27.
Interventions	BWSTT: Walking with 20% bodyweight support (BWS) for 12 minutes, then 4.5 minutes rest, followed by 10% BWS for 12 minutes, another 4.5 minutes rest and finally 12 minutes walking with 0% BWS. Walking speed started at 0.5 km/hr and was ramped up in 0.5 km/hr increments to 3.0 km/hr as tolerated.
	PT: General conditioning, range of motion exercise, activities of daily living training and gait training.
	Changes in medication not allowed. Participants who changed their medication were excluded.
Outcomes	UPDRS
	UPDRS subscales (mental, ADL, motor and complications)
	Gait speed



Miyai 2002 (Continued)

No. steps taken for 10 metre walk

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Low risk	Inclusion and exclusion criteria stated
Randomisation Method	Unclear risk	Method not stated
Concealment of Allocation	High risk	Envelope method used
Similarity at Baseline	Low risk	Group characteristics similar at baseline
Intention to Treat Analysis	Unclear risk	Analysed on an intention-to-treat basis
Withdrawals Described	High risk	Withdrawals at 17%
Cointerventions Constant	Low risk	Changes in medication not allowed. Participants who changed their medication were excluded
Comparable arms	Low risk	Same time and attention given to participants in both arms
Blinded Assessors	High risk	Assessors were not blinded

Morris 2009

Methods	Parallel group design
	Randomised using generated random number sequences by an independent university source
	Method of analysis not stated
	Treated as inpatients for 2 weeks in 16 sessions each lasting 45 minutes
	Assessed at baseline, immediately after treatment and at 3 months
	Assessors were blinded
Participants	14 participants in movement strategies (MS) group and 14 in exercise (Ex) group. 2 participants missing from follow-up in exercise group.
	Participants mean age 68 years (MS), 66 years (Ex); Male/female ratio, disease severity and duration for groups were not stated.
	Inclusion criteria: Aged 21 -80 years, medically stable, diagnosis of IPD confirmed by a neurologist, >23 MMSE with a minimum of 2 out of 3 on the recall question, Hoehn & Yahr II or III, able to walk 10 metres 3 times without assistance. Exclusion criteria: Unsafe to participate in the therapy programs
Interventions	MS: Learn how to use cognitive strategies such as focusing their attention on movement and responding to external cues to enhance walking, turning, standing up from a chair and obstacle negotiation. Based on principles of Victorian comprehensive Parkinson's disease program.
	Ex: Lower limb and trunk strengthening exercises, spinal and lower limb flexibility exercises and receiving feedback on optimal postural alignment for a range positions.



Morris 2009	(Continued)
-------------	-------------

Not stated whether the drug therapy was constant during treatment.

Outcomes

UPDRS Motor

UPDRS ADL

10 m walk test

Timed up and go

2 min walk test

Balance-shoulder tug

PDQ-39

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Low risk	Inclusion and exclusion criteria stated
Randomisation Method	Low risk	Randomised using generated random number sequences by an independent university source
Concealment of Allocation	Low risk	Randomised using generated random number sequences by an independent university source
Similarity at Baseline	Unclear risk	Only average age of groups given
Intention to Treat Analysis	Unclear risk	Method of analysis not stated
Withdrawals Described	Low risk	Withdrawals at 7%
Cointerventions Constant	Unclear risk	Not stated whether the drug therapy was constant during treatment
Comparable arms	Low risk	Same time and attention given to participants in both arms
Blinded Assessors	Low risk	Assessors were blinded

Palmer 1986

Methods	Parallel group design		
Randomised in pairs matched for stage of disease, sex and age, method not stated			
	Method of analysis not stated		
	Treated as outpatients for 12 weeks in 1 hour sessions 3 times per week		
	Assessed at baseline and immediately after therapy		
	Assessors were blinded		
Participants	7 participants in United Parkinson Foundation (UPF) exercise program, 7 in upper body karate (UBK) training program. No dropouts were described.		



Palmer :	L986	(Continued)
----------	------	-------------

Participants mean age 63.9 years (UPF), 65.9 years (UBK); Male/female ratio not stated; Hoehn and Yahr 2.4 (UPF), 2.4 (UBK); Duration of condition not stated.

Inclusion criteria: IPD, stabilization on a regimen of pharmacologic therapy, ability to attend the scheduled evaluation and exercise sessions. Exclusion criteria: Physical problems that might cause them to risk injury during exercises.

Interventions

UPF: Stretch exercises from the UPF exercise program led by a corrective therapist.

UBK: Trained in upper body Karate techniques by a rehabilitation nursing student who had a black belt in karate. All karate was done in a seated position. Each session consisted of approximately 15 mins warm-up stretching exercises, 35 mins karate training and 10 mins cool down stretching exercises.

Drug therapy was stable during therapy period.

Outcomes

Forearm pronation/supination rate

Pursuit score walk index

Degree of activated rigidity

Degree of arm tremor

Activated rigidity

Grip strength

9-hole peg test

Minnesota placing and turning test

Arm swings test

Rapid alternating arm movement test

Button board

Putting shirt on and off

Putting shoes and socks on and off

Getting up from chair

Long latency stretch response

Notes

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Low risk	Inclusion and exclusion criteria stated
Randomisation Method	Unclear risk	Method not stated
Concealment of Allocation	Unclear risk	Randomisation method not stated
Similarity at Baseline	Low risk	Group characteristics similar at baseline
Intention to Treat Analysis	Unclear risk	Method of analysis not stated
Withdrawals Described	Low risk	No withdrawals



Palmer 1986	(Continued)
-------------	-------------

Cointerventions Constant	Low risk	Drug therapy was stable during therapy period
Comparable arms	Low risk	Same time and attention given to participants in both arms
Blinded Assessors	Low risk	Assessors were blinded

Pelosin 2010	
Methods	Parallel group design
	Randomised by computerised random-number generator performed by an independent researcher
	Method of analysis not stated
	Treated as outpatients for 4 weeks with hour long sessions delivered 3 times per week
	Assessed at baseline, 2 days after completion of all therapy sessions and 4 weeks later
	Assessors were blinded
Participants	9 participants in action group, 9 in landscape group. 2 additional patients were recruited but excluded due to past history of neurological conditions other than PD. No dropouts described.
	Participants mean age 68.8 years (action), 70.2 years (landscape); Male/female ratio not stated; Hoehn & Yahr 2.1 (action), 2.2 (landscape); Duration of condition 11.6 years (action), 9.5 years (landscape).
	Inclusion criteria: IPD according to the UK PDS BB criteria. All on stable medication regime. Mobile despite occurrence of freezing at least once a week (minimum score of 2 on item 3 of the FOG questionnaire) and for at least 2s (minimum score of 1 on item 4 of FOG-Q). MMSE > 24. Exclusion criteria not stated.
Interventions	Action: Instructed to carefully watch 6 video clips (each clip lasting 6 mins) showing strategies useful in circumventing FOG episodes. During each training session 2 video clips (with different sequences of actions) were presented twice. The complexity of movements progressively increased from simple actions to more complex movements. All actions shown in the video clips were performed by a physical therapist. To ensure proper attention during the video presentation, patients were explicitly asked to concentrate on how the actions were performed and were not allowed to imitate any movement. After video clip observation, patients were asked to practice (for the remaining time of the session-36 min) the observed actions repetitively and accurately according to the instructions of the physical therapist.
	Landscape: Matched the experimental protocol, with the exception that during each training sessions they watched 2 video clips (presented twice) containing sequences of static pictures of mountains and seaside, countryside, and desert scenes without any living (human or animal) representations. During training sessions, patients in the landscape group performed the same movements/ actions used

for the Action group following the physical therapist's instructions, in the exact order and for the same

Outcomes

FOG Questionnaire and FOG diary

Drug therapy was stable during trial period.

Timed up and go

amount of time.

10 metre walking test

Tinetti scale part I and II

Berg balance scale



Pelos	in 2010	(Continued)
--------------	---------	-------------

PDQ-39

Notes Only data for FOG could be extracted from paper

Risk of bias

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Low risk	Inclusion criteria stated
Randomisation Method	Low risk	Randomised by computerised random-number generator
Concealment of Allocation	Low risk	Randomisation performed by an independent researcher
Similarity at Baseline	Low risk	Group characteristics similar at baseline
Intention to Treat Analysis	Unclear risk	Method of analysis not stated. ITT was not used
Withdrawals Described	Low risk	Withdrawals at 10%
Cointerventions Constant	Low risk	Drug therapy was stable during trial period
Comparable arms	Low risk	Same time and attention given to participants in both arms
Blinded Assessors	Low risk	Assessors were blinded

Picelli 2012

Methods	Parallel group design
	Participants randomised using list, generated by www.randomisation.com kept in locked desk draw
	Analysed on a per protocol basis
	Treated as outpatients for 45 minute sessions, 3 times per week for 4 weeks
	Assessed at baseline, post intervention, and 1 month after treatment
	Assessors were blinded
Participants	21 participants in Robot assisted gait-training group (RAGT) and 20 participants in physiotherapy group (PT). 3 dropouts in RAGT group and 2 in PT group.
	Participants' mean age 68.1 years (RAGT), 68.7 years (PT). Male/female ratio, 10/8 (RAGT), 6/12 (PT). Hoehn and Yahr 2.7 (RAGT), 2.7 (PT). Duration of PD 6.6 years (RAGT), 7.4 years (PT). Baseline data not given for drop-outs.
	Inclusion criteria: Confirmed PD, Hoehn and Yahr score 2.5 or 3, MMSE score > 23. Exclusion criteria: Severe dyskinesias of "on-off" phases, change of PD medications during the study, deficits of somatic sensation in the lower limbs, vestibular disorders or paroxysmal vertigo, other neurological orthopaedic conditions involving the lower limbs and cardiovascular comorbidity.
Interventions	RAGT: 2 motor driven footplates positioned on 2 bars that provide a Robot assisted propulsion by means of a planetary gear system, simulating stance and swing with a ratio of 60% to 40% between the two phases. A progressive reduction of body weight combined with an increase in gait speed. Each training session consisted of 3 parts (each one lasting 10 minutes), with a 5-minute rest after each of

them. Patients were first trained at 20% of body weight supported and at a speed of 1 km/h for 10 minutes; then at 10% of body weight supported and a speed of 1.3 km/hr for 10 mins; and finally, at 0% of



Picelli 2012 (Continued)

body weight supported and a speed of 1.6 km/hr for 10 mins. Patients were instructed to "help" the gait trainer (GT1) gait-like movement during training. Any patient unable to maintain the chosen pace was excluded from the study.

PT: Program included active joint mobilization and conventional gait training. Each treatment session consisted of 2 parts with a 5-minute rest between them. First patients performed active joint mobilization of the lower limbs (hip, knee and ankle) in the supine and prone positions (10 repetitions of 6 exercises) for 10 minutes (5 minutes per position). Then they performed conventional gait therapy based on the proprioceptive neuromuscular facilitation (PNF) concept for 30 minutes. Among the PNF techniques we facilitated pelvic motion to improve control of the pelvis as a "key point of control" for maintaining a gait pattern. Conventional gait therapy consisted of 10 minutes each of rhythmic initiation, slow reversal and agonistic reversal exercises applied to the pelvic region. The time spent in PNF was equal to that spent in RAGT. The same trained therapist treated all the patients in this group and standardized the duration of each part of the treatment.

No changes in medications as this was a criterion for exclusion.

Outcomes

10 m walk test

6 min walk test

Spatiotemporal gait parameters including stride length

Parkinson's fatigue scale

UPDRS Total

Notes

Risk of bias

tated
mization.com
ſ
n groups
s was a criterion for exclusion
b both groups
r

Poliakoff 2009

Methods

Parallel group design

Randomisation method was done by drawing lots and placing allocations in two sets of sealed envelopes for high and low severity of Parkinson's disease



Poliakoff 2009 (Continued)			
,	Method of analysis was	sunclear	
	Treatment delivered in	two 1 hour sessions per week for 10 or 20 weeks	
	Assessed at baseline, a	t 10 weeks and 20 weeks	
	Assessors were not blir	nded	
Participants	16 participants in 20 we group and 6 dropouts f	eek gym group (gym), 16 in 10 week gym group (control). 4 dropouts from gym rom control group.	
		66.6 years (gym), 63.7 years (control); Male/female ratio, 11/5 (gym), 10/6 (contot stated; Duration of condition 7.4 years (gym), 4.7 years (control).	
		to moderate PD. Exclusion criteria: diagnosed with dementia, attendance of for Parkinson's disease, other neurodegenerative disease, >2weeks holiday dy period.	
Interventions	20 Week: 20 week biweekly gym training programme at local leisure complex. 1 hours weekly train in the studio and 1 hour in the gym, each run by gym staff with experience of working with PD. Gyn sions consisted of mainly cardiovascular activity, studio sessions emphasised on gait and agility. T patients used external stimuli (such as music) and team working was encouraged.		
	10 Week: 10 week prog	ramme as above starting 10 weeks after baseline assessment.	
	Not clear whether the o	drug therapy was constant during treatment.	
Outcomes	Simple, choice and serial reaction time		
	Videotaped motor performance		
	PDQ-39		
	UPDRS III		
	Illness perceptions (BIF	PQ)	
	Questionnaire assessin	g experiences of programme	
Notes	Abstract and manuscript accepted for publication in Neurorehabilitation		
	Data used from assessi	ments at baseline and 20 weeks	
Risk of bias			
Bias	Authors' judgement	Support for judgement	
Eligibility Criteria	Unclear risk	Characteristics of included patients described but not stated whether these items were used for the selection of patients for trial	
Randomisation Method	High risk	Drawing lots	
Concealment of Allocation	High risk	Drawing lots	
Similarity at Baseline	Unclear risk	Limited baseline data	
Intention to Treat Analysis	Unclear risk	Attempted to assess all participants regardless of whether they had started or completed intervention. ITT not used	

Withdrawals at 31%

High risk

Withdrawals Described



Poliakoff 2009 (Continued)		
Cointerventions Constant	Unclear risk	Not clear whether the drug therapy was constant during treatment
Comparable arms	Unclear risk	Half the amount of time and attention spent with delayed start group
Blinded Assessors	High risk	Assessors were not blinded

Reuter 2011

Methods	Parallel group design		
	Randomisation was conducted using a computer generated sequence		
	Method of analysis not stated		
	Treatment delivered in 70 minute sessions 3 times per week for 6 months		
	Assessed at baseline and post-intervention		
	Assessors were blinded		
Participants	30 participants in Nordic walking group (NW), 30 in walking group (W) and 30 in flexibility and relaxation group (FR). No dropouts from any group.		
	Participant mean age 62.0 years (NW), 63.0 years (W), 62.1 years (FR); Male/female ratio not stated; Hoehn & Yahr 2.5 (NW), 2.5 (W), 2.4 (FR); Duration of condition 5.3 years (NW), 6.0 years (W), 5.2 years (FR).		
	Inclusion criteria: PD diagnosed using UKBB criteria assessed by movement disorder specialist, Hoehn and Yahr stage II-III. Exclusion criteria: Severe concomitant disease which limit physical performance, second neurological disease.		
Interventions	NW: Consisted of warming up including some flexibility and strength exercises with and without the poles. One session per week was dedicated to practising NW technique, the other sessions focused on endurance training. Participants were encouraged to increase the intensity of the training by walking faster or uphill and to increase the distance walked. Each training session finished with a cooling down programme. Training sessions took place in a park and a forest near to the university hospital.		
	W: Consisted of warming up, technique training, endurance training and cooling down. Instructors emphasised on arm swing and coordination of upper and lower limbs. One session per week included walking uphill to improve muscle strength.		
	FR: Performed flexibility exercises and relaxation training. The training focused on stretching, improving balance and range of movements. The flexibility and relaxation programme did not include aerobic exercises.		
	Medical treatment was optimised prior to study and minor changes were allowed in 5 patients during trial period.		
Outcomes	Max walking speed on treadmill		
	12 m and 24 m walking test		
	Stride length		
	Gait variability		
	UPDRS		
	PDQ-39		



Reuter 2011 (Continued)

Physical activity in everyday life

Adverse effects

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Low risk	Inclusion and exclusion criteria stated
Randomisation Method	Low risk	Computer generated sequence
Concealment of Allocation	Unclear risk	Method not stated
Similarity at Baseline	Low risk	No significant difference in baseline demographic data
Intention to Treat Analysis	Unclear risk	Analysis method not stated
Withdrawals Described	Low risk	No dropouts
Cointerventions Constant	Unclear risk	Minor changes in 5 patients
Comparable arms	Low risk	Same time and attention given to participants in both arms
Blinded Assessors	Low risk	Assessors were blinded

Ridgel 2009

Methods	Parallel group design		
	Randomisation method not stated		
	Method of analysis not stated		
	Treatment delivered in 1 hour sessions, 3 times per week for 8 weeks		
	Assessed at baseline, immediately after completion of treatment, and four weeks after completion of treatment		
	Assessors blinded for UPDRS outcome only		
Participants	5 participants in forced exercise group, 5 in voluntary exercise. No dropouts described.		
	Participants mean age 58 years (forced), 64 years (voluntary); Male/female ratio not stated; Hoehn and Yahr stage not stated; Duration of condition 7.9 years (forced), 4.4 years (voluntary).		
	Inclusion criteria: IPD. Exclusion criteria not stated.		
Interventions	Forced: Tandem stationary bicycle with trainer. Lower extremity forced exercise (FE) intervention using a stationary tandem bicycle. Patient's pedaling rate was increased to approximately 30% more than their preferred rate.		
	Voluntary: Exercise on a stationary single bicycle.		
	Drug therapy was stable during trial period.		



Ridgel 2009 (Continued)

Outcomes UPDRS part III

Manual functional dexterity

Bimanual dexterity

Centre of pressure (CoP)

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Unclear risk	Criteria not stated
Randomisation Method	Unclear risk	Method not stated
Concealment of Allocation	Unclear risk	Method of randomisation not stated
Similarity at Baseline	Unclear risk	Hoehn and Yahr not stated
Intention to Treat Analysis	Unclear risk	Method of analysis not stated
Withdrawals Described	Low risk	Withdrawals at 10%
Cointerventions Constant	Low risk	Drug therapy was stable during trial period
Comparable arms	Low risk	Same time and attention given to participants in both arms
Blinded Assessors	High risk	Assessors blinded for UPDRS outcome only

Robichaud 2012

Methods	Parallel group design		
	Method of randomisation not stated		
	Method of analysis not stated		
	Treatment was delivered in 1 hour sessions twice per week for 24 months		
	Assessed at baseline, 6, 12, 18 and 24 months		
	Assessors were blinded		
Participants	Number randomised not stated. 48 patients assessed at 6 months, down to 38 at 24 months. Group split not stated.		
	Group baseline characteristics not stated. Total mean age 59 years, total mean duration of condition 7 years.		
	Eligibility criteria not stated.		
Interventions	Progressive resistance exercise: Weight lifting program.		
	Fitness counts: Flexibility, balance and strengthening program.		



Dahishaad 2012 (s. a. a. a.				
Robichaud 2012 (Continued)	Not stated whether the drug therapy was constant during treatment.			
Outcomes	UPDRS motor			
	Timed up and go			
	Berg balance scale	Berg balance scale		
	Modified physical perfo	ormance test		
Notes	Abstracts and trial registrations only			
Risk of bias				
Bias	Authors' judgement	Support for judgement		
Eligibility Criteria	Unclear risk	Eligibility criteria not stated		
Randomisation Method	Unclear risk	Method not stated		
Concealment of Allocation	Unclear risk	Method of randomisation not stated		
Similarity at Baseline	Unclear risk	Group baseline characteristics not stated		
Intention to Treat Analysis	Unclear risk	Method of analysis not stated		
Withdrawals Described	High risk	21% lost between 6 and 24 months		
Cointerventions Constant	Unclear risk	Not stated whether the drug therapy was constant during treatment		
Comparable arms	Low risk	Same time and attention given to both groups		
Blinded Assessors	Low risk	Assessors blinded		

Schenkman 2012a

Methods	Parallel group design		
	Randomised using computer generated assignments kept in opaque, sealed envelopes, unsealed by research assistant after baseline assessment		
	Analysed on an intention-to-treat basis		
	Treatment was delivered in 1 hour sessions, 3 times per week for the first 4 months then tapered for 1 month to once monthly sessions out to 16 months. Home exercise group received just once monthly supervised sessions throughout		
	Assessed at baseline, 4 months, 10 months and 16 months		
	Assessors blinded		
Participants	39 participants in Flexibilty/balance/function exercise group (FBF), 41 in Aerobic exercise group (AE) and 41 in Home exercise group (HE).		
	Participant mean age 64.5 years (FBF), 63.4 years (AE), 66.3 years (HE); Male/female ratio 24/15 (FBF), 26/15 (AE), 26/15 (HE); Hoehn and Yahr 2.3 (FBF), 2.2 (AE), 2.3 (HE); Duration of condition 4.9 years (FBF), 3.9 years (AE), 4.5 years (HE).		



Schenkman 2012a (Continued)

Inclusion criteria: Primary PD diagnosed by a movement disorders specialist using the UKBBC, lived in the community, ambulated independently, Hoehn & Yahr stage II-III. Exclusion criteria: Uncontrolled hypertension, on-state freezing or exercise limitations from other disorders, MMSE<24.

Interventions

FBF: Individualised spinal and extremity flexibility exercises followed by group balance/functional training. 2 months of flexibility training one-on-one with a physical therapist followed by 2 months of small group exercise (up to 6 participants) that included flexibility, balance and functional exercise.

AE: Treadmill, bike and/or elliptical trainer. Included 5-10 min of warm-up, 30 minutes exercise at 65% to 80% HR max and 5-10min of cool down. Participants were encouraged to use a treadmill but were permitted to use a stationary bicycle or elliptical trainer.

HE: Exercised at home using the National PD Foundation 'Fitness counts' program. Consisted of exercises in the home setting utilizing Fitness Counts with a single monthly group exercise session supervised by a physical therapist. Flexibility and strengthening exercises in sitting and standing. Daily walking (no specific guidelines).

All participants regardless of group assignment were assisted to develop long term exercise habits. After randomisation and before beginning to exercise, participants met with their trainer to discuss motivation to exercise, potential barriers and strategies to develop exercise habits.

Participants were asked to record supervised and home exercise throughout the 16 months.

Drug therapy was kept constant for the first 4 months.

Outcomes

Overall physical function

Balance - functional reach

Walking economy - VO₂

UPDRS ADL

UPDRS Motor

PDQ-39

Notes

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Low risk	Inclusion and exclusion criteria stated
Randomisation Method	Low risk	Computer generated assignment
Concealment of Allocation	High risk	Sealed envelopes
Similarity at Baseline	Low risk	Group characteristics similar at baseline
Intention to Treat Analysis	Unclear risk	Intention-to-treat analysis used
Withdrawals Described	High risk	Withdrawals at 21%
Cointerventions Constant	Unclear risk	Drug therapy kept constant for first four months only
Comparable arms	Unclear risk	HE group received less time and attention than experimental groups
Blinded Assessors	Low risk	Assessors were blinded



Shankar 2009

Methods	Parallel group design
	Randomisation method not stated
	Method of analysis not stated
	Treatment was delivered in 30 min sessions, twice a week for 8 weeks
	Assessed at baseline and at 2 months
	Assessors were blinded
Participants	10 participants in Treadmill with Cueing (TwC) group, 9 in Treadmill without Cueing (T) group and 10 in Cueing only (C) group. No dropouts described.
	Baseline characteristics not stated for groups.
	Inclusion criteria: Moderate Parkinson's disease. No exclusion criteria.
Interventions	TwC: Walking on the treadmill with music for 30 min twice a week. Music was selected based upon participant input and cadence-matched to the participant's preferred walking speed.
	T: Walking on the treadmill without music for 30 min twice a week.
	C: Listening to music for 30 min twice a week.
	Not stated whether the drug therapy was constant during trial.
Outcomes	Gait and Balance Scale
	UPDRS III
	PDQ-39
	Not stated when examinations took place
Notes	Abstract only

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Unclear risk	Criteria not stated
Randomisation Method	Unclear risk	Method not stated
Concealment of Allocation	Unclear risk	Method of randomisation not stated
Similarity at Baseline	Unclear risk	Baseline characteristics not stated
Intention to Treat Analysis	Unclear risk	Method of analysis not stated
Withdrawals Described	Unclear risk	Withdrawals not described
Cointerventions Constant	Unclear risk	Not stated whether the drug therapy was constant during trial
Comparable arms	Low risk	Same time and attention given to participants in both T and TwC arms



Shankar 2009 (Continued)

Blinded Assessors Low risk Assessors were blinded

Shen 2011

Parallel group design		
Randomisation method not stated		
Method of analysis not stated		
Treatment delivered over 12 weeks		
Assessed at baseline and immediately after training and at 12 weeks follow-up		
Assessors were blinded		
23 participants in balance group, 22 in control group. No dropouts described.		
Participants baseline characteristics not stated.		
Inclusion criteria: PD patients. Exclusion criteria not stated.		
Balance: Task specific to facilitate reaction time and length of compensatory steps during activities with self-induced perturbation and in response to external perturbation.		
Control: Strength training of lower limb muscles with moderate intensity.		
Not stated whether the drug therapy was constant during trial.		
Limit of stability		
Walking speed		
One leg stance time		
Activities-specific balance confidence scale		
UPDRS III		
Abstract only		

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Unclear risk	Criteria not stated
Randomisation Method	Unclear risk	Method not stated
Concealment of Allocation	Unclear risk	Method of randomisation not stated
Similarity at Baseline	Unclear risk	Group characteristics not stated
Intention to Treat Analysis	Unclear risk	Method of analysis not stated
Withdrawals Described	Unclear risk	Withdrawals not described
Cointerventions Constant	Unclear risk	Not stated whether the drug therapy was constant during trial



Shen 2011 (Continued)				
Comparable arms	Low risk	Same time and attention given to participants in both arms		
Blinded Assessors	Low risk	Assessors were blinded		

Shiba 1999

Bias	Authors' judgement Support for judgement			
Risk of bias				
Notes	Abstract only. No numerical data available			
Outcomes	Stride length			
	Auditory stimulation: Patients walked to a rhythm that was 30% of their comfortable walking rhythm. Not stated whether the drug therapy was constant during trial.			
Interventions	Visual training: Patients walked over parallel lines at 90 degrees to the direction of travel. Distance apart of lines dependant on patients normal stride length.			
	Baseline characteristics not stated for groups. Total mean age 65 years; total male/female ratio 3/5. Inclusion criteria: Stable mild to moderate Parkinson's disease. Exclusion criteria not stated.			
Participants	8 participants took part in the study. No dropouts described.			
	Not stated whether the assessors were blinded			
	Assessed at baseline and immediately after treatment			
	Treated as outpatients for an unknown period of time. 1 week between each training regimen.			
	Method of analysis of the data not stated			
Methods	Cross-over group design Method of randomisation not stated			
niba 1999				

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Unclear risk	Criteria not stated
Randomisation Method	Unclear risk	Method not stated
Concealment of Allocation	Unclear risk	Method of randomisation not stated
Similarity at Baseline	Unclear risk	Baseline characteristics not stated
Intention to Treat Analysis	Unclear risk	Method of analysis not stated
Withdrawals Described	Unclear risk	Dropouts not described
Cointerventions Constant	Unclear risk	Not stated whether the drug therapy was constant during trial
Comparable arms	Unclear risk	Treatment schedules not stated
Blinded Assessors	Unclear risk	Not stated whether the assessors were blinded



Si	gu	rge	irsso	n 2	009

Methods	Parallel group design
	Method of randomisation not stated
	Method of analysis not stated
	Treatment delivered in 30 min sessions, 4 sessions per week for 4 weeks
	Assessed at baseline, immediately after treatment and 3 months after treatment
	Not stated whether the assessors were blinded
Participants	26 participants randomised, group split not stated. No dropouts described.
	Baseline characteristics of groups not stated. Total mean Hoehn Yahr 2.1.
	Inclusion and exclusion criteria not stated.
Interventions	Exercise: Walking with visual cues.
	Control: Walking without cues.
	Not stated whether the drug therapy was constant during trial.
Outcomes	Timed up and go
	PDQ-39
Notes	Abstract only

Risk of bias

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Unclear risk	Criteria not stated
Randomisation Method	Unclear risk	Method not described
Concealment of Allocation	Unclear risk	Method of randomisation not described
Similarity at Baseline	Unclear risk	Baseline characteristics of groups not stated
Intention to Treat Analysis	Unclear risk	Method of analysis not stated
Withdrawals Described	Unclear risk	Withdrawals not described
Cointerventions Constant	Unclear risk	Not stated whether the drug therapy was constant during trial
Comparable arms	Unclear risk	Same time and attention given to participants in both arms
Blinded Assessors	Unclear risk	Not stated whether the assessors were blinded

Smania 2010

Methods Parallel group design

Randomisation list used



Sman	ia 20)10 (Continued)
------	-------	--------------	------------

Method of analysis not stated

Treated as outpatients in 50 min sessions, 3 times per week for 7 weeks

Assessed at baseline, immediately post treatment and 1 month post-treatment

Assessors were blinded

Participants

33 participants in balance group, 31 in control group. 5 dropouts in balance group, 4 in control group.

Participants mean age 67.6 years (balance), 67.3 years (control); Male/female 14/14 (balance), 15/12 (control); Hoehn and Yahr not stated; Duration of condition 10.4 years (balance), 8.6 years (control).

Inclusion criteria: Idiopathic PD and postural instability, Hoehn & Yahr 3-4, did not require assistance to rise from chairs or beds. Exclusion criteria: Unstable cardiovascular disease or other chronic conditions that could interfere with their safety during testing or training procedures, other neurological conditions or mental deterioration (MMSE <23), severe dyskinesias or 'on-off' phases.

Interventions

Balance: Each patient was submitted to balance training consisting of exercises aimed at improving both feedforward and feedback postural reactions. Patients were required to repeat exercises from 3 different groups. Group 1: self-destabilisation of the centre-of-body mass (feedforward), group 2: externally induced destabilisation of the centre-of-body mass (feedback), group 3: emphasis of coordination between leg and arm movements during walking as well as locomotor dexterity over an obstacle course and other potentially destabilising activities (continuous feedback and feedforward adjustments). During each session 10 exercises were undertaken, 4 from group 1, 4 from group 2 and 2 from group 3. 5 – 10 repeats of each for 5 mins. Complexity of exercises was increased with patient's ability. Support was provided by therapist at pelvis or chest when required.

Control: Exercises not specifically designed to improve postural reactions. Active joint mobilisation, muscle stretching and motor coordination exercises. During each session 10 exercises were undertaken, 6 in supine position, 2 in the sitting position and 2 in the standing position. 5 – 10 repeats of each for 5 mins. Complexity of exercises was increased with patient's ability.

Drug therapy was stable during trial period.

Outcomes

Berg balance scale

Activities specific balance confidence

Postural transfer test

Self destabilization of the centre of foot pressure test

Falls diary

UPDRS Total

Modified Hoehn and Yahr

Geriatric depression scale

Notes

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Unclear risk	Charateristics of included patients described but not stated whether these items were used for the selection of patients for trial
Randomisation Method	Unclear risk	Randomisation list used

Low risk

Low risk



Smania 2010 (Continued)		
Concealment of Allocation	High risk	Randomisation list kept in locked desk of principal investigator
Similarity at Baseline	Unclear risk	Hoehn and Yahr scores not stated
Intention to Treat Analysis	Unclear risk	Method of analysis not stated
Withdrawals Described	High risk	Withdrawals at 14%
Cointerventions Constant	Low risk	Drug therapy was stable during trial period

Assessors were blinded

Same time and attention given to participants in both arms

Talakkad 2011

Comparable arms

Blinded Assessors

	Method of randomisation not stated Method of analysis not described Treated for 8 hours over 4 weeks		
	Treated for 8 hours over 4 weeks		
	Assessed at baseline and after 4 weeks of intervention		
	Not stated whether assessors were blinded		
Participants	60 participants were randomised into this trial. Dropouts were not described.		
	Baseline characteristics of participants were not stated.		
	Eligibility criteria not stated.		
Interventions	Conventional gait training (CGT).		
	Partial weight supported treadmill training: 20% unweighting.		
	Control: No specific intervention.		
	Drug therapy not described.		
Outcomes	Dynamic posturography		
	UPDRS (total and motor subscore)		
	Beat-to-beat finger blood pressure		
Notes	Abstract only		
	Control arm not included in this review		
Risk of bias			
Bias	Authors' judgement Support for judgement		
Eligibility Criteria	Unclear risk Eligibility criteria not stated		



Talakkad 2011 (Continued)		
Randomisation Method	Unclear risk	Method of randomisation not stated
Concealment of Allocation	Unclear risk	Method of randomisation not stated
Similarity at Baseline	Unclear risk	Baseline characteristics not stated
Intention to Treat Analysis	Unclear risk	Method of analysis not described
Withdrawals Described	Unclear risk	Dropouts not described
Cointerventions Constant	Unclear risk	Drug therapy not described
Comparable arms	Low risk	Same time and attention given to participants in both arms
Blinded Assessors	Unclear risk	Not stated whether assessors were blinded

Thaut 1996

KISK OT DIAS			
Risk of bias			
Notes	3 arms to trial; RAS, SPT and no treatment. SPT versus no treatment are compared in 'Physiotherapy for patients with Parkinson's disease' Cochrane review. New correspondence with author on randomisation method		
Outcomes	Stride velocity Stride cadence Stride length EMG analysis on leg muscles		
	NT: No treatment. Drug therapy was stable during trial period.		
Interventions	RAS: Individual. 30 min/day walking to 3 different tempos of music. For 1st week; normal tempo = pretest cadence, quick = 5-10% faster, fast = an additional 5-10% faster. After each week each tempo was increased by 5-10% to a maximum pace of 130 steps/min. SPT: Individual. 30 min/day walking at normal, quick and fast speeds.		
	Patients mean age 69 years (RAS), 74 years (SPT), 71 (NT); Male/female 10/5 (RAS), 8/3 (SPT), 8/3 (NT); Hoehn and Yahr 2.4 (RAS), 2.5 (SPT), 2.6 (NT); Duration of condition 7.2 years (RAS), 5.4 years (SPT), 8.5 years (NT). Inclusion criteria: IPD with significant gait deficits but able to walk without physical assistance. Exclusion criteria not stated.		
Participants	15 patients in novel rhythmic auditory stimulation group (RAS), 11 in standard self paced training group (SPT) and 11 in no treatment group (NT). No dropouts noted.		
	Not stated whether assessors were blinded		
Methods	Parallel group design Randomised using a computerised random number generator Analysed on an intention-to-treat basis Treated at home or in the community, for 10.5 hours over 3 weeks. Assessed in laboratory, at baseline and immediately after treatment		



Thaut 1996 (Continued)		
Eligibility Criteria	Low risk	Inclusion and exclusion criteria confirmed with author correspondence
Randomisation Method	Low risk	Randomised using a computerised random number generator
Concealment of Allocation	Unclear risk	Method not stated
Similarity at Baseline	Low risk	Group characteristics similar at baseline
Intention to Treat Analysis	Low risk	Analysed on an intention-to-treat basis
Withdrawals Described	Low risk	No withdrawals
Cointerventions Constant	Low risk	Drug therapy was stable during trial period
Comparable arms	Low risk	Same time and attention given to participants in both arms
Blinded Assessors	Unclear risk	Not stated whether assessors were blinded

Toole 2005

Methods	Parallel group design
	Randomised using table of random numbers
	Method of analysis not stated
	Treatment delivered in 20 mins sessions, 3 times per week for 6 weeks
	Assessed at baseline, within 1 week of completing intervention and 4 weeks later
	Not stated whether assessors were blinded
Participants	23 participants randomised, group split not stated. No dropouts described.
	Participants mean age 75.4 years standard treadmill group (ST), 76.4 years unweighted treadmill group (UT), 72.0 years weighted treadmill group (WT); Male/female ratio not stated; Hoehn & Yahr 4.8 (ST), 3.6 (UT), 3.4 (WT); Duration of condition not stated.
	Inclusion criteria: Parkinsonism, Hoehn and Yahr 1-4. Exclusion criteria: Uncompensated cardiovascular disease, uncontrolled high blood pressure, leg claudication, significant dementia, other disorders of comprehension and/or other medical conditions that would interfere with the participants safety and comfort during submaximal exercise.
Interventions	ST: Treadmill training with no loading or unloading.
	UT: Treadmill training assisted by the biodex unweighting system at a 25% body weight reduction for 15 mins then 5 mins with system removed.
	WT: Treadmill training wearing weighted scuba-diving belt which increased normal body weight by 5% for 15 mins then 5 mins without belt.
	Not stated whether the drug therapy was constant during trial.
Outcomes	Balance from dynamic posturography
	Berg balance scale
	UPDRS



Toole 2005 (Continued)

Biomechanical assessment of strength and range of motion

Gait

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement	
Eligibility Criteria	Low risk	Criteria stated	
Randomisation Method	Unclear risk	Table of random numbers, unclear how generated	
Concealment of Allocation	Unclear risk	Randomised using table of random numbers	
Similarity at Baseline	Low risk	Group characteristics similar at baseline	
Intention to Treat Analysis	Unclear risk	Method of analysis not stated	
Withdrawals Described	Unclear risk	Withdrawals not described	
Cointerventions Constant	Unclear risk	Not stated whether the drug therapy was constant during trial	
Comparable arms	Low risk	Same time and attention given to participants in both arms	
Blinded Assessors	Unclear risk	Not stated whether assessors were blinded	

Vivas 2011

Methods	Parallel group design
	Randomisation method not stated
	Method of analysis not stated
	Treated as outpatients with twice weekly 45 minute sessions for 4 weeks
	Assessed at baseline, after 4 weeks treatment and at 17 days follow-up
	Assessors were not blinded
Participants	6 participants in water based exercise group (Water), 6 in land based exercise group (Land). 1 dropout in Water group.
	Participant mean age 65.7 years (Water), 68.3 years (Land); Male/female ratio 3/3 (Water), 4/2 (Land); Hoehn and Yahr 2.7 (Water), 2.4 (Land); Duration of condition 4.2 (Water), 7.8 (Land).
	Inclusion criteria: IPD, ability to follow a stable medication schedule, Hoehn & Yahr stage II-III (Off medication), lack of dementia (MMSE ≥ 24). Exclusion criteria: unable to walk independently, undergone surgical treatment for PD.
Interventions	Water: Warm-up exercises, trunk mobility exercises, postural stability exercises, transferring oneself and changing body positions, all carried out in water. Progression was encouraged with introduction of more complex exercises as appropriate.



Vivas 2011	(Continued)
------------	-------------

Land: Warm-up exercises, trunk mobility exercises, postural stability exercises, transferring oneself and changing body positions, all carried out on land. Progression was encouraged with introduction of more complex exercises as appropriate.

 $\label{lem:medication} \textit{Medication withheld for 12 hours before evaluations, for OFF-dose performance.}$

Outcomes Functional reach test

Berg balance scale

Gait - turn time, velocity, cadence, step amplitude

Timed up and go

UPDRS

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Low risk	Inclusion and exclusion criteria stated
Randomisation Method	Unclear risk	Method not stated
Concealment of Allocation	Unclear risk	Randomisation method not stated
Similarity at Baseline	Low risk	Group characteristics similar at baseline
Intention to Treat Analysis	Unclear risk	Analysis method not stated
Withdrawals Described	Low risk	Withdrawals at 8%
Cointerventions Constant	Unclear risk	Not stated whether the drug therapy was constant during trial
Comparable arms	Low risk	Same time and attention given to both groups
Blinded Assessors	High risk	Assessors not blinded

Werner 2010

Methods	Parallel group design		
	Randomised by individual not involved in testing or treatment using a computer generated number list for allocation		
	Analysed on an Intention-to-treat basis		
	Treatment delivered in 90 min sessions, 2 times per week for 2 weeks		
	Assessed at baseline (4-5 days prior to start of intervention), 1 week after training and at 3, 6 and 12 months		
	Assessors were blinded		
Participants	6 participants in Verbal instruction with augmented feedback group (VIAF), 6 in Verbal instruction only group (VI). 5 dropouts at follow-up time-points, group split not stated.		



Werner 2010 (Continued)

Participant mean age 72.8 years (VIAF), 69.3 years (VI); Male/female ratio 5/1 (VIAF), 5/1 (VI); Hoehn and Yahr 2.3 (VIAF), 2.5 (VI); Duration of condition not stated.

Inclusion criteria: Adequate vision. Exclusion criteria: MMSE<21, marked dyskinesia or other neurological impairments in addition to PD. Cardiac or musculoskeletal pathology that might impact gait, required an assistive device to walk, had video feedback or gait training previously or had difficulty understanding English.

Interventions

VIAB: 15 trials, of walking 7.5 m then returning to starting position with instruction to 'Walk as well as you can', per session. Verbal instruction and augmented feedback before each trial and 3 minutes of seated rest between each trial. Verbal instructions were to 'take big step' prior to each trial, during rest period participants were given 3 viewings of video playback of their walking performance from prior trial. On each viewing they were asked to a) focus on step length and comment on what they observed, b) indicate what they planned to do on next attempt. If subject did not comment accurately after viewing video or did not verbalise what they planned to do next, additional verbal 'knowledge of performance' (KP) or suggestions/corrections ('transitional information' (TI)) were provided by the experimenter.

VI: Trials conducted in same manner as for VIAB group but during rest period the group engaged in discussion or were given something to read.

Drug therapy was stable during trial period.

Outcomes

Stride length

Cadence

Gait velocity

Shoulder excursion

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Low risk	Inclusion and exclusion criteria stated
Randomisation Method	Low risk	Computer generated randomisation
Concealment of Allocation	Low risk	Randomised by individual not involved in testing or treatment
Similarity at Baseline	Low risk	Group characteristics similar at baseline
Intention to Treat Analysis	Low risk	Analysed on an intention-to-treat basis
Withdrawals Described	Low risk	No withdrawals at post-intervention time point. Withdrawals at 42% for follow up
Cointerventions Constant	Low risk	Drug therapy was stable during trial period
Comparable arms	Low risk	Same time and attention given to both groups
Blinded Assessors	Low risk	Assessors were blinded



Yaı	าฮ	20	110
	.0		

Methods Parallel group design

Randomised by independent arbiter who picked out a sealed envelopes 30 mins before the start of the

intervention

Method of analysis not stated

Treatment delivered in 30 mins sessions, 3 times per week for 4 weeks

Assessed at baseline, within 7 days of completion of treatment, 1 month after treatment

Assessors were not blinded

Participants

16 participants in downhill walking group (DW), 17 in physiotherapy group (PT). 2 dropouts in DW group, 5 dropouts in PT group.

Participants mean age 68.1 years (DW), 66.3 years (PT); Male/female 9/6 (DW), 7/8 (PT); Hoehn and Yahr 2.2 (DW), 2.2 (PT); Duration of condition 4.8 years (DW), 5.3 years (PT).

Inclusion criteria: Diagnosed with IPD (as defined by the UK Brain Bank Criteria) by a neurologist, Hoehn and Yahr stages I through III, ability to walk independently, stable medication usage, freedom from any other problems that might affect training and ability to understand instructions and follow commands. Exclusion criteria not stated.

Interventions

DW: Downhill walking training using a treadmill. Subjects walked on a motorized treadmill with a body weight support (BWS) system under the close supervision of a physical therapist. Initially the downhill grade was set at 3%. If subjects could walk with correct erect posture and large strides without stumbling during the training period, the downhill grade was then increased by 1% per training period. Treadmill speed was set at a level that was comfortable for each participant. A BWS of <40% of the body weight was provided and decreased to the maximum extent possible. Rest periods were provided as needed.

PT:Conventional therapy training program consisted of flexibility exercises (5 mins), strengthening exercises (7 mins), proprioceptive neuromuscular facilitation, coordination training, balance training (8 mins) and overground walking training (10 mins).

Drug therapy was stable during trial period.

Outcomes

Speed

Cadence

Stride length

Thoriacic kyphosis

Muscle strength

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Eligibility Criteria	Low risk	Inclusion criteria stated
Randomisation Method	High risk	Randomised by independent arbiter who picked out a sealed envelopes
Concealment of Allocation	High risk	Randomised by sealed envelopes
Similarity at Baseline	Low risk	Group characteristics similar at baseline



Yang 2010 (Conti

Intention to Treat Analysis	Unclear risk	Method of analysis not stated
Withdrawals Described	High risk	Withdrawals at 21%
Cointerventions Constant	Low risk	Drug therapy was stable during trial period
Comparable arms	Low risk	Same time and attention given to participants in both arms
Blinded Assessors	High risk	Assessors were not blinded

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Byl 2009	After contacting the author it was found that the study was not properly randomised
Cheon 2012	Not stated whether patients were randomised into trial. Author contacted but no response
Chouza 2011	Whole body vibration technique not usually used by physiotherapists
Cianci 2010	Excluded as confounded due to use of rolling walker
Donovan 2011	Quasi randomised using alternate allocation
Farley 2005	Insufficient information available from abstract for review and no author correspondence was achieved
Filippin 2010	Single arm study, so not randomised trial
Frazzitta 2012	Comparator arm consists of suggested home exercise and usual care only
Gobbi 2009	Insufficient information available from abstract for review and not stated whether randomisation was carried out
Haase 2011	Single short session of intervention; treatment under one day. Not rehabilitative therapy
Kamsma 1995	Partially randomised with a portion of patients placed in a convenient group based on practical grounds such as availability of transport
Klassen 2007	Difference between the two physiotherapy arms is multidisciplinary education for which the physiotherapy component is unknown
Knobl 2011	Not properly randomised, placed in groups
Lee 2011	Outcomes not relevant, stroke outcomes used and patients trained on one side of the body only
Ma 2009	Crossover trial carried out over one day only
Marchese 2000	Not properly randomised, pseudo-random number list used
Modugno 2010	Trial included a theatre training arm which could not be considered solely as physiotherapy
Munneke 2010	Randomised trial of systems of care for delivery of physiotherapy, not comparing different physiotherapy techniques



Study	Reason for exclusion
Pacchetti 2000	Two arm trial included active music therapy arm, not considered to be solely physiotherapy intervention
Pohl 2003	Randomised multiple intervention cross-over, over 4 consecutive days. Randomisation was of the sequence of the interventions, therefore not RCT
Rochester 2011	Excluded as the study was a randomised crossover over a couple of hours
Sage 2009a	After contacting the author it was found that the study was not properly randomised
Sage 2009b	After contacting the author it was found that the study was not properly randomised
Sage 2010	After contacting the author it was found that the study was not properly randomised
Stallibrass 2001	The method of therapy used - Alexander Technique - is not used by physiotherapists. Therefore this trial was excluded
Tamir 2007	Quasi-randomised trial using alternate allocation of patients to arms
Tickle-Degnen 2010	Multidisciplinary rehabilitation trail. Percentage component of physiotherapy was not specified, therefore unable to differentiate the contribution of physiotherapy to any change in the outcome measures
Wulf 2009	Multiple cross-over trial over a short period of time
Yen 2011	No outcome measures relevant to our review

Characteristics of ongoing studies [ordered by study ID]

Brauer 2011

Brauer 2011	
Trial name or title	Single and dual task gait training in people with Parkinson's disease: A protocol for a randomised controlled trial
Methods	Parallel group design
	Randomised with opaque envelopes containing allocations from a computer generated random number sequence prepared by offsite investigator not involved in recruitment, intervention or data collection
	Analysed on an intention-to-treat basis
	Treated as outpatients for 40 - 60 minutes session 3 times a week for 4 weeks, setting not stated
	Assessed at baseline, immediately after treatment and after 6 months follow-up
	Assessors blinded
Participants	Planning to recruit 60 participants.
	Inclusion criteria: 18 years or older, IPD diagnosed using UKBB criteria, able to walk 100 m independently with or without gait aids, report reduced step length or slowed gait speed, confirmed by clinical examination, Hoehn and Yahr stage I-IV. Exclusion criteria: Neurological condition other than PD, musculoskeletal or cardiopulmonary conditions that affect the ability to safely walk, had surgery for PD such as deep brain stimulation, score <24 on the mini-mental status examination.



Brauer 2011 (Continued)

Interventions

Single task training: One-to-one therapy sessions with therapist. Individually progressed program of gait training aimed at improving step length via repeated practice of straight line walking, turning, obstacle negotiation and challenging gait tasks such as increasing speed and altering surface challenges. External cues to increase step length will be used when needed. Instructions will not be given while participant is walking to avoid dual tasking. A home program will be incorporated at week 2 for 6 months, which includes a walking program and a range of balance, strengthening and postural exercises.

Dual task training: One-to-one therapy sessions with therapist. Aim to improve step length under dual task conditions that is when concurrently performing added cognitive or motor tasks. Participants will undertake repeated practice of walking aiming to improve step length using external cueing techniques including verbal, visual or auditory approaches. Progressing to internal concurrent cueing of appropriate step length. The gait tasks undertaken will be progressed from simple to more complex tasks as outlined for the single task group. In addition a variety of added tasks will be progressively integrated into the training program. These include tasks such as listening, speaking, conversing, generation of simple and complex lists, language, calculation and motor tasks increasing in complexity. Tasks will include those designed to reflect functional everyday activities such as carrying bags, getting keys out of a pocket, counting money, recalling directions or making a shopping list. Complexity will be progressively integrated as more complex tasks result in greater dual task interference with gait in people with PD. If able, participants will be progressed to performing increasingly complex cognitive tasks while concurrently walking. Motor tasks such as carrying and manipulation will also be included as added tasks.

Outcomes

Step length

Gait speed

Cadence

Stride length

Step length coefficient of variation

Double support time

Trail making A&B tests

Strop colour-word interference test

Digit span test

Timed up and go

6-minute walk test

Measures of community mobility questionnaire

Activity measured using ActivPAL

Quality of life

PDQ-39

EQ-5D

Hoehn and Yahr

UPDRS III

Freezing of Gait questionnaire

Ambulatory self confidence questionnaire

Compliance



Brauer 2011 (Continued)

Sta		

Contact information	s.brauer@uq.edu.au	
Notes	Protocol only	

Protas 2005

Trial name or title	Gait and step training to prevent falls in Parkinson's disease		
Methods	Parallel group design		
	Treatment delivered for 1 hour per day, 3 times per week for 8 weeks		
	Assessed at baseline, at 8 weeks, at 1 month follow-up and 5 month follow-up		
	Assessors blinded		
Participants	Planning to recruit 90 participants.		
	Inclusion criteria: Diagnosis of PD, postural instability - gait disorder predominant PD, history of falls, gait freezing or a positive pull test, stable regimen of medications, ability to stand and walk 3 m without assistance, stage 2 or 3 of the Hoehn and Yahr disability scale and moderate or higher cognitive scores.		
Interventions	GSP: Gait and step perturbation. Gait training on a treadmill and in a safety harness while walking in four directions: frontwards, backwards, left sideways and right sideways. Training will start with a treadmill speed that is equivalent to fastest overground walking speed for forward walking and fastest possible for other directions and will increase during training. This group will also receive step training while positioned in 4 directions consisting of suddenly turning the treadmill on/off. The subjects will be required to maintain their balance during perturbations.		
	SPT: Seated exercise. This group will receive seated active range of motion, and upper and lower extremity aerobic training.		
Outcomes	Usual and fastest gait speed		
	5-step test		
	UPDRS		
	Gait and balance scale		
	Center for Epidemiological Studies depression scale		
	Cognistat - a co-morbidity scale		
	Activities balance confidence		
	Physical activity scale for the elderly		
	Limits of stability		
	Falls frequency		
Starting date	25th September 2006		
Contact information	shinowara@nih.gov		



Protas 2005 (Continued)

Notes

_			_		-	
	han	kman		nı	- 7 P	٩
36	11611	RIIIAII	_	LU		э,

Trial name or title	Endurance exercise in Parkinson's disease		
Methods	Parallel group design		
	Randomisation method not stated		
	Treatment was delivered four times a week for 6 months		
	Assessors were blinded		
Participants	Estimated total enrolment, 126 participants. Three arm trial: No intervention control group (NT), vigorous exercise group (VE), moderate exercise group (ME).		
	Inclusion Criteria: Clinical diagnosis of primary Parkinson's disease, in a Hoehn and Yahr stage less than stage III, disease duration is less than 5 years, not likely to require dopaminergic therapy within 6 months. Exclusion Criteria: Use of any PD medication within 60 days prior to the beginning the study, in-		
	cluding levodopa, direct dopamine agonists, amantadine, Rasagiline (Azilect), Selegiline (Eldepryl), Artane (trihexyphenidyl). Duration of previous use of medications for PD that exceeds 90 days. Expected to require dopaminergic therapy in the next 6 months. Poorly controlled or unstable cardiovascular disease. Uncontrolled hypertension. Hypo- or hyperthyroidism, abnormal liver function, abnormal renal function. Mild cognitive impairment (Montreal Cognitive Assessment score<26/30). Depression that precludes ability to exercise (Beck depression score>13). Disorders that interfere with ability to perform endurance exercises. Regular participation in vigorous endurance exercise. Evidence of serious arrhythmias or ischemic heart disease. Any clinically significant medical condition, psychiatric condition, drug or alcohol abuse, or laboratory abnormality that would, in the judgment of the investigators, interfere with the ability to participate in the study.		
Interventions	NT: Control Group. Wait listed to moderate or vigorous exercise after 6 months of no exercise.		
	VE: Endurance exercise at 80-85% HR max.		
	ME: Endurance exercise at 60 - 65% HR max.		
Outcomes	Adherence to exercise		
	UPDRS Motor		
	Adverse events		
Starting date	April 2012		
Contact information	lindsey.pederson@ucdenver.edu		

Smith 2009

Trial name or title Strength training and medication effects in Parkinson's disease: effects on hypokinesia in Parkinson's disease: effects on hypokinesia in Parkinson's disease



Smit	h 2	009	(Continued)
------	-----	-----	-------------

Methods	Treated for 12 weeks
Participants	Inclusion criteria: Male or female at least 40 years of age, neurologist diagnosed idiopathic PD (using UK Brain Bank Criteria), ambulatory and medically cleared by their physician to participate in an exercise regimen, clinical signs of hypokinesia or postural instability, Folstein Mini-Mental State Examination score > 23, currently taking dopamine replacement medication.
	Exclusion criteria: Previous surgical management of PD (pallidotomy, DBS), motor fluctuations and/or dyskinesias uncontrolled by medications, central nervous system disorder (other than PD), myopathic disease (e.g. focal myopathy) that affects skeletal muscle structure/function rheumatological disease that has an effect on muscle and/or mobility, unstable cardiovascular disease that limits exercise abilities, impaired knee flexion, <90°, extreme claustrophobia (secondary to the inability to perform the MRI scans) regular (2-3 times/week) aerobic or resistance exercise performed over the past 6 months.
Interventions	High force resistance training
	Standard care
Outcomes	Muscle structure
	Muscle force output
	Hypokinesia
Starting date	August 2007
Contact information	sheldon.smith@hsc.utah.edu
Notes	

Uc 2010

Trial name or title	Effects of aerobic exercise in Parkinson's disease
Methods	Assessors blinded
Participants	100 participants to be enrolled.
	Inclusion criteria: Veteran or non-veteran, presence of all 3 cardinal features of Parkinson's disease (resting tremor, bradykinesia and rigidity) which must be asymmetrical, Hoehn & Yahr stage I-III, men or women aged 50-80 capable of performing the planned exercise programs, intention to remain in the local area over the study period, stable dopaminergic treatment regimen for at least 4 weeks prior to baseline without any clinical need for medication adjustment at the time of screening.
	Exclusion criteria: Secondary parkinsonism, Parkinson-plus syndromes, MMSE score<24, participating in an aerobic exercise program, an unstable dosage of drugs active in the central nervous system during the 60 days before the baseline visit, participation in drug studies or the use of investigational drugs within 30 days before screening, structural brain disease, active epilepsy, acute illness or active confounding medical, neurological or musculoskeletal conditions, alcoholism or other forms of drug addiction, inability to complete the graded exercise test, lack of medical clearance from our pulmonologist, intention to move or take > 1 month vacation during the study period, contraindications to MRI or claustrophobia requiring sedation.
Interventions	Aerobic exercise in the form of brisk walking using different training formats, four arms: group/continuous, group/interval, single/continuous and single/interval.



Uc 2010 (Co	ntinued)
--------------------	----------

(**************************************							
Outcomes	Cycle ergonometry						
	Functional and morphometric MRI						
	Trophic factor						
	Biomarker assays						
	Safety exams						
	Lab results (ECG, biochemistry, CBC)						
	Tests for Parkinsonism and cognition						
	Questionnaires about quality of life, mood and activities of daily living						
Starting date	February 2009						
Contact information	ergun-uc@uiowa.edu						
Notes							

van Nimwegen 2010

Trial name or title	ParkFit study: A randomised controlled trial evaluating the effectiveness of a multifaceted behavioural program to increase physical activity in Parkinson patients			
Methods	Parallel group design			
	Randomised assigned using minimisation algorithm			
	Treatment delivered over 2 years, offered a maximum of 35 sessions per year, each session lasting 30 mins			
	Assessed at baseline, at 6 months, 12 months 18 months and 24 months			
	Assessors were blinded			
Participants	586 participants randomised, 299 in ParkFit group and 287 in ParkSafe group. No dropouts described.			
	Baseline characteristics for groups not stated			
	Inclusion criteria: PD according to UKBB criteria, aged between 40 and 75 years, sedentary lifestyle (<3 times a week vigorous-intensity physical activity for <60 mins or <3 times a week moderate-intensity physical activity for <150 mins), Hoehn & Yahr score ≤3. Exclusion criteria: Unclear diagnosis (no gratifying and sustained response to dopaminergic therapy), MMSE score <24, unable to complete dutch questionnaires, severe co-morbidity interfering with daily functioning, daily institutionalised care, deep brain surgery.			
Interventions	Both interventions were delivered exclusively by experienced therapists who participate in the Dutch ParkinsonNet.			
	ParkFit: Brochure provided, <i>ParkFit</i> , covering specific strategies to promote behavioural change. Physical therapists serve as personal activity coaches who guide patients towards a more active lifestyle, during specific coaching sessions. Patient and coach create activity goals in order to obtain the 6-month goals. Patients receive a personal ambulatory monitor with automated visual feedback showing the amount of actually delivered daily physical activity, recorded by a triaxial accelerometer. The ParkFit program also included a maximum of 19 physical therapy sessions in year			



van Nimwegen 2010 (Continued)	1 and 23 in year 2. Based on individual disabilities, therapist and patient jointly formulate treatment aims based on the evidence-based guideline of physical therapy for PD. ParkSafe: Patients receive a brochure, <i>ParkSafe</i> with information about the benefits of physical therapy. Specific emphasis is given to the importance of safety when performing daily activities. Patients receive an individualised physical therapy program. Total of 35 sessions per year: 19 physiotherapy plus 16 coach sessions). Physical therapist and patient jointly formulate the aims of the projected treatment plan, based on individual problems and disabilities. Not stated whether the drug therapy was constant during trial.					
Outcomes	Level of physical activity					
	6 minute walk test					
	PDQ-39					
Starting date	September 2008					
Contact information	m.munneke@neuro.umcn.nl					
Notes	Subgroup analysis of those who succeeded in increasing their activity levels versus those who n't, assessed for disease progression and physical fitness					


ADDITIONAL TABLES

atts at post test	
	Cochi Libra

Table 1. Included trials: summary table of results

Stu dly - Grocap- tion	Interventions	terventions Withdrawals at post-test (with reasons)		post-test (with rea-		post-test (with rea-		Outcomes	Summary of results at post-test
_	Arm 1	Arm 2	Arm 3	Arm 4	Arm 2	Arm 3	Arm 4		
da ter- 201 k oo,	Visual cueing on the ground (n=14; 30 minutes, 3 times per aweek, 6 weeks)	Visual cueing on the treadmill (n=14; 30 minutes, 3 times per week, 6 weeks)	Con- trol (n=14)		0	0	Primary: Step length Secondary: UPDRS III, timed up and go, gait speed, cadence, double support time, step time, step-to-step variability, step time variability, 30 second chair stand	There was no difference between treatment arms	
	nethysiotherapy with smental practice (n=25; 1 hour (or 2 x 30 minutes) weekly, 6 weeks)	Physiotherapy with relaxation (n=25; 1 hour (or 2 x 30 minutes) weekly, 6 weeks)		3 (n=2; hos-pi-talised with re-lapse, n=1; too con-frontin	4 (n=3; hos- pi- talised with re- lapse, n=1 died)		Visual analogue scale, timed up and go, 10 m walk test	There was no difference between treatment arms	
ni cona	Aerobic training a,(n=13; 45 minutes, 3 times per week, 7 weeks)	Qigong group (n=13; 45 minutes, 3 times per week, 7 weeks)		2 (n=1; poor com- pli- ance, n=1; back pain after 3rd ses- sion)	2 (n=1; poor com- pli- ance, n=1; fall- relat- ed frac- ture)		Primary: UPDRS III, UPDRS II, Brown's disability scale (BDS), 6 minute walk test, Borg scale for breathlessness, Beck depression in- ventory (BDI), PDQ-39. Secondary: Spirometry test, Maximum cardiopulmonary exercise test	There was no difference between treatment arms for primary outcomes	

Table 1. Included trials: summary table of results (Continued)



of 6

Cochrane
Library

Trusted evidence.
Informed decisions.
Better health.

Table 1. Included trials: sun	u. y tubite on resu	ses- sions over 8 weeks)		Sit-to-stand test, Transcranial magnetic stimulation (subset)	
FraMon-Treadmill with audi- it- tes- tory and visual cues ta cano (n=20; 20 minutes 200 Raly daily, 4 weeks)	Auditory and visual cues (n=20; 20 min- utes daily, 4 weeks)	0	0	UPDRS III, Gait speed, Freezing of gait questionnaire, Stride length, 6 minute walking test	The treadmill with auditory and visual cues had significant improvement in gait speed, freezing of gait questionnaire, stride length and 6 minute walking test when compared to the auditory and visual cues group. There were no differences between treatment arms for UPDRS III
Hac\$t. Tango (n=9; 1 hour, neyLouis20 sessions within 13 200Mis- weeks) souri, USA	Exercise (n=10; 1 hour, 20 sessions within 13 weeks)	0	0	UPDRS III, Berg balance scale, Freezing of gait, Timed up and go, Velocity of walking and dual-task walking	There was no difference between treatment arms
Hac\$t. Tango group (n=19; neyLouis] hour, twice weekly, 200Mis- total 20 sessions in souri,13 weeks) USA	Waltz/foxtrot group (n=19; 1 hour, twice weekly, total 20 sessions in 13 weeks)	Tai 5 of The Chi fly of Tai 1 of The Chi fly of Tai 1 of	(n=1; up in- 20; jury b- at b- at ex;s) home, ; n=1; re infre- n, quent ; at- ns- ten- ta- dance n (un- b- known is, ; nge	4 PDQ-39, UPDRS III, (n=2; (n=1; trans Resg balance scale, Tandem stance portal st, Timed up and go test, One leg tion thence test probilems, sance test n=1; tion hos- (un- pi- re- tal- lat- i- ed), sa- n=1;) tion an- (un- kle re- in- lat- jury, ed), n=1; n=1; death in- in suf- fam- fi- i- ciently) ly in-	PDQ-39 significantly improved in the tango arm compared to the waltz/foxtrot and Tai Chi arms. Timed up and go test was significantly improved in the tango arm compared to the waltz/foxtrot and Tai Chi arms. There was no difference between treatment arms in UPDRS III

Cochrane Library

tense ex-

er-

cise)

percentage

Hac\$t. Partnered tango neyLouis(n=19; 1 hour, twice 201Mis- weekly, 10 weeks) souri, USA

Non-partnered tango (n=20; 1 hour, twice weekly, 10 weeks)

7 5 (n=1; (n=1; exprogrespressed lack sive deof incline ter-

in est, menn=1; tal new

stajob intertus, n=2; fered with excesclass, sive n=1; unretrav-

elling lated dismedical tance, n=1; probfelt lems,

classn=1; es work were com-

mittoo faments, tigun=1; ing, un-

n=3; able to reunable turn

to refor folturn low-up for fol-

mealow-up sures) measures)

Tandem stance, one leg stance, Timed up and go, 6 minute walk test, Gait velocity, Cadence, Stride length, Swing percentage, Double support

There was no difference between treatment

Trusted evidence.
Informed decisions.
Better health.

Table 1. Included trials: sur	nmary table of results	(Continued)			
HasElori-Tai Chi (n=unknown, 200da, total n=23; 1 hour, USA twice weekly, 16 weeks)	Qi-gong (n=un- known, total n=23; 1 hour, twice week- ly, 16 weeks)	Un- known	Un- known	Gait initiation, Gait velocity, Stride length, Stance, Double limb support, Step duration	There was no difference between treatment arms
HirsNorthCombined balance	Balance training	0	0	Balance, Muscle strength (subset	Combined balance and resistance training

HasElori-Tai Chi (n=unknown, 200 d a, total n=23; 1 hour, USA twice weekly, 16 weeks)	Qi-gong (n=un- known, total n=23; 1 hour, twice week- ly, 16 weeks)	Un- known	Un- known	Gait initiation, Gait velocity, Stride length, Stance, Double limb support, Step duration	There was no difference between treatment arms
Hirshbrtl Combined balance 1996ar- and resistance train- oli- ing (n=6; 45 minutes, na, 3 times per week, 10 USA weeks)	Balance training group (n=9; 30 min- utes, 3 times per week, 10 weeks)	0	0	Balance, Muscle strength (subset group): knee extensors, knee flexors, ankle plantar flexors. Latency to fall, % of trials resulting in falls	Combined balance and resistance training improved balance scores significantly more than the balance training group. There were no differences between treatment arms for the falls outcomes
Jou Crétil Asymmetric motor 201Erandraining program (n=unknown, total n=50; 1 hour, 3 times per week, 8 weeks)	Broad program (n=unknown, to- tal n=50; 1 hour, 3 times per week, 8 weeks)	Un- known	Un- known	UPDRS III, GMT score, Rapid alternating movements, Handwriting and spiralography, PDQ-39, Dpression (GDS-15), Video recording of 8 activities of daily living and biomechanical evaluations	Abatract describing methodology. No result data
JunGeor-Aerobic exercise cosgia, (n=unknown, total 2000SA n=56; 6 months)	Tai Chi (n=un- known, total n=56; 6 months)	Qi- Un- gong known (n=un- (total known,n=16: to- n=2; tal seri- n=56; ous 6 ad- month syerse events, n=14; unre- lated med- ical or lo- gis- tical prob- lems)	Un- known (total n=16: n=2; seri- ous ad- verse events, n=14; unre- lated med- ical or lo- gis- tical prob- lems)	Un- UPDRS total, UPDRS motor, UPDRS know/hDL, PDQ-39, Clinical global impres- (to- sion, Walking speed, Falls tal n=16: n=2; se- ri- ous ad- verse events, n=14; un- re- lat- ed med- ical or lo- gis- ti- cal	UPDRS ADL scores improved significantly more with Qi-gong than aerobic exercise. There was no difference between treatment arms for PDQ-39 and UPDRS total and motor subscores. There was insufficient information on clinical global impression, walking speed and falls


Cochran

 Table 1. Included trials: summary table of results (Continued)

problems)

				lems)	
Kha\$au- Physiotherapy and laf di treadmill (n=15) 201Ara- bia	Physiotherapy (n=15)	Un- known	Un- known	UPDRS II & III, Hamilton rating scale of depression, Walking speed, Walking distance	Both treatment arms showed significant improvement in walking distance, speed and ADL. A significant improvement in depression only observed in physiotherapy and treadmill group. Unclear if there were any differences between treatment arms
Li Ore- Tai Chi (n=65; 1 hour, 201ĝon, twice weekly, 24 USA weeks)	Resistance training (n=65; 1 hour, twice weekly, 24 weeks)	Con- 9 trol, (n=4; stretchhealth ing prob- group lem, (n=65; n=3; 1 non- hour, com- twice mit- week- tal/time ly, con- 24 flict, weeks)n=2; relo- cat- ing)	6 (n=4; health problem, n=1; non- com- mit- etal/time con- flict, n=1; relo- cat- ing)	4 Primary: Two indicators of postural (n=3;stability: maximum excursion and dihealthectional control. problem, Secondary: Stride length, Walking n=1; velocity, Strength of bilateral knee non-extensors and flexors, Functional com-reach test, Timed up and go, UPDRS mit-III, Number of falls etal/time conflict)	The Tai Chi group performed significantly better than those in the resistance training and stretching groups on the primary outcomes. The Tai Chi group had significantly better performance/scores in many outcomes compared to the stretching group. The Tai Chi group out performed the resistance training group on the stride length and functional reach
LouCaitólConventional physi- 2010a cal therapy (n=6, ses- do sions, twice weekly, Parartá,tal 12 sessions) Brazil	Complementary activities (n=6, ses- sions, twice week- ly, total 12 ses- sions)	Un- known	Un- known	Timed up and go, Anterior functional reach	There was no difference between treatment arms
MalHongAudio-visual cued 200Kongtask-specific training Chi- (n=21; 20 minutes, na 3 times per week, 4 weeks)	Conventional exercise (n=21; 45 minutes, twice weekly, 4 weeks)	Con- 2 trol (n=1; (n=18; change 4 of weeks)med- ica- tion, n=1; heel pain)	2 (n=1; fall with frac- ture, n=1; went over- seas)	4 Peak horizontal velocity, Peak ver- (n=2;tical velocity, Movement time,3 D de- Kinematics data of sit-to-stand clined to come back, n=2; went over- seas)	The audio-visual cued task-specific training group increased both peak horizontal and vertical velocities and reduced time taken to complete sit-to-stand. These improvements were greater than those of the conventional exercise group and control

Table 1. Included trials: summary table of results (Continued)



Progressive strength training (n=70; 2 hours + 2 hours home practice program, once a week, 8 weeks)

Life 2 skills (n=1; con- untrol able (n=71; or 2 unhours will-

> а week,

weeks)

1 (n=1;health reasons) + 2 ing hours to athome tend, prac- n=1; tice death) program, once

12 Primary: Falls: no. of fallers per (n=2; group, no. of multiple fallers per un- group, falls rate over 12 months in able each group. or

Secondary: Number of injurious falls, unwill- Walking speed, 6 minute walk test, Timed up and go, UPDRS II and III, PDQ-39, EuroQol-5D to at-

health, n=1; preference for exercise group, n=1; death, n=2; health reasons, n=1; unspecified, n=1; group was "depressing", n=2, not exercis-

tend,

n=2;

poor

Time to first fall during the intervention phase did not differ across groups. Full trial results not yet published



41	11-
Library	Cochrane

Trusted evidence.
Informed decisions.
Better health.

	Table 1.	Included	trials:	summary	table	of results	(Continued)
--	----------	----------	---------	---------	-------	------------	-------------

	Tuble 1. Illetadea tilats. sui	initially table of results (co	nunueu)			
				ing or		
				re- cei\	,	
				ing		
:				fall: ed-		
				u-		
:				ca- tior		
	Miy@s- Body weight sup- 200@- ported treadmill ka, training (n=5; 45 min- Japamtes, three times per week, 4 weeks)	Physical therapy (n=5; 45 minutes, three times per week, 4 weeks)	0	0	UPDRS, UPDRS subscales (mental, ADL, motor and complications), Overground ambulation endurance, Gait speed, No. steps taken for 10 metre walk	Cross-over trial, combined data presented. Body weight-supported treadmill training showed greater improvement in UPDRS total, ambulation speed and number of steps than physical therapy arm. Additional data supplied by author allowed data to be separated and there was no difference between arms
·	Miy@s- Body weight sup- 200@- ported treadmill ka, training (n=11; 45 Japamninutes, three times per week, 4 weeks)	Physical therapy (n=9; 45 minutes, three times per week, 4 weeks)	1 (n=1, med- ica- tion change	3 (n=3, med- ica- tion ed):hanged)	Primary: UPDRS, Gait speed, No. steps taken for 10 metre walk. Secondary: UPDRS subscales (mental, ADL, motor and complications)	Body weight-supported treadmill training had significantly greater improvement than the physical therapy group in gait speed at 1 month; and in the number of steps at 1, 3 and 4 months
	MorMel- Movement strateris bourgey, training (n=14; 45 200Aus- minutes, max 16 sestraliasions over 2 weeks)	Musculoskeletal exercise (n=14; 45 minutes, max 16 sessions over 2 weeks)	0	2 (n=2; lost to fol- low up, no de- tails)	Primary: UPDRS Motor and ADL (combined). Secondary: 10 m walk test, Timed up and go, 2 min walk test, Balance-shoulder tug, PDQ-39	The movement strategy training group had a significant improvement in balance compared to the musculoskeletal arm. There were no other differences between treatment arms
	Palmin- United Parkinson 1986eap 6 oundation exerlis, cise program (n=7; 1 Min- hour, three times per neso-week, 12 weeks) ta, USA	Upper body karate training program (n=7; 1 hour, three times per week, 12 weeks)	0	0	Forearm pronation/supination rate, Pursuit score walk index, Degree of activated rigidity, Degree of arm tremor, Activated rigidity, Grip strength, 9-hole peg test, Minnesota placing and turning test, Arm swings test, Rapid alternating arm movement test, Button board, Putting	Study did not compare differences between treatment arms

 Table 1. Included trials: summary table of results (Continued)

shirt on and off, Putting shoes and

				socks on and off, Getting up from chair, Long latency stretch response	
Pel@en- Action plus physical 2016- therapy group (n=9; 1 va, hour, three times per Italy week, 4 weeks)	Landscape plus physical therapy group (n=9; 1 hour, three times per week, 4 weeks)	1 (n=1; found to have past his- to- ry of neu- ro- log- ical con- di- tions other than PD or im- plan- ta- tion for deep brain stim- ula- tion)	1 (n=1; found to have past his- to- ry of neu- ro- log- ical con- di- tions other than PD or im- plan- ta- tion for deep brain stim- ula- tion)	FOG Questionnaire and FOG diary, Timed up and go, 10 metre walking test, Tinetti scale part I and II, Berg balance scale, PDQ-39	At post-test there was no difference between treatment arms
Pi- Veror pot assisted gait- cel-Italy training group (n=21; li 45 minutes, three 2012 times per week, 4 weeks)	Physiotherapy group (n=20; 45 minutes, three times per week, 4 weeks)	3 (n=3; lack of co- op- era- tion)	2 (n=2; lack of co- op- era- tion)	Primary: 10 m walk test, 6 min walk test. Secondary: Spatiotemporal gait parameters including stride length, Parkinson's fatigue scale, UPDRS Total	Robot assisted gait-training significantly improved 10 m walk test, 6 min walk test, Parkinson's fatigue scale, UPDRS Total compared to the physiotherapy group
Po-Man-20 week gym group li- ches-(n=16; 1 hour, twice a week, 20 weeks)	10 week gym group (n=16; 1 hour, twice a week, 10 weeks)	4 (n=1; ran-	6 (n=1; ran-	Simple, choice and serial reaction time, Videotaped motor perfor- mance, PDQ-39, UPDRS III, Illness	No significant differences between arms, except for the learning sequence RT outcome



perceptions (BIPQ), Questionnaire assessing experiences of programme

Table 1.	Included trials: summary table of results	(Continued)
ako fé r,		don

ako fér ,	domiseddomised	
200 0 K	but	but
	did	did
	not	not
	start	start
	inter-	inter-
	ven-	ven-
	tion,	tion,
	n=3;	n=5;
	did	did
	not	not
	com-	com-
	plete	plete
	or in-	or in-

ReuGiess Mordic walking 201Ger- group (n=30; 70 minmanyutes, three times per week, 6 months)

Walking group (n=30; 70 minutes, three times per week, 6 months)

ibility
and
relaxation
group
(n=30;
70
minutes,
three

sufficient

sessions)

Flex- 0

times per

week,

months)

suffi-

cient ses-

sions)

0

Max walking speed on treadmill, 12 m and 24 m walking test, Stride length, Gait variability, UPDRS, PDQ-39, Physical activity in everyday life, Adverse effects Main reported differences were that the Nordic walking group was superior to the walking and flexibility and relaxation programme in improving postural stability, stride length, gait pattern, and gait variability

Rid & eveForced exercise 200 and, group (n=5; 1 hour, Ohio,3 times per week, 8 USA weeks) Voluntary exercise group (n=5; 1 hour, 3 times per week, 8 weeks) 0

0

UPDRS part III

Manual functional dexterity, Bimanual dexterity, Centre of pressure (CoP)

UPDRS motor scores showed a significantly greater improvement in the forced exercise arm compared to the voluntary exercise arm. Only forced exercise resulted in significant improvements in bimanual dexterity

Table 1. Included trials: summary table of results (Continued)



Fitness counts (n=unknown (total n=48 at 6 month time point); 1 hour, twice per week, 24 months) Un- Unknown known UPDRS motor, Timed up and go, Berg balance scale, Modified physical performance test There was no difference between treatment

SchDekr	r Ele xibilty/bal-
201 2e r,	ance/function ex-
Col-	ercise group (n=39;
orad	d, hour, 3 times per
USA	week for the first 4
	months then tapered
	for 1 month to once
	monthly sessions out
	to 16 months)

Aerobic exercise
group (n=41; 1
hour, 3 times per
week for the first
4 months then tad pered for 1 month
to once monthly
sessions out to 16
months)

exer-(n=1; health cise group prob-(n=41; lems, once n=2; month-perly sonsual ispersues) vised sessions)

Home 3 at 4 7 at 4 6 Primary: Overall physical function, Balance - functional reach, Walking months months at (n=1;economy - VO₂ 4 months not (n=3; Secondary: hapру not hap- UPDRS ADL, UPDRS Motor, PDQ-39 with the ру prowith

Overall physical function: improvement at 4 months was greater in the Flexibilty/balance/function exercise group than the aerobic and home exercise group. Functional reach was not different between groups. Walking economy: aerobic exercise improved compared to Flexibilty/balance/function exercise group. The only secondary outcome that showed significant differences was UPDRS ADL: Flexibilty/balance/function exercise group performed better than home exercise group at 4 months

pointdid ment, not n=2; rehealth turn probcalls, lems, n=1 n=1; unmoved, able n=1; to decomceased) mit the time, n=1; missed ар-

0

pointment)

gram,

missed

n=2;

ap-

the

pro-

gram,

n=1;

Sha@adar Treadmill with cue-200@ary, ing group (n=10; 30 Canaminutes, twice a

da week, 8 weeks)

Treadmill without cueing (n=9; 30 minutes, twice a week, 8 weeks) Cue- 0 ing on- ly group (n=10;

0 Gait and Balance Scale, UPDRS III, PDO-39

Limited data. There was no difference between treatment arms

- Juli
Cochrane Library

Table 1.	Included trials: summary table of results (Continued)
	20

30 minutes, twice ek, eks)

0

0

		a wee 8 wee
ShelflongBalance group (n=23;	Strength training	

leg stance time, Activities-specific	Limited data. The balance group improved significantly more than the strength group in movement velocity and one leg stance

tion gait training

Sig-Reyk-Walking with visual

urgiatun cues (n=unknown,

sondur, total n=26; 30 min-

200Rce- utes, 4 sessions per

land week, 4 weeks)

201Kongtreatment delivered

Chi- over 12 weeks)

na

Auditory stimulation (n=unknown, total n=8; treated for an unknown period of time)

Walking without

cues (n=unknown,

total n=26; 30 min-

utes, 4 sessions per

week, 4 weeks)

(n=22; treatment

delivered over 12

weeks)

Un-Unknown known

0

0

Stride length

Secondary:

Timed up and go, PDO-39

Limited data. There was no difference between treatment arms for Timed up and go

Stride length was significantly greater after vi-

sual stimulation than after auditory stimula-

Smalero	nBaalance training
nia Italy	(n=33; 50 minutes,
2010	3 times per week, 7
	weeks)

General physical exercise (n=31; 50 minutes, 3 times per week, 7 weeks) 5 4 (n=2; (n=2;ununcocooperoperativeative-

Primary: Berg balance scale, Activities-specific balance confidence, Postural transfer test, Self destabilization of the centre of foot pressure test, Falls diary.

There was a significant difference in favour of the balance training arm in the Berg Balance Scale and the self destabilization of the centre of foot pressure test when compared to the general physical exercise group

ness, ness, n=3; n=2; medmedical ical comcomplicaplications) tions)

UPDRS Total, Modified Hoehn and Yahr, Geriatric depression scale

Ta-Ban-	Conventional gait
lakl ga d	training (n=un-
201lbre,	known, total n=60; 8
In-	hours over 4 weeks)
dia	

Partial weight supported treadmill training: 20% unweighting (n=unknown, total n=60;

Con-Un-Untrol: known spe-

No

cif-

Un- Dynamic posturography, UPDRS (toknown knowtral and motor subscore), Beat-tobeat finger blood pressure

Limited data. Partial weight supported (-20%) treadmill training had a significantly greater improvement in UPDRS motor score compared to conventional gait training

Cochrane
Library

Table 1. Included trials: sum	8 hours over 4 weeks)	ic in- ter- ven- tion (n=un- known, to- tal n=60)			
ThaQol- Novel rhythmic au- 1996radoditory stimulation USA group (n=15; 30 min- utes per day, total 10.5 hours over 3 weeks)	Standard self paced training group (n=11; 30 minutes per day, total 10.5 hours over 3 weeks)	No 0 treat- ment group (n=11)	0	Stride velocity, Stride cadence Stride length, EMG analysis o muscles	
TooFeori-Standard treadmill 200da, group (n=unknown, USA total n=23; 20 min- utes, 3 times per week, 6 weeks)	Unweighted treadmill group (n=un-known, total n=23; 20 minutes, 3 times per week, 6 weeks)	WeightUn- ed known tread- mill group (n=un- known, to- tal n=23; 20 min- utes, 3 times per week, 6 weeks)	Un- known	Un- Balance from dynamic postu knowphy, Berg balance scale, UPD mechanical assessment of st and range of motion, Gait	RS, Bio- arms for function and stability in gait and dy-
Vi- A Water based exercise vas Coru寶和pup (n=6; 45 min- 201\$painutes, twice weekly, 4 weeks)	Land based exercise group (n=6; 45 minutes, twice weekly, 4 weeks)	1 (n=1; in- fluen- za -	0	Functional reach test, Berg b scale, Gait -turn time, velocit dence, step amplitude, Time and go, UPDRS	y, ca- ly improved Berg Balance Scale and UPDRS

Cochrane
Library

Trusted evidence.
Informed decisions.
Better health.

Table 1. Included trials: sur	nmary table of results (did not re- ceive inter- ven- tion)			
WeiNew Verbal instruction er York, with augmented 2010SA feedback group (n=6; 90 minutes, 2 times per week, 2 weeks)	Verbal instruction only group (n=6; 90 minutes, 2 times per week, 2 weeks)	0	0	Stride length, Cadence, Gait velocity, Shoulder excursion	There was no difference between treatment arms
YangaipeDownhill walking 201Tai- group (n=16; 30 min- wan utes, 3 times per week, 4 weeks)	Physiotherapy group (n=17; 30 minutes, 3 times per week, 4 weeks)	1 (n=1; low mo- tiva- tion)	2 (n=1; con- flict- ing work sched- ule, n=1; trans- port prob- lem)	Speed, Candence, Stride length, Thoriacic kyphosis, Muscle strength	Downhill walking significantly improved gait speed, stride length and muscle strength of knee extensors compared to physiotherapy



APPENDICES

Appendix 1. Search strategies

Cochrane Movement Disorders Specialised Register (search end of January 2012).

- #1. SR-Movement
- #2. physiotherapy
- #3. exercise
- #4. physical therapy
- #5. rehabilitation
- #6. #2 or #3 or #4 or #5
- #7. Parkinson
- #8. Parkinsons disease
- #9. parkinsonism
- #10. #7 or #8 or #9
- #11. #1 and #6 and #10

The Cochrane Library and The Cochrane Controlled Trials Register (Wiley online library) (Issue 1, 2012).

- #1. MeSH descriptor: [Parkinson Disease] explode all trees
- #2. parkinson*
- #3. parkinsonism
- #4. {or #1-#3}
- #5. MeSH descriptor Physical Therapy (Speciality) explode all trees
- #6. MeSH descriptor Physical Therapy Modalities explode all trees
- #7. physical therapy
- #8. physiotherapy
- #9. MeSH descriptor Exercise explode all trees
- #10. exercise
- #11. MeSH descriptor Rehabilitation explode all trees
- #12. rehabilitation
- #13. {or #5-#13}
- #14. #4 and #13

MEDLINE (Ovid) 1966 - Jauary (Week 4) 2012.

- 1. randomized controlled trial.pt
- 2. controlled clinical trial.pt
- 3. randomized.ab.
- 4. placebo.ab.



- 5. drug therapy.fs.
- 6. randomly.ab.
- 7. trial.ab.
- 8. groups.ab.
- 9. 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8
- 10. exp animals/ not humans.sh.
- 11.9 not 10
- 12. exp Parkinson Disease/
- 13. "parkinson*".ab.ti.
- 14. 12 or 13
- 15.exp "Physical Therapy (Specialty)"/
- 16. physiotherapy.mp.
- 17. exp Exercise/
- 18. exp Rehabilitation/
- 19. exercise*.mp. [mp=title, original title, abstract, name of substance word, subject heading word, unique identifier]
- 20. rehabilitation*.mp. [mp=title, original title, abstract, name of substance word, subject heading word, unique identifier]
- 21. 15 or 16 or 17 or 18 or 19 or 20
- 22. 11 and 14 and 21
- 23. limit 22 to yr="2001 -Current"

EMBASE (Ovid) 1974 - January (Week 4) 2012.

- 1. random\$.mp. [mp=title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer name]
- 2. factorial\$.mp. [mp=title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer name]
- 3. (crossover\$ or cross-over\$).mp. [mp=title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer name]
- 4. placebo\$.mp. [mp=title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer name]
- 5. (doubl\$ adj blind\$).mp. [mp=title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer name]
- 6. (singl\$ adj blind\$).mp. [mp=title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer name]
- 7. assign\$.mp. [mp=title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer name]
- 8. allocat\$.mp. [mp=title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer name]
- 9. volunteer\$.mp. [mp=title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer name]
- 10. crossover procedure.sh.
- 11. Double-blind Procedure.sh.
- 12. Randomized Controlled Trial.sh.
- 13. Single-blind Procedure.sh.
- 14. 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13



- 15. exp Parkinson Disease/ 16. parkinson*.mp. 17. 15 or 16 18. exp physiotherapy/ 19. physical therapy.mp. 20. exp Exercise/ 21. exp Rehabilitation/ 22. physiotherapy.mp. [mp=title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer name] 23. exercise.mp. [mp=title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer name] 24. rehabilitation.mp. [mp=title, abstract, subject headings, heading word, drug trade name, original title, device manufacturer name] 25. 18 or 19 or 20 or 21 or 22 or 23 or 24 26. 14 and 17 and 25 27. limit 26 to yr="2001 - Current" CINAHL (EBSCO) (1982-2012). S1. (MH "Random Assignment") S2. (MH "Comparative Studies") S3. (MH "Clinical Research+") S4. (MH "Clinical Trials+") S5. (MH "Evaluation Research+") S6. TX ((control* or clinic* or prospective*) adj5 (trial* or study or studies)) S7. TX cross*over* S8. TX (compar* adj5 (trial* or study* or studies)) S9. "random\$" S10. "placebo" S11. "RCT" S12. (S1 or S2 or S3 or S4 or S5 or S6 or S7 or S8 or S9 or S10 or S11)
- S14. "Parkinson*"
- S15. "physiotherapy"
- S16. (MH "Physical Therapy+")

S13. (MH "Parkinson Disease")

- S17. (MH "Exercise+")
- S18. rehabilitation
- S19. S13 or S14
- S20. S15 or S16 or S17 or S18
- S21. S12 and S19 and S20



ISI Web of Science: Science Citation Index Expanded (SCI-EXPANDED) (1981 to January 2012), ISI Web of Science: Conference Proceedings Citation Index-Science (CPCI-S) (1982 to January 2012).

- #1. TS=clinical trial*
- #2. TS=research design
- #3. TS=comparative stud*
- #4. TS=evaluation stud*
- #5. TS=controlled trial*
- #6. TS=follow-up stud*
- #7. TS=prospective stud*
- #8. TS=random*
- #9. TS=placebo*
- #10. TS=(single blind*)
- #11. TS=(double blind*)
- #12. #11 OR #10 OR #9 OR #8 OR #7 OR #6 OR #5 OR #4 OR #3 OR #2 OR #1
- #13. TS=(parkinson disease)
- #14. TS=parkinson*
- #15. #13 OR #14
- #16. TS=physiotherap*
- #17. TS=(physical therap*)
- #18. TS=exercise*
- #19. TS=rehabilitation*
- #20. #19 OR #18 OR #17 OR #16
- #21. #20 AND #15 AND #12

AMED (EBSCO) (1985-2012).

- S1. (MH "Random Assignment")
- S2. (MH "Comparative Studies")
- S3. (MH "Clinical Research+")
- S4. (MH "Clinical Trials+")
- S5. (MH "Evaluation Research+")
- S6. TX ((control* or clinic* or prospective*) adj5 (trial* or study or studies))
- S7. TX cross*over*
- S8. TX (compar* adj5 (trial* or study* or studies))
- S9. "random\$"
- S10. "placebo"
- S11. "RCT"



S12. (S1 or S2 or S3 or S4 or S5 or S6 or S7 or S8 or S9 or S10 or S11)

S13. (MH "Parkinson Disease")

S14. "Parkinson*"

S15. "physiotherapy"

S16. (MH "Physical Therapy+")

S17. (MH "Exercise+")

S18. rehabilitation

S19. S13 or S14

S20. S15 or S16 or S17 or S18

S21. S12 and S19 and S20

REHABDATA (1995-2012). Searched using the term Parkinson*

REHADAT (1990 - 2012). Searched using the term Parkinson.

PEDro (1929-2012). Searched using the term Parkinson.

GEROLIT (1979-2012). Searched using the terms "parkinson*" AND "physi*", "parkinson*" AND "exercise*", "parkinson* AND rehabilitation*"

LILACS (Virtual Health Library) (1982-2012); MedCarib (Virtual Health Library) (17th Century-2012); IMEMR (1984-2012). Searched using the terms "Parkinson\$" AND "physi\$", "Parkinson\$" AND "exercise\$", "Parkinson\$" AND "rehabilitat\$", "Parkinson\$" AND "physiotherap\$".

The Cochrane Controlled Trials Register. See the Cochrane Library.

The CentreWatch Clinical Trials listing service. Screened all entries under the Medical Condition 'Parkinson's Disease'.

The metaRegister of Controlled Trial; NIDRR. Searched using the term Parkinson.

Clinical Trials.gov. Searched using the terms "parkinson" AND "(physical therapy OR physiotherapy OR exercise OR rehabilitation)".

RePORT. Searched using the terms "Parkinson" and "Rehabilitation", "Parkinson" and "Exercise", "Parkinson" and "Physical Therapy", "Parkinson" and "Physiotherapy".

NRR. Searched using the terms "Parkinson" and "Exercise", "Parkinson" and "Exercising", "Parkinson" and "Physiotherapy", "Parkinson" and "Physiotherapies", "Parkinson" and "Physical", "Parkinson" and "rehabilitation", "Parkinson" and "Rehabilitating", "Parkinson" and "Rehabilitate".

Conference Proceedings Citation Index (1982-2012). See ISI Web of Science.

DISSABS (DISSertation ABStracts) (1999-2012). Search using the term "parkinson".

Conference Papers Index (ProQuest) (1982-2012); Index to Theses (1970-2012); ProQuest dissertations and theses databases (1861-2012).

Searched using the terms (all(Physical therapy*) OR all(physiotherap*) OR all(exercise) OR all(rehabilitation*)) AND all(parkinson*).

Electronic Theses Online Service (EThOS) (16th Century-2012). Searched using the terms "parkinson" AND "physical therapy", "parkinson" AND "physiotherapy", "parkinson" AND "exercise", "parkinson" AND "rehabilitation".

WHAT'S NEW



Date	Event	Description
13 May 2014	New citation required but conclusions have not changed	Conclusions not changed
13 May 2014	New search has been performed	Search updated to 31 January 2012
		New studies added, conclusions unchanged

HISTORY

Protocol first published: Issue 4, 2000 Review first published: Issue 1, 2001

Date	Event	Description
29 November 2000	New citation required and conclusions have changed	Substantive amendment

CONTRIBUTIONS OF AUTHORS

Claire Tomlinson was involved in searching and selection of studies, data extraction, analysis and interpretation of the review.

Clare Herd was involved in searching and selection of studies, data extraction, analysis and interpretation of the review.

Smitaa Patel was involved in selection of studies, data extraction.

Charmaine Meek was involved in selection of studies, data extraction and provided expert physiotherapy input into the interpretation of the review.

Carl Clarke contributed to the design of the protocol and was involved in the interpretation of the review providing clinical input.

Rebecca Stowe contributed to the design of the protocol and was involved in searching and selection of studies and interpretation of the review.

Laila Shah was involved in searching and selection of studies for the review.

Catherine Sackley contributed to the design of the protocol and provided expert physiotherapy input into the interpretation of the review.

Katherine Deane undertook the 2001 Cochrane Review, and was involved in the interpretation of this review.

Keith Wheatley contributed to the design of the protocol and was involved in the interpretation of the review.

Natalie Ives contributed to the design of the protocol and was involved in the analysis and interpretation of the review.

DECLARATIONS OF INTEREST

Carl Clarke, Natalie Ives, Charmaine Meek, Smitaa Patel, Catherine Sackley and Keith Wheatley are either recruiting for or involved in the running of the UK PD REHAB trial.

SOURCES OF SUPPORT

Internal sources

• No sources of support supplied

External sources

· Parkinson's UK, UK.



• Department of Health, UK.

INDEX TERMS

Medical Subject Headings (MeSH)

*Physical Therapy Modalities; Gait [physiology]; Parkinson Disease [*rehabilitation]; Randomized Controlled Trials as Topic

MeSH check words

Humans