Regression Models for Epidemiology

Ezra Morrison

Last modified: 2024-06-05: 12:02:43 (PM)

| Pr | eface | 1 |
|----|--|-----|
| | Using these lecture notes | 1 |
| | Other resources | 5 |
| | License | 8 |
| 1. | Introduction | 10 |
| | 1.1. Introduction to Epi 204 | 12 |
| | 1.2. Regression models | |
| | | |
| I. | Generalized Linear Models | 27 |
| 2. | Linear (Gaussian) Models | 31 |
| | 2.1. Overview | 33 |
| | 2.2. Understanding Gaussian Linear Regression Models | 34 |
| | 2.3. Estimating Linear Models via Maximum Likelihood | 52 |
| | 2.4. Inference about Gaussian Linear Regression Models | 59 |
| | 2.5. Goodness of fit | 65 |
| | 2.6. Rescaling | 73 |
| | 2.7. Prediction | 75 |
| | 2.8. Diagnostics | 79 |
| | 2.9. Model selection | 114 |
| | 2.10. Categorical covariates with more than two levels | 122 |
| | 2.11. Ordinal covariates | |
| 3. | Models for Binary Outcomes | 132 |
| | 3.1 Introduction | 134 |

| | 2.0 | D:-1- E-t:t: 1 D1:-t: | 196 |
|-----|-------|--|-------------|
| | 3.2. | Risk Estimation and Prediction | |
| | 3.3. | Comparing Probabilities | |
| | 3.4. | Odds and Odds Ratios | |
| | 3.5. | The logit and expit functions | |
| | 3.6. | Introduction to logistic regression | |
| | 3.7. | Multiple logistic regression | |
| | 3.8. | Fitting logistic regression models | |
| | 3.9. | Model comparisons for logistic models | |
| | | Residual-based diagnostics | |
| | | Other link functions for Bernoulli outcomes | |
| | | Quasibinomial | |
| | 3.13 | Further reading | 240 |
| 4. | Mod | lels for Count Outcomes | 241 |
| ₹. | | nowledgements | |
| | 4.1. | Introduction | |
| | 4.2. | Inference for count regression models | |
| | 4.3. | Prediction | |
| | 4.4. | Diagnostics | |
| | 4.5. | Zero-inflation | |
| | 4.6. | Over-dispersion | |
| | 4.7. | Example: needle-sharing | |
| | 4.1. | Example: needle-sharing | 200 |
| | | | |
| II. | Tir | ne to Event Models | 26 1 |
| 5. | Intro | oduction to Survival Analysis | 263 |
| | 5.1. | Overview | 265 |
| | 5.2. | Time-to-event outcome distributions | |
| | 5.3. | Distribution functions for time-to-event variables | |
| | 5.4. | Parametric Models for Time-to-Event Outcomes | |
| | 5.5. | Nonparametric Survival Analysis | |
| | 5.6. | Example: clinical trial for pediatric acute leukemia | |
| | 5.7. | The Kaplan-Meier Product Limit Estimator | |

| | 5.8. Using | g the survival package in R | 303 |
|----|-------------|--|-----|
| | 5.9. Two-s | sample tests | 308 |
| | 5.10. Exam | nple: Bone Marrow Transplant Data | 312 |
| | 5.11. Nelso | on-Aalen Estimates of Cumulative Hazard and Survival | 321 |
| 6. | Proportion | nal Hazards Models | 325 |
| | 6.1. Intro | duction | 327 |
| | 6.2. Cox's | s Proportional Hazards Model | 329 |
| | 6.3. Cox 1 | Model for the bmt data | 341 |
| | 6.4. Adjus | stment for Ties (optional) | 351 |
| | 6.5. Build | ling Cox Proportional Hazards models | 354 |
| | 6.6. Diagr | nostic graphs for proportional hazards assumption | 357 |
| | 6.7. Predi | ictions and Residuals | 366 |
| | 6.8. Good | lness of Fit using the Cox-Snell Residuals | 375 |
| | 6.9. Mart | ingale Residuals | 377 |
| | | king for Outliers and Influential Observations | |
| | 6.11. Strat | ified survival models | 392 |
| | | e-varying covariates | |
| | 6.13. Recu | rrent Events | 420 |
| | 6.14. Age a | as the time scale | 426 |
| 7. | Parametri | c survival models | 427 |
| | 7.1. Parai | metric Survival Models | 429 |
| | 7.2. Comb | bining left-truncation and interval-censoring | 444 |
| Re | ferences | | 445 |
| Αŗ | pendices | | 451 |
| Α. | Mathemat | tics | 451 |
| | A.1. Elem | entary Algebra | 451 |
| | | onentials and Logarithms | |
| | | vatives | |

| | A.4. Vector Calculus | 458 |
|----|--|-----|
| | A.5. Additional resources | 461 |
| В. | Probability | 463 |
| | B.1. Statistical events | 465 |
| | B.2. Random variables | 465 |
| | B.3. Key probability distributions | 470 |
| | B.4. Characteristics of probability distributions | 479 |
| C. | Estimation | 489 |
| | C.1. Probabilistic models | 489 |
| | C.2. Estimands, estimates, and estimators | 490 |
| | C.3. Accuracy of estimators | 493 |
| D. | Inference | 500 |
| | D.1. Interpretation of Negative Findings | 500 |
| Ε. | Introduction to Maximum Likelihood Inference | 501 |
| | E.1. Overview of maximum likelihood estimation | 503 |
| | E.2. Example: Maximum likelihood for Tropical Cyclones in | |
| | Australia | 515 |
| | E.3. Maximum likelihood inference for univariate Gaussian models | 544 |
| | E.4. Example: hormone therapy study | 549 |
| F. | Common Mistakes | 567 |
| | F.1. Parameters versus random variables | 569 |
| | F.2. Quarto | |
| | F.3. LaTeX | 570 |
| G. | Notation | 571 |
| | G.1. Information matrices | 572 |
| | G.2. The percent sign | |
| н. | Statistical computing in R | 573 |
| | H 1 Functions | 573 |

| | H.2. data.frames and tibbles | 574 |
|----|-------------------------------------|-----|
| | $	ext{H.3. The tidyverse}$ | 574 |
| | H.4. Piping | 575 |
| | H.5. Quarto | 576 |
| | H.6. Packages | 576 |
| | H.7. Git | 577 |
| | H.8. Spatial data science | 577 |
| | H.9. Shiny apps | 577 |
| | H.10.Other resources for learning R | 578 |
| I. | Contributing to rme | 579 |
| | I.1. Style guide | 579 |
| | I.2. Fixing typos | |
| | I.3. Bigger changes | 580 |
| | I.4. Code of Conduct | |
| | I.5. Additional references | |

Preface

This web-book is derived from my lecture slides for Epidemiology 204: "Quantitative Epidemiology III: Statistical Models", at UC Davis.

I have drawn these materials from many sources, including but not limited to:

- David Rocke¹'s materials from the 2021 edition of this course²
- • Hua Zhou³'s materials from the 2020 edition of Biostat 200C at UCLA⁴
- Vittinghoff et al. (2012)
- Dobson and Barnett (2018)
- Harrell (2015)

Using these lecture notes

These lecture notes are available online at https://d-morrison.github.io/rme/. The online notes are searchable and are currently being iteratively updated⁵. A pdf version of the notes is also downloadable from https://d-

¹https://dmrocke.ucdavis.edu/

²https://dmrocke.ucdavis.edu/Class/EPI204-Spring-2021/EPI204-Spring-2021.html

³https://hua-zhou.github.io/

⁴https://ucla-biostat-200c-2020spring.github.io/schedule/schedule.html

⁵see the source file repository for recent changes: https://github.com/d-morrison/rme

morrison.github.io/rme/Regression-Models-for-Epidemiology.pdf, and the source files are available at https://github.com/d-morrison/rme.

Compiling chapters as lecture slide decks

Each chapter's source file can also be compiled as a lecture slide deck, using the _quarto-revealjs.yml⁶ Quarto profile⁷ included in the git repository on Github⁸.

For example, to compile Chapter 3 as a slide deck:

- 1) install quarto⁹
- 2) clone the project repository from Github¹⁰
- 3) Install the project dependencies using devtools:

```
library(devtools) # install from CRAN if needed
devtools::install_deps()
```

4) Render the chapter using the revealjs profile using the following terminal shell command:

```
quarto render logistic-regression.qmd --profile=revealjs
```

You can also render all the chapters listed in the _quarto-revealjs.yml¹¹ Quarto profile¹² as slide decks simultaneously:

⁶https://github.com/d-morrison/rme/blob/main/_quarto-revealjs.yml

⁷https://quarto.org/docs/projects/profiles.html

⁸https://github.com/d-morrison/rme

⁹https://quarto.org/docs/get-started/

¹⁰https://github.com/d-morrison/rme

¹¹https://github.com/d-morrison/rme/blob/main/ quarto-revealjs.yml

¹²https://quarto.org/docs/projects/profiles.html

quarto render --profile=revealjs

Extracting LaTeX commands from the online version of the notes

If you want to extract the LaTeX commands for any math expressions in the online lecture notes, you should be able to right-click and get this pop-up menu:



Figure 1.: Pop-up menu produced by right-clicking on math in online notes

If you select "TeX commands", you will get a window with LaTeX code. 13

¹³MathJax¹⁴ is more or less a dialect of LaTeX



Figure 2.: LaTeX source code window

You can also grab the TeX commands from the quarto source files on github, but those files use custom macros (defined in https://github.com/d-morrison/rme/blob/main/macros.qmd), so it's a little harder to reuse code from the source files.

Dark Mode

The online notes have two color palette themes: light and dark. You can toggle between them using the oval button near the top-left corner:



Figure 3.: Palette toggle

Other resources

These notes represent my still-developing perspective on regression models in epidemiology. Many other statisticians and epidemiologists have published their own perspectives, and I encourage you to explore your many options and find ones that resonate with you. I have attempted to cite my sources throughout these notes.

Here are some additional resources that I've come across; I haven't had time to read some of them thoroughly yet, but they're all on my to-do list. I'll add my thoughts on them over time.

• Dobson and Barnett (2018) is a classic textbook on GLMs. It was used in UCLA Biostatistics's MS-level GLMs course (Biostat 200C) when I took it, and it helped me a lot. It is fairly mathematically

Other resources

rigorous and concise, bordering on terse. It covers GLMs in detail, and survival analysis briefly, and it also has helpful chapters on Bayesian methods. I have adapted examples and explanations from it extensively in these notes.

- Hosmer, Lemeshow, and Sturdivant (2013) is a classic text on logistic regression. I haven't read it yet.
- Agresti (2012) is another classic text for GLMs. I haven't read it yet.
- Agresti (2018) appears to be a more applied version of Agresti (2012). I haven't read it yet. There are extra exercises ¹⁵ and other resources available on the Student Companion Site ¹⁶
- Agresti (2015) has "More than 400 exercises for readers to practice and extend the theory, methods, and data analysis"; might be more theoretical?
- Dunn, Smyth, et al. (2018) is a recent textbook on GLMs. It doesn't cover time-to-event models, and it doesn't use the modern tidyverse¹⁷ packages (ggplot2¹⁸, dplyr¹⁹, etc.), but otherwise it seems great.
- Moore (2016) is a recent textbook on survival analysis. It also doesn't use the tidyverse, but otherwise seems great.
- Klein and Moeschberger (2003) is a classic text for survival analysis. I read most of it in grad school, and it was very helpful. Examples and explanations from it are borrowed extensively in the second half of these notes (partially filtered through David Rocke's course notes.)

¹⁵https://bcs.wiley.com/he-bcs/Books?action=resource&bcsId=11293&itemId= 1119405262&resourceId=44770

¹⁶https://bcs.wiley.com/he-bcs/Books?action=index&itemId=1119405262&bcsId= 11293

¹⁷https://tidyverse.org/

¹⁸https://ggplot2.tidyverse.org/

¹⁹https://dplvr.tidvverse.org/

Other resources

- Kalbfleisch and Prentice (2011) is another classic survival analysis text; I haven't read it yet.
- Kleinbaum and Klein (2010) is a mostly applied-level "self-learning" text for logistic regression; I read it cover-to-cover before grad school, and found it very helpful.
- Kleinbaum and Klein (2012) is the corresponding "self-learning" text for survival analysis; I read it cover-to-cover before grad school, and found it very helpful.
- Harrell (2015) is another popular textbook. It uses ggplot2²⁰ but not dplyr²¹, and covers logistic regression and survival analysis (no Poisson or NB models?). An abbreviated but continuously updated version with audio clips is available at https://hbiostat.org/rmsc/.
- Fox (2015) is another standard text. ²²
- McCullagh and Nelder (1989) is a classic, theoretical textbook on GLMs $^{23}\,$
- Dalgaard (2008) covers GLMs and survival analysis at an applied level, using base R
- Vittinghoff et al. (2012) covers GLMs, survival analysis, and causal inference, using Stata. The authors are UCSF professors, and it is used for the core Epi PhD courses there. I read this book nearly cover-to-cover before grad school, and it was hugely helpful for me, both for statistical modeling and for causal inference (I think it provided my first exposure to DAGs).
- Faraway (2016) has GLMs but not survival analysis

²⁰https://ggplot2.tidyverse.org/

²¹https://dplyr.tidyverse.org/

²²I don't have anything to say about this book, because I haven't opened it yet, but I've heard it's great!

 $^{^{23}}$ haven't opened it either

License

- Selvin (2001) provides worked-out examples of applications for a wide range of statistical analysis techniques. The Author²⁴ is a retired UC Berkeley Biostatistics professor; he used it in a graduate-level biostat/epi course.
- Jewell (2003) is by another UC Berkeley professor²⁵; it mostly covers logistic regression, with one chapter on survival analysis.
- https://ucla-biostat-200c-2020spring.github.io/schedule/schedule.
 html provides course notes for "Biostat 200C Methods in Biostatistics C" at UCLA, which is at the Biostatistics MS level.
- https://online.stat.psu.edu/stat504/book/ provides course notes for "STAT 504 Analysis of Discrete Data" at Penn State University. It includes logistic regression and Poisson regression, as well as 2-way tables and other related topics, and includes SAS code.
- Nahhas (2024) is currently in-development
- Clayton and Hills (2013) covers binary regression, count regression, and survival analysis. Haven't started it yet.
- https://thomaselove.github.io/2020-432-book/index.html is another set of lecture notes.
- Woodward (2013) covers GLMs and survival; haven't read it yet, but it looks comprehensive.

License

This book is licensed to you under Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License²⁶.

²⁴https://publichealth.berkelev.edu/people/steve-selvin

²⁵https://publichealth.berkeley.edu/people/nicholas-jewell

²⁶http://creativecommons.org/licenses/by-nc-nd/4.0/

License

The code samples in this book are licensed under Creative Commons CC0 1.0 Universal (CC0 1.0) 27 , i.e. public domain.

 $[\]overline{^{27} \rm https://creative commons.org/public domain/zero/1.0/$

Configuring R

Functions from these packages will be used throughout this document:

```
library(conflicted) # check for conflicting function definitions
# library(printr) # inserts help-file output into markdown output
library(rmarkdown) # Convert R Markdown documents into a variety of formats.
library(pander) # format tables for markdown
library(ggplot2) # graphics
library(ggeasy) # help with graphics
library(ggfortify) # help with graphics
library(dplyr) # manipulate data
library(tibble) # `tibble`s extend `data.frame`s
library(magrittr) # `%>%` and other additional piping tools
library(haven) # import Stata files
library(knitr) # format R output for markdown
library(tidyr) # Tools to help to create tidy data
library(plotly) # interactive graphics
library(dobson) # datasets from Dobson and Barnett 2018
library(parameters) # format model output tables for markdown
library(haven) # import Stata files
```

```
library(latex2exp) # use LaTeX in R code (for figures and tables)
library(fs) # filesystem path manipulations
library(survival) # survival analysis
library(survminer) # survival analysis graphics
library(KMsurv) # datasets from Klein and Moeschberger
library(parameters) # format model output tables for
library(webshot2) # convert interactive content to static for pdf
library(forcats) # functions for categorical variables ("factors")
library(stringr) # functions for dealing with strings
library(lubridate) # functions for dealing with dates and times
```

Here are some R settings I use in this document:

```
rm(list = ls()) # delete any data that's already loaded into R

conflicts_prefer(dplyr::filter)
ggplot2::theme_set(
    ggplot2::theme_bw() +
        # ggplot2::labs(col = "") +
        ggplot2::theme(
        legend.position = "bottom",
        text = ggplot2::element_text(size = 12, family = "serif")))

knitr::opts_chunk$set(message = FALSE)
options('digits' = 4)

panderOptions("big.mark", ",")
pander::panderOptions("table.emphasize.rownames", FALSE)
pander::panderOptions("table.split.table", Inf)
conflicts_prefer(dplyr::filter) # use the `filter()` function from dplyr() by defaul
legend_text_size = 9
```

1.1. Introduction to Epi 204

Welcome to Epidemiology 204: Quantitative Epidemiology III (Statistical Models).

In this course, we will start where Epi 203 left off: with linear regression models.

Note

Epi 203/STA 130B/STA 131B is a prerequisite for this course. If you haven't passed one of these courses, please talk to me ASAP.

1.1.1. What you should already know

Epi 202: probability models for different data types

- Probability distributions
 - binomial
 - Poisson
 - Gaussian
 - exponential
- Characteristics of probability distributions
 - Mean, median, mode, quantiles
 - Variance, standard deviation, overdispersion
- Characteristics of samples
 - independence, dependence, covariance, correlation
 - ranks
 - identical vs nonidentical distribution (homogeneity vs heterogeneity)
 - Laws of Large Numbers

- Central Limit Theorem for the mean of an iid sample

Epi 203: inference for one or several homogenous populations

- the maximum likelihood inference framework:
 - likelihood functions
 - log-likelihood functions
 - score functions
 - estimating equations
 - information matrices
 - point estimates
 - standard errors
 - confidence intervals
 - hypothesis tests
 - p-values
- Hypothesis tests for one, two, and >2 groups:
 - t-tests/ANOVA for Gaussian models
 - chi-square tests for binomial and Poisson models
 - nonparametric tests:
 - * Wilcoxon signed-rank test for matched pairs
 - * Mann–Whitney/Kruskal-Wallis rank sum test for 2 independent samples
 - * Fisher's exact test for contingency tables
 - * Cochran–Mantel–Haenszel-Cox log-rank test
- Some linear regression

For all of the quantities above, and especially for confidence intervals and p-values, you should know how **both**: - how to compute them - how to interpret them

Stat 108: linear regression models

• building models for Gaussian outcomes

- multiple predictors
- interactions
- regression diagnostics
- fundamentals of R programming; e.g.:
 - Wickham, Çetinkaya-Rundel, and Grolemund (2023)
 - Dalgaard (2008)
- RMarkdown or Quarto for formatting homework¹
 - LaTeX for writing math in RMarkdown/Quarto

1.1.2. What we will cover in this course

- Linear (Gaussian) regression models (review and more details)
- Regression models for non-Gaussian outcomes
 - binary
 - count
 - time to event
- Statistical analysis using R

1.2. Regression models

Why do we need them?

- continuous predictors
- not enough data to analyze some subgroups individually

¹https://r4ds.hadley.nz/quarto

1.2.1. Example: Adelie penguins



Figure 1.1.: Palmer penguins

1.2.2. Linear regression



Figure 1.2.: Palmer penguins with linear regression fit

1.2.3. Curved regression lines

```
ggpenguins2 = ggpenguins +
  stat_smooth(
   method = "lm",
   formula = y ~ log(x),
   geom = "smooth") +
  xlab("Bill length (mm)") +
  ylab("Body mass (g)")
ggpenguins2
```



Figure 1.3.: Palmer penguins - curved regression lines

1.2.4. Multiple regression

```
ggpenguins =
  palmerpenguins::penguins |>
  ggplot(
   aes(x = bill_length_mm ,
        y = body_mass_g,
        color = species
   )
) +
  geom_point() +
  stat_smooth(
  method = "lm",
```

```
formula = y ~ x,
  geom = "smooth") +
  xlab("Bill length (mm)") +
  ylab("Body mass (g)")
ggpenguins |> print()
```



Figure 1.4.: Palmer penguins - multiple groups

1.2.5. Modeling non-Gaussian outcomes

```
library(glmx)
data(BeetleMortality)
beetles = BeetleMortality |>
```

```
mutate(
    pct = died/n,
    survived = n - died
)

plot1 =
    beetles |>
    ggplot(aes(x = dose, y = pct)) +
    geom_point(aes(size = n)) +
    xlab("Dose (log mg/L)") +
    ylab("Mortality rate (%)") +
    scale_y_continuous(labels = scales::percent) +
    # xlab(bquote(log[10]), bquote(CS[2])) +
    scale_size(range = c(1,2))

print(plot1)
```



Figure 1.5.: Mortality rates of adult flour beetles after five hours' exposure to gaseous carbon disulphide (Bliss 1935)

1.2.6. Why don't we use linear regression?

```
beetles_long =
  beetles |>
  reframe(.by = everything(),
      outcome = c(
      rep(1, times = died),
      rep(0, times = survived))
)
lm1 =
```

```
beetles_long |>
lm(
   formula = outcome ~ dose,
   data = _)

range1 = range(beetles$dose) + c(-.2, .2)

f.linear = function(x) predict(lm1, newdata = data.frame(dose = x))

plot2 =
   plot1 +
   geom_function(fun = f.linear, aes(col = "Straight line")) +
   labs(colour="Model", size = "")

print(plot2)
```



Figure 1.6.: Mortality rates of adult flour beetles after five hours' exposure to gaseous carbon disulphide (Bliss 1935)

1.2.7. **Zoom out**




Figure 1.7.: Mortality rates of adult flour beetles after five hours' exposure to gaseous carbon disulphide (Bliss 1935)

1.2.8. log transformation of dose?

```
lm2 =
  beetles_long |>
  lm(formula = outcome ~ log(dose), data = _)

f.linearlog = function(x) predict(lm2, newdata = data.frame(dose = x))

plot3 = plot2 +
```

```
expand_limits(x = c(1.6, 2)) +
geom_function(fun = f.linearlog, aes(col = "Log-transform dose"))
print(plot3 + expand_limits(x = c(1.6, 2)))
```



Figure 1.8.: Mortality rates of adult flour beetles after five hours' exposure to gaseous carbon disulphide (Bliss 1935)

1.2.9. Logistic regression

```
glm1 = beetles |>
   glm(formula = cbind(died, survived) ~ dose, family = "binomial")
```

```
f = function(x) predict(glm1, newdata = data.frame(dose = x), type = "response")
plot4 = plot3 + geom_function(fun = f, aes(col = "Logistic regression"))
print(plot4)
```



Figure 1.9.: Mortality rates of adult flour beetles after five hours' exposure to gaseous carbon disulphide (Bliss 1935)

1.2.10. Three parts to regression models

- What distribution does the outcome have for a specific subpopulation defined by covariates? (**outcome model**)
- How does the combination of covariates relate to the mean? (link function)

• How do the covariates combine? (linear predictor/linear component)

$$\boldsymbol{\eta} \stackrel{\mathrm{def}}{=} \tilde{\boldsymbol{x}}^{\top} \tilde{\boldsymbol{\beta}} = \boldsymbol{\beta}_0 + \boldsymbol{\beta}_1 \boldsymbol{x}_1 + \boldsymbol{\beta}_2 \boldsymbol{x}_2 + \dots$$

Part I. Generalized Linear Models

This section is primarily adapted starting from the textbook "An Introduction to Generalized Linear Models" (4th edition, 2018) by Annette J. Dobson and Adrian G. Barnett:

https://doi.org/10.1201/9781315182780

The type of predictive model one uses depends on several issues; one is the type of response.

- Measured values such as quantity of a protein, age, weight usually can be handled in an ordinary linear regression model, possibly after a log transformation.
- Patient survival, which may be censored, calls for a different method (survival analysis, Cox regression).
- If the response is binary, then can we use logistic regression models
- If the response is a count, we can use Poisson regression
- If the count has a higher variance than is consistent with the Poisson, we can use a negative binomial or over-dispersed Poisson
- Other forms of response can generate other types of generalized linear models

We need a linear predictor of the same form as in linear regression βx . In theory, such a linear predictor can generate any type of number as a prediction, positive, negative, or zero

We choose a suitable distribution for the type of data we are predicting (normal for any number, gamma for positive numbers, binomial for binary responses, Poisson for counts)

We create a link function which maps the mean of the distribution onto the set of all possible linear prediction results, which is the whole real line $(-\infty, \infty)$. The inverse of the link function takes the linear predictor to the actual prediction.

- Ordinary linear regression has identity link (no transformation by the link function) and uses the normal distribution
- If one is predicting an inherently positive quantity, one may want to use the log link since ex is always positive.
- An alternative to using a generalized linear model with a log link, is to transform the data using the log. This is a device that works well with measurement data and may be usable in other cases, but it cannot be used for 0/1 data or for count data that may be 0.

Table 1.1.: R glm() Families

| Family | Links |
|------------------|--|
| gaussian | identity, log, inverse |
| binomial | logit, probit, cauchit, log, cloglog |
| gamma | inverse , identity, log |
| inverse.gaussian | 1/mu^2, inverse, identity, log |
| Poisson | log, identity, sqrt |
| quasi | identity, logit, probit, cloglog, |
| | inverse, \log , $1/mu^2$ and $sqrt$ |
| quasibinomial | logit, probit, identity, cloglog, |
| | inverse, log, 1/mu ² and sqrt |
| quasipoisson | log, identity, logit, probit, cloglog, |
| | inverse, $1/\text{mu}^2$ and sqrt |

Table 1.2.: R glm() Link Functions; $\eta = X\beta = g(\mu)$

| | | | | - 0 / |
|----------|---------------------|---------------------|---|--|
| | | | | Inverse Link |
| Name | Domain | Range | Link Function | Function |
| identity | $y(-\infty,\infty)$ | $(-\infty, \infty)$ | $\eta = \mu$. | $\mu = \eta$ |
| \log | $(0, \infty)$ | $(-\infty, \infty)$ | $\eta = \log \{\mu\}$ | $\mu = \exp\left\{\eta\right\}$ |
| inverse | $(0, \infty)$ | $(0, \infty)$ | $\eta = 1/\mu$ | $\mu = 1/\eta$ |
| logit | (0,1) | $(-\infty, \infty)$ | $\eta =$ | $\mu = \exp\left\{\eta\right\}/(1 +$ |
| | | | $\log\left\{\mu/(1-\mu)\right\}$ | $\exp\left\{\eta ight\}$ |
| probit | (0, 1) | $(-\infty, \infty)$ | $\eta = \Phi^{-1}(\mu)$ | $\mu = \Phi(\eta)$ |
| cloglog | (0, 1) | $(-\infty, \infty)$ | $\eta =$ | $\mu =$ |
| | | | $\log\left\{-\log\left\{1-\mu\right\}\right]$ | $\} 1 - \exp\left\{-\exp\left\{\eta\right\}\right\}$ |
| 1/mu^2 | $2(0,\infty)$ | $(0, \infty)$ | $\eta = 1/\mu^2$ | $\mu = 1/\sqrt{\eta}$ |
| sqrt | $(0,\infty)$ | $(0,\infty)$ | $\eta = \sqrt{\mu}$ | $\mu = \eta^2$ |

Configuring R

Functions from these packages will be used throughout this document:

```
library(conflicted) # check for conflicting function definitions
# library(printr) # inserts help-file output into markdown output
library(rmarkdown) # Convert R Markdown documents into a variety of formats.
library(pander) # format tables for markdown
library(ggplot2) # graphics
library(ggeasy) # help with graphics
library(ggfortify) # help with graphics
library(dplyr) # manipulate data
library(tibble) # `tibble`s extend `data.frame`s
library(magrittr) # `%>%` and other additional piping tools
library(haven) # import Stata files
library(knitr) # format R output for markdown
library(tidyr) # Tools to help to create tidy data
library(plotly) # interactive graphics
library(dobson) # datasets from Dobson and Barnett 2018
library(parameters) # format model output tables for markdown
library(haven) # import Stata files
```

```
library(latex2exp) # use LaTeX in R code (for figures and tables)
library(fs) # filesystem path manipulations
library(survival) # survival analysis
library(survminer) # survival analysis graphics
library(KMsurv) # datasets from Klein and Moeschberger
library(parameters) # format model output tables for
library(webshot2) # convert interactive content to static for pdf
library(forcats) # functions for categorical variables ("factors")
library(stringr) # functions for dealing with strings
library(lubridate) # functions for dealing with dates and times
```

Here are some R settings I use in this document:

Note

This content is adapted from:

- Dobson and Barnett (2018), Chapters 2-6
- Dunn, Smyth, et al. (2018), Chapters 2-3
- Vittinghoff et al. (2012), Chapter 4

There are numerous textbooks specifically for linear regression, including:

- Kutner et al. (2005): used for UCLA Biostatistics MS level linear models class
- Chatterjee and Hadi (2015): used for Stanford MS-level linear models class
- Seber and Lee (2012): used for UCLA Biostatistics PhD level linear models class
- Kleinbaum et al. (2014): same first author as Kleinbaum and Klein (2010) and Kleinbaum and Klein (2012)

2.1. Overview

2.1.1. Why this course includes linear regression

- This course is about *generalized linear models* (for non-Gaussian outcomes)
- UC Davis STA 108 ("Applied Statistical Methods: Regression Analysis") is a prerequisite for this course, so everyone here should have some understanding of linear regression already.
- We will review linear regression to:

- make sure everyone is caught up
- to provide an epidemiological perspective on model interretation.

2.1.2. Chapter overview

- Section 2.2: how to interpret linear regression models
- Section 2.3: how to estimate linear regression models
- Section 2.4: how to quantify uncertainty about our estimates
- Section 2.8: how to tell if your model is insufficiently complex

2.2. Understanding Gaussian Linear Regression Models

2.2.1. Motivating example: birthweights and gestational age

Suppose we want to learn about the distributions of birthweights ($outcome\ Y$) for (human) babies born at different gestational ages ($covariate\ A$) and with different chromosomal sexes ($covariate\ S$) (Dobson and Barnett (2018) Example 2.2.2).

2.2.1.1. Data as table

```
library(dobson)
data("birthweight", package = "dobson")
birthweight |> knitr::kable()
```

Table 2.1.: birthweight data (Dobson and Barnett (2018) Example 2.2.2)

| boys gestational age | boys weight | girls gestational age | girls weight |
|----------------------|-------------|-----------------------|--------------|
| 40 | 2968 | 40 | 3317 |
| 38 | 2795 | 36 | 2729 |
| 40 | 3163 | 40 | 2935 |
| 35 | 2925 | 38 | 2754 |
| 36 | 2625 | 42 | 3210 |
| 37 | 2847 | 39 | 2817 |
| 41 | 3292 | 40 | 3126 |
| 40 | 3473 | 37 | 2539 |
| 37 | 2628 | 36 | 2412 |
| 38 | 3176 | 38 | 2991 |
| 40 | 3421 | 39 | 2875 |
| 38 | 2975 | 40 | 3231 |

2.2.1.2. Reshape data for graphing

```
bw =
  birthweight |>
  pivot_longer(
    cols = everything(),
    names_to = c("sex", ".value"),
    names_sep = "s "
) |>
  rename(age = `gestational age`) |>
  mutate(
    sex = sex |>
    case_match(
        "boy" ~ "male",
        "girl" ~ "female") |>
    factor(levels = c("female", "male")))
```

bw

Table 2.2.: birthweight data reshaped

| sex | age | weight |
|--------|-----|--------|
| male | 40 | 2968 |
| female | 40 | 3317 |
| male | 38 | 2795 |
| female | 36 | 2729 |
| male | 40 | 3163 |
| female | 40 | 2935 |
| male | 35 | 2925 |
| female | 38 | 2754 |
| male | 36 | 2625 |
| female | 42 | 3210 |
| male | 37 | 2847 |
| female | 39 | 2817 |
| male | 41 | 3292 |
| female | 40 | 3126 |
| male | 40 | 3473 |
| female | 37 | 2539 |
| male | 37 | 2628 |
| female | 36 | 2412 |
| male | 38 | 3176 |
| female | 38 | 2991 |
| male | 40 | 3421 |
| female | 39 | 2875 |
| male | 38 | 2975 |
| female | 40 | 3231 |
| | | |

2.2.1.3. Data as graph

```
plot1 = bw |>
    ggplot(aes(
        x = age,
        y = weight,
        linetype = sex,
        shape = sex,
        col = sex)) +
    theme_bw() +
    xlab("Gestational age (weeks)") +
    ylab("Birthweight (grams)") +
    theme(legend.position = "bottom") +
    # expand_limits(y = 0, x = 0) +
    geom_point(alpha = .7)
print(plot1 + facet_wrap(~ sex))
```



Figure 2.1.: birthweight data (Dobson and Barnett (2018) Example 2.2.2)

2.2.1.4. Data notation

Let's define some notation to represent this data.

- Y: birthweight (measured in grams)
- S: chromosomal sex: "male" (XY) or "female" (XX)

- M: indicator variable for $S = \text{``male''}^1$
- M = 0 if female (XX)
- M = 1 if male (XY)
- F: indicator variable for S = "female"²
- F = 1 if female (XX)
- F = 0 if male (XY)
- A: estimated gestational age at birth (measured in weeks).

Note

Female is the **reference level** for the categorical variable S (chromosomal sex) and corresponding indicator variable M. The choice of a reference level is arbitrary and does not limit what we can do with the resulting model; it only makes it more computationally convenient to make inferences about comparisons involving that reference group.

2.2.2. Parallel lines regression

We don't have enough data to model the distribution of birth weight separately for each combination of gestational age and sex, so let's instead consider a (relatively) simple model for how that distribution varies with gestational age and sex:

$$p(Y = y|A = a, S = s) \sim_{\text{iid}} N(\mu(a, s), \sigma^2)$$

 $^{^{1}}M$ is implicitly a deterministic function of S

 $^{^{2}}F$ is implicitly a deterministic function of S

$$\mu(a,s) \stackrel{\text{def}}{=} \mathbb{E}\left[Y|A=a,S=s\right]$$
$$= \beta_0 + \beta_A a + \beta_M m$$
 (2.1)

Table 2.3 shows the parameter estimates from R. Figure 2.2 shows the estimated model, superimposed on the data.

```
bw_lm1 = lm(
  formula = weight ~ sex + age,
  data = bw)

bw_lm1 |>
  parameters() |>
  print_md(
    include_reference = TRUE,
    # show_sigma = TRUE,
    select = "{estimate}")
```

Table 2.3.: Estimate of Model 2.1 for birthweight data

| Parameter | Estimate |
|--------------|----------|
| (Intercept) | -1773.32 |
| sex (female) | 0.00 |
| sex (male) | 163.04 |
| age | 120.89 |

```
bw =
  bw |>
  mutate(`E[Y|X=x]` = fitted(bw_lm1)) |>
  arrange(sex, age)

plot2 =
```

```
plot1 %+% bw +
geom_line(aes(y = `E[Y|X=x]`))
print(plot2)
```



Figure 2.2.: Parallel-slopes model of birthweight

2.2.2.1. Model assumptions and predictions

To learn what this model is assuming, let's plug in a few values.

Exercise 2.1. According to this model, what's the mean birthweight for a female born at 36 weeks?

Table 2.4.: Estimated coefficients for model 2.1

Solution.

$$\begin{split} E[Y|A=0, A=36] &= \beta_0 + \beta_M \cdot 0 + \beta_A \cdot 36 \\ &= 2578.8739 \end{split}$$

Exercise 2.2. What's the mean birthweight for a male born at 36 weeks?

```
coef(bw_lm1)
#> (Intercept) sexmale age
#> -1773.3 163.0 120.9
```

Solution.

$$\begin{split} E[Y|M=1, A=36] &= \beta_0 + \beta_M \cdot 1 + \beta_A \cdot 36 \\ &= 2741.9132 \end{split}$$

Exercise 2.3. What's the difference in mean birthweights between males born at 36 weeks and females born at 36 weeks?

```
coef(bw_lm1)
#> (Intercept) sexmale age
#> -1773.3 163.0 120.9
```

Solution.

$$E[Y|M=1, A=36] - E[Y|M=0, A=36]$$

= 2741.9132 - 2578.8739
= 163.0393

Shortcut:

$$\begin{split} E[Y|M=1,A=36] - E[Y|M=0,A=36] \\ &= (\beta_0 + \beta_M \cdot 1 + \beta_A \cdot 36) - (\beta_0 + \beta_M \cdot 0 + \beta_A \cdot 36) \\ &= \beta_M \\ &= 163.0393 \end{split}$$

Note that age doesn't show up in this difference: in other words, according to this model, the difference between females and males with the same gestational age is the same for every age.

That's an assumption of the model; it's built-in to the parametric structure, even before we plug in the estimated values of those parameters.

That's why the lines are parallel.

2.2.3. Interactions

What if we don't like that parallel lines assumption?

Then we need to allow an "interaction" between age A and sex S:

$$E[Y|A=a,S=s] = \beta_0 + \beta_A a + \beta_M m + \beta_{AM}(a \cdot m) \tag{2.2}$$

Now, the slope of mean birthweight E[Y|A,S] with respect to gestational age A depends on the value of sex S.

```
bw_lm2 = lm(weight ~ sex + age + sex:age, data = bw)
bw_lm2 |>
  parameters() |>
  print_md(
    include_reference = TRUE,
    # show_sigma = TRUE,
    select = "{estimate}")
```

Table 2.5.: Birthweight model with interaction term

| Parameter | Estimate |
|---|----------|
| (Intercept) | -2141.67 |
| sex (female) | 0.00 |
| sex (male) | 872.99 |
| age | 130.40 |
| $\mathrm{sex}\ (\mathrm{male})\times\mathrm{age}$ | -18.42 |

```
bw =
  bw |>
  mutate(
    predlm2 = predict(bw_lm2)
) |>
  arrange(sex, age)

plot1_interact =
  plot1 %+% bw +
  geom_line(aes(y = predlm2))

print(plot1_interact)
```



Figure 2.3.: Birthweight model with interaction term

Now we can see that the lines aren't parallel.

Here's another way we could rewrite this model (by collecting terms involving S):

$$E[Y|A,M] = \beta_0 + \beta_M M + (\beta_A + \beta_{AM} M) A$$

Note

If you want to understand a coefficient in a model with interactions, collect terms for the corresponding variable, and you will see what other variables are interacting with the variable you are interested in.

In this case, the coefficient S is interacting with A. So the slope of Y with respect to A depends on the value of M.

According to this model, there is no such thing as "the slope of birthweight with respect to age". There are two slopes, one for each sex.³ We can only talk about "the slope of birthweight with respect to age among males" and "the slope of birthweight with respect to age among females".

Then: that coefficient is the difference in means per unit change in its corresponding coefficient, when the other collected variables are set to 0.

To learn what this model is assuming, let's plug in a few values.

Exercise 2.4. According to this model, what's the mean birthweight for a female born at 36 weeks?

Solution.

³using the definite article "the" would mean there is only one slope.

$$E[Y|A=0,X_2=36] = \beta_0 + \beta_M \cdot 0 + \beta_A \cdot 36 + \beta_{AM} \cdot (0*36) = 2552.7333$$

Exercise 2.5. What's the mean birthweight for a male born at 36 weeks?

Solution.

$$\begin{split} E[Y|A=0, X_2=36] &= \beta_0 + \beta_M \cdot 1 + \beta_A \cdot 36 + \beta_{AM} \cdot 1 \cdot 36 \\ &= 2762.7069 \end{split}$$

Exercise 2.6. What's the difference in mean birthweights between males born at 36 weeks and females born at 36 weeks?

48

Solution.

$$\begin{split} E[Y|M=1,A=36] - E[Y|M=0,A=36] \\ &= (\beta_0 + \beta_M \cdot 1 + \beta_A \cdot 36 + \beta_{AM} \cdot 1 \cdot 36) \\ &- (\beta_0 + \beta_M \cdot 0 + \beta_A \cdot 36 + \beta_{AM} \cdot 0 \cdot 36) \\ &= \beta_S + \beta_{AM} \cdot 36 \\ &= 209.9736 \end{split}$$

Note that age now does show up in the difference: in other words, according to this model, the difference in mean birthweights between females and males with the same gestational age can vary by gestational age.

That's how the lines in the graph ended up non-parallel.

2.2.4. Stratified regression

We could re-write the interaction model as a stratified model, with a slope and intercept for each sex:

$$\mathbb{E}[Y|A=a,S=s] = \beta_M m + \beta_{AM}(a \cdot m) + \beta_F f + \beta_{AF}(a \cdot f) \qquad (2.3)$$

Compare this stratified model with our interaction model, Equation 2.2:

$$\mathbb{E}[Y|A=a,S=s] = \beta_0 + \beta_A a + \beta_M m + \beta_{AM}(a \cdot m)$$

In the stratified model, the intercept term β_0 has been relabeled as β_F .

```
bw_lm2 = lm(weight ~ sex + age + sex:age, data = bw)
bw_lm2 |>
  parameters() |>
  print_md(
   include_reference = TRUE,
```

```
# show_sigma = TRUE,
select = "{estimate}")
```

Table 2.6.: Birthweight model with interaction term

| Parameter | Estimate |
|-------------------------|----------|
| (Intercept) | -2141.67 |
| sex (female) | 0.00 |
| sex (male) | 872.99 |
| age | 130.40 |
| $sex (male) \times age$ | -18.42 |

```
bw_lm_strat =
  bw |>
  lm(
    formula = weight ~ sex + sex:age - 1,
    data = _)

bw_lm_strat |>
  parameters() |>
  print_md(
    # show_sigma = TRUE,
    select = "{estimate}")
```

Table 2.7.: Birthweight model - stratified betas

| Parameter | Estimate |
|---------------------------|----------|
| sex (female) | -2141.67 |
| sex (male) | -1268.67 |
| $sex (female) \times age$ | 130.40 |
| $sex (male) \times age$ | 111.98 |

Table 2.7.: Birthweight model - stratified betas

| Parameter | Estimate |
|-----------|----------|
| | |

2.2.5. Curved-line regression

If we transform some of our covariates (Xs) and plot the resulting model on the original covariate scale, we end up with curved regression lines:

```
bw_lm3 = lm(weight ~ sex:log(age) - 1, data = bw)
library(palmerpenguins)
ggpenguins <-
 palmerpenguins::penguins |>
  dplyr::filter(species == "Adelie") |>
  ggplot(
    aes(x = bill_length_mm , y = body_mass_g)) +
  geom_point() +
  xlab("Bill length (mm)") +
 ylab("Body mass (g)")
ggpenguins2 = ggpenguins +
  stat_smooth(
   method = "lm",
    formula = y \sim log(x),
    geom = "smooth") +
  xlab("Bill length (mm)") +
  ylab("Body mass (g)")
ggpenguins2 |> print()
```



Figure 2.4.: palmerpenguins model with bill_length entering on log scale

2.3. Estimating Linear Models via Maximum Likelihood

2.3.1. Likelihood, log-likelihood, and score functions for linear regression

In EPI 203 and Appendix E, we learned how to fit outcome-only models of the form $p(X=x|\theta)$ to iid data $\tilde{x}=(x_1,...,x_n)$ using maximum likelihood estimation.

Now, we apply the same procedure to linear regression models:

$$\mathcal{L}(\tilde{y}|\mathbf{x}, \beta, \sigma^2) = \prod_{i=1}^{n} (2\pi\sigma^2)^{-1/2} \exp\left\{-\frac{1}{2\sigma^2} (y_i - \tilde{x_i}'\beta)^2\right\}$$
(2.4)

$$\ell(\tilde{\boldsymbol{y}}|\mathbf{x},\boldsymbol{\beta},\sigma^2) = -\frac{n}{2}\log\left\{\sigma^2\right\} - \frac{1}{2\sigma^2}\sum_{i=1}^{n}(y_i - \tilde{\boldsymbol{x_i}}'\boldsymbol{\beta})^2 \tag{2.5}$$

$$\ell_{\beta}'(\tilde{\boldsymbol{y}}|\mathbf{x},\beta,\sigma^2) = -\frac{1}{2\sigma^2} \frac{\partial}{\partial \beta} \left(\sum_{i=1}^n (y_i - \tilde{\boldsymbol{x}_i}^\top \beta)^2 \right) \tag{2.6}$$

Let's switch to matrix-vector notation:

$$\sum_{i=1}^n (y_i - \tilde{x}_i^\top \tilde{\beta})^2 = (\tilde{y} - \mathbf{X} \tilde{\beta})' (\tilde{y} - \mathbf{X} \tilde{\beta})$$

So

$$\begin{split} (\tilde{y} - \mathbf{X}\tilde{\beta})'(\tilde{y} - \mathbf{X}\tilde{\beta}) &= (\tilde{y}' - \tilde{\beta}'X')(\tilde{y} - \mathbf{X}\tilde{\beta}) \\ &= y'y - \tilde{\beta}'X'y - y'\mathbf{X}\tilde{\beta} + \tilde{\beta}'\mathbf{X}'\mathbf{X}\beta \\ &= y'y - 2y'\mathbf{X}\beta + \beta'\mathbf{X}'\mathbf{X}\beta \end{split}$$

2.3.2. Deriving the linear regression score function

We will use some results from vector calculus:

$$\begin{split} \frac{\partial}{\partial \beta} \left(\sum_{i=1}^{n} (y_i - x_i'\beta)^2 \right) &= \frac{\partial}{\partial \beta} (\tilde{y} - X\beta)' (\tilde{y} - X\beta) \\ &= \frac{\partial}{\partial \beta} (y'y - 2y'X\beta + \beta'X'X\beta) \\ &= (-2X'y + 2X'X\beta) \\ &= -2X'(y - X\beta) \\ &= -2X'(y - \mathbb{E}[y]) \\ &= -2X'\varepsilon(y) \end{split} \tag{2.7}$$

So if $\ell(\beta, \sigma^2) = 0$, then

$$0 = (-2X'y + 2X'X\beta)$$

$$2X'y = 2X'X\beta$$

$$X'y = X'X\beta$$

$$(X'X)^{-1}X'y = \beta$$

The second derivative matrix $\ell''_{\beta,\beta'}(\beta,\sigma^2;\mathbf{X},\tilde{y})$ is negative definite at $\beta=(X'X)^{-1}X'y$, so $\hat{\beta}_{ML}=(X'X)^{-1}X'y$ is the MLE for β .

54

Similarly (not shown):

$$\hat{\sigma}_{ML}^2 = \frac{1}{n} (Y - X \hat{\beta})' (Y - X \hat{\beta})$$

And

$$\begin{split} \mathcal{I}_{\beta} &= E[-\ell_{\beta,\beta'}'(Y|X,\beta,\sigma^2)] \\ &= \frac{1}{\sigma^2} X' X \end{split}$$

So:

$$Var(\hat{\beta}) \approx (\mathcal{I}_{\beta})^{-1} = \sigma^2 (X'X)^{-1}$$

and

$$\hat{\beta}\dot{\sim}N(\beta,\mathcal{I}_{\beta}^{-1})$$

These are all results you have hopefully seen before.

In the Gaussian linear regression case, we also have exact results:

$$\frac{\hat{\beta}_j}{\hat{\mathrm{se}}\left(\hat{\beta}_j\right)} \; \sim \; t_{n-p}$$

In our model 2 above, $\hat{\mathcal{I}}(\beta)$ is:

bw_lm2 |> parameters() |> print_md()

```
bw_lm2 |> vcov()
#>
                (Intercept)
                             sexmale
                                           age sexmale:age
#> (Intercept)
                    1353968 -1353968 -34871.0
                                                    34871.0
#> sexmale
                   -1353968
                             2596387
                                       34871.0
                                                   -67211.0
                                                     -899.9
#> age
                     -34871
                                34871
                                         899.9
#> sexmale:age
                      34871
                               -67211
                                        -899.9
                                                     1743.5
```

If we take the square roots of the diagonals, we get the standard errors listed in the model output:

```
bw_lm2 |> vcov() |> diag() |> sqrt()
#> (Intercept) sexmale age sexmale:age
#> 1163.60 1611.33 30.00 41.76
```

Table 2.8.: Estimated model for birthweight data with interaction term

| Parameter | Coefficient | SE | 95% CI | t(20) | p |
|---|-------------|---------|------------------------|-------|--------|
| (Intercept) | -2141.67 | 1163.60 | (-4568.90, 285.56) | -1.84 | 0.081 |
| sex (male) | 872.99 | 1611.33 | (-2488.18, 4234.17) | 0.54 | 0.594 |
| age | 130.40 | 30.00 | (67.82, 192.98) | 4.35 | < .001 |
| $\begin{array}{c} \text{sex (male)} \times \\ \text{age} \end{array}$ | -18.42 | 41.76 | (-105.52, 68.68) | -0.44 | 0.664 |

So we can do confidence intervals, hypothesis tests, and p-values exactly as in the one-variable case we looked at previously.

2.3.3. Residual Standard Deviation

 $\hat{\sigma}$ represents an *estimate* of the *Residual Standard Deviation* parameter, σ . We can extract $\hat{\sigma}$ from the fitted model, using the sigma() function:

```
sigma(bw_lm2)
#> [1] 180.6
```

2.3.3.1. σ is NOT "Residual standard error"

In the summary.lm() output, this estimate is labeled as "Residual standard error":

```
summary(bw_lm2)
#>
#> Call:
#> lm(formula = weight ~ sex + age + sex:age, data = bw)
#> Residuals:
     Min
          1Q Median
                          3Q
                                Max
#> -246.7 -138.1 -39.1 176.6 274.3
#>
#> Coefficients:
              Estimate Std. Error t value Pr(>|t|)
#> (Intercept) -2141.7
                        1163.6
                                  -1.84 0.08057 .
#> sexmale
               873.0
                          1611.3
                                 0.54 0.59395
                130.4
                                   4.35 0.00031 ***
#> age
                            30.0
#> sexmale:age
              -18.4
                            41.8
                                  -0.44 0.66389
#> ---
#> Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
```

```
#>
#> Residual standard error: 181 on 20 degrees of freedom
#> Multiple R-squared: 0.643, Adjusted R-squared: 0.59
#> F-statistic: 12 on 3 and 20 DF, p-value: 0.000101
```

However, this is a misnomer:

```
library(printr) # captures ? documentation
?stats::sigma
#> Extract Residual Standard Deviation 'Sigma'
#>
#> Description:
#>
#>
        Extract the estimated standard deviation of the errors, the
#>
        "residual standard deviation" (misnamed also "residual standard
        error", e.g., in 'summary.lm()''s output, from a fitted model).
#>
#>
#>
        Many classical statistical models have a _scale parameter_,
#>
        typically the standard deviation of a zero-mean normal (or
        Gaussian) random variable which is denoted as sigma. 'sigma(.)'
#>
#>
        extracts the _estimated_ parameter from a fitted model, i.e.,
#>
        sigma^.
#>
#> Note:
#>
#>
        The misnomer "Residual standard *error*" has been part of too many
        R (and S) outputs to be easily changed there.
#>
```

2.4. Inference about Gaussian Linear Regression Models

2.4.1. Motivating example: birthweight data

Research question: is there really an interaction between sex and age?

$$\begin{split} H_0: \beta_{AM} &= 0 \\ H_A: \beta_{AM} &\neq 0 \\ P(|\hat{\beta}_{AM}| > |-18.4172| \mid H_0) &= ? \end{split}$$

2.4.2. Wald tests and CIs

R can give you Wald tests for single coefficients and corresponding CIs:

```
bw_lm2 |>
  parameters() |>
  print_md(
   include_reference = TRUE)
```

| Parameter | Coefficient | SE | 95% CI | t(20) | p |
|---|-------------|---------|------------------------|-------|--------|
| (Intercept) | -2141.67 | 1163.60 | (-4568.90, 285.56) | -1.84 | 0.081 |
| sex (female) | 0.00 | | , | | |
| sex (male) | 872.99 | 1611.33 | (-2488.18, 4234.17) | 0.54 | 0.594 |
| age | 130.40 | 30.00 | (67.82, 192.98) | 4.35 | < .001 |
| $\begin{array}{l} \text{sex (male)} \times \\ \text{age} \end{array}$ | -18.42 | 41.76 | (-105.52, 68.68) | -0.44 | 0.664 |

To understand what's happening, let's replicate these results by hand for the interaction term.

2.4.3. P-values

```
bw_lm2 |>
  parameters(keep = "sexmale:age") |>
  print_md(
   include_reference = TRUE)
```

| Parameter | Coefficient | SE | 95% CI | t(20) | p |
|--------------|-------------|-------|------------------|-------|-------|
| sex (male) × | -18.42 | 41.76 | (-105.52, 68.68) | -0.44 | 0.664 |
| age | | | | | |

```
beta_hat = coef(summary(bw_lm2))["sexmale:age", "Estimate"]
se_hat = coef(summary(bw_lm2))["sexmale:age", "Std. Error"]
dfresid = bw_lm2$df.residual
t_stat = abs(beta_hat)/se_hat
pval_t =
   pt(-t_stat, df = dfresid, lower.tail = TRUE) +
   pt(t_stat, df = dfresid, lower.tail = FALSE)
```

$$\begin{split} &P\left(|\hat{\beta}_{AM}| > |-18.4172| \Big| H_0\right) \\ &= \Pr\left(\left|\frac{\hat{\beta}_{AM}}{\hat{SE}(\hat{\beta}_{AM})}\right| > \left|\frac{-18.4172}{41.7558}\right| \middle| H_0\right) \\ &= \Pr\left(|T_{20}| > 0.4411| H_0\right) \\ &= 0.6639 \end{split}$$

This matches the result in the table above.

2.4.4. Confidence intervals

```
bw_lm2 |>
  parameters(keep = "sexmale:age") |>
  print_md(
   include_reference = TRUE)
```

| Parameter | Coefficient | SE | 95% CI | t(20) | р |
|--------------|-------------|-------|------------------|-------|-------|
| sex (male) × | -18.42 | 41.76 | (-105.52, 68.68) | -0.44 | 0.664 |
| age | | | | | |

```
q_t = qt(
  p = 0.975,
  df = dfresid,
  lower.tail = TRUE)

q_t = qt(
  p = 0.025,
  df = dfresid,
  lower.tail = TRUE)

confint_radius_t =
  se_hat * q_t

confint_t = beta_hat + c(-1,1) * confint_radius_t

print(confint_t)
#> [1] 68.68 -105.52
```

This also matches.

2.4.5. Gaussian approximations

Here are the asymptotic (Gaussian approximation) equivalents:

2.4.6. P-values

```
bw_lm2 |>
  parameters(keep = "sexmale:age") |>
  print_md(
   include_reference = TRUE)
```

| Parameter | Coefficient | SE | 95% CI | t(20) | p |
|---------------------|-------------|-------|------------------|-------|-------|
| $sex (male) \times$ | -18.42 | 41.76 | (-105.52, 68.68) | -0.44 | 0.664 |
| age | | | | | |

```
pval_z = pnorm(abs(t_stat), lower = FALSE) * 2
print(pval_z)
#> [1] 0.6592
```

2.4.7. Confidence intervals

```
bw_lm2 |>
  parameters(keep = "sexmale:age") |>
  print_md(
   include_reference = TRUE)
```

| Parameter | Coefficient | SE | 95% CI | t(20) | р |
|--------------|-------------|-------|------------------|-------|-------|
| sex (male) × | -18.42 | 41.76 | (-105.52, 68.68) | -0.44 | 0.664 |
| age | | | | | |

```
confint_radius_z = se_hat * qnorm(0.975, lower = TRUE)
confint_z =
  beta_hat + c(-1,1) * confint_radius_z
print(confint_z)
#> [1] -100.26 63.42
```

2.4.8. Likelihood ratio statistics

```
logLik(bw_lm2)
#> 'log Lik.' -156.6 (df=5)
logLik(bw_lm1)
#> 'log Lik.' -156.7 (df=4)

LR = (logLik(bw_lm2) - logLik(bw_lm1)) |> as.numeric()
delta_df = (bw_lm1$df.residual - df.residual(bw_lm2))

x_max = 1
```

```
d_lLR = function(x, df = delta_df) dchisq(x, df = df)

chisq_plot =
    ggplot() +
```

```
geom_function(fun = d_llR) +
stat_function( fun = d_llR, xlim = c(llR, x_max), geom = "area", fill = "gray") +
geom_segment(aes(x = llR, xend = llR, y = 0, yend = d_llR(llR)), col = "red") +
xlim(0.0001,x_max) +
ylim(0,4) +
ylab("p(X=x)") +
xlab("log(likelihood ratio) statistic [x]") +
theme_classic()
chisq_plot |> print()
```



Figure 2.5.: Chi-square distribution

Now we can get the p-value:

```
pchisq(
    q = 2*lLR,
    df = delta_df,
    lower = FALSE) |>
    print()
#> [1] 0.6298
```

In practice you don't have to do this by hand; there are functions to do it for you:

```
# built in
library(lmtest)
lrtest(bw_lm2, bw_lm1)
```

| #Df | LogLik | Df | Chisq | Pr(>Chisq) |
|-----|--------|----|--------|------------|
| 5 | -156.6 | NA | NA | NA |
| 4 | -156.7 | -1 | 0.2323 | 0.6298 |

2.5. Goodness of fit

2.5.1. AIC and BIC

When we use likelihood ratio tests, we are comparing how well different models fit the data.

Likelihood ratio tests require "nested" models: one must be a special case of the other.

If we have non-nested models, we can instead use the Akaike Information Criterion (AIC) or Bayesian Information Criterion (BIC):

- AIC = $-2 * \ell(\hat{\theta}) + 2 * p$
- BIC = $-2 * \ell(\hat{\theta}) + p * \log(n)$

where ℓ is the log-likelihood of the data evaluated using the parameter estimates $\hat{\theta}$, p is the number of estimated parameters in the model (including $\hat{\sigma}^2$), and n is the number of observations.

You can calculate these criteria using the logLik() function, or use the built-in R functions:

2.5.1.1. AIC in R

```
-2 * logLik(bw_lm2) |> as.numeric() +
    2*(length(coef(bw_lm2))+1) # sigma counts as a parameter here
#> [1] 323.2

AIC(bw_lm2)
#> [1] 323.2
```

2.5.1.2. BIC in R

```
-2 * logLik(bw_lm2) |> as.numeric() +
  (length(coef(bw_lm2))+1) * log(nobs(bw_lm2))
#> [1] 329

BIC(bw_lm2)
#> [1] 329
```

Large values of AIC and BIC are worse than small values. There are no hypothesis tests or p-values associated with these criteria.

2.5.2. (Residual) Deviance

Let q be the number of distinct covariate combinations in a data set.

```
bw.X.unique =
  bw |>
  count(sex, age)

n_unique.bw = nrow(bw.X.unique)
```

For example, in the birthweight data, there are q=12 unique patterns (Table 2.15).

```
bw.X.unique
```

Table 2.15.: Unique covariate combinations in the birthweight data, with replicate counts

| sex | age | n |
|--------|-----|---|
| female | 36 | 2 |
| female | 37 | 1 |
| female | 38 | 2 |
| female | 39 | 2 |
| female | 40 | 4 |
| female | 42 | 1 |
| male | 35 | 1 |
| male | 36 | 1 |
| male | 37 | 2 |
| | | |

Table 2.15.: Unique covariate combinations in the birthweight data, with replicate counts

| sex | age | n |
|------|-----|---|
| male | 38 | 3 |
| male | 40 | 4 |
| male | 41 | 1 |

Definition 2.1 (Replicates). If a given covariate pattern has more than one observation in a dataset, those observations are called **replicates**.

Example 2.1 (Replicates in the birthweight data). In the birthweight dataset, there are 2 replicates of the combination "female, age 36" (Table 2.15).

Exercise 2.7 (Replicates in the birthweight data). Which covariate pattern(s) in the birthweight data has the most replicates?

Solution 2.1 (Replicates in the birthweight data). Two covariate patterns are tied for most replicates: males at age 40 weeks and females at age 40 weeks. 40 weeks is the usual length for human pregnancy (Polin, Fox, and Abman (2011)), so this result makes sense.

```
bw.X.unique |> dplyr::filter(n == max(n))
```

| sex | age | n |
|--------|-----|---|
| female | 40 | 4 |
| male | 40 | 4 |

2.5.2.1. Saturated models

The most complicated model we could fit would have one parameter (a mean) for each covariate pattern, plus a variance parameter:

```
lm_max =
  bw |>
  mutate(age = factor(age)) |>
  lm(
    formula = weight ~ sex:age - 1,
    data = _)

lm_max |>
  parameters() |>
  print_md()
```

Table 2.17.: Saturated model for the birthweight data

| Parameter | Coefficient | SE | 95% CI | t(12) | p |
|-----------------------|-------------|--------|-----------|-------|------|
| sex (male) × | 2925.00 | 187.92 | (2515.55, | 15.56 | < |
| age35 | | | 3334.45) | | .001 |
| sex (female) \times | 2570.50 | 132.88 | (2280.98, | 19.34 | < |
| age36 | | | 2860.02) | | .001 |
| $sex (male) \times$ | 2625.00 | 187.92 | (2215.55, | 13.97 | < |
| age36 | | | 3034.45) | | .001 |
| sex (female) \times | 2539.00 | 187.92 | (2129.55, | 13.51 | < |
| age37 | | | 2948.45) | | .001 |
| sex (male) \times | 2737.50 | 132.88 | (2447.98, | 20.60 | < |
| age37 | | | 3027.02) | | .001 |
| sex (female) \times | 2872.50 | 132.88 | (2582.98, | 21.62 | < |
| age38 | | | 3162.02) | | .001 |
| $sex (male) \times$ | 2982.00 | 108.50 | (2745.60, | 27.48 | < |
| age38 | | | 3218.40) | | .001 |
| sex (female) \times | 2846.00 | 132.88 | (2556.48, | 21.42 | < |
| age39 | | | 3135.52) | | .001 |
| sex (female) \times | 3152.25 | 93.96 | (2947.52, | 33.55 | < |
| age40 | | | 3356.98) | | .001 |
| $sex (male) \times$ | 3256.25 | 93.96 | (3051.52, | 34.66 | < |
| age40 | | | 3460.98) | | .001 |
| $sex (male) \times$ | 3292.00 | 187.92 | (2882.55, | 17.52 | < |
| age41 | | | 3701.45) | | .001 |
| sex (female) \times | 3210.00 | 187.92 | (2800.55, | 17.08 | < |
| age42 | | | 3619.45) | | .001 |

We call this model the **full**, **maximal**, or **saturated** model for this dataset.

We can calculate the log-likelihood of this model as usual:

```
logLik(lm_max)
#> 'log Lik.' -151.4 (df=13)
```

We can compare this model to our other models using chi-square tests, as usual:

lrtest(lm_max, bw_lm2)

| #Df | LogLik | Df | Chisq | Pr(>Chisq) |
|-----|--------|----|-------|------------|
| 13 | -151.4 | NA | NA | NA |
| 5 | -156.6 | -8 | 10.36 | 0.241 |

The likelihood ratio statistic for this test is

$$\lambda = 2 * (\ell_{\text{full}} - \ell) = 10.3554$$

where:

- $\ell_{\rm max}$ is the log-likelihood of the full model: -151.4016
- ℓ is the log-likelihood of our comparison model (two slopes, two intercepts): -156.5793

This statistic is called the **deviance** or **residual deviance** for our two-slopes and two-intercepts model; it tells us how much the likelihood of that model deviates from the likelihood of the maximal model.

The corresponding p-value tells us whether there we have enough evidence to detect that our two-slopes, two-intercepts model is a worse fit for the data than the maximal model; in other words, it tells us if there's evidence that we missed any important patterns. (Remember, a nonsignificant p-value could mean that we didn't miss anything and a more complicated model is unnecessary, or it could mean we just don't have enough data to tell the difference between these models.)

2.5.3. Null Deviance

Similarly, the least complicated model we could fit would have only one mean parameter, an intercept:

$$\mathrm{E}[Y|X=x] = \beta_0$$

We can fit this model in R like so:

| Parameter | Coefficient | SE | 95% CI | t(23) | р |
|-------------|-------------|-------|--------------------|-------|--------|
| (Intercept) | 2967.67 | 57.58 | (2848.56, 3086.77) | 51.54 | < .001 |

This model also has a likelihood:

```
logLik(lm0)
#> 'log Lik.' -169 (df=2)
```

And we can compare it to more complicated models using a likelihood ratio test:

| #Df | LogLik | Df | Chisq | Pr(>Chisq) |
|-----|--------|----|-------|------------|
| 5 | -156.6 | NA | NA | NA |
| 2 | -169.0 | -3 | 24.75 | 0 |

The likelihood ratio statistic for the test comparing the null model to the maximal model is

$$\lambda=2*(\ell_{\mathrm{full}}-\ell_0)=35.1067$$

where:

- + ℓ_0 is the log-likelihood of the null model: -168.955
- $\ell_{\rm full}$ is the log-likelihood of the maximal model: -151.4016

In R, this test is:

| #Df | LogLik | Df | Chisq | Pr(>Chisq) |
|-----|--------|-----|-------|------------|
| 13 | -151.4 | NA | NA | NA |
| 2 | -169.0 | -11 | 35.11 | 2e-04 |

This log-likelihood ratio statistic is called the **null deviance**. It tells us whether we have enough data to detect a difference between the null and full models.

2.6. Rescaling

2.6.1. Rescale age

```
bw =
  bw |>
  mutate(
    `age - mean` = age - mean(age),
    `age - 36wks` = age - 36
```

```
lm1c = lm(weight ~ sex + `age - 36wks`, data = bw)
lm2c = lm(weight ~ sex + `age - 36wks` + sex: `age - 36wks`, data = bw)
parameters(lm2c, ci_method = "wald") |> print_md()
```

| Parameter | Coefficient | SE | 95% CI | t(20) | p |
|---|-------------|--------|---------------------|-------|--------|
| (Intercept) | 2552.73 | 97.59 | (2349.16, 2756.30) | 26.16 | < .001 |
| sex (male) | 209.97 | 129.75 | (-60.68, 480.63) | 1.62 | 0.121 |
| age - 36 wks | 130.40 | 30.00 | (67.82, 192.98) | 4.35 | < .001 |
| $\begin{array}{l} {\rm sex~(male)} \times {\rm age} \; \text{-} \\ {\rm 36wks} \end{array}$ | -18.42 | 41.76 | (-105.52, 68.68) | -0.44 | 0.664 |

Compare with what we got without rescaling:

```
parameters(bw_lm2, ci_method = "wald") |> print_md()
```

| Parameter | Coefficient | SE | 95% CI | t(20) | p |
|-------------|-------------|---------|------------------------|-------|--------|
| (Intercept) | -2141.67 | 1163.60 | (-4568.90, 285.56) | -1.84 | 0.081 |
| sex (male) | 872.99 | 1611.33 | (-2488.18, 4234.17) | 0.54 | 0.594 |
| age | 130.40 | 30.00 | (67.82, 192.98) | 4.35 | < .001 |

| Parameter | Coefficient | SE | 95% CI | t(20) | p |
|---------------------------------------|-------------|-------|------------------|-------|-------|
| $\overline{\text{sex (male)}} \times$ | -18.42 | 41.76 | (-105.52, 68.68) | -0.44 | 0.664 |
| age | | | | | |

2.7. Prediction

2.7.1. Prediction for linear models

Definition 2.2 (Predicted value). In a regression model p(y|x), the **predicted value** of y given x is the estimated mean of Y given X:

$$\hat{y} \stackrel{\text{def}}{=} \hat{\mathbf{E}} \left[Y | X = x \right]$$

For linear models, the predicted value can be straightforwardly calculated by multiplying each predictor value x_j by its corresponding coefficient β_j and adding up the results:

$$\begin{split} \hat{Y} &= \hat{E}[Y|X=x] \\ &= x'\hat{\beta} \\ &= \hat{\beta}_0 \cdot 1 + \hat{\beta}_1 x_1 + \ldots + \hat{\beta}_p x_p \end{split}$$

2.7.2. Example: prediction for the birthweight data

```
X = c(1,1,40)
sum(X * coef(bw_lm1))
#> [1] 3225
```

R has built-in functions for prediction:

```
x = tibble(age = 40, sex = "male")
bw_lm1 |> predict(newdata = x)
#> 1
#> 3225
```

If you don't provide **newdata**, R will use the covariate values from the original dataset:

```
predict(bw_lm1)
      1
                 3
                                                  9
                                                      10
                                                                 12
                                                                       13
                            5
                                                            11
                                                                            14
                                                                                       16
#> 3225 3062 2984 2579 3225 3062 2621 2821 2742 3304 2863 2942 3346 3062 3225 2700
     17
           18
                19
                     20
                           21
                                22
                                      23
                                           24
#> 2863 2579 2984 2821 3225 2942 2984 3062
```

These special predictions are called the *fitted values* of the dataset:

Definition 2.3. For a given dataset (\tilde{Y}, \mathbf{X}) and corresponding fitted model $p_{\hat{\beta}}(\tilde{y}|\mathbf{x})$, the **fitted value** of y_i is the predicted value of y when $\tilde{X} = \tilde{x}_i$ using the estimate parameters $\hat{\beta}$.

R has an extra function to get these values:

```
fitted(bw_lm1)
      1
            2
                 3
                                            8
                                                  9
                                                      10
                                                            11
                                                                 12
                                                                       13
                                                                            14
                                                                                  15
                                                                                       16
  3225 3062 2984 2579 3225
                              3062 2621 2821 2742 3304 2863 2942 3346 3062 3225 2700
           18
                19
                     20
                           21
                                 22
                                      23
                                           24
     17
#> 2863 2579 2984 2821 3225 2942 2984 3062
```

2.7.3. Quantifying uncertainty in predictions

```
bw_lm1 |>
    predict(
        newdata = x,
        se.fit = TRUE)

#> $fit

#> 1

#> 3225

#>

#> $se.fit

#> [1] 61.46

#>

#> $df

#> [1] 21

#>

#> $residual.scale

#> [1] 177.1
```

This is a list(); you can extract the elements with \$ or magrittr::use_series():

```
bw_lm1 |>
  predict(
    newdata = x,
    se.fit = TRUE) |>
  use_series(se.fit)
#> [1] 61.46
```

You can get **confidence intervals** for $\mathbb{E}[Y|X=x]$:

```
bw_lm1 |> predict(
  newdata = x,
  interval = "confidence")
```

| fit | lwr | upr |
|------|------|------|
| 3225 | 3098 | 3353 |

You can also get **prediction intervals** for the value of an individual outcome Y:

```
bw_lm1 |>
predict(newdata = x, interval = "predict")

fit lwr upr
```

2836

3615

3225

The warning from the last command is: "predictions on current data refer to *future* responses" (since you already know what happened to the current data, and thus don't need to predict it).

See ?predict.lm for more.

2.8. Diagnostics



Tip

This section is adapted from Dobson and Barnett (2018, secs. 6.2-6.3) and Dunn, Smyth, et al. (2018) Chapter 3^a .

^ahttps://link.springer.com/chapter/10.1007/978-1-4419-0118-7_3

2.8.1. Assumptions in linear regression models

$$Y|\tilde{X} \sim N(\tilde{X}'\beta, \sigma^2)$$

- 1. Normality: The distribution conditional on a given X value is normal
- 2. Correct Functional Form: The conditional means have the structure

$$E[Y|\tilde{X} = \tilde{x}] = \tilde{x}'\beta$$

- 3. Homoskedasticity: The variance σ^2 is constant (with respect to \tilde{x})
 - 4. Independence: The observations are statistically independent

2.8.2. Direct visualization

The most direct way to examine the fit of a model is to compare it to the raw observed data.

```
bw =
  bw |>
  mutate(
    predlm2 = predict(bw_lm2)
) |>
  arrange(sex, age)

plot1_interact =
  plot1 %+% bw +
  geom_line(aes(y = predlm2))

print(plot1_interact)
```



Figure 2.6.: Birthweight model with interaction term

It's not easy to assess these assumptions from this model. If there are multiple continuous covariates, it becomes even harder to visualize the raw data.

2.8.3. Residuals

Maybe we can transform the data and model in some way to make it easier to inspect.

Definition 2.4 (Residual noise). The **residual noise** in a probabilistic model p(Y) is the difference between an observed value y and its distributional mean:

$$\varepsilon(y) \stackrel{\text{def}}{=} y - \mathbb{E}[Y] \tag{2.8}$$

We use the same notation for residual noise that we used for errors. $\mathbb{E}[Y]$ can be viewed as an estimate of Y, before y is observed. Conversely, each observation y can be viewed as an estimate of $\mathbb{E}[Y]$ (albeit an imprecise one, individually, since n = 1).

We can rearrange Equation 2.8 to view y as the sum of its mean plus the residual noise:

$$y = \mathbb{E}\left[Y\right] + \varepsilon y$$

Theorem 2.1 (Residuals in Gaussian models). If Y has a Gaussian distribution, then $\varepsilon(Y)$ also has a Gaussian distribution, and vice versa.

Proof. Left to the reader. \Box

Definition 2.5 (Residual errors of a fitted model value). The **residual** of a fitted value \hat{y} (shorthand: "residual") is its error:

$$\begin{aligned} e(\hat{y}) &\stackrel{\text{def}}{=} \varepsilon \left(\hat{y} \right) \\ &= y - \hat{y} \end{aligned}$$

 $e(\hat{y})$ can be seen as the maximum likelihood estimate of the residual noise:

$$\begin{split} e(\hat{y}) &= y - \hat{y} \\ &= \hat{\varepsilon}_{ML} \end{split}$$

2.8.3.1. General characteristics of residuals

Theorem 2.2. For unbiased estimators $\hat{\theta}$:

$$\mathbb{E}\left[e(y)\right] = 0\tag{2.9}$$

$$Var(e(y)) \approx \sigma^2$$
 (2.10)

Proof.

Equation 2.9:

$$\begin{split} \mathbb{E}\left[e(y)\right] &= \mathbb{E}\left[y - \hat{y}\right] \\ &= \mathbb{E}\left[y\right] - \mathbb{E}\left[\hat{y}\right] \\ &= \mathbb{E}\left[y\right] - \mathbb{E}\left[y\right] \\ &= 0 \end{split}$$

Equation 2.10:

$$\begin{split} \operatorname{Var}\left(e(y)\right) &= \operatorname{Var}\left(y - \hat{y}\right) \\ &= \operatorname{Var}\left(y\right) + \operatorname{Var}\left(\hat{y}\right) - 2\operatorname{Cov}\left(y, \hat{y}\right) \\ & \approx \operatorname{Var}\left(y\right) + 0 - 2 \cdot 0 \\ &= \operatorname{Var}\left(y\right) \\ &= \sigma^2 \end{split}$$

2.8.3.2. Characteristics of residuals in Gaussian models

With enough data and a correct model, the residuals will be approximately Guassian distributed, with variance σ^2 , which we can estimate using $\hat{\sigma}^2$: that is:

$$e_i \sim_{\mathrm{iid}} N(0, \hat{\sigma}^2)$$

Example 2.2 (residuals in birthweight data). R provides a function for residuals:

```
resid(bw_lm2)
          1
                  2
                           3
                                    4
                                             5
                                                     6
                                                                       8
                                                                                       10
#>
    176.27 -140.73 -144.13
                              -59.53
                                       177.47 -126.93
                                                        -68.93
                                                                 242.67 -139.33
                                                                                    51.67
#>
                          13
                                   14
                                            15
                                                    16
                                                                      18
                                                                                       20
         11
                 12
                                                             17
                                                                               19
#>
    156.67 -125.13
                     274.28
                             -137.71
                                       -27.69 -246.69 -191.67
                                                                 189.33
                                                                          -11.67 -242.64
#>
        21
                 22
                          23
                                   24
    -47.64
           262.36
                     210.36
```

Exercise 2.8. Check R's output by computing the residuals directly.

Solution.

```
bw$weight - fitted(bw_lm2)
        1
                 2
                        3
                                 4
                                         5
                                                                 8
                                                                                 10
   176.27 -140.73 -144.13 -59.53
#>
                                   177.47 -126.93
                                                   -68.93
                                                            242.67 -139.33
                                                                             51.67
#>
                12
                        13
                               14
                                        15
                                                16
                                                        17
                                                                18
                                                                                 20
       11
                                                                        19
#>
   156.67 -125.13 274.28 -137.71
                                   -27.69 -246.69 -191.67
                                                            189.33 -11.67 -242.64
       21
#>
                22
                        23
                                24
   -47.64 262.36 210.36 -30.62
```

This matches R's output!

2.8.3.3. Graph the residuals

```
xlab("Gestational age (weeks)") +
ylab("residuals (grams)") +
theme(legend.position = "bottom") +
# expand_limits(y = 0, x = 0) +
geom_point(alpha = .7)
print(plot_bw_resid + facet_wrap(~ sex))
```



Figure 2.7.: Residuals of interaction model for birthweight data

Definition 2.6 (Standardized residuals).

$$r_i = \frac{e_i}{\widehat{SD}(e_i)}$$

Hence, with enough data and a correct model, the standardized residuals will be approximately standard Gaussian; that is,

$$r_i \; \sim_{\mathrm{iid}} \; N(0,1)$$

2.8.4. Marginal distributions of residuals

To look for problems with our model, we can check whether the residuals e_i and standardized residuals r_i look like they have the distributions that they are supposed to have, according to the model.

2.8.4.1. Standardized residuals in R

| rst | tandard(bw_lm2) | | | | | | | | | | |
|-----------------------------|-----------------|----------|----------|----------|----------|----------|---------|------------|--|--|--|
| #> | 1 | 2 | 3 | 4 | 5 | 6 | | 7 8 | | | |
| #> | 1.15982 | -0.92601 | -0.87479 | -0.34723 | 1.03507 | -0.73473 | -0.3990 | 1 1.43752 | | | |
| #> | 9 | 10 | 11 | 12 | 13 | 14 | 1 | 5 16 | | | |
| #> | -0.82539 | 0.30606 | 0.92807 | -0.87616 | 1.91428 | -0.86559 | -0.1643 | 0 -1.46376 | | | |
| #> | 17 | 18 | 19 | 20 | 21 | 22 | 2 | 3 24 | | | |
| #> | -1.11016 | 1.09658 | -0.06761 | -1.46159 | -0.28696 | 1.58040 | 1.2671 | 7 -0.19805 | | | |
| resid(bw_lm2)/sigma(bw_lm2) | | | | | | | | | | | |
| #> | 1 | 2 | 3 | 4 | 5 | 6 | | 7 8 | | | |
| #> | 0.97593 | -0.77920 | -0.79802 | -0.32962 | 0.98258 | -0.70279 | -0.3816 | 6 1.34357 | | | |
| #> | 9 | 10 | 11 | 12 | 13 | 14 | 1 | 5 16 | | | |
| #> | -0.77144 | 0.28606 | 0.86741 | -0.69282 | 1.51858 | -0.76244 | -0.1533 | 1 -1.36584 | | | |
| #> | 17 | 18 | 19 | 20 | 21 | 22 | 2 | 3 24 | | | |
| #> | -1.06123 | 1.04825 | -0.06463 | -1.34341 | -0.26376 | 1.45262 | 1.1647 | 1 -0.16954 | | | |

These are not quite the same, because R is doing something more complicated and precise to get the standard errors. Let's not worry about those details for now; the difference is pretty small in this case:

```
rstandard_compare_plot =
  tibble(
    x = resid(bw_lm2)/sigma(bw_lm2),
    y = rstandard(bw_lm2)) |>
  ggplot(aes(x = x, y = y)) +
  geom_point() +
  theme_bw() +
  coord_equal() +
  xlab("resid(bw_lm2)/sigma(bw_lm2)") +
 ylab("rstandard(bw_lm2)") +
  geom_abline(
    aes(
      intercept = 0,
      slope = 1,
      col = "x=y")) +
  labs(colour="") +
  scale_colour_manual(values="red")
print(rstandard_compare_plot)
```



Let's add these residuals to the tibble of our dataset:

```
bw =
  bw |>
  mutate(
    fitted_lm2 = fitted(bw_lm2),

    resid_lm2 = resid(bw_lm2),
    # resid_lm2 = weight - fitted_lm2,

std_resid_lm2 = resid_lm2 / sigma(bw_lm2),
    # std_resid_lm2 = resid_lm2 / sigma(bw_lm2)
)
```

```
bw |>
  select(
   sex,
   age,
   weight,
   fitted_lm2,
   resid_lm2,
   std_resid_lm2
)
```

| sex | age | weight | fitted_lm2 | resid_lm2 | std_resid_lm2 |
|--------------------|-----|--------|------------|-----------|---------------|
| female | 36 | 2729 | 2553 | 176.27 | 1.1598 |
| female | 36 | 2412 | 2553 | -140.73 | -0.9260 |
| female | 37 | 2539 | 2683 | -144.13 | -0.8748 |
| female | 38 | 2754 | 2814 | -59.53 | -0.3472 |
| female | 38 | 2991 | 2814 | 177.47 | 1.0351 |
| female | 39 | 2817 | 2944 | -126.93 | -0.7347 |
| female | 39 | 2875 | 2944 | -68.93 | -0.3990 |
| female | 40 | 3317 | 3074 | 242.67 | 1.4375 |
| female | 40 | 2935 | 3074 | -139.33 | -0.8254 |
| female | 40 | 3126 | 3074 | 51.67 | 0.3061 |
| female | 40 | 3231 | 3074 | 156.67 | 0.9281 |
| female | 42 | 3210 | 3335 | -125.13 | -0.8762 |
| $_{\mathrm{male}}$ | 35 | 2925 | 2651 | 274.28 | 1.9143 |
| $_{ m male}$ | 36 | 2625 | 2763 | -137.71 | -0.8656 |
| $_{\mathrm{male}}$ | 37 | 2847 | 2875 | -27.69 | -0.1643 |
| $_{\mathrm{male}}$ | 37 | 2628 | 2875 | -246.69 | -1.4638 |
| $_{\mathrm{male}}$ | 38 | 2795 | 2987 | -191.67 | -1.1102 |
| $_{\mathrm{male}}$ | 38 | 3176 | 2987 | 189.33 | 1.0966 |
| $_{ m male}$ | 38 | 2975 | 2987 | -11.67 | -0.0676 |
| $_{\mathrm{male}}$ | 40 | 2968 | 3211 | -242.64 | -1.4616 |
| | | | | | |

2. Linear (Gaussian) Models

| sex | age | weight | $fitted_lm2$ | ${\rm resid_lm2}$ | std_resid_lm2 |
|------|-----|--------|---------------|--------------------|-------------------|
| male | 40 | 3163 | 3211 | -47.64 | -0.2870 |
| male | 40 | 3473 | 3211 | 262.36 | 1.5804 |
| male | 40 | 3421 | 3211 | 210.36 | 1.2672 |
| male | 41 | 3292 | 3323 | -30.62 | -0.1981 |

Now let's build histograms:

```
resid_marginal_hist =
  bw |>
  ggplot(aes(x = resid_lm2)) +
  geom_histogram()

print(resid_marginal_hist)
```



Figure 2.8.: Marginal distribution of (nonstandardized) residuals

Hard to tell with this small amount of data, but I'm a bit concerned that the histogram doesn't show a bell-curve shape.

```
std_resid_marginal_hist =
  bw |>
  ggplot(aes(x = std_resid_lm2)) +
  geom_histogram()

print(std_resid_marginal_hist)
```



Figure 2.9.: Marginal distribution of standardized residuals

This looks similar, although the scale of the x-axis got narrower, because we divided by $\hat{\sigma}$ (roughly speaking).

Still hard to tell if the distribution is Gaussian.

2.8.5. QQ plot of standardized residuals

Another way to assess normality is the QQ plot of the standardized residuals versus normal quantiles:

```
library(ggfortify)
# needed to make ggplot2::autoplot() work for `lm` objects

qqplot_lm2_auto =
  bw_lm2 |>
  autoplot(
    which = 2, # options are 1:6; can do multiple at once
    ncol = 1) +
  theme_classic()

print(qqplot_lm2_auto)
```



If the Gaussian model were correct, these points should follow the dotted line.

Fig 2.4 panel (c) in Dobson and Barnett (2018) is a little different; they

didn't specify how they produced it, but other statistical analysis systems do things differently from R.

See also Dunn, Smyth, et al. (2018) §3.5.4⁴.

2.8.5.1. QQ plot - how it's built

Let's construct it by hand:

```
bw = bw |>
 mutate(
   p = (rank(std_resid_lm2) - 1/2)/n(), # "Blom's method"
    expected_quantiles_lm2 = qnorm(p)
  )
qqplot_lm2 =
  bw |>
  ggplot(
    aes(
      x = expected_quantiles_lm2,
      y = std_resid_lm2,
      col = sex,
      shape = sex)
  ) +
  geom_point() +
  theme_classic() +
  theme(legend.position='none') + # removing the plot legend
  ggtitle("Normal Q-Q") +
```

 $^{^4} https://link.springer.com/chapter/10.1007/978-1-4419-0118-7_3\#Sec14:~: text=3.5.4\%20Q\%E2\%80\%93Q\%20Plots\%20and\%20Normality$

```
xlab("Theoretical Quantiles") +
ylab("Standardized residuals")

# find the expected line:

ps <- c(.25, .75)  # reference probabilities
a <- quantile(rstandard(bw_lm2), ps) # empirical quantiles
b <- qnorm(ps)  # theoretical quantiles

qq_slope = diff(a)/diff(b)
qq_intcpt = a[1] - b[1] * qq_slope

qqplot_lm2 =
   qqplot_lm2 +
   geom_abline(slope = qq_slope, intercept = qq_intcpt)

print(qqplot_lm2)</pre>
```



2.8.6. Conditional distributions of residuals

If our Gaussian linear regression model is correct, the residuals e_i and standardized residuals r_i should have:

- an approximately Gaussian distribution, with:
- a mean of 0
- a constant variance

This should be true for every value of x.

If we didn't correctly guess the functional form of the linear component of the mean,

$$\mathrm{E}[Y|X=x] = \beta_0 + \beta_1 X_1 + \ldots + \beta_p X_p$$

Then the the residuals might have nonzero mean.

Regardless of whether we guessed the mean function correctly, ther the variance of the residuals might differ between values of x.

2.8.6.1. Residuals versus fitted values

To look for these issues, we can plot the residuals e_i against the fitted values \hat{y}_i (Figure 2.10).

Residuals vs Fitted 300 18 22 200 100 Residuals 0 -100-20016 2600 2800 3000 3200 Fitted values

Figure 2.10.: birthweight model (Equation 2.2): residuals versus fitted values

If the model is correct, the blue line should stay flat and close to 0, and the cloud of dots should have the same vertical spread regardless of the fitted value.

If not, we probably need to change the functional form of linear component of the mean,

$$\mathrm{E}[Y|X=x] = \beta_0 + \beta_1 X_1 + \ldots + \beta_p X_p$$

2.8.6.2. Example: PLOS Medicine title length data

(Adapted from Dobson and Barnett (2018), §6.7.1)

```
data(PLOS, package = "dobson")
library(ggplot2)
fig1 =
  PLOS |>
  ggplot(
    aes(x = authors,
        y = nchar)
  ) +
  geom_point() +
  theme(legend.position = "bottom") +
  labs(col = "") +
  guides(col=guide_legend(ncol=3))
fig1
```



Figure 2.11.: Number of authors versus title length in $PLOS\ Medicine$ articles

Linear fit

```
lm_PLOS_linear = lm(
  formula = nchar ~ authors,
  data = PLOS)
```

```
fig2 = fig1 +
  geom_smooth(
   method = "lm",
        fullrange = TRUE,
```

```
aes(col = "lm(y ~ x)"))
fig2
library(ggfortify)
autoplot(lm_PLOS_linear, which = 1, ncol = 1)
```



Figure 2.12.: Number of authors versus title length in $PLOS\ Medicine,$ with linear model fit

Quadratic fit

method = "lm",

```
lm_PLOS_quad = lm(
  formula = nchar ~ authors + I(authors^2),
  data = PLOS)

fig3 =
  fig2 +
geom_smooth(
```

```
fullrange = TRUE,
  formula = y ~ x + I(x ^ 2),
  aes(col = "lm(y ~ x + I(x^2))")
)
fig3
autoplot(lm_PLOS_quad, which = 1, ncol = 1)
```



Figure 2.13.: Number of authors versus title length in $PLOS\ Medicine,$ with quadratic model fit

Linear versus quadratic fits

```
library(ggfortify)
autoplot(lm_PLOS_linear, which = 1, ncol = 1)
autoplot(lm_PLOS_quad, which = 1, ncol = 1)
```



Figure 2.14.: Residuals versus fitted plot for linear and quadratic fits to PLOS data

Cubic fit

```
lm_PLOS_cub = lm(
  formula = nchar ~ authors + I(authors^2) + I(authors^3),
  data = PLOS)
```

```
fig4 =
  fig3 +
geom_smooth(
    method = "lm",
    fullrange = TRUE,
    formula = y ~ x + I(x ^ 2) + I(x ^ 3),
    aes(col = "lm(y ~ x + I(x^2) + I(x ^ 3))")
)
fig4
autoplot(lm_PLOS_cub, which = 1, ncol = 1)
```



Figure 2.15.: Number of authors versus title length in PLOS Medicine, with cubic model fit

Logarithmic fit

```
lm_PLOS_log = lm(nchar ~ log(authors), data = PLOS)
```

```
fig5 = fig4 +
  geom_smooth(
    method = "lm",
    fullrange = TRUE,
    formula = y ~ log(x),
    aes(col = "lm(y ~ log(x))")
)
fig5
autoplot(lm_PLOS_log, which = 1, ncol = 1)
```



Figure 2.16.: logarithmic fit

Model selection

anova(lm_PLOS_linear, lm_PLOS_quad)

Table 2.27.: linear vs quadratic

| Res.Df | RSS | Df | Sum of Sq | F | Pr(>F) |
|--------|--------|----|-----------|------|--------|
| 876 | 947502 | NA | NA | NA | NA |
| 875 | 880950 | 1 | 66552 | 66.1 | 0 |

anova(lm_PLOS_quad, lm_PLOS_cub)

Table 2.28.: quadratic vs cubic

| Res.Df | RSS | Df | Sum of Sq | F | Pr(>F) |
|--------|--------|----|-----------|-------|--------|
| 875 | 880950 | NA | NA | NA | NA |
| 874 | 865933 | 1 | 15018 | 15.16 | 1e-04 |

AIC/BIC

```
AIC(lm_PLOS_quad)
#> [1] 8568
AIC(lm_PLOS_cub)
#> [1] 8555

AIC(lm_PLOS_cub)
#> [1] 8555
AIC(lm_PLOS_log)
#> [1] 8544

BIC(lm_PLOS_cub)
```

BIC(lm_PLOS_cub)

#> [1] 8578

BIC(lm_PLOS_log)

#> [1] 8558

Extrapolation is dangerous

```
fig_all = fig5 +
   xlim(0, 60)
fig_all
```



Figure 2.17.: Number of authors versus title length in $PLOS\ Medicine$

2.8.6.3. Scale-location plot

We can also plot the square roots of the absolute values of the standardized residuals against the fitted values (Figure 2.18).

autoplot(bw_lm2, which = 3, ncol = 1) |> print()



Figure 2.18.: Scale-location plot of birthweight data

Here, the blue line doesn't need to be near 0, but it should be flat. If not, the residual variance σ^2 might not be constant, and we might need to transform our outcome Y (or use a model that allows non-constant variance).

2.8.6.4. Residuals versus leverage

We can also plot our standardized residuals against "leverage", which roughly speaking is a measure of how unusual each x_i value is. Very

unusual x_i values can have extreme effects on the model fit, so we might want to remove those observations as outliers, particularly if they have large residuals.





Figure 2.19.: birthweight model with interactions (Equation 2.2): residuals versus leverage

The blue line should be relatively flat and close to 0 here.

2.8.7. Diagnostics constructed by hand

```
bw =
  bw |>
  mutate(
    predlm2 = predict(bw_lm2),
    residlm2 = weight - predlm2,
    std_resid = residlm2 / sigma(bw_lm2),
    # std_resid_builtin = rstandard(bw_lm2), # uses leverage
    sqrt_abs_std_resid = std_resid |> abs() |> sqrt()
)
```

Residuals vs fitted

```
resid_vs_fit = bw |>
    ggplot(
       aes(x = predlm2, y = residlm2, col = sex, shape = sex)
) +
    geom_point() +
    theme_classic() +
    geom_hline(yintercept = 0)
```

```
print(resid_vs_fit)
```



Standardized residuals vs fitted

```
bw |>
  ggplot(
   aes(x = predlm2, y = std_resid, col = sex, shape = sex)
) +
  geom_point() +
  theme_classic() +
  geom_hline(yintercept = 0)
```



Standardized residuals vs gestational age

```
bw |>
    ggplot(
    aes(x = age, y = std_resid, col = sex, shape = sex)
) +
    geom_point() +
    theme_classic() +
    geom_hline(yintercept = 0)
```



sqrt(abs(rstandard())) vs fitted

Compare with autoplot(bw_lm2, 3)

```
bw |>
    ggplot(
    aes(x = predlm2, y = sqrt_abs_std_resid, col = sex, shape = sex)
) +
    geom_point() +
    theme_classic() +
    geom_hline(yintercept = 0)
```



2.9. Model selection

(adapted from Dobson and Barnett (2018) §6.3.3; for more information on prediction, see James et al. (2013) and Harrell (2015)).

If we have a lot of covariates in our dataset, we might want to choose a small subset to use in our model.

There are a few possible metrics to consider for choosing a "best" model.

2.9.1. Mean squared error

We might want to minimize the **mean squared error**, $E[(y-\hat{y})^2]$, for new observations that weren't in our data set when we fit the model.

Unfortunately,

$$\frac{1}{n} \sum_{i=1}^{n} (y_i - \hat{y}_i)^2$$


gives a biased estimate of $\mathrm{E}[(y-\hat{y})^2]$ for new data. If we want an unbiased estimate, we will have to be clever.

2.9.1.1. Cross-validation

```
results_reduced =
  tibble(
    model = "wgt+protein",
    errs = cv_reduced$reps[])
results_full =
  tibble(model = "wgt+age+protein",
        errs = cv_full$reps[])

cv_results =
  bind_rows(results_reduced, results_full)

cv_results |>
  ggplot(aes(y = model, x = errs)) +
  geom_boxplot()
```



comparing metrics

```
compare_results = tribble(
    ~ model, ~ cvRMSE, ~ r.squared, ~adj.r.squared, ~ trainRMSE, ~loglik,
    "full", cv_full$cv, summary(full_model)$r.squared, summary(full_model)$adj.r.squared,
    "reduced", cv_reduced$cv, summary(reduced_model)$r.squared, summary(reduced_model)
compare_results
```

| model | cvRMSE | r.squared | adj.r.squared | trainRMSE | loglik |
|---------|--------|-----------|---------------|-----------|--------|
| full | 7.079 | 0.4805 | 0.3831 | 5.956 | -61.84 |
| reduced | 6.524 | 0.4454 | 0.3802 | 5.971 | -62.49 |

anova(full_model, reduced_model)

| Res.Df | RSS | Df | Sum of Sq | F | Pr(>F) |
|--------|-------|----|-----------|-------|--------|
| 16 | 567.7 | NA | NA | NA | NA |
| 17 | 606.0 | -1 | -38.36 | 1.081 | 0.3139 |

2.9.1.2. stepwise regression

| :> | ols_step_both_ai | c(full_mode | 1) | | | |
|---|---|---|-------------------------------|--------|---|---------|
| ;> ;> | | Ste | pwise Summa | ry | | |
| _ | Variable | AIC | SBC | SBIC | R2 | Adj. R2 |
| > 1 | Base Model protein (+) weight (+) | 137.950 | 140.937 | 80.438 | 0.21427 | 0.17061 |
| | Model Output | | | | | |
| :> :> | | Model S | • | | | |
| > R > R-Square > Adj. 1 | ared R-Squared R-Squared | 0.667 0.445 0.380 | RMSE MSE Coef. V AIC | ar | 5.505 35.648 15.879 132.981 136.964 | |
| > > RMSE > MSE: > MAE: > AIC: | | are Error for Crror cion Criteri | | | 100.004 | |
| :> | | A | NOVA | | | |
| ;> ;> | Sum c | | | | | |

| #> | | Squares | DF M | lean Square | F | Sig. | |
|----|-------------|----------|------------|---------------|--------|--------|--------|
| #> | | | | | | | |
| #> | Regression | 486.778 | 2 | 243.389 | 6.827 | 0.0067 | |
| #> | Residual | 606.022 | 17 | 35.648 | | | |
| #> | Total | 1092.800 | 19 | | | | |
| #> | | | | | | | |
| #> | | | | | | | |
| #> | | | Par | ameter Estima | ates | | |
| #> | | | | | | | |
| #> | model | Beta | Std. Error | Std. Beta | t | Sig | lower |
| #> | | | | | | | |
| #> | (Intercept) | 33.130 | 12.572 | | 2.635 | 0.017 | 6.607 |
| #> | protein | 1.824 | 0.623 | 0.534 | 2.927 | 0.009 | 0.509 |
| #> | weight | -0.222 | 0.083 | -0.486 | -2.662 | 0.016 | -0.397 |
| #> | | | | | | | |

2.9.1.3. Lasso

$$\arg \min_{\theta} \ell(\theta) + \lambda \sum_{j=1}^p |\beta_j|$$


```
library(glmnet)
y = carbohydrate$carbohydrate
x = carbohydrate |>
    select(age, weight, protein) |>
    as.matrix()
fit = glmnet(x,y)
```

autoplot(fit, xvar = 'lambda')



Figure 2.20.: Lasso selection

```
cvfit = cv.glmnet(x,y)
plot(cvfit)
```



2.10. Categorical covariates with more than two levels

2.10.1. Example: birthweight

In the birthweight example, the variable sex had only two observed values:

```
unique(bw$sex)
#> [1] female male
#> Levels: female male
```

If there are more than two observed values, we can't just use a single variable with 0s and 1s.

2.10.2.

For example, Table 2.31 shows the (in)famous⁵ iris data (Anderson (1935)), and Table 2.32 provides summary statistics. The data include three species: "setosa", "versicolor", and "virginica".

head(iris)

Table 2.31.: The iris data

| Sepal.Length | Sepal.Width | Petal.Length | Petal.Width | Species |
|--------------|-------------|--------------|-------------|---------|
| 5.1 | 3.5 | 1.4 | 0.2 | setosa |
| 4.9 | 3.0 | 1.4 | 0.2 | setosa |
| 4.7 | 3.2 | 1.3 | 0.2 | setosa |

⁵https://www.meganstodel.com/posts/no-to-iris/

Table 2.31.: The iris data

| Sepal.Length | Sepal.Width | Petal.Length | Petal.Width | Species |
|--------------|-------------|--------------|-------------|---------|
| 4.6 | 3.1 | 1.5 | 0.2 | setosa |
| 5.0 | 3.6 | 1.4 | 0.2 | setosa |
| 5.4 | 3.9 | 1.7 | 0.4 | setosa |

```
library(table1)
table1(
  x = ~ . | Species,
  data = iris,
  overall = FALSE
)
```

If we want to model Sepal.Length by species, we could create a variable X that represents "setosa" as X=1, "virginica" as X=2, and "versicolor" as X=3.

data(iris) # this step is not always necessary, but ensures you're starting
from the original version of a dataset stored in a loaded package

```
iris =
  iris |>
  tibble() |>
  mutate(
    X = case_when(
        Species == "setosa" ~ 1,
        Species == "virginica" ~ 2,
        Species == "versicolor" ~ 3
```

Table 2.32.: Summary statistics for the ${\tt iris}$ data

| | setosa | versicolor | virginica |
|-------------------|----------------------------|-------------------|-------------------|
| | (N=50) | (N=50) | (N=50) |
| Sepal.Length | | | |
| Mean (SD) | $5.01 \ (0.352)$ | 5.94 (0.516) | 6.59 (0.636) |
| Median [Min, Max] | 5.00 [4.30, 5.80] | 5.90 [4.90, 7.00] | 6.50 [4.90, 7.90] |
| Sepal.Width | | | |
| Mean (SD) | 3.43(0.379) | 2.77(0.314) | 2.97(0.322) |
| Median [Min, Max] | 3.40 [2.30, 4.40] | 2.80 [2.00, 3.40] | 3.00 [2.20, 3.80] |
| Petal.Length | | | |
| Mean (SD) | 1.46 (0.174) | 4.26 (0.470) | 5.55 (0.552) |
| Median [Min, Max] | 1.50 [1.00, 1.90] | 4.35 [3.00, 5.10] | 5.55 [4.50, 6.90] |
| Petal.Width | | | |
| Mean (SD) | $0.246 \ (0.105)$ | 1.33(0.198) | $2.03 \ (0.275)$ |
| Median [Min, Max] | $0.200 \ [0.100, \ 0.600]$ | 1.30 [1.00, 1.80] | 2.00 [1.40, 2.50] |

```
iris |>
  distinct(Species, X)
```

Table 2.33.: iris data with numeric coding of species

| Species | X |
|------------|---|
| setosa | 1 |
| versicolor | 3 |
| virginica | 2 |

Then we could fit a model like:

```
iris_lm1 = lm(Sepal.Length ~ X, data = iris)
iris_lm1 |> parameters() |> print_md()
```

Table 2.34.: Model of iris data with numeric coding of Species

| Parameter | Coefficient | SE | 95% CI | t(148) | p |
|-------------|-------------|------|--------------|--------|--------|
| (Intercept) | 4.91 | 0.16 | (4.60, 5.23) | 30.83 | < .001 |
| X | 0.47 | 0.07 | (0.32, 0.61) | 6.30 | < .001 |

2.10.3. Let's see how that model looks:

```
iris_plot1 = iris |>
  ggplot(
```

```
aes(
    x = X,
    y = Sepal.Length)
) +
geom_point(alpha = .1) +
geom_abline(
    intercept = coef(iris_lm1)[1],
    slope = coef(iris_lm1)[2]) +
theme_bw(base_size = 18)
print(iris_plot1)
```



Figure 2.21.: Model of iris data with numeric coding of Species

We have forced the model to use a straight line for the three estimated means. Maybe not a good idea?

2.10.4. Let's see what R does with categorical variables by default:

```
iris_lm2 = lm(Sepal.Length ~ Species, data = iris)
iris_lm2 |> parameters() |> print_md()
```

Table 2.35.: Model of iris data with Species as a categorical variable

| Parameter | Coefficient | SE | 95% CI | t(147) | p |
|---------------------|-------------|------|--------------|--------|--------|
| (Intercept) | 5.01 | 0.07 | (4.86, 5.15) | 68.76 | < .001 |
| Species | 0.93 | 0.10 | (0.73, 1.13) | 9.03 | < .001 |
| (versicolor) | | | | | |
| Species (virginica) | 1.58 | 0.10 | (1.38, 1.79) | 15.37 | < .001 |

2.10.5. Re-parametrize with no intercept

If you don't want the default and offset option, you can use "-1" like we've seen previously:

```
iris.lm2b = lm(Sepal.Length ~ Species - 1, data = iris)
iris.lm2b |> parameters() |> print_md()
```

| Parameter | Coefficient | SE | 95% CI | t(147) | p |
|---------------------|-------------|------|--------------|--------|--------|
| Species (setosa) | 5.01 | 0.07 | (4.86, 5.15) | 68.76 | < .001 |
| Species | 5.94 | 0.07 | (5.79, 6.08) | 81.54 | < .001 |
| (versicolor) | | | | | |
| Species (virginica) | 6.59 | 0.07 | (6.44, 6.73) | 90.49 | < .001 |

2.10.6. Let's see what these new models look like:

```
iris_plot2 =
  iris |>
  mutate(
    predlm2 = predict(iris_lm2)) |>
  arrange(X) |>
  ggplot(aes(x = X, y = Sepal.Length)) +
  geom_point(alpha = .1) +
  geom_line(aes(y = predlm2), col = "red") +
  geom_abline(
    intercept = coef(iris_lm1)[1],
    slope = coef(iris_lm1)[2]) +
  theme_bw(base_size = 18)

print(iris_plot2)
```



Figure 2.22.

2.10.7. Let's see how R did that:

```
formula(iris_lm2)
#> Sepal.Length ~ Species
model.matrix(iris_lm2) |> as_tibble() |> unique()
```

| (Intercept) | Speciesversicolor | Speciesvirginica |
|-------------|-------------------|------------------|
| 1 | 0 | 0 |
| 1 | 1 | 0 |
| 1 | 0 | 1 |

This is called a "corner point parametrization".

```
formula(iris.lm2b)
#> Sepal.Length ~ Species - 1
model.matrix(iris.lm2b) |> as_tibble() |> unique()
```

| Speciessetosa | Speciesversicolor | Speciesvirginica |
|---------------|-------------------|------------------|
| 1 | 0 | 0 |
| 0 | 1 | 0 |
| 0 | 0 | 1 |

This can be called a "group point parametrization".

There are more options; see Dobson and Barnett (2018) §6.4.1 and the codingMatrices package⁶ vignette⁷ (Venables (2023)).

2.11. Ordinal covariates

(c.f. Dobson and Barnett (2018) §2.4.4)

We can create ordinal variables in R using the ordered() function⁸.

Example 2.3.

⁶https://CRAN.R-project.org/package=codingMatrices

 $^{^{7}} https://cran.r-project.org/web/packages/codingMatrices/vignettes/codingMatrices. \\pdf$

⁸or equivalently, factor(ordered = TRUE)

```
url = paste0(
   "https://regression.ucsf.edu/sites/g/files/tkssra6706/",
   "f/wysiwyg/home/data/hersdata.dta")
library(haven)
hers = read_dta(url)
hers |> head()
```

Table 2.39.: HERS dataset

| Н | Tagera | ae | płak | хифі | kidas | l g ad | a igl | ealpo | ted | cebali | iosch | adi | heb | eitp | ishkiightalatyllavoightalatyllatokaltHIIIG cHo IHIIIGS IBDB3ge1(|
|---|--------|----|------|------|-------|---------------|------------------|-------|-----|--------|-------|-----|-----|------|---|
| 0 | 702 | 1 | 0 | 0 | 0 | 5 | 3 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 73 23 9600 98273 23 9630 992189 23 2473 2013 72677 1388 7.0 |
| 0 | 622 | 1 | 0 | 0 | 0 | 1 | 3 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 70 28 5520 96473 28 3500 978430 2 444610 2 165458711 8 06.2 |
| 1 | 691 | 0 | 0 | 0 | 0 | 3 | 3 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 10 2/205110.7829 6/40. 733.738 2546 6.7 215 2 545 6.6 016 0 3 4 86.9 |
| 0 | 641 | 0 | 1 | 1 | 0 | 1 | 3 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 64 24 \$ 7 00\$ 7 4758 2 2. 77 20\$ 9 0 2 2041 6 £ 1 5 2 0 7 2 3 . 7 13 7 5 7 26.4 |
| 0 | 651 | 0 | 0 | 0 | 0 | 2 | 3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 57 2 0.7900.790458 2 2.72805.79572145042610 7 23 5 7 3 2513 9 7 9 56.5 |
| 1 | 682 | 1 | 0 | 1 | 0 | 3 | 3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 60 2 9 9 510000 5 7 2 7 3 620.9110 2 12 3 7 2 8112022 53 511 2 7 4 86.8 |

```
# C(contr = codingMatrices::contr.diff)
```

Logistic regression and variations



Configuring R

Functions from these packages will be used throughout this document:

```
library(conflicted) # check for conflicting function definitions
# library(printr) # inserts help-file output into markdown output
library(rmarkdown) # Convert R Markdown documents into a variety of formats.
library(pander) # format tables for markdown
library(ggplot2) # graphics
library(ggeasy) # help with graphics
library(ggfortify) # help with graphics
library(dplyr) # manipulate data
library(tibble) # `tibble`s extend `data.frame`s
library(magrittr) # `%>%` and other additional piping tools
library(haven) # import Stata files
library(knitr) # format R output for markdown
```

```
library(tidyr) # Tools to help to create tidy data
library(plotly) # interactive graphics
library(dobson) # datasets from Dobson and Barnett 2018
library(parameters) # format model output tables for markdown
library(haven) # import Stata files
library(latex2exp) # use LaTeX in R code (for figures and tables)
library(fs) # filesystem path manipulations
library(survival) # survival analysis
library(survminer) # survival analysis graphics
library(KMsurv) # datasets from Klein and Moeschberger
library(parameters) # format model output tables for
library(webshot2) # convert interactive content to static for pdf
library(forcats) # functions for categorical variables ("factors")
library(stringr) # functions for dealing with strings
library(lubridate) # functions for dealing with dates and times
```

Here are some R settings I use in this document:

```
pander::panderOptions("table.split.table", Inf)
conflicts_prefer(dplyr::filter) # use the `filter()` function from dplyr() by defaul
legend_text_size = 9
```

Acknowledgements

This content is adapted from:

- Dobson and Barnett (2018), Chapter 7
- Vittinghoff et al. (2012), Chapter 5
- David Rocke¹'s materials from the 2021 edition of Epi 204²
- Nahhas (2024) Chapter 6^3

3.1. Introduction

3.1.1. What is logistic regression?

Logistic regression is a framework for modeling binary outcomes, conditional on one or more *predictors* (a.k.a. *covariates*).

Exercise 3.1 (Examples of binary outcomes). What are some examples of binary outcomes in the health sciences?

¹https://dmrocke.ucdavis.edu/

²https://dmrocke.ucdavis.edu/Class/EPI204-Spring-2021/EPI204-Spring-2021.html

³https://www.bookdown.org/rwnahhas/RMPH/blr.html

Solution. Examples of binary outcomes include:

- exposure (exposed vs unexposed)
- disease (diseased vs healthy)
- recovery (recovered vs unrecovered)
- relapse (relapse vs remission)
- return to hospital (returned vs not)
- vital status (dead vs alive)

Logistic regression uses the Bernoulli distribution to model the outcome variable, conditional on one or more covariates.

Exercise 3.2. Write down a mathematical definition of the Bernoulli distribution.

Solution. The **Bernoulli distribution** family for a random variable X is defined as:

$$\begin{split} \Pr(X = x) &= \mathbb{1}_{x \in \{0,1\}} \pi^x (1 - \pi)^{1 - x} \\ &= \left\{ \begin{matrix} \pi, x = 1 \\ 1 - \pi, x = 0 \end{matrix} \right. \end{split}$$

3.1.2. Logistic regression versus linear regression

Logistic regression differs from linear regression, which uses the Gaussian ("normal") distribution to model the outcome variable, conditional on the covariates.

Exercise 3.3. Recall: what kinds of outcomes is linear regression used for?

Solution. Linear regression is typically used for numerical outcomes that aren't event counts or waiting times for an event. Examples of outcomes that are often analyzed using linear regression include include weight, height, and income.

3.2. Risk Estimation and Prediction

In Epi 203, you have already seen methods for modeling binary outcomes using one covariate that is also binary (such as exposure/non-exposure). In this section, we review one-covariate analyses, with a special focus on risk ratios and odds ratios, which are important concepts for interpreting logistic regression.

Example 3.1 (Oral Contraceptive Use and Heart Attack).

• Research question: how does oral contraceptive (OC) use affect the risk of myocardial infarction (MI; a.k.a. heart attack)?

This was an issue when oral contraceptives were first developed, because the original formulations used higher concentrations of hormones. Modern OCs don't have this issue.

Table 3.1 contains simulated data for an imaginary follow-up (a.k.a. *prospective*) study in which two groups are identified, one using OCs and another not using OCs, and both groups are tracked for three years to determine how many in each groups have MIs.

Exercise 3.4. Review: estimate the probabilities of MI for OC users and non-OC users in Example 3.1.

Solution.

$$\hat{\mathbf{p}}(MI|OC) = \frac{13}{5000} = 0.0026$$

$$\hat{\mathbf{p}}(MI|\neg OC) = \frac{7}{10000} = 7 \times 10^{-4}$$

Table 3.1.: Simulated data from study of oral contraceptive use and heart attack risk

```
library(dplyr)
oc_mi =
 tribble(
   ~OC, ~MI, ~Total,
   "OC use", 13, 5000,
   "No OC use", 7, 10000
 ) |>
 mutate(`No MI` = Total - MI) |>
 relocate(`No MI`, .after = MI)
totals =
 oc_mi |>
 summarize(across(c(MI, `No MI`, Total), sum)) |>
 mutate(OC = "Total")
tbl_oc_mi = bind_rows(oc_mi, totals)
tbl_oc_mi
#> # A tibble: 3 x 4
#>
   OC
          MI `No MI` Total
#> <chr>
            <dbl> <dbl> <dbl>
#> 1 OC use 13
                    4987 5000
#> 2 No OC use
                7
                     9993 10000
#> 3 Total 20 14980 15000
```

3.2.0.1. Controls

Two meanings of "controls"

Depending on context, "controls" can mean either individuals who don't experience an exposure of interest, or individuals who don't experience an outcome of interest.

Definition 3.1 (cases and controls in retrospective studies). In *retrospective studies*, participants who experience the outcome of interest are called **cases**, while participants who don't experience that outcome are called **controls**.

Definition 3.2 (treatment groups and control groups in prospective studies). In *prospective studies*, the group of participants who experience the treatment or exposure of interest is called the **treatment group**, while the participants who receive the baseline or comparison treatment (for example, clinical trial participants who receive a placebo or a standard-of-care treatment rather than an experimental treatment) are called **controls**.

3.3. Comparing Probabilities

3.3.1. Risk differences

The simplest comparison of two probabilities, π_1 , and π_2 , is the difference of their values:

Definition 3.3 (Risk difference). The **risk difference** of two probabilities, π_1 , and π_2 , is the difference of their values:

$$\delta(\pi_1,\pi_2) \stackrel{\mathrm{def}}{=} \pi_1 - \pi_2$$

Example 3.2 (Difference in MI risk). In Example 3.1, the maximum likelihood estimate of the difference in MI risk between OC users and OC non-users is:

$$\begin{split} \hat{\delta}(\pi(OC),\pi(\neg OC)) &= \delta(\hat{\pi}(OC),\hat{\pi}(\neg OC)) \\ &= \hat{\pi}(OC) - \hat{\pi}(\neg OC) \\ &= 0.0026 - 7 \times 10^{-4} \\ &= 0.0019 \end{split}$$

3.3.2. Risk ratios

Definition 3.4 (Relative risk ratios).

The **relative risk** of probability π_1 compared to another probability π_2 , also called the **risk ratio**, **relative risk ratio**, **probability ratio**, or **rate ratio**, is the ratio of those probabilities:

$$\rho(\pi_1,\pi_2) = \frac{\pi_1}{\pi_2}$$

Example 3.3.

Above, we estimated that:

$$\hat{p}(MI|OC) = 0.0026$$

$$\hat{\mathbf{p}}(MI|\neg OC) = 7 \times 10^{-4}$$

So we might estimate that the $relative\ risk$ of MI for OC versus non-OC is:

rr = (13/5000)/(7/10000)

$$\begin{split} \hat{\rho}(OC, \neg OC) &= \frac{\hat{\mathbf{p}}(MI|OC)}{\hat{\mathbf{p}}(MI|\neg OC)} \\ &= \frac{0.0026}{7 \times 10^{-4}} \\ &= 3.7143 \end{split}$$

We might summarize this result by saying that "the estimated probability of MI among OC users was 3.7143 as high as the estimated probability among OC non-users.

3.3.3. Relative risk differences

Definition 3.5 (Relative risk difference).

Sometimes, we divide the risk difference by the comparison probability; the result is called the **relative risk difference**:

$$\xi(\pi_1,\pi_2) \stackrel{\mathrm{def}}{=} \frac{\delta(\pi_1,\pi_2)}{\pi_2}$$

Theorem 3.1 (Relative risk difference equals risk ratio minus 1).

$$\xi(\pi_1, \pi_2) = \rho(\pi_1, \pi_2) - 1$$

Proof.

$$\begin{split} \xi(\pi_1, \pi_2) & \stackrel{\text{def}}{=} \frac{\delta(\pi_1, \pi_2)}{\pi_2} \\ & = \frac{\pi_1 - \pi_2}{\pi_2} \\ & = \frac{\pi_1}{\pi_2} - 1 \\ & = \rho(\pi_1, \pi_2) - 1 \end{split}$$

3.3.4. Changing reference groups in risk comparisons

Risk differences, risk ratios, and relative risk differences are defined by two probabilities, plus a choice of which probability is the **baseline** or **reference** probability (i.e., which probability is the subtrahend of the risk difference or the denominator of the risk ratio).

$$\delta(\pi_2,\pi_1) = -\delta(\pi_1,\pi_2)$$

$$\begin{split} \rho(\pi_2,\pi_1) &= \left(\rho(\pi_1,\pi_2)\right)^{-1} \\ \xi(\pi_2,\pi_1) &= \left(\xi(\pi_2,\pi_1) + 1\right)^{-1} - 1 \end{split}$$

Exercise 3.5. Prove the relationships above.

Example 3.4 (Switching the reference group in a risk ratio). Above, we estimated that the risk ratio of OC versus non-OC is:

$$\rho(OC, \neg OC) = 3.7143$$

In comparison, the risk ratio for non-OC versus OC is:

$$\begin{split} \rho(\neg OC, OC) &= \frac{\hat{\mathbf{p}}(MI|\neg OC)}{\hat{\mathbf{p}}(MI|OC)} \\ &= \frac{7\times 10^{-4}}{0.0026} \\ &= 0.2692 \\ &= \frac{1}{\rho(OC, \neg OC)} \end{split}$$

3.4. Odds and Odds Ratios

3.4.1. Odds and probabilities

In logistic regression, we will make use of a mathematically-convenient transformation of probability, called *odds*.

Definition 3.6 (Odds). The **odds** of an outcome A, which we will represent using ω ("omega"), is the probability that the outcome occurs, divided by the probability that it doesn't occur:

$$\omega(A) \stackrel{\text{def}}{=} \frac{\Pr(A)}{\Pr(\neg A)}$$

Theorem 3.2. If the probability of an outcome A is $Pr(A) = \pi$, then the corresponding odds of A is:

$$odds\{\pi\} = \frac{\pi}{1-\pi} \tag{3.1}$$

Proof.

$$\begin{aligned} \Pr(\neg A) &= 1 - \Pr(A) \\ &= 1 - \pi \end{aligned}$$

Function 3.1, which transforms probabilities into odds, can be called the **odds function**. Figure 3.1 graphs the shape of this function.

```
odds = function(pi) pi / (1 - pi)
library(ggplot2)
ggplot() +
  geom_function(
   fun = odds,
   arrow = arrow(ends = "last"),
   mapping = aes(col = "odds function")
 xlim(0, .99) +
 xlab("Probability") +
  ylab("Odds") +
  geom_abline(aes(
   intercept = 0,
    slope = 1,
    col = "y=x"
  )) +
  theme_bw() +
  labs(colour = "") +
  theme(legend.position = "bottom")
```



Figure 3.1.: Odds versus probability

Example 3.5 (Computing odds from probabilities). In Exercise 3.4, we estimated that the probability of MI, given OC use, is $\pi(OC) \stackrel{\text{def}}{=} \Pr(MI|OC) = 0.0026$. If this estimate is correct, then the odds of MI, given OC use, is:

$$\begin{split} \omega(OC) &\stackrel{\text{def}}{=} \frac{\Pr(MI|OC)}{\Pr(\neg MI|OC)} \\ &= \frac{\Pr(MI|OC)}{1 - \Pr(MI|OC)} \\ &= \frac{\pi(OC)}{1 - \pi(OC)} \\ &= \frac{0.0026}{1 - 0.0026} \\ &\approx 0.002607 \end{split}$$

Exercise 3.6 (Computing odds from probabilities). Estimate the odds of MI, for non-OC users.

Solution.

$$\omega(\neg OC) = 7.0049 \times 10^{-4}$$

Theorem 3.3 (One-sample MLE for odds). Let $X_1,...X_n$ be a set of n iid Bernoulli trials, and let $X = \sum_{i=1}^n X_i$ be their sum.

Then the maximum likelihood estimate of the odds of X = 1, ω , is:

$$\hat{\omega} = \frac{x}{n-x}$$

Proof.

$$1 - \hat{\pi} = 1 - \frac{x}{n}$$
$$= \frac{n}{n} - \frac{x}{n}$$
$$= \frac{n - x}{n}$$

Thus, the estimated odds is:

$$\frac{\hat{\pi}}{1 - \hat{\pi}} = \frac{\left(\frac{x}{n}\right)}{\left(\frac{n-x}{n}\right)}$$
$$= \frac{x}{n-x}$$

That is, odds can be calculated directly as "# events" divided by "# nonevents" (without needing to calculate $\hat{\pi}$ and $1 - \hat{\pi}$ first).

Example 3.6 (calculating odds using the shortcut). In Example 3.5, we calculated

$$\omega(OC) = 0.0026$$

Let's recalculate this result using our shortcut.

Solution 3.1.

$$\omega(OC) = \frac{13}{5000 - 13} \\ = 0.0026$$

Same answer as in Example 3.5!

$3.\ \, \textit{Models for Binary Outcomes}$

Theorem 3.4 (Simplified expression for odds function).

An equivalent expression for the odds function is

$$odds\{\pi\} = \left(\left(\pi\right)^{-1} - 1\right)^{-1}$$
 (3.2)

Exercise 3.7. Prove that Equation 3.2 is equivalent to Definition 3.6.

Theorem 3.5 (Derivative of odds function).

$$odds'\left\{\pi\right\} = \frac{1}{\left(1 - \pi\right)^2}$$

Proof.

$$\operatorname{odds}' \{\pi\} = \frac{\partial}{\partial \pi} \left(\frac{\pi}{1 - \pi} \right)$$

$$= \frac{\frac{\partial}{\partial \pi} \pi}{1 - \pi} - \left(\frac{\pi}{(1 - \pi)^2} \cdot \frac{\partial}{\partial \pi} (1 - \pi) \right)$$

$$= \frac{1}{1 - \pi} - \frac{\pi}{(1 - \pi)^2} \cdot (-1)$$

$$= \frac{1}{1 - \pi} + \frac{\pi}{(1 - \pi)^2}$$

$$= \frac{1 - \pi}{(1 - \pi)^2} + \frac{\pi}{(1 - \pi)^2}$$

$$= \frac{1 - \pi + \pi}{(1 - \pi)^2}$$

$$= \frac{1}{(1 - \pi)^2}$$

3.4.1.1. Odds of rare events

For rare events (small π), odds and probabilities are nearly equal, because $1-\pi\approx 1$ (see Figure 3.1).

For example, in Example 3.5, the probability and odds differ by 6.7776×10^{-6} .

Exercise 3.8. What odds value corresponds to the probability $\pi = 0.2$, and what is the numerical difference between these two values?

Solution.

$$\omega = \frac{\pi}{1-\pi} = \frac{.2}{.8} = .25$$

Theorem 3.6. Let $\omega = \frac{\pi}{1-\pi}$. Then:

$$\omega - \pi = \omega \cdot \pi$$

Exercise 3.9. Prove Theorem 3.6.

Solution.

$$\begin{split} \omega - \pi &= \frac{\pi}{1 - \pi} - \pi \\ &= \frac{\pi}{1 - \pi} - \frac{\pi(1 - \pi)}{1 - \pi} \\ &= \frac{\pi}{1 - \pi} - \frac{\pi - \pi^2}{1 - \pi} \\ &= \frac{\pi - (\pi - \pi^2)}{1 - \pi} \\ &= \frac{\pi - \pi + \pi^2}{1 - \pi} \\ &= \frac{\pi^2}{1 - \pi} \\ &= \frac{\pi}{1 - \pi} \\ &= \omega \pi \end{split}$$

Lemma 3.1 (Odds of a non-event). If π is the odds of event A and ω is the corresponding odds of A, then the odds of $\neg A$ are:

$$\omega(\neg A) = \frac{1-\pi}{\pi}$$

Proof. Left to the reader.

3.4.2. The inverse odds function

Definition 3.7 (inverse odds function). The inverse odds function,

invodds
$$\{\omega\} \stackrel{\text{def}}{=} \frac{\omega}{1+\omega}$$

converts odds into their corresponding probabilities (Figure 3.2).

The inverse-odds function takes an odds as input and produces a probability as output. Its domain of inputs is $[0, \infty)$ and its range of outputs is [0, 1].

I haven't seen anyone give the inverse-odds function a concise name; maybe prob()?

```
odds_inv = function(omega) (1 + omega^-1)^-1
ggplot() +
   geom_function(fun = odds_inv, aes(col = "inverse-odds")) +
   xlab("Odds") +
   ylab("Probability") +
   xlim(0,5) +
   ylim(0,1) +
   geom_abline(aes(intercept = 0, slope = 1, col = "x=y"))
```



Figure 3.2.: The inverse odds function, invodds $\{\omega\}$

Exercise 3.10. What probability corresponds to an odds of $\omega = 1$, and what is the numerical difference between these two values?

Solution.

$$\pi(1) = \frac{1}{1+1} = \frac{1}{2} = .5$$
$$1 - \pi(1) = 1 - .5 = .5$$

Lemma 3.2 (Simplified expression for inverse odds function).

An equivalent expression for the inverse odds function is

$$\pi(\omega) = (1 + \omega^{-1})^{-1} \tag{3.3}$$

Exercise 3.11. Prove that Equation 3.3 is equivalent to Definition 3.7.

Lemma 3.3 (One minus inverse-odds).

$$1 - invodds\{\omega\} = \frac{1}{1+\omega}$$

Proof.

$$\begin{aligned} 1 - \text{invodds} \left\{ \omega \right\} &= 1 - \frac{\omega}{1 + \omega} \\ &= \frac{1 + \omega}{1 + \omega} - \frac{\omega}{1 + \omega} \\ &= \frac{(1 + \omega) - \omega}{1 + \omega} \\ &= \frac{1 + \omega - \omega}{1 + \omega} \\ &= \frac{1}{1 + \omega} \end{aligned}$$

Theorem 3.7. If ω is the odds of event A, then the probability that A does not occur is:

$$\Pr(\neg A) = \frac{1}{1+\omega}$$

Proof.

Use Lemma 3.3:

$$\begin{aligned} \Pr(\neg A) &= 1 - \Pr(A) \\ &= 1 - \text{invodds} \{\omega\} \\ &= \frac{1}{1 + \omega} \end{aligned}$$

Theorem 3.8 (Derivative of inverse odds function).

$$invodds' \{\omega\} = \frac{1}{(1+\omega)^2}$$

Proof.

Use the quotient rule:

$$\operatorname{invodds}'(\omega) = \frac{\partial}{\partial \omega} \operatorname{invodds} \{\omega\}$$

$$= \frac{\partial}{\partial \omega} \frac{\omega}{1 + \omega}$$

$$= \frac{\frac{\partial}{\partial \omega} \omega}{1 + \omega} - \frac{\omega}{(1 + \omega)^2} \cdot \frac{\partial}{\partial \omega} (1 + \omega)$$

$$= \frac{1}{1 + \omega} - \frac{\omega}{(1 + \omega)^2} \cdot 1$$

$$= \frac{1}{1 + \omega} - \frac{\omega}{(1 + \omega)^2}$$

$$= \frac{1 + \omega}{(1 + \omega)^2} - \frac{\omega}{(1 + \omega)^2}$$

$$= \frac{1 + \omega - \omega}{(1 + \omega)^2}$$

$$= \frac{1}{(1 + \omega)^2}$$

Corollary 3.1.

$$\mathit{invodds}'\left\{\omega\right\} = \left(1 - \mathit{invodds}\left\{\omega\right\}\right)^2$$

3.4.3. Odds ratios

Now that we have defined odds, we can introduce another way of comparing event probabilities: odds ratios.

Definition 3.8 (Odds ratio). The **odds ratio** for two odds ω_1 , ω_2 is their ratio:

$$\theta(\omega_1,\omega_2) = \frac{\omega_1}{\omega_2}$$

Example 3.7 (Calculating odds ratios). In Example 3.1, the odds ratio for OC users versus OC-non-users is:

$$\theta(\omega(OC), \omega(\neg OC)) = \frac{\omega(OC)}{\omega(\neg OC)}$$
$$= \frac{0.0026}{7 \times 10^{-4}}$$
$$= 3.7143$$

3.4.3.1. A shortcut for calculating odds ratio estimates

The general form of a two-by-two table is shown in Table 3.2.

Table 3.2.: A generic 2x2 table

| | Event | Non-Event | Total |
|-------------|--------------|-----------|---------|
| Exposed | a | b | a+b |
| Non-exposed | \mathbf{c} | d | c+d |
| Total | a+c | b+d | a+b+c+d |

From this table, we have:

- $\bullet \ \ \widehat{\pi}(Event|Exposed) = a/(a+b)$
- $\hat{\pi}(\neg Event|Exposed) = b/(a+b)$
- $\hat{\omega}(Event|Exposed) = \frac{\left(\frac{a}{a+b}\right)}{\left(\frac{b}{a+b}\right)} = \frac{a}{b}$
- $\hat{\omega}(Event|\neg Exposed) = \frac{c}{d}$ (see Exercise 3.12)
- $\theta(Exposed, \neg Exposed) = \frac{\frac{a}{b}}{\frac{c}{d}} = \frac{ad}{bc}$

Exercise 3.12. Given Table 3.2, show that $\hat{\omega}(Event|\neg Exposed) = \frac{c}{d}$.

158

3.4.3.2. Properties of odds ratios

Odds ratios have a special property: we can swap a covariate with the outcome, and the odds ratio remains the same.

Theorem 3.9 (Odds ratios are reversible). For any two events A, B:

$$\theta(A|B) = \theta(B|A)$$

Proof.

$$\theta(A|B) \stackrel{\text{def}}{=} \frac{\omega(A|B)}{\omega(A|\neg B)}$$

$$= \frac{\left(\frac{p(A|B)}{p(\neg A|B)}\right)}{\left(\frac{p(A|\neg B)}{p(\neg A|\neg B)}\right)}$$

$$= \left(\frac{p(A|B)}{p(\neg A|B)}\right) \left(\frac{p(A|\neg B)}{p(\neg A|\neg B)}\right)^{-1}$$

$$= \left(\frac{p(A|B)}{p(\neg A|B)}\right) \left(\frac{p(\neg A|\neg B)}{p(A|\neg B)}\right)$$

$$= \left(\frac{p(A|B)}{p(\neg A|B)} \cdot \frac{p(B)}{p(B)}\right) \left(\frac{p(\neg A|\neg B)}{p(A|\neg B)} \cdot \frac{p(\neg B)}{p(\neg B)}\right)$$

$$= \left(\frac{p(A,B)}{p(\neg A,B)}\right) \left(\frac{p(\neg A,\neg B)}{p(A,\neg B)}\right)$$

$$= \left(\frac{p(B,A)}{p(B,\neg A)}\right) \left(\frac{p(\neg B,\neg A)}{p(\neg B,A)}\right)$$

$$= \left(\frac{p(B,A)}{p(\neg B,A)}\right) \left(\frac{p(\neg B,\neg A)}{p(B,\neg A)}\right)$$

$$= [\text{reverse the preceding steps}]$$

$$= \theta(B|A)$$

Example 3.8. In Example 3.1, we have:

160

$$\begin{split} \theta(MI;OC) &\stackrel{\text{def}}{=} \frac{\omega(MI|OC)}{\omega(MI|\neg OC)} \\ &\stackrel{\text{def}}{=} \frac{\left(\frac{\Pr(MI|OC)}{\Pr(\neg MI|OC)}\right)}{\left(\frac{\Pr(MI|\neg OC)}{\Pr(\neg MI, \neg OC)}\right)} \\ &= \frac{\left(\frac{\Pr(MI,OC)}{\Pr(\neg MI,OC)}\right)}{\left(\frac{\Pr(MI,OC)}{\Pr(\neg MI,OC)}\right)} \\ &= \left(\frac{\Pr(MI,OC)}{\Pr(\neg MI,OC)}\right) \left(\frac{\Pr(\neg MI,\neg OC)}{\Pr(MI,\neg OC)}\right) \\ &= \left(\frac{\Pr(MI,OC)}{\Pr(MI, \neg OC)}\right) \left(\frac{\Pr(\neg MI, \neg OC)}{\Pr(\neg MI,OC)}\right) \\ &= \left(\frac{\Pr(OC,MI)}{\Pr(\neg OC,MI)}\right) \left(\frac{\Pr(\neg OC,\neg MI)}{\Pr(OC,\neg MI)}\right) \\ &= \left(\frac{\Pr(OC|MI)}{\Pr(\neg OC|MI)}\right) \left(\frac{\Pr(\neg OC|\neg MI)}{\Pr(OC|\neg MI)}\right) \\ &= \frac{\left(\frac{\Pr(OC|MI)}{\Pr(\neg OC|MI)}\right)}{\left(\frac{\Pr(OC|\neg MI)}{\Pr(\neg OC|\rightarrow MI)}\right)} \\ &\stackrel{\text{def}}{=} \frac{\omega(OC|MI)}{\omega(OC|\neg MI)} \\ &\stackrel{\text{def}}{=} \theta(OC;MI) \end{split}$$

Exercise 3.13. For Table 3.2, show that $\hat{\theta}(Exposed, Unexposed) =$ $\hat{\theta}(Event, \neg Event).$

Conditional odds ratios have the same reversibility property:

Theorem 3.10 (Conditional odds ratios are reversible). For any three events A, B, C:

$$\theta(A|B,C) = \theta(B|A,C)$$

Proof. Apply the same steps as for Theorem 3.9, inserting C into the conditions (RHS of |) of every expression.

3.4.3.3. Odds Ratios vs Probability (Risk) Ratios

When the outcome is rare (i.e., its probability is small) for both groups being compared in an odds ratio, the odds of the outcome will be similar to the probability of the outcome, and thus the risk ratio will be similar to the odds ratio.

Case 1: rare events

For rare events, odds ratios and probability (a.k.a. risk, a.k.a. prevalence) ratios will be close:

$$\pi_1 = .01~\pi_2 = .02$$

```
pi1 = .01
pi2 = .02
pi2/pi1
#> [1] 2
odds(pi2)/odds(pi1)
#> [1] 2.02
```

Example 3.9. In Example 3.1, the outcome is rare for both OC and non-OC participants, so the odds for both groups are similar to the corresponding probabilities, and the odds ratio is similar the risk ratio.

Case 2: frequent events

```
\pi_1 = .4 \ \pi_2 = .5
```

For more frequently-occurring outcomes, this won't be the case:

```
pi1 = .4
pi2 = .5
pi2/pi1
#> [1] 1.25
odds(pi2)/odds(pi1)
#> [1] 1.5
```

3.4.3.4. Odds Ratios in Case-Control Studies

Table 3.1 simulates a follow-up study in which two populations were followed and the number of MI's was observed. The risks are P(MI|OC) and $P(MI|\neg OC)$ and we can estimate these risks from the data.

But suppose we had a case-control study in which we had 100 women with MI and selected a comparison group of 100 women without MI (matched as groups on age, etc.). Then MI is not random, and we cannot compute

P(MI|OC) and we cannot compute the risk ratio. However, the odds ratio however can be computed.

The disease odds ratio is the odds for the disease in the exposed group divided by the odds for the disease in the unexposed group, and we cannot validly compute and use these separate parts.

We can still validly compute and use the exposure odds ratio, which is the odds for exposure in the disease group divided by the odds for exposure in the non-diseased group (because exposure can be treated as random):

$$\hat{\theta}(OC|MI) = \frac{\hat{\omega}(OC|MI)}{\hat{\omega}(OC|\neg MI)}$$

And these two odds ratios, $\hat{\theta}(MI|OC)$ and $\hat{\theta}(OC|MI)$, are mathematically equivalent, as we saw in Section 3.4.3.2:

$$\hat{\theta}(MI|OC) = \hat{\theta}(OC|MI)$$

Exercise 3.14. Calculate the odds ratio of MI with respect to OC use, assuming that Table 3.1 comes from a case-control study. Confirm that the result is the same as in Example 3.7.

Solution.

•
$$\omega(OC|MI) = P(OC|MI)/(1 - P(OC|MI) = \frac{13}{7} = 1.8571$$

•
$$\omega(OC|\neg MI) = P(OC|\neg MI)/(1-P(OC|\neg MI) = \frac{4987}{9993} = 0.499$$

•
$$\theta(OC, MI) = \frac{\omega(OC|MI)}{\omega(OC|-MI)} = \frac{13/7}{4987/9993} = 3.7214$$

This is the same estimate we calculated in Example 3.7.

3.4.3.5. Odds Ratios in Cross-Sectional Studies

- If a cross-sectional study is a probability sample of a population (which it rarely is) then we can estimate risks.
- If it is a sample, but not an unbiased probability sample, then we need to treat it in the same way as a case-control study.
- We can validly estimate odds ratios in either case.
- But we can usually not validly estimate risks and risk ratios.

3.5. The logit and expit functions

3.5.1. The logit function

Definition 3.9 (logit function).

The **logit function** of a probability π is the natural logarithm of the odds function of π :

$$\operatorname{logit}(\pi) \stackrel{\text{def}}{=} \log \left\{ \omega(\pi) \right\}$$

Figure 3.3 shows the shape of the logit() function.

```
logit = function(p) log(odds(p))

logit_plot =
    ggplot() +
    geom_function(
        fun = logit,
        arrow = arrow(ends = "both")) +
    xlim(.001, .999) +
    ylab("logit(p)") +
    xlab("p") +
    theme_bw()
print(logit_plot)
```




Figure 3.3.: The logit function

Theorem 3.11.

$$logit(\pi) = log\left\{\frac{\pi}{1-\pi}\right\} \tag{3.4}$$

Exercise 3.15 (Compose the logit function). Prove Theorem 3.11.

Proof. Apply Definition 3.9 and then Definition 3.6 (details left to the reader). $\hfill\Box$

Theorem 3.12 (Derivative of logit function).

$$logit'(\pi) = \frac{1}{(\pi)(1-\pi)}$$

Proof.

$$\log it'(\pi) = \frac{\partial}{\partial \pi} \log it(\pi)$$

$$= \frac{\partial}{\partial \pi} \log \{\omega(\pi)\}$$

$$= \frac{\omega'(\pi)}{\omega(\pi)}$$

$$= \omega'(\pi) \frac{1}{\omega(\pi)}$$

$$= \frac{1}{(1-\pi)^2} \frac{1-\pi}{\pi}$$

$$= \frac{1}{(\pi)(1-\pi)}$$

3.5.2. The expit function

Definition 3.10 (expit, logistic, inverse-logit). The **expit function** (Figure 3.4) of a log-odds η , also known as the **inverse-logit function** or **logistic function**, is the inverse-odds of the exponential of η :

$$\operatorname{expit}(\eta) \stackrel{\operatorname{def}}{=} \operatorname{invodds} \left\{ \exp \left\{ \eta \right\} \right\}$$



Figure 3.4.: The expit function

Theorem 3.13 (logit and expit are each others' inverses).

$$logit\{expit\{\eta\}\}=\eta$$

$$expit\{logit\{\pi\}\}=\pi$$

Proof. Left to the reader.

Theorem 3.14 (Expressions for expit function).

$$\begin{split} \mathit{expit}(\eta) &= \frac{\mathit{exp}\left\{\eta\right\}}{1 + \mathit{exp}\left\{\eta\right\}} \\ &= (1 + \mathit{exp}\left\{-\eta\right\})^{-1} \end{split}$$

Proof. Apply definitions and Lemma 3.2. Details left to the reader. \Box

Lemma 3.4.

$$1-\exp{\{\eta\}}=\left(1+\exp{\{\eta\}}\right)^{-1}$$

Proof. Using Lemma 3.3:

$$\begin{aligned} 1 - \operatorname{expit} \left\{ \eta \right\} &= 1 - \operatorname{invodds} \left\{ \exp \left\{ \eta \right\} \right\} \\ &= \frac{1}{1 + \exp \left\{ \eta \right\}} \\ &= \left(1 + \exp \left\{ \eta \right\} \right)^{-1} \end{aligned}$$

Lemma 3.5.

$$\mathit{expit}'\left\{\eta\right\} = (\mathit{expit}\left\{\eta\right\})(1-\mathit{expit}\left\{\eta\right\})$$

Proof. Using Theorem 3.8:

$$\begin{split} & \operatorname{expit}'\left\{\eta\right\} = \frac{\partial}{\partial \eta} \operatorname{expit}\left\{\eta\right\} \\ & = \frac{\partial}{\partial \eta} \operatorname{invodds}\left\{\exp\left\{\eta\right\}\right\} \\ & = \operatorname{invodds}'\left\{\exp\left\{\eta\right\}\right\} \frac{\partial}{\partial \eta} \exp\left\{\eta\right\} \\ & = \frac{1}{\left(1 + \exp\left\{\eta\right\}\right)^2} \exp\left\{\eta\right\} \\ & = \frac{\exp\left\{\eta\right\}}{\left(1 + \exp\left\{\eta\right\}\right)^2} \\ & = \frac{\exp\left\{\eta\right\}}{1 + \exp\left\{\eta\right\}} \frac{1}{1 + \exp\left\{\eta\right\}} \\ & = \operatorname{expit}\left\{\eta\right\} \left(1 - \operatorname{expit}\left\{\eta\right\}\right) \end{split}$$

Proof. Alternatively, we can use Theorem 3.14:

$$\begin{split} & \exp \operatorname{it}' \left\{ \eta \right\} = \frac{\partial}{\partial \eta} \operatorname{expit} \left\{ \eta \right\} \\ & = \frac{\partial}{\partial \eta} (1 + \exp \left\{ -\eta \right\})^{-1} \\ & = -(1 + \exp \left\{ -\eta \right\})^{-2} \frac{\partial}{\partial \eta} (1 + \exp \left\{ -\eta \right\}) \\ & = -(1 + \exp \left\{ -\eta \right\})^{-2} (-\exp \left\{ -\eta \right\}) \\ & = (1 + \exp \left\{ -\eta \right\})^{-2} (\exp \left\{ -\eta \right\}) \\ & = (1 + \exp \left\{ -\eta \right\})^{-1} \frac{\exp \left\{ -\eta \right\}}{1 + \exp \left\{ -\eta \right\}} \\ & = (1 + \exp \left\{ -\eta \right\})^{-1} \frac{1}{1 + \exp \left\{ \eta \right\}} \\ & = \operatorname{expit} \left\{ \eta \right\} (1 - \operatorname{expit} \left\{ \eta \right\}) \end{split}$$

3.5.3. Diagram of expit and logit

$$\left[\pi \stackrel{\text{def}}{=} \Pr(Y=1)\right] \underbrace{\xrightarrow[1+\omega]{\frac{\pi}{1-\pi}}}_{\text{expit}(\eta)} \left[\omega \stackrel{\text{def}}{=} \operatorname{odds}(Y=1)\right] \xrightarrow[\exp \operatorname{it}(\eta)]{\log\{\omega\}}}_{\text{expit}(\eta)} \left[\eta \stackrel{\text{def}}{=} \operatorname{log-odds}(Y=1)\right]$$

3.6. Introduction to logistic regression

- In Example 3.1, we estimated the risk and the odds of MI for two groups, defined by oral contraceptive use.
- If the predictor is quantitative (dose) or there is more than one predictor, the task becomes more difficult.

• In this case, we will use logistic regression, which is a generalization of the linear regression models you have been using that can account for a binary response instead of a continuous one.

3.6.1. Binary outcomes models - one group, no covariates

$$\begin{split} \mathbf{P}(Y=1) &= \pi \\ \mathbf{P}(Y=0) &= 1 - \pi \\ \mathbf{P}(Y=y) &= \pi^y (1-\pi)^{1-y} \\ \mathbf{y} &= (y_1, ..., y_n) \\ \mathcal{L}(\pi; \mathbf{y}) &= \pi^{\sum y_i} (1-\pi)^{n-\sum y_i} \\ \ell(\pi, \mathbf{y}) &= \left(\sum y_i\right) \log\left\{\pi\right\} + \left(n - \sum y_i\right) \log\left\{1 - \pi\right\} \\ &= \left(\sum y_i\right) \left(\log\left\{\pi\right\} - \log\left\{1 - \pi\right\}\right) + n \cdot \log\left\{1 - \pi\right\} \\ &= \left(\sum y_i\right) \log\left\{\frac{\pi}{1-\pi}\right\} + n \cdot \log\left\{1 - \pi\right\} \\ &= \left(\sum y_i\right) \operatorname{logit}(\pi) + n \cdot \log\left\{1 - \pi\right\} \end{split}$$

3.6.2. Binary outcomes - general

$$\begin{split} \mathbf{P}(Y_i = 1) &= \pi_i \\ \mathbf{P}(Y_i = 0) &= 1 - \pi_i \end{split}$$

$$\mathbf{P}(Y_i = y_i) &= (\pi_i)_i^y (1 - \pi_i)^{1 - y_i}$$

$$\mathcal{L}_i(\pi_i) &= \mathbf{P}(Y_i = y_i)$$

$$\ell_i(\pi_i) = \log \left\{ \mathcal{L}_i(\pi_i) \right\} \\ &= y_i \log \left\{ \pi_i \right\} + (1 - y_i) \log \left\{ 1 - \pi_i \right\} \end{split}$$

For iid data $\tilde{y}=(y_1,...,y_n)$:

$$\begin{split} \mathcal{L}(\pi; \tilde{y}) &= \mathbf{P}(Y_1 = y_1, \dots, Y_n = y_n) \\ &= \prod_{i=1}^n \mathbf{P}(Y_i = y_i) \\ &= \prod_{i=1}^n \mathcal{L}_i(\pi_i) \end{split}$$

3.6.3. Modeling $\boldsymbol{\pi}_i$ as a function of \boldsymbol{X}_i

If there are only a few distinct X_i values, we can model π_i separately for each value of X_i .

Otherwise, we need regression.

$$\pi(x) \equiv \mathbf{E}(Y = 1|X = x)$$
$$= f(x^{\top}\beta)$$

Typically, we use the expit inverse-link:

$$\pi(\tilde{x}) = \operatorname{expit}(\tilde{x}'\beta) \tag{3.5}$$

3.6.4. Meet the beetles

```
library(glmx)
data(BeetleMortality, package = "glmx")
beetles = BeetleMortality |>
 mutate(
   pct = died/n,
   survived = n - died
 )
plot1 =
 beetles |>
 ggplot(aes(x = dose, y = pct)) +
 geom_point(aes(size = n)) +
 xlab("Dose (log mg/L)") +
 ylab("Mortality rate (%)") +
  scale_y_continuous(labels = scales::percent) +
  scale_size(range = c(1,2)) +
 theme_bw(base_size = 18)
print(plot1)
```



Figure 3.5.: Mortality rates of adult flour beetles after five hours' exposure to gaseous carbon disulphide (Bliss 1935)

3.6.5. Why don't we use linear regression?

```
beetles_long = beetles |>
  reframe(
    .by = everything(),
  outcome = c(
    rep(1, times = died),
    rep(0, times = survived))) |>
  as_tibble()

lm1 = beetles_long |> lm(formula = outcome ~ dose)
```

```
f.linear = function(x) predict(lm1, newdata = data.frame(dose = x))
range1 = range(beetles$dose) + c(-.2, .2)

plot2 =
  plot1 +
  geom_function(
   fun = f.linear,
    aes(col = "Straight line")) +
  labs(colour="Model", size = "")

plot2 |> print()
```




Figure 3.6.: Mortality rates of adult flour beetles after five hours' exposure to gaseous carbon disulphide (Bliss 1935)

3.6.6. Zoom out






Figure 3.7.: Mortality rates of adult flour beetles after five hours' exposure to gaseous carbon disulphide (Bliss 1935)

3.6.7. log transformation of dose?

```
lm2 = beetles_long |> lm(formula = outcome ~ log(dose))
f.linearlog = function(x) predict(lm2, newdata = data.frame(dose = x))
plot3 = plot2 +
```

```
expand_limits(x = c(1.6, 2)) +
geom_function(fun = f.linearlog, aes(col = "Log-transform dose"))
(plot3 + expand_limits(x = c(1.6, 2))) |> print()
```




Figure 3.8.: Mortality rates of adult flour beetles after five hours' exposure to gaseous carbon disulphide (Bliss 1935)

3.6.8. Logistic regression

```
#| label: fig-beetles_5
#| fig-cap: "Mortality rates of adult flour beetles after five hours' exposure to ga
beetles_glm_grouped = beetles |>
```

```
glm(formula = cbind(died, survived) ~ dose, family = "binomial")
f = function(x)
  beetles_glm_grouped |>
  predict(newdata = data.frame(dose = x), type = "response")
plot4 = plot3 + geom_function(fun = f, aes(col = "Logistic regression"))
plot4 |> print()
```



- 56
- 58
- 60
- 62

Model

- Log-transform dose
- Logistic regression
- Straight line

3.6.9. Three parts to regression models

• What distribution does the outcome have for a specific subpopulation defined by covariates? (outcome model)

- How does the combination of covariates relate to the mean? (link function)
- How do the covariates combine? (linear predictor, interactions)

3.6.10. Logistic regression in R

```
beetles_glm_grouped =
  beetles |>
  glm(
    formula = cbind(died, survived) ~ dose,
    family = "binomial")

library(parameters)
beetles_glm_grouped |>
  parameters() |>
  print_md()
```

| Parameter | Log-Odds | SE | 95% CI | ${f z}$ | p |
|-------------|----------|------|------------------|---------|--------|
| (Intercept) | -60.72 | | (-71.44, -51.08) | | |
| dose | 34.27 | 2.91 | (28.85, 40.30) | 11.77 | < .001 |

Fitted values:

3.6.11. Individual observations

```
beetles_glm_ungrouped =
  beetles_long |>
  glm(
    formula = outcome ~ dose,
    family = "binomial")

beetles_glm_ungrouped |> parameters() |> print_md()
```

Table 3.4.: beetles data in long format

| 1 | | _ 7 | _ | | | | | |
|----|-----|-------------|-------------|-------------|-------------|------------------|-------------|--|
| | | s_long | | | | | | |
| #> | # A | tibbl | Le: 48 | 1 x 6 | | | | |
| #> | | dose | died | n | pct | ${\tt survived}$ | outcome | |
| #> | | <dbl></dbl> | <int></int> | <int></int> | <dbl></dbl> | <int></int> | <dbl></dbl> | |
| #> | 1 | 1.69 | 6 | 59 | 0.102 | 53 | 1 | |
| #> | 2 | 1.69 | 6 | 59 | 0.102 | 53 | 1 | |
| #> | 3 | 1.69 | 6 | 59 | 0.102 | 53 | 1 | |
| #> | 4 | 1.69 | 6 | 59 | 0.102 | 53 | 1 | |
| #> | 5 | 1.69 | 6 | 59 | 0.102 | 53 | 1 | |
| #> | 6 | 1.69 | 6 | 59 | 0.102 | 53 | 1 | |
| #> | 7 | 1.69 | 6 | 59 | 0.102 | 53 | 0 | |
| #> | 8 | 1.69 | 6 | 59 | 0.102 | 53 | 0 | |
| #> | 9 | 1.69 | 6 | 59 | 0.102 | 53 | 0 | |
| #> | 10 | 1.69 | 6 | 59 | 0.102 | 53 | 0 | |
| #> | # i | 471 n | nore r | OWS | | | | |

Table 3.5.: logistic regression model for beetles data with individual Bernoulli data

| Parameter | Log-Odds | SE | 95% CI | Z | p |
|------------------|-----------------|----|------------------------------------|---|---|
| (Intercept) dose | -60.72 34.27 | | (-71.44, -51.08) (28.85, 40.30) | | |

Here's the previous version again:

```
beetles_glm_grouped |> parameters() |> print_md()
```

Table 3.6.: logistic regression model for beetles data with grouped (binomial) data

| Parameter | Log-Odds | SE | 95% CI | ${f z}$ | p |
|------------------|-----------------|----|------------------------------------|---------|---|
| (Intercept) dose | -60.72 34.27 | | (-71.44, -51.08) (28.85, 40.30) | | |

They seem the same! But not quite:

```
logLik(beetles_glm_grouped)
#> 'log Lik.' -18.72 (df=2)
logLik(beetles_glm_ungrouped)
#> 'log Lik.' -186.2 (df=2)
```

The difference is due to the binomial coefficient $\binom{n}{x}$ which isn't included in the individual-observations (Bernoulli) version of the model.

3.7. Multiple logistic regression

3.7.1. Coronary heart disease (WCGS) study data

Let's use the data from the Western Collaborative Group Study (WCGS) (Rosenman et al. (1975)) to explore multiple logistic regression:

From Vittinghoff et al. (2012):

"The Western Collaborative Group Study (WCGS) was a large epidemiological study designed to investigate the association between the "type A" behavior pattern and coronary heart disease (CHD)".

From Wikipedia, "Type A and Type B personality theory":

"The hypothesis describes Type A individuals as outgoing, ambitious, rigidly organized, highly status-conscious, impatient, anxious, proactive, and concerned with time management....

The hypothesis describes Type B individuals as a contrast to those of Type A. Type B personalities, by definition, are noted to live at lower stress levels. They typically work steadily and may enjoy achievement, although they have a greater tendency to disregard physical or mental stress when they do not achieve."

3.7.1.1. Study design

from ?faraway::wcgs:

3154 healthy young men aged 39-59 from the San Francisco area were assessed for their personality type. All were free from coronary heart disease at the start of the research. Eight and a half years later change in CHD status was recorded.

Details (from faraway::wcgs)

The WCGS began in 1960 with 3,524 male volunteers who were employed by 11 California companies. Subjects were 39 to 59 years old and free of heart disease as determined by electrocardiogram. After the initial screening, the study population dropped to 3,154 and the number of companies to 10 because of various exclusions. The cohort comprised both blue- and white-collar employees.

3.7.2. Baseline data collection

socio-demographic characteristics:

- age
- education
- marital status
- income
- occupation
- physical and physiological including:
- height
- weight
- blood pressure
- electrocardiogram
- corneal arcus;

biochemical measurements:

- cholesterol and lipoprotein fractions;
- medical and family history and use of medications;

behavioral data:

- Type A interview,
- smoking,
- exercise
- alcohol use.

Later surveys added data on:

- anthropometry
- triglycerides
- Jenkins Activity Survey
- caffeine use

Average follow-up continued for 8.5 years with repeat examinations.

3.7.3. Load the data

Here, I load the data:

```
### load the data directly from a UCSF website:
library(haven)
url = paste0(
    # I'm breaking up the url into two chunks for readability
    "https://regression.ucsf.edu/sites/g/files/",
    "tkssra6706/f/wysiwyg/home/data/wcgs.dta")
wcgs = haven::read_dta(url)
```

Table 3.7.: wcgs data

```
wcgs |> head()
#> # A tibble: 6 x 22
       age arcus behpat
                          bmi chd69 chol
                                            dbp dibpat height
                                                                  id lnsbp lnwght
#>
     <dbl> <lgl> <fct> <dbl> <fct> <dbl> <fct> <dbl> <fct>
                                                         <dbl> <dbl> <dbl>
                                                                            <dbl>
#> 1
        50 TRUE A1
                                                            67 2343
                                                                     4.88
                                                                             5.30
                         31.3 No
                                      249
                                             90 Type A
#> 2
       51 FALSE A1
                         25.3 No
                                      194
                                             74 Type A
                                                           73
                                                               3656 4.79
                                                                             5.26
#> 3
       59 TRUE A1
                         28.7 No
                                      258
                                             94 Type A
                                                            70 3526 5.06
                                                                             5.30
#> 4
     51 TRUE A1
                                                            69 22057 4.84
                                                                             5.01
                         22.1 No
                                      173
                                             80 Type A
#> 5
       44 FALSE A1
                         22.3 No
                                      214
                                             80 Type A
                                                           71 12927 4.84
                                                                             5.08
                                             76 Type A
#> 6
     47 FALSE A1
                         27.1 No
                                      206
                                                            64 16029 4.75
                                                                             5.06
#> # i 10 more variables: ncigs <dbl>, sbp <dbl>, smoke <fct>, t1 <dbl>,
       time169 <dbl>, typchd69 <fct>, uni <dbl>, weight <dbl>, wghtcat <fct>,
#> #
       agec <fct>
```

3.7.4. Data cleaning

Now let's do some data cleaning

```
library(arsenal) # provides `set_labels()`
library(forcats) # provides `as_factor()`
library(haven)
library(plotly)
wcgs = wcgs |>
    mutate(
    age = age |>
        arsenal::set_labels("Age (years)"),

arcus =
    arcus |>
    as.logical() |>
```

```
arsenal::set_labels("Arcus Senilis"),
time169 =
  time169 |>
  as.numeric() |>
  arsenal::set_labels("Observation (follow up) time (days)"),
dibpat =
  dibpat |>
  as_factor() |>
  relevel(ref = "Type B") |>
  arsenal::set_labels("Behavioral Pattern"),
typchd69 = typchd69 |>
 labelled(
    label = "Type of CHD Event",
    labels =
      c(
        "None" = 0,
        "infdeath" = 1,
        "silent" = 2,
        "angina" = 3)),
# turn stata-style labelled variables in to R-style factors:
across(
  where (is.labelled),
  haven::as_factor)
```

3.7.5. What's in the data

Here's a table of the data:

```
wcgs |>
select(-c(id, uni, t1)) |>
tableby(chd69 ~ ., data = _) |>
summary(
    pfootnote = TRUE,
    title =
        "Baseline characteristics by CHD status at end of follow-up")
```

Table 3.8.: Baseline characteristics by CHD status at end of follow-up

| | No | Yes | Total | p |
|--------------------|--------------|----------------|------------|-------------|
| | (N=2897) | (N=257) | (N=3154) | value |
| Age (years) | | | | < |
| , , | | | | 0.001^{1} |
| Mean (SD) | 46.082 | 48.490 | 46.279 | |
| | (5.457) | (5.801) | (5.524) | |
| Range | 39.000 - | 39.000 - | 39.000 - | |
| | 59.000 | 59.000 | 59.000 | |
| Arcus Senilis | | | | < |
| | | | | 0.001^{2} |
| N-Miss | 0 | 2 | 2 | |
| FALSE | 2058 | 153 | 2211 | |
| | (71.0%) | (60.0%) | (70.1%) | |
| TRUE | 839 | 102 | 941 | |
| | (29.0%) | (40.0%) | (29.9%) | |
| Behavioral Pattern | , | , | , , | < |
| | | | | 0.001^{2} |
| A1 | 234 (8.1%) | 30 (11.7%) | 264 (8.4%) | |
| A2 | $1\dot{1}77$ | 148 | 1325 | |
| | (40.6%) | (57.6%) | (42.0%) | |
| B3 | 1155 | $61\ (23.7\%)$ | 1216 | |
| | (39.9%) | , | (38.6%) | |

| | No (N=2897) | Yes (N=257) | Total (N=3154) | p value |
|--------------------|-----------------|----------------|-------------------|-------------|
| B4 | 331 | 18 (7.0%) | 349 | |
| | (11.4%) | _ ((,,,,,, | (11.1%) | |
| Body Mass Index | (' ' ' ' ' ' ' | | (| < |
| (kg/m2) | | | | 0.001^{1} |
| Mean (SD) | 24.471 | 25.055 | 24.518 | |
| , | (2.561) | (2.579) | (2.567) | |
| Range | 11.191 - | 19.225 - | 11.191 - | |
| J | 37.653 | 38.947 | 38.947 | |
| Total Cholesterol | | | | < |
| | | | | 0.001^{1} |
| N-Miss | 12 | 0 | 12 | |
| Mean (SD) | 224.261 | 250.070 | 226.372 | |
| , | (42.217) | (49.396) | (43.420) | |
| Range | 103.000 - | 155.000 - | 103.000 - | |
| | 400.000 | 645.000 | 645.000 | |
| Diastolic Blood | | | | < |
| Pressure | | | | 0.001^{1} |
| Mean (SD) | 81.723 | 85.315 | 82.016 | |
| , , | (9.621) | (10.311) | (9.727) | |
| Range | 58.000 - | 64.000 - | 58.000 - | |
| _ | 150.000 | 122.000 | 150.000 | |
| Behavioral Pattern | | | | < |
| | | | | 0.001^{2} |
| Type B | 1486 | 79 (30.7%) | 1565 | |
| | (51.3%) | , | (49.6%) | |
| Type A | 1411 | 178 | 1589 | |
| • | (48.7%) | (69.3%) | (50.4%) | |
| Height (inches) | , , | , | , | 0.290^{1} |
| Mean (SD) | 69.764 | 69.938 | 69.778 | |
| . , | (2.539) | (2.410) | (2.529) | |

| | No (N=2897) | Yes (N=257) | Total (N=3154) | p value |
|--------------------|----------------|----------------|-------------------|-------------|
| Range | 60.000 - | 63.000 - | 60.000 - | |
| | 78.000 | 77.000 | 78.000 | |
| Ln of Systolic | | | | < |
| Blood Pressure | | | | 0.001^{1} |
| Mean (SD) | 4.846 | 4.900 | 4.850 | |
| , , | (0.110) | (0.125) | (0.112) | |
| Range | 4.585 - | 4.605 - | 4.585 - | |
| J | 5.438 | 5.298 | 5.438 | |
| Ln of Weight | | | | < |
| · · | | | | 0.001^{1} |
| Mean (SD) | 5.126 | 5.155 | 5.128 | |
| , | (0.123) | (0.118) | (0.123) | |
| Range | 4.357 - | 4.868 - | 4.357 - | |
| J | 5.670 | 5.768 | 5.768 | |
| Cigarettes per day | | | | < |
| - · | | | | 0.001^{1} |
| Mean (SD) | 11.151 | 16.665 | 11.601 | |
| , , | (14.329) | (15.657) | (14.518) | |
| Range | 0.000 - | 0.000 - | 0.000 - | |
| J | 99.000 | 60.000 | 99.000 | |
| Systolic Blood | | | | < |
| Pressure | | | | 0.001^{1} |
| Mean (SD) | 128.034 | 135.385 | 128.633 | |
| , , | (14.746) | (17.473) | (15.118) | |
| Range | 98.000 - | 100.000 - | 98.000 - | |
| | 230.000 | 200.000 | 230.000 | |
| Current smoking | | | | < |
| o . | | | | 0.001^{2} |
| No | 1554 | 98 (38.1%) | 1652 | |
| | (53.6%) | , | (52.4%) | |
| | ` / | | ` / | |

| | No (N. 2007) | Yes | Total | p |
|---------------------|--------------|-------------|------------|-------------|
| | (N=2897) | (N=257) | (N=3154) | value |
| Yes | 1343 | 159 | 1502 | |
| | (46.4%) | (61.9%) | (47.6%) | |
| Observation (follow | | | | < |
| up) time (days) | | | | 0.001^{1} |
| Mean (SD) | 2775.158 | 1654.700 | 2683.859 | |
| | (562.205) | (859.297) | (666.524) | |
| Range | 238.000 - | 18.000 - | 18.000 - | |
| | 3430.000 | 3229.000 | 3430.000 | |
| Type of CHD | | | | |
| Event | | | | |
| None | 0 (0.0%) | 0 (0.0%) | 0 (0.0%) | |
| infdeath | 2897 | 0(0.0%) | 2897 | |
| | (100.0%) | , | (91.9%) | |
| silent | 0 (0.0%) | 135 | 135 (4.3%) | |
| | , | (52.5%) | , | |
| angina | 0 (0.0%) | 71(27.6%) | 71 (2.3%) | |
| 4 | 0(0.0%) | 51 (19.8%) | 51 (1.6%) | |
| Weight (lbs) | , | , | , | < |
| 3 () | | | | 0.001^{1} |
| Mean (SD) | 169.554 | 174.463 | 169.954 | |
| , | (21.010) | (21.573) | (21.096) | |
| Range | 78.000 - | 130.000 - | 78.000 - | |
| G. | 290.000 | 320.000 | 320.000 | |
| Weight Category | | | | < |
| | | | | 0.001^2 |
| < 140 | 217 (7.5%) | 15 (5.8%) | 232 (7.4%) | 0.00 |
| 140-170 | 1440 | 98 (38.1%) | 1538 | |
| 110 110 | (49.7%) | 00 (00.170) | (48.8%) | |
| 170-200 | 1049 | 122 | 1171 | |
| =. | (36.2%) | (47.5%) | (37.1%) | |
| > 200 | 191 (6.6%) | 22 (8.6%) | 213 (6.8%) | |

| | No | Yes | Total | p |
|---------------|-----------------|----------------|----------------|-------------|
| | (N=2897) | (N=257) | (N=3154) | value |
| RECODE of age | | | | < |
| (Age) | | | | 0.001^{2} |
| 35-40 | 512 | $31\ (12.1\%)$ | 543 | |
| | (17.7%) | | (17.2%) | |
| 41-45 | 1036 | 55~(21.4%) | 1091 | |
| | (35.8%) | | (34.6%) | |
| 46-50 | 680 | 70~(27.2%) | 750 | |
| | (23.5%) | | (23.8%) | |
| 51-55 | 463 | 65~(25.3%) | 528 | |
| | (16.0%) | | (16.7%) | |
| 56-60 | $206 \ (7.1\%)$ | $36\ (14.0\%)$ | $242\ (7.7\%)$ | |

- 1. Linear Model ANOVA
- 2. Pearson's Chi-squared test

3.7.6. Data by age and personality type

For now, we will look at the interaction between age and personality type (dibpat). To make it easier to visualize the data, we summarize the event rates for each combination of age:

```
chd_grouped_data =
  wcgs |>
  summarize(
    .by = c(age, dibpat),
    n = sum(chd69 %in% c("Yes", "No")),
    x = sum(chd69 == "Yes")) |>
  mutate(
    `n - x` = n - x,
    `p(chd)` = (x / n) |>
```

```
labelled(label = "CHD Event by 1969"),
   \circ dds(chd) = p(chd) / (1 - p(chd)),
   `logit(chd)` = log(`odds(chd)`)
 )
chd_grouped_data
#> # A tibble: 42 x 8
      #>
    <dbl> <fct> <int> <int> <int> <dbl+lbl>
#>
                                            <dbl>
                                                       <dbl>
                                                       -2.14
#>
   1
      50 Type A
                76 8
                            68 0.105
                                            0.118
  2
#>
       51 Type A
                 67 11
                            56 0.164
                                            0.196
                                                       -1.63
                 30
                       7
                            23 0.233
#>
      59 Type A
                                            0.304
                                                       -1.19
                113 9
#> 4 44 Type A
                           104 0.0796
                                            0.0865
                                                       -2.45
#> 5 47 Type A
                72
                      7
                            65 0.0972
                                            0.108
                                                       -2.23
                          124 0.0677
101 0.0648
#> 6
      40 Type A
                133
                      9
                                            0.0726
                                                       -2.62
#> 7
      41 Type A
                108
                      7
                                            0.0693
                                                       -2.67
#> 8 43 Type A
                       7
                 97
                            90 0.0722
                                            0.0778
                                                       -2.55
                       7
#> 9
       54 Type A
                 53
                            46 0.132
                                            0.152
                                                       -1.88
#> 10
                       12
                           68 0.15
                                            0.176
                                                       -1.73
      48 Type A
                 80
#> # i 32 more rows
```

3.7.7. Graphical exploration

```
library(ggplot2)
library(ggeasy)
library(scales)
chd_plot_probs =
   chd_grouped_data |>
   ggplot(
   aes(
    x = age,
```

```
y = p(chd),
      col = dibpat)
  ) +
 geom_point(aes(size = n), alpha = .7) +
 scale_size(range = c(1,4)) +
 geom_line() +
 theme_bw() +
 ylab("P(CHD Event by 1969)") +
 scale_y_continuous(
   labels = scales::label_percent(),
   sec.axis = sec_axis(
     ~ odds(.),
     name = "odds(CHD Event by 1969)")) +
  ggeasy::easy_labs() +
 theme(legend.position = "bottom")
print(chd_plot_probs)
```



Figure 3.9.: CHD rates by age group, probability scale

3.7.7.1. Odds scale

```
trans_odds = trans_new(
  name = "odds",
  transform = odds,
  inverse = odds_inv)

chd_plot_odds = chd_plot_probs +
  scale_y_continuous(
    trans = trans_odds, # this line changes the vertical spacing
```




Figure 3.10.: CHD rates by age group, odds spacing

3.7.7.2. Log-odds (logit) scale

```
trans_logit = trans_new(
   name = "logit",
   transform = logit,
   inverse = expit)

chd_plot_logit =
   chd_plot_probs +
   scale_y_continuous(
     trans = trans_logit, # this line changes the vertical spacing
   name = chd_plot_probs$labels$y,
   breaks = c(seq(.01, .1, by = .01), .15, .2),
   minor_breaks = NULL,
   sec.axis = sec_axis(
     ~ logit(.),
     name = "log(odds(CHD Event by 1969))"))

print(chd_plot_logit)
```



Figure 3.11.: CHD data (logit-scale)

3.7.8. Logistic regression models for CHD data

Here, we fit stratified models for CHD by personality type.

```
chd_glm_strat = glm(
   "formula" = chd69 == "Yes" ~ dibpat + dibpat:age - 1,
   "data" = wcgs,
   "family" = binomial(link = "logit")
)
chd_glm_strat |> parameters() |> print_md()
```

Table 3.9.: CHD model, stratified parametrization

| | Log- | | | | |
|--------------------------|-------|------|----------------|---------|--------|
| Parameter | olds | SE | 95% CI | ${f z}$ | p |
| dibpat (Type B) | -5.80 | 0.98 | (-7.73, -3.90) | -5.95 | < .001 |
| dibpat (Type A) | -5.50 | 0.67 | (-6.83, -4.19) | -8.18 | < .001 |
| dibpat (Type B) \times | 0.06 | 0.02 | (0.02, 0.10) | 3.01 | 0.003 |
| age | | | | | |
| dibpat (Type A) \times | 0.07 | 0.01 | (0.05, 0.10) | 5.24 | < .001 |
| age | | | | | |

We can get the corresponding odds ratios $(e^{\beta}s)$ by passing exponentiate = TRUE to parameters():

```
chd_glm_strat |>
  parameters(exponentiate = TRUE) |>
  print_md()
```

Table 3.10.: Odds ratio estimates for CHD model

| | Odds | | | | |
|-----------------|----------|--------|--------------|---------|-------|
| Parameter | Ratio | SE | 95% CI | ${f z}$ | p |
| dibpat (Type B) | 3.02e-03 | 2.94e- | (4.40e-04, | -5.95 | < |
| | | 03 | 0.02) | | .001 |
| dibpat (Type A) | 4.09e-03 | 2.75e- | (1.08e-03, | -8.18 | < |
| | | 03 | 0.02) | | .001 |
| dibpat (Type B) | 1.06 | 0.02 | (1.02, 1.11) | 3.01 | 0.003 |
| \times age | | | | | |
| dibpat (Type A) | 1.07 | 0.01 | (1.05, 1.10) | 5.24 | < |
| \times age | | | | | .001 |


3.7.9. Models superimposed on data

We can graph our fitted models on each scale (probability, odds, log-odds).

3.7.9.1. probability scale

```
curve_type_A = function(x)
  chd_glm_strat |> predict(
    type = "response",
    newdata = tibble(age = x, dibpat = "Type A"))
}
curve_type_B = function(x)
  chd_glm_strat |> predict(
    type = "response",
    newdata = tibble(age = x, dibpat = "Type B"))
}
chd_plot_probs_2 =
  chd_plot_probs +
 geom_function(
   fun = curve_type_A,
    aes(col = "Type A")
  ) +
  geom_function(
    fun = curve_type_B,
```

```
aes(col = "Type B")
)
print(chd_plot_probs_2)
```



3.7.9.2. odds scale

```
# curve_type_A = function(x)
# {
# chd_glm_strat |> predict(
# type = "link",
# newdata = tibble(age = x, dibpat = "Type A")) |> exp()
```

```
# }
# curve_type_B = function(x)
# {
# chd_glm_strat |> predict(
     type = "link",
     newdata = tibble(age = x, dibpat = "Type B")) |> exp()
# }
chd_plot_odds_2 =
 chd_plot_odds +
 geom_function(
   fun = curve_type_A,
   aes(col = "Type A")
  ) +
 geom_function(
   fun = curve_type_B,
   aes(col = "Type B")
print(chd_plot_odds_2)
```



Figure 3.12.

3.7.9.3. log-odds (logit) scale

```
chd_plot_logit_2 =
  chd_plot_logit +
  geom_function(
    fun = curve_type_A,
    aes(col = "Type A")
) +
  geom_function(
  fun = curve_type_B,
```

```
aes(col = "Type B")
)
print(chd_plot_logit_2)
```




Figure 3.13.

3.7.10. reference-group and contrast parametrization

We can also use the corner-point parametrization (with reference groups and contrasts):

```
chd_glm_contrasts =
  wcgs |>
```

```
glm(
    "data" = _,
    "formula" = chd69 == "Yes" ~ dibpat*I(age - 50),
    "family" = binomial(link = "logit")
chd_glm_contrasts |>
  parameters() |>
  print_md()
```

Table 3.11.: CHD model (corner-point parametrization)

| | Log- | | | | |
|--------------------------|-------|------|----------------|---------|-------|
| Parameter | Odds | SE | 95% CI | ${f z}$ | p |
| (Intercept) | -2.73 | 0.13 | (-2.98, -2.49) | -21.45 | < |
| | | | | | .001 |
| dibpat (Type A) | 0.82 | 0.15 | (0.53, 1.13) | 5.42 | < |
| | | | | | .001 |
| age - 50 | 0.06 | 0.02 | (0.02, 0.10) | 3.01 | 0.003 |
| dibpat (Type A) \times | 0.01 | 0.02 | (-0.04, 0.06) | 0.42 | 0.674 |
| age - 50 | | | | | |

Compare with Table 3.10.

Exercise 3.16. If I give you model 1, how would you get the coefficients of model 2?

Theorem 3.15. For the logistic regression model:

$$\begin{array}{ll} \bullet & Y_i | \tilde{X}_i & \sim & \mathit{Ber}(\pi(\tilde{X}_i)) \\ \bullet & \pi(\tilde{x}) = \mathit{expit}\left\{ \tilde{x}' \tilde{\beta} \right\} \end{array}$$

Consider two covariate patterns, \tilde{x} and $\tilde{x^*}$.

The odds ratio comparing these covariate patterns is:

$$\omega(\tilde{\boldsymbol{x}}, \tilde{\boldsymbol{x^*}}) = \exp\left\{(\tilde{\boldsymbol{x}} - \tilde{\boldsymbol{x^*}})^\top \tilde{\boldsymbol{\beta}}\right\}$$

Proof.

$$\begin{split} \omega(\tilde{x}, \tilde{x^*}) &= \frac{\omega(Y = 1 | \tilde{X} = \tilde{x})}{\omega(Y = 1 | \tilde{X} = \tilde{x^*})} \\ &= \frac{\exp\left\{\tilde{x}^\top \tilde{\beta}\right\}}{\exp\left\{\tilde{x^*}^\top \tilde{\beta}\right\}} \\ &= \exp\left\{\tilde{x}^\top \tilde{\beta} - \tilde{x^*}^\top \tilde{\beta}\right\} \\ &= \exp\left\{(\tilde{x}^\top - \tilde{x^*}^\top) \tilde{\beta}\right\} \\ &= \exp\left\{(\tilde{x} - \tilde{x^*})^\top \tilde{\beta}\right\} \end{split}$$

3.8. Fitting logistic regression models

3.8.1. Maximum likelihood estimation for ciid data

Assume:

- $\begin{array}{ll} \bullet & Y_i | \tilde{X}_i \ \sim & \mathrm{Ber}(\pi(X_i)) \\ \bullet & \pi(\tilde{x}) = \mathrm{expit} \left\{ \tilde{x}' \tilde{\beta} \right\} \end{array}$

3.8.1.1. log-likelihood function

$$\begin{split} \ell(\tilde{\beta}, \tilde{y}) &= \log \left\{ \mathcal{L}(\tilde{\beta}, \tilde{y}) \right\} \\ &= \sum_{i=1}^{n} \underline{\ell_i}(\pi(\tilde{x}_i)) \end{split} \tag{3.6}$$

$$\begin{split} & \ell_i(\pi) = y_i \log \left\{ \pi \right\} + (1 - y_i) \log \left\{ 1 - \pi \right\} \\ & = y_i \log \left\{ \pi \right\} + (1 \cdot \log \left\{ 1 - \pi \right\} - y_i \cdot \log \left\{ 1 - \pi \right\}) \\ & = y_i \log \left\{ \pi \right\} + (\log \left\{ 1 - \pi \right\} - y_i \log \left\{ 1 - \pi \right\}) \\ & = y_i \log \left\{ \pi \right\} + \log \left\{ 1 - \pi \right\} - y_i \log \left\{ 1 - \pi \right\} \\ & = y_i \log \left\{ \pi \right\} - y_i \log \left\{ 1 - \pi \right\} + \log \left\{ 1 - \pi \right\} \\ & = (y_i \log \left\{ \pi \right\} - y_i \log \left\{ 1 - \pi \right\}) + \log \left\{ 1 - \pi \right\} \\ & = y_i (\log \left\{ \pi \right\} - \log \left\{ 1 - \pi \right\}) + \log \left\{ 1 - \pi \right\} \\ & = y_i (\log \left\{ \frac{\pi}{1 - \pi} \right\}) + \log \left\{ 1 - \pi \right\} \\ & = y_i (\log (\pi)) + \log \left\{ 1 - \pi \right\} \end{split}$$

3.8.1.2. score function

$$\begin{split} \ell'(\tilde{\beta}) & \stackrel{\text{def}}{=} \frac{\partial}{\partial \tilde{\beta}} \ell(\tilde{\beta}) \\ & = \frac{\partial}{\partial \tilde{\beta}} \sum_{i=1}^n \ell_i(\tilde{\beta}) \\ & = \sum_{i=1}^n \frac{\partial}{\partial \tilde{\beta}} \ell_i(\tilde{\beta}) \\ & = \sum_{i=1}^n \ell_i'(\tilde{\beta}) \end{split}$$

$$\begin{split} \ell_i'(\tilde{\beta}) &= \frac{\partial}{\partial \tilde{\beta}} y_i \left(\operatorname{logit} \left\{ \pi_i \right\} \right) + \operatorname{log} \left\{ 1 - \pi_i \right\} \\ &= \frac{\partial}{\partial \tilde{\beta}} \left\{ y_i \left(\tilde{x}_i' \tilde{\beta} \right) + \operatorname{log} \left\{ 1 - \pi_i \right\} \right\} \\ &= \left\{ y_i \frac{\partial}{\partial \tilde{\beta}} \left(\tilde{x}_i' \tilde{\beta} \right) + \frac{\partial}{\partial \tilde{\beta}} \operatorname{log} \left\{ 1 - \pi_i \right\} \right\} \\ &= \left\{ \tilde{x}_i y_i + \frac{\partial}{\partial \tilde{\beta}} \operatorname{log} \left\{ 1 - \operatorname{expit} (\tilde{x}_i' \tilde{\beta}) \right\} \right\} \\ &= \left\{ \tilde{x}_i y_i + \frac{\partial}{\partial \tilde{\beta}} \operatorname{log} \left\{ \left(1 + \operatorname{exp} \left\{ \tilde{x}_i' \tilde{\beta} \right\} \right)^{-1} \right\} \right\} \\ &= \left\{ \tilde{x}_i y_i - \frac{\partial}{\partial \tilde{\beta}} \operatorname{log} \left\{ 1 + \operatorname{exp} \left\{ \tilde{x}_i' \tilde{\beta} \right\} \right\} \right\} \end{split}$$

Now we need to apply the chain rule:

$$\begin{split} \frac{\partial}{\partial \beta} \log \left\{ 1 + \exp \left\{ \tilde{x}_i' \beta \right\} \right\} &= \frac{1}{1 + \exp \left\{ \tilde{x}_i' \beta \right\}} \frac{\partial}{\partial \beta} \left\{ 1 + \exp \left\{ \tilde{x}_i' \beta \right\} \right\} \\ &= \frac{\partial}{\partial \beta} \left\{ 1 + \exp \left\{ \tilde{x}_i' \beta \right\} \right\} = \exp \left\{ \tilde{x}_i' \beta \right\} \\ &= \tilde{x}_i \exp \left\{ \tilde{x}_i' \beta \right\} \end{split}$$

So:

$$\begin{split} \frac{\partial}{\partial \beta} \log \left\{ 1 + \exp \left\{ \tilde{x}_i' \beta \right\} \right\} &= \frac{1}{1 + \exp \left\{ \tilde{x}_i' \beta \right\}} \exp \left\{ \tilde{x}_i' \beta \right\} \tilde{x}_i \\ &= \frac{\exp \left\{ \tilde{x}_i' \beta \right\}}{1 + \exp \left\{ \tilde{x}_i' \beta \right\}} \tilde{x}_i \\ &= \tilde{x}_i \mathrm{expit} \left\{ \tilde{x}_i' \beta \right\} \end{split}$$

So:

$$\begin{split} \ell_i'(\tilde{\beta}) &= \tilde{x}_i y_i - \tilde{x}_i \mathrm{expit} \left\{ \tilde{x}_i' \beta \right\} \\ &= \tilde{x}_i (y_i - \mathrm{expit} \left\{ \tilde{x}_i' \beta \right\}) \\ &= \tilde{x}_i (y_i - \pi_i) \\ &= \tilde{x}_i (y_i - \mathbb{E}[Y_i | \tilde{X}_i = \tilde{x}_i]) \\ &= \tilde{x}_i \; \varepsilon(y_i | \tilde{X}_i = \tilde{x}_i) \end{split}$$

This last expression is essentially the same as we found in linear regression.

Putting the pieces of $\ell'(\tilde{\beta})$ back together, we have:

$$\ell'(\tilde{\beta}) = \sum_{i=1}^n \left\{ \tilde{x}_i(y_i - \operatorname{expit}\left\{\tilde{x}_i'\beta\right\}) \right\}$$

Setting $\ell'(\tilde{\beta}; \tilde{y}) = 0$ gives us:

$$\sum_{i=1}^{n} \{ \tilde{x}_i(y_i - \text{expit} \{ \tilde{x}_i' \beta \}) \} = 0$$
 (3.7)

In general, the estimating equation $\ell'(\tilde{\beta};\tilde{y})=0$ cannot be solved analytically.

Instead, we can use the Newton-Raphson method:

$$\hat{\theta}^* \leftarrow \hat{\theta}^* - \left(\ell''\left(\tilde{y}; \hat{\theta}^*\right)\right)^{-1} \ell'\left(\tilde{y}; \hat{\theta}^*\right)$$

We make an iterative series of guesses, and each guess helps us make the next guess better (i.e., higher log-likelihood). You can see some information about this process like so:

```
options(digits = 8)
temp =
  wcgs |>
  glm(
    control = glm.control(trace = TRUE),
    data = _,
  formula = chd69 == "Yes" ~ dibpat*age,
  family = binomial(link = "logit")
)
```

```
#> Deviance = 1775.7899 Iterations - 1
#> Deviance = 1708.5396 Iterations - 2
#> Deviance = 1704.0434 Iterations - 3
#> Deviance = 1703.9833 Iterations - 4
#> Deviance = 1703.9832 Iterations - 5
#> Deviance = 1703.9832 Iterations - 6
```

After each iteration of the fitting procedure, the deviance $(2(\ell_{\text{full}} - \ell(\hat{\beta})))$ is printed. You can see that the algorithm took six iterations to converge to a solution where the likelihood wasn't changing much anymore.

3.9. Model comparisons for logistic models

3.9.1. Deviance test

We can compare the maximized log-likelihood of our model, $\ell(\hat{\beta}; \mathbf{x})$, versus the log-likelihood of the full model (aka saturated model aka maximal model), ℓ_{full} , which has one parameter per covariate pattern. With enough data, $2(\ell_{\text{full}} - \ell(\hat{\beta}; \mathbf{x})) \dot{\sim} \chi^2(N-p)$, where N is the number of distinct covariate patterns and p is the number of β parameters in our model. A significant p-value for this **deviance** statistic indicates that there's some detectable pattern in the data that our model isn't flexible enough to catch.

Caution

The deviance statistic needs to have a large amount of data for each covariate pattern for the χ^2 approximation to hold. A guideline from Dobson is that if there are q distinct covariate patterns $x_1...,x_q$, with $n_1,...,n_q$ observations per pattern, then the expected frequencies $n_k \cdot \pi(x_k)$ should be at least 1 for every pattern $k \in 1:q$.

If you have covariates measured on a continuous scale, you may not be able to use the deviance tests to assess goodness of fit.

3.9.2. Hosmer-Lemeshow test

If our covariate patterns produce groups that are too small, a reasonable solution is to make bigger groups by merging some of the covariate-pattern groups together.

Hosmer and Lemeshow (1980) proposed that we group the patterns by their predicted probabilities according to the model of interest. For example, you could group all of the observations with predicted probabilities of 10% or less together, then group the observations with 11%-20% probability together, and so on; g = 10 categories in all.

Then we can construct a statistic

$$X^2 = \sum_{c=1}^g \frac{(o_c - e_c)^2}{e_c}$$

where o_c is the number of events observed in group c, and e_c is the number of events expected in group c (based on the sum of the fitted values $\hat{\pi}_i$ for observations in group c).

If each group has enough observations in it, you can compare X^2 to a χ^2 distribution; by simulation, the degrees of freedom has been found to be approximately g-2.

For our CHD model, this procedure would be:

```
wcgs =
wcgs |>
mutate(
   pred_probs_glm1 = chd_glm_strat |> fitted(),
   pred_prob_cats1 =
```

```
pred_probs_glm1 |>
    cut(breaks = seq(0, 1, by = .1),
        include.lowest = TRUE))

HL_table =
    wcgs |>
    summarize(
        .by = pred_prob_cats1,
        n = n(),
        o = sum(chd69 == "Yes"),
        e = sum(pred_probs_glm1)
    )

library(pander)
HL_table |> pander()
```

| pred_prob_cats1 | n | 0 | е |
|-----------------|-----------|-----|-------|
| (0.1, 0.2] | 785 | 116 | 108 |
| (0.2, 0.3] | 64 | 12 | 13.77 |
| [0,0.1] | $2,\!305$ | 129 | 135.2 |

Our statistic is $X^2=1.11028711;\ p(\chi^2(1)>1.11028711)=0.29201955,$ which is our p-value for detecting a lack of goodness of fit.

Unfortunately that grouping plan left us with just three categories with any observations, so instead of grouping by 10% increments of predicted probability, typically analysts use deciles of the predicted probabilities:

```
wcgs =
  wcgs |>
  mutate(
    pred_probs_glm1 = chd_glm_strat |> fitted(),
    pred_prob_cats1 =
      pred_probs_glm1 |>
      cut(breaks = quantile(pred_probs_glm1, seq(0, 1, by = .1)),
          include.lowest = TRUE))
HL_table =
  wcgs |>
  summarize(
    .by = pred_prob_cats1,
   n = n(),
    o = sum(chd69 == "Yes"),
    e = sum(pred_probs_glm1)
HL_table |> pander()
```

| pred_prob_cats1 | n | О | e |
|------------------|-----|----|-------|
| (0.114, 0.147] | 275 | 48 | 36.81 |
| (0.147, 0.222] | 314 | 51 | 57.19 |
| (0.0774, 0.0942] | 371 | 27 | 32.56 |
| (0.0942, 0.114] | 282 | 30 | 29.89 |
| (0.0633, 0.069] | 237 | 17 | 15.97 |

| n | 0 | е |
|-----|--------------------------|--------------------------------------|
| 306 | 20 | 22.95 |
| 413 | 27 | 24.1 |
| 310 | 14 | 14.15 |
| 407 | 16 | 13.91 |
| 239 | 7 | 9.48 |
| | 306 413 310 407 | 306 20 413 27 310 14 407 16 |


Now we have more evenly split categories. The p-value is 0.56041994, still not significant.

Graphically, we have compared:

```
HL_plot =
HL_table |>
ggplot(aes(x = pred_prob_cats1)) +
geom_line(aes(y = e, x = pred_prob_cats1, group = "Expected", col = "Expected")) +
geom_point(aes(y = e, size = n, col = "Expected")) +
geom_point(aes(y = o, size = n, col = "Observed")) +
geom_line(aes(y = o, col = "Observed", group = "Observed")) +
scale_size(range = c(1,4)) +
```

```
theme_bw() +
ylab("number of CHD events") +
theme(axis.text.x = element_text(angle = 45))
```

print(HL_plot)



pred_prob_cats1

3.9.3. Comparing models

- AIC = $-2 * \ell(\hat{\theta}) + 2 * p$ [lower is better]
- BIC = $-2 * \ell(\hat{\theta}) + p * \log(n)$ [lower is better]
- likelihood ratio [higher is better]

3.10. Residual-based diagnostics

3.10.1. Logistic regression residuals only work for grouped data

Residuals only work if there is more than one observation for most covariate patterns.

Here we will create the grouped-data version of our CHD model from the WCGS study:

```
wcgs_grouped =
wcgs |>
summarize(
    .by = c(dibpat, age),
    n = n(),
    chd = sum(chd69 == "Yes"),
    `!chd` = sum(chd69 == "No")
)

chd_glm_strat_grouped = glm(
    "formula" = cbind(chd, `!chd`) ~ dibpat + dibpat:age - 1,
    "data" = wcgs_grouped,
    "family" = binomial(link = "logit")
)

chd_glm_strat_grouped |> parameters() |> print_md()
```

| Parameter | Log- Odds | SE | 95% CI | Z | р |
|-----------------|--------------|------|----------------|-------|--------|
| dibpat (Type B) | -5.80 | 0.98 | (-7.73, -3.90) | -5.95 | < .001 |
| dibpat (Type A) | -5.50 | 0.67 | (-6.83, -4.19) | -8.18 | < .001 |

| | Log- | | | | |
|--|------|------|--------------|---------|--------|
| Parameter | Odds | SE | 95% CI | ${f z}$ | p |
| $\overline{\text{dibpat (Type B)} \times}$ | 0.06 | 0.02 | (0.02, 0.10) | 3.01 | 0.003 |
| age dibpat (Type A) \times age | 0.07 | 0.01 | (0.05, 0.10) | 5.24 | < .001 |

3.10.2. (Response) residuals

$$e_k \stackrel{\mathrm{def}}{=} \bar{y}_k - \hat{\pi}(x_k)$$

(k indexes the covariate patterns)

We can graph these residuals e_k against the fitted values $\hat{\pi}(x_k)$:

```
wcgs_grouped =
  wcgs_grouped |>
 mutate(
    fitted = chd_glm_strat_grouped |> fitted(),
    fitted_logit = fitted |> logit(),
    response_resids =
      chd_glm_strat_grouped |> resid(type = "response")
  )
wcgs_response_resid_plot =
  wcgs_grouped |>
  ggplot(
   mapping = aes(
     x = fitted,
      y = response_resids
    )
  ) +
```

```
geom_point(
   aes(col = dibpat)
) +
geom_hline(yintercept = 0) +
geom_smooth(
   se = TRUE,
   method.args = list(
      span=2/3,
      degree=1,
      family="symmetric",
      iterations=3),
   method = stats::loess)
```

① Don't worry about these options for now; I chose them to match autoplot() as closely as I can. plot.glm and autoplot use stats::lowess instead of stats::lowess; stats::lowess is older, hard to use with geom_smooth, and hard to match exactly with stats::lowess; see https://support.bioconductor.org/p/2323/.]

```
wcgs_response_resid_plot |> print()
```



We can see a slight fan-shape here: observations on the right have larger variance (as expected since $var(\bar{y}) = \pi(1-\pi)/n$ is maximized when $\pi=0.5$).

3.10.3. Pearson residuals

The fan-shape in the response residuals plot isn't necessarily a concern here, since we haven't made an assumption of constant residual variance, as we did for linear regression.

However, we might want to divide by the standard error in order to make the graph easier to interpret. Here's one way to do that:

The Pearson (chi-squared) residual for covariate pattern k is:

$$X_k = \frac{\bar{y}_k - \hat{\pi}_k}{\sqrt{\hat{\pi}_k (1 - \hat{\pi}_k)/n_k}}$$

where

$$\begin{split} \hat{\pi}_k & \stackrel{\text{def}}{=} \hat{\pi}(x_k) \\ & \stackrel{\text{def}}{=} \hat{P}(Y = 1 | X = x_k) \\ & \stackrel{\text{def}}{=} \text{expit}(x_i' \hat{\beta}) \\ & \stackrel{\text{def}}{=} \text{expit}(\hat{\beta}_0 + \sum_{i=1}^p \hat{\beta}_j x_{ij}) \end{split}$$

Let's take a look at the Pearson residuals for our CHD model from the WCGS data (graphed against the fitted values on the logit scale):

```
library(ggfortify)
```

autoplot(chd_glm_strat_grouped, which = 1, ncol = 1) |> print()

Residuals vs Fitted



The fan-shape is gone, and these residuals don't show any obvious signs of model fit issues.

3.10.3.1. Pearson residuals plot for beetles data

If we create the same plot for the beetles model, we see some strong evidence of a lack of fit:

autoplot(beetles_glm_grouped, which = 1, ncol = 1) |> print()

Residuals vs Fitted



3.10.3.2. Pearson residuals with individual (ungrouped) data

What happens if we try to compute residuals without grouping the data by covariate pattern?

library(ggfortify)

autoplot(chd_glm_strat, which = 1, ncol = 1) |> print()

Residuals vs Fitted



Meaningless.

3.10.3.3. Residuals plot by hand (optional section)

If you want to check your understanding of what these residual plots are, try building them yourself:

```
wcgs_grouped =
 wcgs_grouped |>
 mutate(
   fitted = chd_glm_strat_grouped |> fitted(),
   fitted_logit = fitted |> logit(),
   resids = chd_glm_strat_grouped |> resid(type = "pearson")
wcgs_resid_plot1 =
 wcgs_grouped |>
 ggplot(
   mapping = aes(
     x = fitted_logit,
     y = resids
    )
  ) +
 geom_point(
   aes(col = dibpat)
  geom_hline(yintercept = 0) +
  geom_smooth(se = FALSE,
              method.args = list(
                span=2/3,
```

```
degree=1,
    family="symmetric",
    iterations=3,
    surface="direct"
    # span = 2/3,
    # iterations = 3
    ),
    method = stats::loess)

# plot.glm and autoplot use stats::lowess, which is hard to use with
# geom_smooth and hard to match exactly;
# see https://support.bioconductor.org/p/2323/
```



3.10.4. Pearson chi-squared goodness of fit test

The Pearson chi-squared goodness of fit statistic is:

$$X^2 = \sum_{k=1}^{m} X_k^2$$

Under the null hypothesis that the model in question is correct (i.e., sufficiently complex), $X^2 \sim \chi^2(N-p)$.

```
X = chd_glm_strat_grouped |>
  resid(type = "pearson")

chisq_stat = sum(X^2)

pval = pchisq(
  chisq_stat,
  lower = FALSE,
  df = length(X) - length(coef(chd_glm_strat_grouped)))
```

For our CHD model, the p-value for this test is 0.26523556; no significant evidence of a lack of fit at the 0.05 level.

3.10.4.1. Standardized Pearson residuals

Especially for small data sets, we might want to adjust our residuals for leverage (since outliers in X add extra variance to the residuals):

$$r_{P_k} = \frac{X_k}{\sqrt{1-h_k}}$$

where h_k is the leverage of X_k . The functions autoplot() and plot.lm() use these for some of their graphs.

3.10.5. Deviance residuals

For large sample sizes, the Pearson and deviance residuals will be approximately the same. For small sample sizes, the deviance residuals from covariate patterns with small sample sizes can be unreliable (high variance).

$$d_k = \mathrm{sign}(y_k - n_k \hat{\pi}_k) \left\{ \sqrt{2[\ell_{\mathrm{full}}(x_k) - \ell(\hat{\beta}; x_k)]} \right\}$$

3.10.5.1. Standardized deviance residuals

$$r_{D_k} = \frac{d_k}{\sqrt{1-h_k}}$$

3.10.6. Diagnostic plots

Let's take a look at the full set of autoplot() diagnostics now for our CHD model:




Figure 3.14.: Diagnostics for CHD model

:::

Things look pretty good here. The QQ plot is still usable; with large

samples; the residuals should be approximately Gaussian.

3.10.6.1. Beetles

Let's look at the beetles model diagnostic plots for comparison:

```
beetles_glm_grouped |> autoplot(which = 1:6) |> print()
```



Figure 3.15.: Diagnostics for logistic model of BeetleMortality data

Hard to tell much from so little data, but there might be some issues here.

3.11. Other link functions for Bernoulli outcomes

If you want risk ratios, you can sometimes get them by changing the link function:

```
data(anthers, package = "dobson")
anthers.sum<-aggregate(
   anthers[c("n","y")],
   by=anthers[c("storage")],FUN=sum)

anthers_glm_log = glm(
   formula = cbind(y,n-y)~storage,
   data=anthers.sum,
   family=binomial(link="log"))

anthers_glm_log |> parameters() |> print_md()
```

| Parameter | Log-Risk | SE | 95% CI | Z | p |
|-------------|--------------|------|--------------------------------|---------------|-----------------|
| (Intercept) | -0.80 0.17 | | (-1.04, -0.58) (0.02, 0.31) | -6.81 2.31 | < .001 0.021 |
| storage | 0.17 | 0.07 | (0.02, 0.31) | 2.31 | 0.021 |

Now $\exp \{\beta\}$ gives us risk ratios instead of odds ratios:

```
anthers_glm_log |> parameters(exponentiate = TRUE) |> print_md()
```

| Parameter | Risk Ratio | SE | 95% CI | ${f z}$ | p |
|-------------|------------|------|--------------|---------|--------|
| (Intercept) | 0.45 | 0.05 | (0.35, 0.56) | -6.81 | < .001 |
| storage | 1.18 | 0.09 | (1.03, 1.36) | 2.31 | 0.021 |

Let's compare this model with a logistic model:

```
anthers_glm_logit = glm(
  formula = cbind(y, n - y) ~ storage,
  data = anthers.sum,
  family = binomial(link = "logit"))
anthers_glm_logit |> parameters(exponentiate = TRUE) |> print_md()
```

| Parameter | Odds Ratio | SE | 95% CI | Z | p |
|------------------------|--------------|----|------------------------------|---|---|
| (Intercept) storage | 0.76 1.49 | | (0.45, 1.27) (1.06, 2.10) | | |

[to add: fitted plots on each outcome scale]

When I try to use link ="log" in practice, I often get errors about not finding good starting values for the estimation procedure. This is likely because the model is producing fitted probabilities greater than 1.

When this happens, you can try to fit Poisson regression models instead (we will see those soon!). But then the outcome distribution isn't quite right, and you won't get warnings about fitted probabilities greater than 1. In my opinion, the Poisson model for binary outcomes is confusing and not very appealing.

3.11.1. WCGS: link functions

```
wcgs_glm_logit_link =
  chd_grouped_data |>
  mutate(type = dibpat |> relevel(ref = "Type B")) |>
  glm(
    "formula" = cbind(x, `n - x`) ~ dibpat * age,
```

```
"data" = _,
    "family" = binomial(link = "logit")
)

wcgs_glm_identity_link =
    chd_grouped_data |>
    mutate(type = dibpat |> relevel(ref = "Type B")) |>
    glm(
        "formula" = cbind(x, `n - x`) ~ dibpat * age,
        "data" = _,
        "family" = binomial(link = "identity")
)

wcgs_glm_identity_link |> coef() |> pander()
```

| | $\operatorname{dibpatType}$ | | ${\text{dibpatType}}$ |
|-------------|-----------------------------|----------|-----------------------|
| (Intercept) | A | age | A:age |
| -0.08257 | -0.1374 | 0.002906 | 0.004194 |

```
library(ggfortify)
wcgs_glm_logit_link |> autoplot(which = c(1), ncol = 1) + facet_wrap(~dibpat)
wcgs_glm_identity_link |> autoplot(which = c(1), ncol = 1) + facet_wrap(~dibpat)
```



Figure 3.16.: Residuals vs Fitted plot for wcgs models

```
beetles_lm =
  beetles_long |>
  lm(formula = died ~ dose)

beetles =
  beetles |> mutate(
    resid_logit = beetles_glm_grouped |> resid(type = "response"))
beetles_glm_grouped |> autoplot(which = c(1), ncol = 1)
beetles_lm |> autoplot(which = c(1), ncol = 1)
```



Figure 3.17.: Residuals vs Fitted plot for BeetleMortality models

3.12. Quasibinomial

See Hua Zhou⁴'s lecture notes⁵

3.13. Further reading

• Hosmer, Lemeshow, and Sturdivant (2013) is a classic textbook on logistic regression

⁴https://hua-zhou.github.io/

 $^{^5} https://ucla-biostat-200c-2020 spring.github.io/slides/04-binomial/binomial.html#: ~:text=0.05%20%27%207%200.1%20%27%20%27%201-,Quasi%2Dbinomial,-Another%20way%20to$

4. Models for Count Outcomes

| Poisson regression and variations |
|--|
| |
| |
| Acknowledgements |
| This content is adapted from: |
| Dobson and Barnett (2018), Chapter 9 Vittinghoff et al. (2012), Chapter 8 |
| |
| |

Configuring R

Functions from these packages will be used throughout this document:

Acknowledgements

```
library(conflicted) # check for conflicting function definitions
# library(printr) # inserts help-file output into markdown output
library(rmarkdown) # Convert R Markdown documents into a variety of formats.
library(pander) # format tables for markdown
library(ggplot2) # graphics
library(ggeasy) # help with graphics
library(ggfortify) # help with graphics
library(dplyr) # manipulate data
library(tibble) # `tibble`s extend `data.frame`s
library(magrittr) # `%>%` and other additional piping tools
library(haven) # import Stata files
library(knitr) # format R output for markdown
library(tidyr) # Tools to help to create tidy data
library(plotly) # interactive graphics
library(dobson) # datasets from Dobson and Barnett 2018
library(parameters) # format model output tables for markdown
library(haven) # import Stata files
library(latex2exp) # use LaTeX in R code (for figures and tables)
library(fs) # filesystem path manipulations
library(survival) # survival analysis
library(survminer) # survival analysis graphics
library(KMsurv) # datasets from Klein and Moeschberger
library(parameters) # format model output tables for
library(webshot2) # convert interactive content to static for pdf
library(forcats) # functions for categorical variables ("factors")
library(stringr) # functions for dealing with strings
library(lubridate) # functions for dealing with dates and times
```

Here are some R settings I use in this document:

```
rm(list = ls()) # delete any data that's already loaded into R
conflicts_prefer(dplyr::filter)
```

Acknowledgements

4.1. Introduction

This chapter presents models for count data outcomes. With covariates, the event rate λ becomes a function of the covariates $\tilde{X}=(X_1,\ldots,X_n)$. Typically, count data models use a log $\{\}$ link function, and thus an exp $\{\}$ inverse-link function. That is:

$$\begin{split} \mathbb{E}[Y|\tilde{X} = \tilde{x}, T = t] &= \mu(\tilde{x}, t) \\ \mu(\tilde{x}, t) &= \lambda(\tilde{x}) \cdot t \\ \lambda(\tilde{x}) &= \exp\left\{\eta(\tilde{x})\right\} \\ \eta(\tilde{x}) &= \tilde{x}'\tilde{\beta} = \beta_0 + \beta_1 x_1 + \dots + \beta_p x_p \end{split}$$

Therefore,

$$\begin{split} \log \left\{ \mathbb{E}[Y|\tilde{X} = \tilde{x}, T = t] \right\} &= \log \left\{ \mu(\tilde{x}) \right\} \\ &= \log \left\{ \lambda(\tilde{x}) \cdot t \right\} \\ &= \log \left\{ \lambda(\tilde{x}) \right\} + \log \left\{ t \right\} \\ &= \log \left\{ \exp \left\{ \eta(\tilde{x}) \right\} \right\} + \log \left\{ t \right\} \\ &= \eta(\tilde{x}) + \log \left\{ t \right\} \\ &= \tilde{x}' \tilde{\beta} + \log \left\{ t \right\} \\ &= (\beta_0 + \beta_1 x_1 + \dots + \beta_p x_p) + \log \left\{ t \right\} \end{split}$$

In contrast with the other covariates (represented by \tilde{X}), T enters this expression with a log $\{\}$ transformation and without a corresponding β coefficient.

Note

Terms that enter the linear component of a model without a coefficient, such as $\log \{t\}$ here, are called **offsets**.

4.1.1. Rate ratios

Differences on the log-rate scale become ratios on the rate scale, because

$$\exp\left\{a - b\right\} = \frac{\exp\left\{a\right\}}{\exp\left\{b\right\}}$$

(recall from Algebra 2)

Therefore, according to this model, differences of δ in covariate x_j correspond to rate ratios of $\exp\{\beta_j \cdot \delta\}$.

That is, letting \tilde{X}_{-j} denote vector \tilde{X} with element j removed:

Acknowledgements

$$\begin{cases} \log\left\{\mathbb{E}[Y|\mathbf{X}_{j}=a,\tilde{X}_{-j}=\tilde{x}_{-j},T=t]\right\} \\ -\log\left\{\mathbb{E}[Y|\mathbf{X}_{j}=b,\tilde{X}_{-j}=\tilde{x}_{-j},T=t]\right\} \end{cases} \\ = \begin{cases} \log\left\{t\right\} + \beta_{0} + \beta_{1}x_{1} + \ldots + \beta_{j}(a) + \ldots + \beta_{p}x_{p} \\ -\log\left\{t\right\} + \beta_{0} + \beta_{1}x_{1} + \ldots + \beta_{j}(b) + \ldots + \beta_{p}x_{p} \right\} \\ = \beta_{j}(a-b) \end{cases}$$

And accordingly,

$$\frac{\mathbb{E}[Y|\boldsymbol{X_j}=a,\tilde{X}_{-j}=\tilde{x}_{-j},T=t]}{\mathbb{E}[Y|\boldsymbol{X_j}=b,\tilde{X}_{-j}=\tilde{x}_{-j},T=t]}=\exp\left\{\beta_j(a-b)\right\}$$

4.2. Inference for count regression models

4.2.1. Confidence intervals for regression coefficients and rate ratios

As usual:

$$\beta \in \left[\hat{\beta} \pm z_{1 - \frac{\alpha}{2}} \cdot \hat{\operatorname{se}} \left(\hat{\beta} \right) \right]$$

Rate ratios: exponentiate CI endpoints

$$\exp\left\{\beta\right\} \in \left[\exp\left\{\hat{\beta} \underline{+} z_{1-\frac{\alpha}{2}} \cdot \hat{\operatorname{se}}\left(\hat{\beta}\right)\right\}\right]$$

Acknowledgements

4.2.2. Hypothesis tests for regression coefficients

$$t = \frac{\hat{\beta} - \beta_0}{\hat{\operatorname{se}}\left(\hat{\beta}\right)}$$

Compare t or |t| to the tails of the standard Gaussian distribution, according to the null hypothesis.

4.2.3. Comparing nested models

log(likelihood ratio) tests, as usual.

4.3. Prediction

$$\begin{split} \hat{y} &\stackrel{\text{def}}{=} \hat{\mathbb{E}}[Y|\tilde{X} = \tilde{x}, T = t] \\ &= \hat{\mu}(\tilde{x}, t) \\ &= \hat{\lambda}(\tilde{x}) \cdot t \\ &= \exp\left\{\hat{\eta}(\tilde{x})\right\} \cdot t \\ &= \exp\left\{\tilde{x}'\hat{\beta}\right\} \cdot t \end{split}$$

4.4. Diagnostics

4.4.1. Residuals

4.4.1.1. Observation residuals

$$e \stackrel{\mathrm{def}}{=} y - \hat{y}$$

4.4.1.2. Pearson residuals

$$r = \frac{e}{\hat{\operatorname{se}}\left(e\right)} \approx \frac{e}{\sqrt{\hat{y}}}$$

4.4.1.3. Standardized Pearson residuals

$$r_p = \frac{r}{\sqrt{1-h}}$$

where h is the "leverage" (which we will continue to leave undefined).

4.4.1.4. Deviance residuals

$$d_k = \mathrm{sign}(y - \hat{y}) \left\{ \sqrt{2[\ell_{\mathrm{full}}(y) - \ell(\hat{\beta}; y)]} \right\}$$

Note

$$\operatorname{sign}(x) \stackrel{\text{def}}{=} \frac{x}{|x|}$$

In other words:

- $\begin{array}{ll} \bullet & \mathrm{sign}(x) = -1 \mathrm{~if~} x < 0 \\ \bullet & \mathrm{sign}(x) = 0 \mathrm{~if~} x = 0 \end{array}$
- $\operatorname{sign}(x) = 1 \text{ if } x > 0$

4.5. Zero-inflation

4.5.1. Models for zero-inflated counts

We assume a latent (unobserved) binary variable, Z, which we model using logistic regression:

$$P(Z = 1|X = x) = \pi(x) = \text{expit}(\gamma_0 + \gamma_1 x_1 + ...)$$

According to this model, if Z=1, then Y will always be zero, regardless of X and T:

$$P(Y = 0|Z = 1, X = x, T = t) = 1$$

Otherwise (if Z = 0), Y will have a Poisson distribution, conditional on X and T, as above.

Even though we never observe Z, we can estimate the parameters γ_0 - γ_p , via maximum likelihood:

$$P(Y = y|X = x, T = t) = P(Y = y, Z = 1|...) + P(Y = y, Z = 0|...)$$

(by the Law of Total Probability)

where

$$P(Y=y,Z=z|\ldots)=P(Y=y|Z=z,\ldots)P(Z=z|\ldots)$$

Exercise 4.1. Expand P(Y=0|X=x,T=t), P(Y=1|X=x,T=t) and P(Y=y|X=x,T=t) into expressions involving P(Z=1|X=x,T=t) and P(Y=y|Z=0,X=x,T=t).

Exercise 4.2. Derive the expected value and variance of Y, conditional on X and T, as functions of P(Z=1|X=x,T=t) and $\mathbb{E}[Y|Z=0,X=x,T=t]$.

4.6. Over-dispersion

The Poisson distribution model **forces** the variance to equal the mean. In practice, many count distributions will have a variance substantially larger than the mean (or occasionally, smaller).

Definition 4.1 (Overdispersion). A random variable X is **overdispersed** relative to a model p(X = x) if if its empirical variance in a dataset is larger than the value is predicted by the fitted model $\hat{p}(X = x)$.

c.f. Dobson and Barnett (2018) §3.2.1, 7.7, 9.8; Vittinghoff et al. (2012) §8.1.5; and https://en.wikipedia.org/wiki/Overdispersion.

When we encounter overdispersion, we can try to reduce the residual variance by adding more covariates.

4.6.1. Negative binomial models

There are alternatives to the Poisson model. Most notably, the negative binomial model.

We can still model μ as a function of X and T as before, and we can combine this model with zero-inflation (as the conditional distribution for the non-zero component).

4.6.2. Quasipoisson

An alternative to Negative binomial is the "quasipoisson" distribution. I've never used it, but it seems to be a method-of-moments type approach rather than maximum likelihood. It models the variance as $\mathrm{Var}\,(Y) = \mu\theta$, and estimates θ accordingly.

See ?quasipoisson in R for more.

4.7. Example: needle-sharing

(adapted from Vittinghoff et al. (2012), §8)

```
library(ggplot2)

needles |>
    ggplot(
    aes(
        x = age,
        y = shared_syr,
        shape = sex,
        col = ethn
    )
    ) +
    geom_point(
        size = 3,
        alpha = .5) +
    facet_grid(
    cols = vars(sex, polydrug),
```

Table 4.1.: Needle-sharing data

```
library(tidyverse)
library(haven)
needles =
  "inst/extdata/needle_sharing.dta" |>
  read dta() |>
  as_tibble() |>
  mutate(
    hivstat =
      hivstat |>
      case_match(
        1 ~ "HIV+",
        0 ~ "HIV-") |>
      factor() |>
      relevel(ref = "HIV-"),
    polydrug =
      polydrug |>
      case_match(
        1 ~ "multiple drugs used",
        0 ~ "one drug used") |>
      factor() |>
      relevel(ref = "one drug used"),
    homeless =
      homeless |>
      case_match(
        1 ~ "homeless",
        0 ~ "not homeless") |>
      factor() |>
      relevel(ref = "not homeless"),
    sex = sex |> factor() |> relevel(ref = "M"))
needles
#> # A tibble: 128 x 17
#>
         id sex ethn
                             age dprsn_dx sexabuse shared_syr hivstat hplsns nivdu
      <dbl> <fct> <chr>
                           <dbl>
                                    <dbl>
                                             <dbl>
                                                         <dbl> <fct>
                                                                        <dbl> <dbl>
#>
#>
   1 2104 M
                White
                              47
                                        5
                                                 0
                                                             1 HIV-
                                                                            6
                                                                                 90
                             25$
                                                                            2
   2 2009 M
                White
                                        1
                                                  0
#>
                                                             1 HIV+
                                                                                  4
                                                             1 HIV-
   3 2032 M
                White
                             52
                                        1
                                                  0
#>
                                                                           18
                                                                                 90
#>
   4 2063 M
                AA
                              47
                                        1
                                                 1
                                                             1 HIV-
                                                                            1
                                                                                120
                Hispanic
   5 2059 M
                              32
                                                             2 HIV-
#>
                                        1
                                                 0
                                                                           12
                                                                                120
#>
   6 2077 M
                Hispanic
                              54
                                        1
                                                 0
                                                             2 HIV-
                                                                           10
                                                                                120
#>
   7 2042 F
                              32
                                        5
                                                 0
                                                             2 HIV-
                                                                            8
                                                                                 15
                  White
                                        5
                                                 0
                                                                                120
#>
   8 2017 M
                  White
                              26
                                                             2 HIV-
                                                                           11
                                                 0
#> 9 2119 M
                  White
                              54
                                        1
                                                             3 HIV-
                                                                            2
                                                                                 90
#> 10 2085 F
                                        5
                                                  0
                              19
                                                             3 HIV-
                                                                                 90
                  White
#> # i 118 more rows
#> # i 7 more variables: shsyryn <dbl>, sqrtnivd <dbl>, logshsyr <dbl>,
       polydrug <fct>, sqrtninj <dbl>, homeless <fct>, shsyr <dbl>
```

```
rows = vars(homeless)) +
theme(legend.position = "bottom")
```




Figure 4.1.: Rates of needle sharing

4.7.0.1. Covariate counts

There's only one individual with sex = Trans, which unfortunately isn't enough data to analyze. We will remove that individual:

{.r .cell-code code-summary="remove singleton observation
with `sex == "Trans"`"} needles = needles |> filter(sex !=
"Trans")

Table 4.2.: Counts of observations in needles dataset by sex, unhoused status, and multiple drug use

```
needles |>
 dplyr::select(sex, homeless, polydrug) |>
 summary()
#>
      sex
                    homeless
                                           polydrug
#> M
             not homeless:63 one drug used
       :97
                                              :109
#> F
        :30
                       :61 multiple drugs used: 19
             homeless
#> Trans: 1 NA's : 4
```

4.7.1. models

```
glm1 = glm(
  data = needles,
  family = stats::poisson,
  shared_syr ~ age + sex + homeless*polydrug
)

library(parameters)
glm1 |> parameters(exponentiate = TRUE) |>
  print_md()
```

Table 4.3.: Poisson model for needle-sharing data

| Parameter | IRR | SE | 95% CI | \mathbf{z} | p |
|--------------------------------|-----------|------------|--------|--------------|-------|
| (Intercept) | 4.52 | 1.15 | (2.74, | 5.90 | < |
| | | | 7.45) | | .001 |
| age | 0.97 | 5.58e- | (0.96, | - | < |
| | | 03 | 0.98) | 5.41 | .001 |
| sex(F) | 1.98 | 0.23 | (1.58, | 5.88 | < |
| | | | 2.49) | | .001 |
| homeless (homeless) | 3.58 | 0.45 | (2.79, | 10.06 | < |
| | | | 4.59) | | .001 |
| polydrug (multiple drugs used) | 1.45e- | 5.82e- | (0.00, | - | 0.969 |
| | 07 | 05 | Inf) | 0.04 | |
| homeless (homeless) \times | 1.27e + 0 | 06.12e + 0 | 0.00, | 0.03 | 0.972 |
| polydrug (multiple drugs used) | | | Inf) | | |

```
library(ggfortify)
autoplot(glm1)
```

_

4.7.1.1. zero-inflation

```
library(glmmTMB)
zinf_fit1 = glmmTMB(
  family = "poisson",
  data = needles,
  formula = shared_syr ~ age + sex + homeless*polydrug,
  ziformula = ~ age + sex + homeless + polydrug # fit won't converge with interaction)
```




Table 4.4.: Diagnostics for Poisson model

Table 4.5.: Negative binomial model for needle-sharing data

```
library(MASS) #need this for glm.nb()
glm1.nb = glm.nb(
 data = needles,
  shared_syr ~ age + sex + homeless*polydrug
summary(glm1.nb)
#>
#> Call:
#> glm.nb(formula = shared_syr ~ age + sex + homeless * polydrug,
      data = needles, init.theta = 0.08436295825, link = log)
#>
#>
#> Coefficients:
#>
                                                Estimate Std. Error z value
#> (Intercept)
                                                9.91e-01 1.71e+00
                                                                      0.58
                                               -2.76e-02 3.82e-02 -0.72
#> age
#> sexF
                                                1.06e+00 8.07e-01 1.32
#> homelesshomeless
                                                1.65e+00 7.22e-01
                                                                      2.29
#> polydrugmultiple drugs used
                                               -2.46e+01 3.61e+04 0.00
#> homelesshomeless:polydrugmultiple drugs used 2.32e+01 3.61e+04 0.00
#>
                                               Pr(>|z|)
#> (Intercept)
                                                  0.563
#> age
                                                  0.469
#> sexF
                                                  0.187
#> homelesshomeless
                                                  0.022 *
#> polydrugmultiple drugs used
                                                  0.999
#> homelesshomeless:polydrugmultiple drugs used
                                                  0.999
#> ---
#> Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
#>
#> (Dispersion parameter for Negative Binomial(0.0844) family taken to be 1)
#>
#>
      Null deviance: 69.193 on 119 degrees of freedom
#> Residual deviance: 57.782 on 114 degrees of freedom
    (7 observations deleted due to missingness)
                            256
#> AIC: 315.5
#>
#> Number of Fisher Scoring iterations: 1
#>
#>
#>
                Theta: 0.0844
            Std. Err.: 0.0197
#>
#>
#> 2 x log-likelihood: -301.5060
```

Table 4.6.: Poisson versus Negative Binomial Regression coefficient estimates

```
tibble(name = names(coef(glm1)), poisson = coef(glm1), nb = coef(glm1.nb))
#> # A tibble: 6 x 3
#>
     name
                                                   poisson
                                                                 nb
#>
     <chr>>
                                                     <dbl>
                                                              <dbl>
#> 1 (Intercept)
                                                    1.51
                                                             0.991
#> 2 age
                                                   -0.0311 -0.0276
#> 3 sexF
                                                    0.684
                                                            1.06
#> 4 homelesshomeless
                                                    1.27
                                                             1.65
#> 5 polydrugmultiple drugs used
                                                           -24.6
                                                  -15.7
#> 6 homelesshomeless:polydrugmultiple drugs used 14.1
                                                            23.2
```

```
zinf_fit1 |>
  parameters(exponentiate = TRUE) |>
  print_md()
```

Table 4.7.: Zero-inflated poisson model

Table 4.7.: # Fixed Effects

| Parameter | IRR | SE | 95% CI | ${f z}$ | р |
|-------------|------|--------|--------|---------|-------|
| (Intercept) | 3.16 | 0.82 | (1.90, | 4.44 | < |
| | | | 5.25) | | .001 |
| age | 1.01 | 5.88e- | (1.00, | 1.74 | 0.081 |
| | | 03 | 1.02) | | |
| sex [F] | 3.43 | 0.44 | (2.67, | 9.68 | < |
| | | | 4.40) | | .001 |

| Parameter | IRR | SE | 95% CI | ${f z}$ | р |
|--------------------------------|---------|------------------|-----------|---------|-------|
| homeless [homeless] | 3.44 | 0.47 | (2.63, | 9.03 | < |
| | | | 4.50) | | .001 |
| polydrug [multiple drugs used] | 1.85e- | 1.21e- | (0.00, | -3.08e- | 0.998 |
| | 09 | 05 | Inf) | 03 | |
| homeless [homeless] \times | 1.38e + | 0 8 .04e+ | 11 (0.00, | 2.87e- | 0.998 |
| polydrug [multiple drugs used] | | | Inf) | 03 | |

Table 4.8.: Zero-inflated poisson model

Table 4.8.: # Zero-Inflation

| | Odds | | | | |
|-----------------------------------|-------|------|--------------|---------|-------|
| Parameter | Ratio | SE | 95% CI | ${f z}$ | p |
| (Intercept) | 0.49 | 0.54 | (0.06, 4.25) | -0.65 | 0.514 |
| age | 1.05 | 0.03 | (1.00, 1.10) | 1.95 | 0.051 |
| sex [F] | 1.44 | 0.84 | (0.46, 4.50) | 0.62 | 0.533 |
| homeless [homeless] | 0.68 | 0.34 | (0.26, 1.80) | -0.78 | 0.436 |
| polydrug [multiple drugs used] | 1.15 | 0.91 | (0.24, 5.43) | 0.18 | 0.858 |

Another R package for zero-inflated models is $pscl^1$ (Zeileis, Kleiber, and Jackman (2008)).

 $^{^{-1} \}rm https://cran.r-project.org/web/packages/pscl/index.html$

4.7.1.2. zero-inflated negative binomial model

```
library(glmmTMB)
zinf_fit1 = glmmTMB(
  family = nbinom2,
  data = needles,
  formula = shared_syr ~ age + sex + homeless*polydrug,
  ziformula = ~ age + sex + homeless + polydrug # fit won't converge with interaction)

zinf_fit1 |>
  parameters(exponentiate = TRUE) |>
  print_md()
```

Table 4.9.: Zero-inflated negative binomial model

Table 4.9.: # Fixed Effects

| Parameter | IRR | SE | 95% CI | ${f z}$ | p |
|--------------------------------|---------|------------|-----------|---------|-------|
| (Intercept) | 1.06 | 1.48 | (0.07, | 0.04 | 0.969 |
| | | | 16.52) | | |
| age | 1.02 | 0.03 | (0.96, | 0.53 | 0.599 |
| | | | 1.08) | | |
| sex [F] | 6.86 | 6.36 | (1.12, | 2.08 | 0.038 |
| | | | 42.16) | | |
| homeless [homeless] | 6.44 | 4.59 | (1.60, | 2.62 | 0.009 |
| | | | 26.01) | | |
| polydrug [multiple drugs used] | 8.25e- | 7.07e- | (0.00, | -2.44e- | 0.998 |
| | 10 | 06 | Inf) | 03 | |
| homeless [homeless] \times | 2.36e + | 08.02e + 1 | 12 (0.00, | 2.25e- | 0.998 |
| polydrug [multiple drugs used] | | | Inf) | 03 | |

Table 4.10.: Zero-inflated negative binomial model

Table 4.10.: # Zero-Inflation

| | Odds | | | | |
|-----------------------------------|-------|------|---------------------|--------------|-------|
| Parameter | Ratio | SE | 95% CI | \mathbf{Z} | p |
| (Intercept) | 0.10 | 0.20 | (1.47e-03, 6.14) | -1.11 | 0.269 |
| age | 1.07 | 0.04 | (0.99, 1.15) | 1.78 | 0.075 |
| sex [F] | 2.72 | 2.40 | (0.48, 15.33) | 1.13 | 0.258 |
| homeless [homeless] | 1.15 | 0.86 | (0.27, 4.96) | 0.19 | 0.853 |
| polydrug [multiple drugs used] | 0.75 | 0.86 | (0.08, 7.12) | -0.25 | 0.799 |

Table 4.11.: Zero-inflated negative binomial model

Table 4.11.: # Dispersion

| Parameter | Coefficient | 95% CI | | |
|-------------|-------------|--------------|--|--|
| (Intercept) | 0.44 | (0.11, 1.71) | | |

Part II. Time to Event Models

In many health sciences applications, binary outcomes are *incompletely observed*. For example, if we are studying whether cancer patients experience a relapse after a initial remission, we may may not be able to follow patients to the end of their lives; instead, we may only know whether each patient has relapsed before the end of the study. If a patient has not relapsed by that point, we might not know if they will relapse at some other date or if they will stay cancer-free for the rest of their lives. ² Their recurrence status at end-of-life is *missing data*. If some study participants withdraw from a study before the end date in the study design, there will be even more missing data. All of this missing data will make logistic regression difficult for this type of data.

However, these outcome observations are not *entirely* missing. We know that those patients stayed relapse free *at least* until the time point when we last saw them. If we also know the *time-to-event* for the participants who did experience events while under study, we can analyze *time-to-event-or-study-exit*, combined with the indicator of which of these two cases occurred, using *survival analysis*. The survival analysis framework is the subject of the rest of these course notes.

²Binary outcomes are typically defined for a specific time-point. It is important to clearly define whether we are interested in outcome status at end of study, at end of life, or at some other time.

Configuring R

Functions from these packages will be used throughout this document:

```
library(conflicted) # check for conflicting function definitions
# library(printr) # inserts help-file output into markdown output
library(rmarkdown) # Convert R Markdown documents into a variety of formats.
library(pander) # format tables for markdown
library(ggplot2) # graphics
library(ggeasy) # help with graphics
library(ggfortify) # help with graphics
library(dplyr) # manipulate data
library(tibble) # `tibble`s extend `data.frame`s
library(magrittr) # `%>%` and other additional piping tools
library(haven) # import Stata files
library(knitr) # format R output for markdown
library(tidyr) # Tools to help to create tidy data
library(plotly) # interactive graphics
library(dobson) # datasets from Dobson and Barnett 2018
library(parameters) # format model output tables for markdown
library(haven) # import Stata files
library(latex2exp) # use LaTeX in R code (for figures and tables)
library(fs) # filesystem path manipulations
```

```
library(survival) # survival analysis
library(survminer) # survival analysis graphics
library(KMsurv) # datasets from Klein and Moeschberger
library(parameters) # format model output tables for
library(webshot2) # convert interactive content to static for pdf
library(forcats) # functions for categorical variables ("factors")
library(stringr) # functions for dealing with strings
library(lubridate) # functions for dealing with dates and times
```

Here are some R settings I use in this document:

```
rm(list = ls()) # delete any data that's already loaded into R

conflicts_prefer(dplyr::filter)
ggplot2::theme_set(
    ggplot2::theme_bw() +
        # ggplot2::labs(col = "") +
    ggplot2::theme(
        legend.position = "bottom",
        text = ggplot2::element_text(size = 12, family = "serif")))

knitr::opts_chunk$set(message = FALSE)
options('digits' = 4)

panderOptions("big.mark", ",")
pander::panderOptions("table.emphasize.rownames", FALSE)
pander::panderOptions("table.split.table", Inf)
conflicts_prefer(dplyr::filter) # use the `filter()` function from dplyr() by defaul legend_text_size = 9
```

5.1. Overview

5.1.1. Time-to-event outcomes

Survival analysis is a framework for modeling *time-to-event* outcomes. It is used in:

- clinical trials, where the event is often death or recurrence of disease.
- engineering reliability analysis, where the event is failure of a device or system.
- insurance, particularly life insurance, where the event is death.

Note

The term *survival analysis* is a bit misleading. Survival outcomes can sometimes be analyzed using binomial models (logistic regression). *Time-to-event models* or *survival time analysis* might be a better name.

5.2. Time-to-event outcome distributions

5.2.1. Distributions of Time-to-Event Data

- The distribution of event times is asymmetric and can be long-tailed, and starts at 0 (that is, P(T < 0) = 0).
- The base distribution is not normal, but exponential.
- There are usually **censored** observations, which are ones in which the failure time is not observed.
- Often, these are **right-censored**, meaning that we know that the event occurred after some known time t, but we don't know the actual event time, as when a patient is still alive at the end of the study.

- Observations can also be **left-censored**, meaning we know the event has already happened at time t, or **interval-censored**, meaning that we only know that the event happened between times t_1 and t_2 .
- Analysis is difficult if censoring is associated with treatment.

5.2.2. Right Censoring

- Patients are in a clinical trial for cancer, some on a new treatment and some on standard of care.
- Some patients in each group have died by the end of the study. We know the survival time (measured for example from time of diagnosis—each person on their own clock).
- Patients still alive at the end of the study are right censored.
- Patients who are lost to follow-up or withdraw from the study may be right-censored.

5.2.3. Left and Interval Censoring

- An individual tests positive for HIV.
- If the event is infection with HIV, then we only know that it has occurred before the testing time t, so this is left censored.
- If an individual has a negative HIV test at time t_1 and a positive HIV test at time t_2 , then the infection event is interval censored.

5.3. Distribution functions for time-to-event variables

5.3.1. The Probability Density Function (PDF)

For a time-to-event variable T with a continuous distribution, the **probability density function** is defined as usual (see Section B.4.1).

In most time-to-event models, this density is assumed to be 0 for all t < 0; that is, $f(t) = 0, \forall t < 0$. In other words, the support of T is typically $[0, \infty)$.

Example 5.1 (exponential distribution). Recall from Epi 202: the pdf of the exponential distribution family of models is:

$$p(T=t) = \mathbb{1}_{t \geq 0} \cdot \lambda \mathrm{e}^{-\lambda t}$$

where $\lambda > 0$.

Here are some examples of exponential pdfs:



5.3.2. The Cumulative Distribution Function (CDF)

The **cumulative distribution function** is defined as:

$$\begin{split} F(t) &\stackrel{\text{def}}{=} \Pr(T \leq t) \\ &= \int_{u=-\infty}^t f(u) du \end{split}$$

Example 5.2 (exponential distribution). Recall from Epi 202: the cdf of the exponential distribution family of models is:

$$P(T \leq t) = \mathbb{1}_{t \geq 0} \cdot (1 - \mathrm{e}^{-\lambda t})$$

where $\lambda > 0$.

Here are some examples of exponential cdfs:



5.3.3. The Survival Function

For survival data, a more important quantity is the **survival function**:

$$S(t) \stackrel{\text{def}}{=} \Pr(T > t)$$

$$= \int_{u=t}^{\infty} p(u) du$$

$$= 1 - F(t)$$

Definition 5.1 (Survival function).

Given a random time-to-event variable T, the **survival function** or **survivor function**, denoted S(t), is the probability that the event time is later than t. If the event in a clinical trial is death, then S(t) is the expected fraction of the original population at time 0 who have survived up to time t and are still alive at time t; that is:

$$S(t) \stackrel{\text{def}}{=} \Pr(T > t) \tag{5.1}$$

Example 5.3 (exponential distribution). Since S(t) = 1 - F(t), the survival function of the exponential distribution family of models is:

$$P(T > t) = \begin{cases} e^{-\lambda t}, t \ge 0\\ 1, t \le 0 \end{cases}$$

where $\lambda > 0$.

Figure 5.1 shows some examples of exponential survival functions.



Figure 5.1.: Exponential Survival Functions

Theorem 5.1. If A_t represents survival status at time t, with $A_t=1$ denoting alive at time t and $A_t=0$ denoting deceased at time t, then:

$$S(t) = \Pr(A_t = 1) = \mathbb{E}[A_t]$$

Theorem 5.2. If T is a nonnegative random variable, then:

$$\mathbb{E}[T] = \int_{t=0}^{\infty} S(t) dt$$

Proof. See https://statproofbook.github.io/P/mean-nnrvar.html or □

5.3.4. The Hazard Function

Another important quantity is the **hazard function**:

Definition 5.2 (Hazard function, hazard rate, hazard rate function).

The hazard function, hazard rate, hazard rate function, for a random variable T at value t, typically denoted as h(t) or $\lambda(t)$, is the conditional density of T at t, given $T \geq t$. That is:

$$h(t) \stackrel{\text{def}}{=} p(T = t | T \ge t)$$

If T represents the time at which an event occurs, then h(t) is the probability that the event occurs at time t, given that it has not occurred prior to time t.

New content for 2025:

Definition 5.3 (Incidence rate). Given a population of individuals indexed by i, each with their own hazard rate $h_i(t)$, the **incidence rate** for that population is the mean hazard rate:

$$\bar{h}(t) \stackrel{\text{def}}{=} \frac{1}{n} \sum_{i=1}^{n} h_i(t)$$

New content for 2025:

Theorem 5.3. If a population of individuals indexed by i all have identical hazard rates $h_i(t) = h(t)$, then the **incidence rate** for that population is equal to the hazard rate:

$$\bar{h}(t) = h(t)$$

(End of new content for 2025)

The hazard function has an important relationship to the density and survival functions, which we can use to derive the hazard function for a given probability distribution (Theorem 5.4).

Lemma 5.1 (Joint probability of a variable with itself).

$$p(T = t, T \ge t) = p(T = t)$$

Proof. Recall from Epi 202: if A and B are statistical events and $A \subseteq B$, then p(A,B)=p(A). In particular, $\{T=t\}\subseteq \{T\geq t\}$, so $p(T=t,T\geq t)=p(T=t)$.

Theorem 5.4.

$$h(t) = \frac{f(t)}{S(t)}$$

Proof.

$$\begin{split} h(t) &= p(T=t|T \geq t) \\ &= \frac{p(T=t,T \geq t)}{p(T \geq t)} \\ &= \frac{p(T=t)}{p(T \geq t)} \\ &= \frac{f(t)}{S(t)} \end{split}$$

Example 5.4 (exponential distribution). The hazard function of the exponential distribution family of models is:

$$\begin{split} P(T=t|T\geq t) &= \frac{f(t)}{S(t)} \\ &= \frac{\mathbbm{1}_{t\geq 0} \cdot \lambda \mathrm{e}^{-\lambda t}}{\mathrm{e}^{-\lambda t}} \\ &= \mathbbm{1}_{t\geq 0} \cdot \lambda \end{split}$$

Figure 5.2 shows some examples of exponential hazard functions.




Figure 5.2.: Examples of hazard functions for exponential distributions

We can also view the hazard function as the derivative of the negative of the logarithm of the survival function:

Theorem 5.5 (transform survival to hazard).

$$h(t) = \frac{\partial}{\partial t} \left\{ -\log \left\{ S(t) \right\} \right\}$$

Proof.

$$h(t) = \frac{f(t)}{S(t)}$$

$$= \frac{-S'(t)}{S(t)}$$

$$= -\frac{S'(t)}{S(t)}$$

$$= -\frac{\partial}{\partial t} \log \{S(t)\}$$

$$= \frac{\partial}{\partial t} \{-\log \{S(t)\}\}$$

5.3.5. The Cumulative Hazard Function

Since $h(t) = \frac{\partial}{\partial t} \{-\log \{S(t)\}\}$ (see Theorem 5.5), we also have:

Corollary 5.1.

$$S(t) = exp\left\{-\int_{u=0}^{t} h(u)du\right\}$$
 (5.2)

The integral in Equation 5.2 is important enough to have its own name: cumulative hazard.

Definition 5.4 (cumulative hazard). The **cumulative hazard function** H(t) is defined as:

$$H(t) \stackrel{\text{def}}{=} \int_{u=0}^{t} h(u) du$$

As we will see below, H(t) is tractable to estimate, and we can then derive an estimate of the hazard function using an approximate derivative of the estimated cumulative hazard.

Example 5.5. The cumulative hazard function of the exponential distribution family of models is:

$$H(t) = \mathbb{1}_{t \geq 0} \cdot \lambda t$$

Figure 5.3 shows some examples of exponential cumulative hazard functions.



Figure 5.3.: Examples of exponential cumulative hazard functions

5.3.6. Some Key Mathematical Relationships among Survival Concepts

5.3.6.1. Diagram:

$$h(t) \xrightarrow{\int_{u=0}^{t} h(u)du} H(t) \xrightarrow{\exp\{-H(t)\}} S(t) \xrightarrow{1-S(t)} F(t)$$
$$h(t) \xleftarrow{\frac{\partial}{\partial t} H(t)} H(t) \xleftarrow{-\log\{S(t)\}} S(t) \xleftarrow{1-F(t)} F(t)$$

5.3.6.2. Identities:

$$S(t) = 1 - F(t)$$

$$= \exp \{-H(t)\}$$

$$S'(t) = -f(t)$$

$$H(t) = -\log \{S(t)\}$$

$$H'(t) = h(t)$$

$$h(t) = \frac{f(t)}{S(t)}$$

$$= -\frac{\partial}{\partial t} \log \{S(t)\}$$

$$f(t) = h(t) \cdot S(t)$$

Some proofs (others left as exercises):

$$S'(t) = \frac{\partial}{\partial t}(1 - F(t))$$
$$= -F'(t)$$
$$= -f(t)$$

$$\begin{split} \frac{\partial}{\partial t} \mathrm{log} \left\{ S(t) \right\} &= \frac{S'(t)}{S(t)} \\ &= -\frac{f(t)}{S(t)} \\ &= -h(t) \end{split}$$

$$\begin{split} H(t) &\stackrel{\text{def}}{=} \int_{u=0}^{t} h(u) du \\ &= \int_{0}^{t} -\frac{\partial}{\partial u} \log \left\{ S(u) \right\} du \\ &= \left[-\log \left\{ S(u) \right\} \right]_{u=0}^{u=t} \\ &= \left[\log \left\{ S(u) \right\} \right]_{u=t}^{u=0} \\ &= \log \left\{ S(0) \right\} - \log \left\{ S(t) \right\} \\ &= \log \left\{ 1 \right\} - \log \left\{ S(t) \right\} \\ &= 0 - \log \left\{ S(t) \right\} \\ &= -\log \left\{ S(t) \right\} \end{split}$$

Corollary:

$$S(t) = \exp\left\{-H(t)\right\}$$

5.3.6.3. Example: Time to death the US in 2004

The first day is the most dangerous:




Figure 5.4.: Daily Hazard Rates in 2004 for US Females

Exercise: hypothesize why these curves differ where they do?




Figure 5.5.: Daily Hazard Rates in 2004 for US Males and Females 1-40 $\,$

Exercise: compare and contrast this curve with the corresponding hazard curve.




Figure 5.6.: Survival Curve in 2004 for US Females

5.3.7. Likelihood with censoring

If an event time T is observed exactly as T=t, then the likelihood of that observation is just its probability density function:

$$\begin{split} \mathcal{L}(t) &= p(T=t) \\ &\stackrel{\text{def}}{=} f_T(t) \\ &= h_T(t) S_T(t) \\ \ell(t) &\stackrel{\text{def}}{=} \log \left\{ \mathcal{L}(t) \right\} \\ &= \log \left\{ h_T(t) S_T(t) \right\} \\ &= \log \left\{ h_T(t) \right\} + \log \left\{ S_T(t) \right\} \\ &= \log \left\{ h_T(t) \right\} - H_T(t) \end{split}$$

If instead the event time T is censored and only known to be after time y, then the likelihood of that censored observation is instead the survival function evaluated at the censoring time:

$$\begin{split} \mathcal{L}(y) &= p_T(T > y) \\ &\stackrel{\text{def}}{=} S_T(y) \\ \ell(y) &\stackrel{\text{def}}{=} \log \left\{ \mathcal{L}(y) \right\} \\ &= \log \left\{ S(y) \right\} \\ &= -H(y) \end{split}$$

What's written above is incomplete. We also observed whether or not the observation was censored. Let C denote the time when censoring would occur (if the event did not occur first); let $f_C(y)$ and $S_C(y)$ be the corresponding density and survival functions for the censoring event.

Let Y denote the time when observation ended (either by censoring or by the event of interest occurring), and let D be an indicator variable

for the event occurring at Y (so D=0 represents a censored observation and D=1 represents an uncensored observation). In other words, let $Y\stackrel{\text{def}}{=} \min(T,C)$ and $D\stackrel{\text{def}}{=} \mathbb{1}\{T <= C\}$.

Then the complete likelihood of the observed data (Y, D) is:

$$\begin{split} \mathcal{L}(y,d) &= p(Y = y, D = d) \\ &= \left[p(T = y, C > y) \right]^d \cdot \left[p(T > y, C = y) \right]^{1-d} \end{split}$$

Typically, survival analyses assume that C and T are mutually independent; this assumption is called "non-informative" censoring.

Then the joint likelihood p(Y,D) factors into the product p(Y),p(D), and the likelihood reduces to:

$$\begin{split} \mathcal{L}(y,d) &= \left[p(T=y,C>y) \right]^d \cdot \left[p(T>y,C=y) \right]^{1-d} \\ &= \left[p(T=y)p(C>y) \right]^d \cdot \left[p(T>y)p(C=y) \right]^{1-d} \\ &= \left[f_T(y)S_C(y) \right]^d \cdot \left[S(y)f_C(y) \right]^{1-d} \\ &= \left[f_T(y)^d S_C(y)^d \right] \cdot \left[S_T(y)^{1-d} f_C(y)^{1-d} \right] \\ &= \left(f_T(y)^d \cdot S_T(y)^{1-d} \right) \cdot \left(f_C(y)^{1-d} \cdot S_C(y)^d \right) \end{split}$$

The corresponding log-likelihood is:

$$\begin{split} \ell(y,d) &= \log \left\{ \mathcal{L}(y,d) \right\} \\ &= \log \left\{ \left(f_T(y)^d \cdot S_T(y)^{1-d} \right) \cdot \left(f_C(y)^{1-d} \cdot S_C(y)^d \right) \right\} \\ &= \log \left\{ f_T(y)^d \cdot S_T(y)^{1-d} \right\} + \log \left\{ f_C(y)^{1-d} \cdot S_C(y)^d \right\} \end{split}$$

Let

- θ_T represent the parameters of $p_T(t)$,
- θ_C represent the parameters of $p_C(c)$,
- $\theta = (\theta_T, \theta_C)$ be the combined vector of all parameters.

The corresponding score function is:

$$\begin{split} \ell'(y,d) &= \frac{\partial}{\partial \theta} \left[\log \left\{ f_T(y)^d \cdot S_T(y)^{1-d} \right\} + \log \left\{ f_C(y)^{1-d} \cdot S_C(y)^d \right\} \right] \\ &= \left(\frac{\partial}{\partial \theta} \log \left\{ f_T(y)^d \cdot S_T(y)^{1-d} \right\} \right) + \left(\frac{\partial}{\partial \theta} \log \left\{ f_C(y)^{1-d} \cdot S_C(y)^d \right\} \right) \end{split}$$

As long as θ_C and θ_T don't share any parameters, then if censoring is non-informative, the partial derivative with respect to θ_T is:

$$\begin{split} \ell_{\theta_T}'(y,d) &\stackrel{\text{def}}{=} \frac{\partial}{\partial \theta_T} \ell(y,d) \\ &= \left(\frac{\partial}{\partial \theta_T} \log \left\{ f_T(y)^d \cdot S_T(y)^{1-d} \right\} \right) + \left(\frac{\partial}{\partial \theta_T} \log \left\{ f_C(y)^{1-d} \cdot S_C(y)^d \right\} \right) \\ &= \left(\frac{\partial}{\partial \theta_T} \log \left\{ f_T(y)^d \cdot S_T(y)^{1-d} \right\} \right) + 0 \\ &= \frac{\partial}{\partial \theta_T} \log \left\{ f_T(y)^d \cdot S_T(y)^{1-d} \right\} \end{split}$$

Thus, the MLE for θ_T won't depend on θ_C , and we can ignore the distribution of C when estimating the parameters of $f_T(t) = p(T = t)$.

Then:

$$\begin{split} \mathcal{L}(y,d) &= f_T(y)^d \cdot S_T(y)^{1-d} \\ &= \left(h_T(y)^d S_T(y)^d \right) \cdot S_T(y)^{1-d} \\ &= h_T(y)^d \cdot S_T(y)^d \cdot S_T(y)^{1-d} \\ &= h_T(y)^d \cdot S_T(y) \\ &= S_T(y) \cdot h_T(y)^d \end{split}$$

That is, if the event occurred at time y (i.e., if d = 1), then the likelihood of (Y, D) = (y, d) is equal to the hazard function at y times the survival function at y. Otherwise, the likelihood is equal to just the survival function at y.

The corresponding log-likelihood is:

$$\begin{split} \ell(y,d) &= \log \left\{ \mathcal{L}(y,d) \right\} \\ &= \log \left\{ S_T(y) \cdot h_T(y)^d \right\} \\ &= \log \left\{ S_T(y) \right\} + \log \left\{ h_T(y)^d \right\} \\ &= \log \left\{ S_T(y) \right\} + d \cdot \log \left\{ h_T(y) \right\} \\ &= -H_T(y) + d \cdot \log \left\{ h_T(y) \right\} \end{split}$$

In other words, the log-likelihood contribution from a single observation (Y, D) = (y, d) is equal to the negative cumulative hazard at y, plus the log of the hazard at y if the event occurred at time y.

5.4. Parametric Models for Time-to-Event Outcomes

5.4.1. Exponential Distribution

- The exponential distribution is the base distribution for survival analysis.
- The distribution has a constant hazard λ
- The mean survival time is λ^{-1}

5.4.1.1. Mathematical details of exponential distribution

$$\begin{split} f(t) &= \lambda \mathrm{e}^{-\lambda t} \\ E(t) &= \lambda^{-1} \\ Var(t) &= \lambda^{-2} \\ F(t) &= 1 - \mathrm{e}^{-\lambda x} \\ S(t) &= \mathrm{e}^{-\lambda x} \\ \ln(S(t)) &= -\lambda x \\ h(t) &= -\frac{f(t)}{S(t)} = -\frac{\lambda \mathrm{e}^{-\lambda t}}{\mathrm{e}^{-\lambda t}} = \lambda \end{split}$$

5.4.1.2. Estimating λ

• Suppose we have m exponential survival times of t_1,t_2,\dots,t_m and k right-censored values at $u_1,u_2,\dots,u_k.$

- A survival time of $t_i = 10$ means that subject i died at time 10. A right-censored time $u_i = 10$ means that at time 10, subject i was still alive and that we have no further follow-up.
- For the moment we will assume that the survival distribution is exponential and that all the subjects have the same parameter λ .

We have m exponential survival times of t_1,t_2,\ldots,t_m and k right-censored values at u_1,u_2,\ldots,u_k . The log-likelihood of an observed survival time t_i is

$$\log\left\{\lambda\mathrm{e}^{-\lambda t_i}\right\} = \log\left\{\lambda\right\} - \lambda t_i$$

and the likelihood of a censored value is the probability of that outcome (survival greater than u_i) so the log-likelihood is

$$\begin{split} \ell_j(\lambda) &= \log \left\{ \lambda \mathrm{e}^{u_j} \right\} \\ &= -\lambda u_j \end{split}$$

_ _ _ _ _ _

Theorem 5.6. Let $T = \sum t_i$ and $U = \sum u_j$. Then:

$$\hat{\lambda}_{ML} = \frac{m}{T + U} \tag{5.3}$$

Proof.

$$\begin{split} \ell(\lambda) &= \sum_{i=1}^m (\ln \lambda - \lambda t_i) + \sum_{j=1}^k (-\lambda u_j) \\ &= m \ln \lambda - (T+U) \lambda \\ \ell'(\lambda) &= m \lambda^{-1} - (T+U) \\ \hat{\lambda} &= \frac{m}{T+U} \end{split}$$

 $\ell'' = -m/\lambda^2$ < 0 $\hat{E}[T] = \hat{\lambda}^{-1}$ $= \frac{T+U}{m}$

5.4.1.3. Fisher Information and Standard Error

$$\begin{split} E[-\ell''] &= m/\lambda^2 \\ \operatorname{Var}\left(\hat{\lambda}\right) &\approx \left(E[-\ell'']\right)^{-1} \\ &= \lambda^2/m \\ \operatorname{SE}\left(\hat{\lambda}\right) &= \sqrt{\operatorname{Var}\left(\hat{\lambda}\right)} \\ &\approx \lambda/\sqrt{m} \end{split}$$

 $\hat{\lambda}$ depends on the censoring times of the censored observations, but Var $(\hat{\lambda})$ only depends on the number of uncensored observations, m, and not on the number of censored observations (k).

5.4.2. Other Parametric Survival Distributions

- Any density on $[0, \infty)$ can be a survival distribution, but the most useful ones are all skew right.
- The most frequently used generalization of the exponential is the Weibull.
- Other common choices are the gamma, log-normal, log-logistic, Gompertz, inverse Gaussian, and Pareto.
- Most of what we do going forward is non-parametric or semiparametric, but sometimes these parametric distributions provide a useful approach.

5.5. Nonparametric Survival Analysis

5.5.1. Basic ideas

- Mostly, we work without a parametric model.
- The first task is to estimate a survival function from data listing survival times, and censoring times for censored data.
- For example one patient may have relapsed at 10 months. Another might have been followed for 32 months without a relapse having occurred (censored).
- The minimum information we need for each patient is a time and a censoring variable which is 1 if the event occurred at the indicated time and 0 if this is a censoring time.

5.6. Example: clinical trial for pediatric acute leukemia

5.6.1. Overview of study

This is from a clinical trial in 1963 for 6-MP treatment vs. placebo for Acute Leukemia in 42 children.

- Pairs of children:
 - matched by remission status at the time of treatment (remstat:
 1 = partial, 2 = complete)
 - randomized to 6-MP (exit times in t2) or placebo (exit times in t1)
- Followed until relapse or end of study.
- All of the placebo group relapsed, but some of the 6-MP group were censored (which means they were still in remission); indicated by relapse variable (0 = censored, 1 = relapse).
- 6-MP = 6-Mercaptopurine (Purinethol) is an anti-cancer ("antineoplastic" or "cytotoxic") chemotherapy drug used currently for Acute lymphoblastic leukemia (ALL). It is classified as an antimetabolite.

5.6.2. Study design

- Clinical trial in 1963 for 6-MP treatment vs. placebo for Acute Leukemia in 42 children.
- Pairs of children:
 - matched by remission status at the time of treatment (remstat)
 - * remstat = 1: partial
 - * remstat = 2: complete

Table 5.1.: drug6mp pediatric acute leukemia data

```
library(KMsurv)
data(drug6mp)
drug6mp = drug6mp |> as_tibble() |> print()
#> # A tibble: 21 x 5
#>
       pair remstat
                         t1
                                t2 relapse
#>
      <int>
               <int> <int> <int>
                                      <int>
    1
           1
                    1
                          1
                                          1
#>
                                10
#>
    2
           2
                    2
                         22
                                7
                                          1
#>
    3
           3
                    2
                          3
                                32
                                          0
    4
           4
                    2
                         12
#>
                                23
                                          1
    5
                    2
#>
           5
                          8
                                22
                                          1
    6
           6
                    1
                         17
                                6
                                          1
#>
#>
    7
           7
                    2
                          2
                                16
                                          1
    8
                    2
           8
                         11
                                34
                                          0
#>
    9
                    2
#>
           9
                          8
                                32
                                          0
#> 10
          10
                         12
                                25
                                          0
#> # i 11 more rows
```

- randomized to 6-MP (exit time: t2) or placebo (t1).
- Followed until relapse or end of study.
 - All of the placebo group relapsed,
 - Some of the 6-MP group were censored.

Table 5.2.: Summary statistics for drug6mp data

| <pre>summary(drug6mp)</pre> | | | | | | | | | |
|-----------------------------|------------|--------------|---------------|--------------|---------------|--|--|--|--|
| #> | pair | remstat | t1 | t2 | relapse | | | | |
| #> | Min. : 1 | Min. :1.00 | Min. : 1.00 | Min. : 6.0 | Min. :0.000 | | | | |
| #> | 1st Qu.: 6 | 1st Qu.:2.00 | 1st Qu.: 4.00 | 1st Qu.: 9.0 | 1st Qu.:0.000 | | | | |
| #> | Median :11 | Median :2.00 | Median: 8.00 | Median :16.0 | Median :0.000 | | | | |
| #> | Mean :11 | Mean :1.76 | Mean : 8.67 | Mean :17.1 | Mean :0.429 | | | | |
| #> | 3rd Qu.:16 | 3rd Qu.:2.00 | 3rd Qu.:12.00 | 3rd Qu.:23.0 | 3rd Qu.:1.000 | | | | |
| #> | Max. :21 | Max. :2.00 | Max. :23.00 | Max. :35.0 | Max. :1.000 | | | | |

5.6.3. Data documentation for drug6mp

```
# library(printr) # inserts help-file output into markdown output
library(KMsurv)
?drug6mp
```

5.6.4. Descriptive Statistics

- The average time in each group is not useful. Some of the 6-MP patients have not relapsed at the time recorded, while all of the placebo patients have relapsed.
- The median time is not really useful either because so many of the 6-MP patients have not relapsed (12/21).
- Both are biased down in the 6-MP group. Remember that lower times are worse since they indicate sooner recurrence.

5.6.5. Exponential model

• We can compute the hazard rate, assuming an exponential model: number of relapses divided by the sum of the exit times (Equation 5.3).

$$\hat{\lambda} = \frac{\sum_{i=1}^{n} D_i}{\sum_{i=1}^{n} Y_i}$$

• For the placebo, that is just the reciprocal of the mean time:

$$\hat{\lambda}_{\text{placebo}} = \frac{\sum_{i=1}^{n} D_i}{\sum_{i=1}^{n} Y_i}$$

$$= \frac{\sum_{i=1}^{n} 1}{\sum_{i=1}^{n} Y_i}$$

$$= \frac{n}{\sum_{i=1}^{n} Y_i}$$

$$= \frac{1}{\bar{Y}}$$

$$= \frac{1}{8.6667}$$

$$= 0.1154$$

• For the 6-MP group, $\hat{\lambda} = 9/359 = 0.025$

$$\hat{\lambda}_{6-\text{MP}} = \frac{\sum_{i=1}^{n} D_i}{\sum_{i=1}^{n} Y_i}$$

$$= \frac{9}{359}$$

$$= 0.0251$$

• The estimated hazard in the placebo group is 4.6 times as large as in the 6-MP group (assuming the hazard is constant over time).

5.7. The Kaplan-Meier Product Limit Estimator

5.7.1. Estimating survival in datasets without censoring

In the drug6mp dataset, the estimated survival function for the placebo patients is easy to compute. For any time t in months, S(t) is the fraction of patients with times greater than t:

5.7.2. Estimating survival in datasets with censoring

- For the 6-MP patients, we cannot ignore the censored data because we know that the time to relapse is greater than the censoring time.
- For any time t in months, we know that 6-MP patients with times greater than t have not relapsed, and those with relapse time less than t have relapsed, but we don't know if patients with censored time less than t have relapsed or not.
- The procedure we usually use is the Kaplan-Meier product-limit estimator of the survival function.
- The Kaplan-Meier estimator is a step function (like the empirical cdf), which changes value only at the event times, not at the censoring times.
- At each event time t, we compute the at-risk group size Y, which is all those observations whose event time or censoring time is at least t.

• If d of the observations have an event time (not a censoring time) of t, then the group of survivors immediately following time t is reduced by the fraction

$$\frac{Y-d}{Y} = 1 - \frac{d}{Y}$$

Definition 5.5 (Kaplan-Meier Product-Limit Estimator of Survival Function). If the event times are t_i with events per time of d_i $(1 \le i \le k)$, then the **Kaplan-Meier Product-Limit Estimator** of the survival function¹ is:

$$\hat{S}(t) = \prod_{t_i < t} \left[\frac{1 - d_i}{Y_i} \right] \tag{5.4}$$

where Y_i is the set of observations whose time (event or censored) is $\geq t_i$, the group at risk at time t_i .

Theorem 5.7 (Kaplan-Meier Estimate with No Censored Observations). If there are no censored data, and there are n data points, then just after (say) the third event time

 $^{^1}$ @def-surv-fn

$$\begin{split} \hat{S}(t) &= \prod_{t_i < t} \left[1 - \frac{d_i}{Y_i} \right] \\ &= \left[\frac{n - d_1}{n} \right] \left[\frac{n - d_1 - d_2}{n - d_1} \right] \left[\frac{n - d_1 - d_2 - d_3}{n - d_1 - d_2} \right] \\ &= \frac{n - d_1 - d_2 - d_3}{n} \\ &= 1 - \frac{d_1 + d_2 + d_3}{n} \\ &= 1 - \hat{F}(t) \end{split}$$

where $\hat{F}(t)$ is the usual empirical CDF estimate.

5.7.3. Kaplan-Meier curve for drug6mp data

Here is the Kaplan-Meier estimated survival curve for the patients who received 6-MP in the drug6mp dataset (we will see code to produce figures like this one shortly):

```
# | echo: false

require(KMsurv)
data(drug6mp)
library(dplyr)
library(survival)

drug6mp_km_model1 =
    drug6mp |>
    mutate(surv = Surv(t2, relapse)) |>
    survfit(formula = surv ~ 1, data = _)

library(ggfortify)
```

```
drug6mp_km_model1 |>
  autoplot(
    mark.time = TRUE,
    conf.int = FALSE) +
  expand_limits(y = 0) +
  xlab('Time since diagnosis (months)') +
  ylab("KM Survival Curve")
```




Figure 5.7.: Kaplan-Meier Survival Curve for 6-MP Patients

5.7.4. Kaplan-Meier calculations

Let's compute these estimates and build the chart by hand:

```
library(KMsurv)
library(dplyr)
data(drug6mp)
drug6mp.v2 =
 drug6mp |>
 as_tibble() |>
  mutate(
   remstat = remstat |>
      case_match(
      1 ~ "partial",
        2 ~ "complete"
      ),
    # renaming to "outcome" while relabeling is just a style choice:
    outcome = relapse |>
      case_match(
       0 ~ "censored",
       1 ~ "relapsed"
  )
km.6mp =
  drug6mp.v2 |>
  summarize(
    .by = t2,
    Relapses = sum(outcome == "relapsed"),
    Censored = sum(outcome == "censored")) |>
  # here we add a start time row, so the graph starts at time 0:
  bind_rows(
    tibble(
      t2 = 0,
      Relapses = 0,
      Censored = 0)
```

```
# sort in time order:
arrange(t2) |>
mutate(
    Exiting = Relapses + Censored,
    `Study Size` = sum(Exiting),
    Exited = cumsum(Exiting) |> dplyr::lag(default = 0),
    `At Risk` = `Study Size` - Exited,
    Hazard = Relapses / `At Risk`,
    `KM Factor` = 1 - Hazard,
    `Cumulative Hazard` = cumsum(`Hazard`),
    `KM Survival Curve` = cumprod(`KM Factor`)
)

library(pander)
pander(km.6mp)
```

| | | | | | | | | | | KM |
|----|------|----------------------|-------------------|----------------|------|---------------------|-------|------------------|---------|----------|
| | | | | | | | | KM | | Sur- |
| | | | | Study | | At | | Fac- | Cumulat | ivevival |
| t2 | Rela | ps & enso | or E okiti | n § ize | Exit | eRisk | Haza | rdtor | Hazard | Curve |
| 0 | 0 | 0 | 0 | 21 | 0 | 21 | 0 | 1 | 0 | 1 |
| 6 | 3 | 1 | 4 | 21 | 0 | 21 | 0.142 | 90.8571 | 0.1429 | 0.8571 |
| 7 | 1 | 0 | 1 | 21 | 4 | 17 | 0.058 | 8 2 .9412 | 0.2017 | 0.8067 |
| 9 | 0 | 1 | 1 | 21 | 5 | 16 | 0 | 1 | 0.2017 | 0.8067 |
| 10 | 1 | 1 | 2 | 21 | 6 | 15 | 0.066 | 6 7 .9333 | 0.2683 | 0.7529 |
| 11 | 0 | 1 | 1 | 21 | 8 | 13 | 0 | 1 | 0.2683 | 0.7529 |
| 13 | 1 | 0 | 1 | 21 | 9 | 12 | 0.083 | 3 3 .9167 | 0.3517 | 0.6902 |
| 16 | 1 | 0 | 1 | 21 | 10 | 11 | 0.090 | 9 0 .9091 | 0.4426 | 0.6275 |
| 17 | 0 | 1 | 1 | 21 | 11 | 10 | 0 | 1 | 0.4426 | 0.6275 |
| 19 | 0 | 1 | 1 | 21 | 12 | 9 | 0 | 1 | 0.4426 | 0.6275 |
| 20 | 0 | 1 | 1 | 21 | 13 | 8 | 0 | 1 | 0.4426 | 0.6275 |

| | | | | | | | | | | KM |
|----|------|----------------------|-------------------|-----------------|------|---------------------|-------|----------|---------|-----------|
| | | | | | | | | KM | | Sur- |
| | | | | Study | | At | | Fac- | Cumulat | tivevival |
| t2 | Rela | ps & enso | or Ec kiti | in § ize | Exit | eRisk | Haza | rdtor | Hazard | Curve |
| 22 | 1 | 0 | 1 | 21 | 14 | 7 | 0.142 | 290.8571 | 0.5854 | 0.5378 |
| 23 | 1 | 0 | 1 | 21 | 15 | 6 | 0.166 | 570.8333 | 0.7521 | 0.4482 |
| 25 | 0 | 1 | 1 | 21 | 16 | 5 | 0 | 1 | 0.7521 | 0.4482 |
| 32 | 0 | 2 | 2 | 21 | 17 | 4 | 0 | 1 | 0.7521 | 0.4482 |
| 34 | 0 | 1 | 1 | 21 | 19 | 2 | 0 | 1 | 0.7521 | 0.4482 |
| 35 | 0 | 1 | 1 | 21 | 20 | 1 | 0 | 1 | 0.7521 | 0.4482 |
| 50 | J | 1 | | - 1 | 20 | 1 | J | 1 | 0.1021 | 0.110 |

5.7.4.1. Summary

For the 6-MP patients at time 6 months, there are 21 patients at risk. At t = 6 there are 3 relapses and 1 censored observations.

The Kaplan-Meier factor is (21-3)/21 = 0.857. The number at risk for the next time (t=7) is 21-3-1=17.

At time 7 months, there are 17 patients at risk. At t=7 there is 1 relapse and 0 censored observations. The Kaplan-Meier factor is (17-1)/17 = 0.941. The Kaplan Meier estimate is $0.857 \times 0.941 = 0.807$. The number at risk for the next time (t=9) is 17-1=16.

Now, let's graph this estimated survival curve using ggplot():

```
library(ggplot2)
conflicts_prefer(dplyr::filter)
km.6mp |>
    ggplot(aes(x = t2, y = `KM Survival Curve`)) +
    geom_step() +
    geom_point(data = km.6mp |> filter(Censored > 0), shape = 3) +
    expand_limits(y = c(0,1), x = 0) +
    xlab('Time since diagnosis (months)') +
    ylab("KM Survival Curve") +
    scale_y_continuous(labels = scales::percent)
```




Figure 5.8.: KM curve for 6MP patients, calculated by hand

5.8. Using the survival package in R

We don't have to do these calculations by hand every time; the survival package and several others have functions available to automate many of these tasks (full list: https://cran.r-project.org/web/views/Survival.html).

5.8.1. The Surv function

To use the survival package, the first step is telling R how to combine the exit time and exit reason (censoring versus event) columns. The Surv() function accomplishes this task.

5.8.1.1. Example: Surv() with drug6mp data

```
library(survival)
   drug6mp.v3 =
     drug6mp.v2 |>
     mutate(
       surv2 = Surv(
         time = t2,
6
         event = (outcome == "relapsed")))
   print(drug6mp.v3)
9
   #> # A tibble: 21 x 7
          pair remstat
                             t1
                                   t2 relapse outcome
                                                          surv2
11
         <int> <chr>
                          <int> <int>
                                                         <Surv>
   #>
                                         <int> <chr>
              1 partial
                             1
                                                            10
   #>
       1
                                   10
                                             1 relapsed
       2
              2 complete
                                   7
                                                             7
                             22
                                             1 relapsed
14
       3
              3 complete
                             3
                                   32
                                             0 censored
                                                            32+
15
   #>
              4 complete
                             12
                                   23
                                             1 relapsed
                                                            23
16
```

| 17 | #> | 5 | 5 | complete | 8 | 22 | 1 relapsed | 22 |
|----|---------------------|----|----|----------|----|----|------------|-----|
| 18 | #> | 6 | 6 | partial | 17 | 6 | 1 relapsed | 6 |
| 19 | #> | 7 | 7 | complete | 2 | 16 | 1 relapsed | 16 |
| 20 | #> | 8 | 8 | complete | 11 | 34 | 0 censored | 34+ |
| 21 | #> | 9 | 9 | complete | 8 | 32 | 0 censored | 32+ |
| 22 | #> | 10 | 10 | complete | 12 | 25 | 0 censored | 25+ |
| 23 | #> # i 11 more rows | | | | | | | |

The output of Surv() is a vector of objects with class Surv. When we print this vector:

- observations where the event was observed are printed as the event time (for example, surv2 = 10 on line 1)
- observations where the event was right-censored are printed as the censoring time with a plus sign (+; for example, surv2 = 32+ on line 3).

5.8.2. The survfit function

Once we have constructed our Surv variable, we can calculate the Kaplan-Meier estimate of the survival curve using the survfit() function.

Note

The documentation for ?survfit isn't too helpful; the survfit.formula documentation is better.

5.8.2.1. Example: survfit() with drug6mp data

Here we use survfit() to create a survfit object, which contains the Kaplan-Meier estimate:

```
drug6mp.km_model = survfit(
  formula = surv2 ~ 1,
  data = drug6mp.v3)
```

print.survfit() just gives some summary statistics:

summary.survfit() shows us the underlying Kaplan-Meier table:

```
summary(drug6mp.km_model)
#> Call: survfit(formula = surv2 ~ 1, data = drug6mp.v3)
#>
#>
    time n.risk n.event survival std.err lower 95% CI upper 95% CI
#>
       6
             21
                       3
                            0.857
                                    0.0764
                                                   0.720
                                                                 1.000
#>
       7
             17
                       1
                            0.807
                                    0.0869
                                                   0.653
                                                                 0.996
#>
      10
             15
                       1
                            0.753
                                    0.0963
                                                   0.586
                                                                 0.968
#>
      13
             12
                       1
                            0.690
                                    0.1068
                                                   0.510
                                                                 0.935
                       1
#>
      16
              11
                            0.627
                                    0.1141
                                                   0.439
                                                                 0.896
#>
      22
              7
                       1
                            0.538
                                    0.1282
                                                   0.337
                                                                 0.858
#>
      23
              6
                            0.448 0.1346
                                                   0.249
                                                                 0.807
```

We can specify which time points we want using the times argument:

```
summary(
  drug6mp.km_model,
  times = c(0, drug6mp.v3$t2))
#> Call: survfit(formula = surv2 ~ 1, data = drug6mp.v3)
#>
    time n.risk n.event survival std.err lower 95% CI upper 95% CI
#>
       0
              21
                                                     1.000
#>
                        0
                              1.000
                                     0.0000
                                                                    1.000
       6
                                     0.0764
              21
                        3
                              0.857
                                                     0.720
                                                                    1.000
#>
       6
#>
              21
                        0
                              0.857
                                     0.0764
                                                     0.720
                                                                    1.000
#>
       6
              21
                        0
                              0.857
                                     0.0764
                                                     0.720
                                                                    1.000
       6
#>
              21
                        0
                              0.857
                                     0.0764
                                                     0.720
                                                                    1.000
       7
              17
                              0.807
                                     0.0869
#>
                        1
                                                     0.653
                                                                   0.996
       9
#>
              16
                        0
                              0.807
                                     0.0869
                                                     0.653
                                                                   0.996
      10
                        1
#>
              15
                              0.753
                                     0.0963
                                                     0.586
                                                                   0.968
      10
              15
                        0
                              0.753
                                     0.0963
                                                                   0.968
#>
                                                     0.586
#>
      11
              13
                        0
                              0.753
                                     0.0963
                                                     0.586
                                                                    0.968
#>
      13
              12
                        1
                              0.690
                                     0.1068
                                                     0.510
                                                                   0.935
                              0.627
                                     0.1141
#>
      16
              11
                        1
                                                     0.439
                                                                   0.896
      17
              10
                        0
                              0.627
                                     0.1141
                                                     0.439
                                                                   0.896
#>
               9
                        0
                              0.627
#>
      19
                                     0.1141
                                                     0.439
                                                                   0.896
      20
               8
                        0
                              0.627
                                     0.1141
#>
                                                     0.439
                                                                   0.896
      22
               7
                        1
                              0.538
                                     0.1282
                                                     0.337
                                                                   0.858
#>
#>
      23
               6
                        1
                              0.448
                                     0.1346
                                                     0.249
                                                                   0.807
#>
      25
               5
                        0
                              0.448
                                     0.1346
                                                     0.249
                                                                   0.807
#>
      32
               4
                        0
                              0.448
                                     0.1346
                                                     0.249
                                                                   0.807
      32
               4
                        0
#>
                              0.448
                                     0.1346
                                                     0.249
                                                                   0.807
#>
      34
               2
                        0
                              0.448
                                     0.1346
                                                     0.249
                                                                   0.807
      35
                        0
                              0.448 0.1346
                                                     0.249
                                                                   0.807
#>
```

?summary.survfit

5.8.3. Plotting estimated survival functions

We can plot survfit objects with plot(), autoplot(), or ggsurvplot():

```
library(ggfortify)
autoplot(drug6mp.km_model)
```




Figure 5.9.: Kaplan-Meier Survival Curve for 6-MP Patients

```
# not shown:
# plot(drug6mp.km_model)
```

$5. \ Introduction \ to \ Survival \ Analysis$

```
# library(survminer)
# ggsurvplot(drug6mp.km_model)
```

5.8.3.1. quantiles of survival curve

We can extract quantiles with quantile():

```
drug6mp.km_model |>
 quantile(p = c(.25, .5)) |>
 as_tibble() |>
 mutate(p = c(.25, .5)) |>
  relocate(p, .before = everything())
#> # A tibble: 2 x 4
         p quantile lower upper
     <dbl>
           <dbl> <dbl> <dbl>
#> 1 0.25
                 13
                       6
                            NA
#> 2 0.5
                 23
                      16
                            NA
```

5.9. Two-sample tests

5.9.1. The survdiff function

```
?survdiff
```

5.9.2. Example: survdiff() with drug6mp data

Now we are going to compare the placebo and 6-MP data. We need to reshape the data to make it usable with the standard survival workflow:


```
library(survival)
library(tidyr)
drug6mp.v4 =
  drug6mp.v3 |>
  select(pair, remstat, t1, t2, outcome) |>
  # here we are going to change the data from a wide format to long:
  pivot_longer(
    cols = c(t1, t2),
    names_to = "treatment",
    values_to = "exit_time") |>
  mutate(
    treatment = treatment |>
      case_match(
        "t1" ~ "placebo",
        "t2" ~ "6-MP"
      ),
    outcome = if_else(
      treatment == "placebo",
      "relapsed",
      outcome
    ),
    surv = Surv(
      time = exit_time,
      event = (outcome == "relapsed"))
```

Using this long data format, we can fit a Kaplan-Meier curve for each treatment group simultaneously:

```
drug6mp.km_model2 =
    survfit(
  formula = surv ~ treatment,
  data = drug6mp.v4)
```

We can plot the curves in the same graph:

drug6mp.km_model2 |> autoplot()



We can also perform something like a t-test, where the null hypothesis is that the curves are the same:

```
survdiff(
 formula = surv ~ treatment,
 data = drug6mp.v4)
#> Call:
#> survdiff(formula = surv ~ treatment, data = drug6mp.v4)
#>
#>
                      N Observed Expected (O-E)^2/E (O-E)^2/V
#> treatment=6-MP
                     21
                                9
                                      19.3
                                                5.46
                                                          16.8
#> treatment=placebo 21
                              21
                                      10.7
                                                9.77
                                                          16.8
#>
    Chisq= 16.8 on 1 degrees of freedom, p= 4e-05
#>
```

By default, survdiff() ignores any pairing, but we can use strata() to perform something similar to a paired t-test:

```
survdiff(
 formula = surv ~ treatment + strata(pair),
 data = drug6mp.v4)
#> Call:
#> survdiff(formula = surv ~ treatment + strata(pair), data = drug6mp.v4)
#>
#>
                      N Observed Expected (O-E)^2/E (O-E)^2/V
#> treatment=6-MP
                     21
                               9
                                      16.5
                                                3.41
                                                          10.7
#> treatment=placebo 21
                              21
                                      13.5
                                                4.17
                                                          10.7
#>
#>
    Chisq= 10.7 on 1 degrees of freedom, p= 0.001
```

Interestingly, accounting for pairing reduces the significant of the difference.

5.10. Example: Bone Marrow Transplant Data

| Data from | Copelan et al. | (1991) | |
|-----------|----------------|--------|--|
| | | | |
| | | | |




Figure 1.1 Recovery Process from a Bone Marrow Transplant

Figure 5.10.: Recovery process from a bone marrow transplant (Fig. 1.1 from Klein and Moeschberger (2003))

313

5.10.1. Study design

- * Treatment
 - allogeneic (from a donor) bone marrow transplant therapy
- * Inclusion criteria
 - acute myeloid leukemia (AML)
 - acute lymphoblastic leukemia (ALL).
- * Possible intermediate events
 - graft vs. host disease (GVHD): an immunological rejection response to the transplant
 - platelet recovery: a return of platelet count to normal levels.

One or the other, both in either order, or neither may occur.

End point events

- relapse of the disease
- death

Any or all of these events may be censored.

5.10.2. KMsurv::bmt data in R

library(KMsurv)
?bmt

5.10.3. Analysis plan

- We concentrate for now on disease-free survival (t2 and d3) for the three risk groups, ALL, AML Low Risk, and AML High Risk.
- We will construct the Kaplan-Meier survival curves, compare them, and test for differences.
- We will construct the cumulative hazard curves and compare them.
- We will estimate the hazard functions, interpret, and compare them.

5.10.4. Survival Function Estimate and Variance

$$\hat{S}(t) = \prod_{t_i < t} \left[1 - \frac{d_i}{Y_i} \right]$$

where Y_i is the group at risk at time t_i .

The estimated variance of $\hat{S}(t)$ is:

Theorem 5.8 (Greenwood's estimator for variance of Kaplan-Meier survival estimator).

$$\widehat{Var}\left(\hat{S}(t)\right) = \hat{S}(t)^2 \sum_{t_i < t} \frac{d_i}{Y_i(Y_i - d_i)} \tag{5.5}$$

We can use Equation 5.5 for confidence intervals for a survival function or a difference of survival functions.

Kaplan-Meier survival curves

```
library(KMsurv)
library(survival)
data(bmt)

bmt =
    bmt |>
    as_tibble() |>
    mutate(
        group =
            group |>
            factor(
                labels = c("ALL","Low Risk AML","High Risk AML")),
        surv = Surv(t2,d3))

km_model1 = survfit(
    formula = surv ~ group,
    data = bmt)
```

```
library(ggfortify)
autoplot(
   km_model1,
   conf.int = TRUE,
   ylab = "Pr(disease-free survival)",
   xlab = "Time since transplant (days)") +
   theme_bw() +
   theme(legend.position="bottom")
```



Figure 5.11.: Disease-Free Survival by Disease Group

5.10.5. Understanding Greenwood's formula (optional)

To see where Greenwood's formula comes from, let $x_i = Y_i - d_i$. We approximate the solution treating each time as independent, with Y_i fixed and ignore randomness in times of failure and we treat x_i as independent binomials $Bin(Y_i, p_i)$. Letting S(t) be the "true" survival function

$$\begin{split} \hat{S}(t) &= \prod_{t_i < t} x_i / Y_i \\ S(t) &= \prod_{t_i < t} p_i \end{split}$$

$$S(t) = \prod_{t_i < t} p_i$$

$$\begin{split} \frac{\hat{S}(t)}{S(t)} &= \prod_{t_i < t} \frac{x_i}{\cdot} p_i Y_i \\ &= \prod_{t_i < t} \frac{\hat{p}_i}{p_i} \\ &= \prod_{t_i < t} \left(1 + \frac{\hat{p}_i - p_i}{p_i} \right) \\ &\approx 1 + \sum_{t_i < t} \frac{\hat{p}_i - p_i}{p_i} \end{split}$$

$$\begin{split} \operatorname{Var}\left(\frac{\hat{S}(t)}{S(t)}\right) &\approx \operatorname{Var}\left(1 + \sum_{t_i < t} \frac{\hat{p}_i - p_i}{p_i}\right) \\ &= \sum_{t_i < t} \frac{1}{p_i^2} \frac{p_i (1 - p_i)}{Y_i} \\ &= \sum_{t_i < t} \frac{(1 - p_i)}{p_i Y_i} \\ &\approx \sum_{t_i < t} \frac{(1 - x_i / Y_i)}{x_i} \\ &= \sum_{t_i < t} \frac{Y_i - x_i}{x_i Y_i} \\ &= \sum_{t_i < t} \frac{d_i}{Y_i (Y_i - d_i)} \\ &: \cdot \operatorname{Var}\left(\hat{S}(t)\right) \approx \hat{S}(t)^2 \sum_{t_i < t} \frac{d_i}{Y_i (Y_i - d_i)} \end{split}$$

5.10.6. Test for differences among the disease groups

Here we compute a chi-square test for assocation between disease group (group) and disease-free survival:

```
survdiff(surv ~ group, data = bmt)
#> Call:
#> survdiff(formula = surv ~ group, data = bmt)
#>
                        N Observed Expected (0-E)^2/E (0-E)^2/V
#>
#> group=ALL
                       38
                                 24
                                        21.9
                                                 0.211
                                                           0.289
#> group=Low Risk AML 54
                                25
                                        40.0
                                                 5.604
                                                          11.012
#> group=High Risk AML 45
                                34
                                        21.2
                                                 7.756
                                                          10.529
#>
   Chisq= 13.8 on 2 degrees of freedom, p= 0.001
```

5.10.7. Cumulative Hazard

$$\begin{split} h(t) &\stackrel{\text{def}}{=} P(T=t|T \geq t) \\ &= \frac{p(T=t)}{P(T \geq t)} \\ &= -\frac{\partial}{\partial t} \log \left\{ S(t) \right\} \end{split}$$

The cumulative hazard (or integrated hazard) function is

$$H(t) \stackrel{\text{def}}{=} \int_0^t h(t)dt$$

Since $h(t) = -\frac{\partial}{\partial t} \mathrm{log} \left\{ S(t) \right\}$ as shown above, we have:

$$H(t) = -\log\{S\}(t)$$

So we can estimate H(t) as:

$$\begin{split} \hat{H}(t) &= -\mathrm{log}\left\{\hat{S}(t)\right\} \\ &= -\mathrm{log}\left\{\prod_{t_i < t} \left[1 - \frac{d_i}{Y_i}\right]\right\} \\ &= -\sum_{t_i < t} \mathrm{log}\left\{1 - \frac{d_i}{Y_i}\right\} \end{split}$$

This is the Kaplan-Meier (product-limit) estimate of cumulative hazard.

5.10.7.1. Example: Cumulative Hazard Curves for Bone-Marrow Transplant (bmt) data

```
autoplot(
  fun = "cumhaz",
  km_model1,
  conf.int = FALSE,
  ylab = "Cumulative hazard (disease-free survival)",
  xlab = "Time since transplant (days)") +
  theme_bw() +
  theme(legend.position="bottom")
```



Figure 5.12.: Disease-Free Cumulative Hazard by Disease Group

5.11. Nelson-Aalen Estimates of Cumulative Hazard and Survival

Definition 5.6 (Nelson-Aalen Cumulative Hazard Estimator).

The point hazard at time t_i can be estimated by d_i/Y_i , which leads to the Nelson-Aalen estimator of the cumulative hazard:

$$\hat{H}_{NA}(t) \stackrel{\text{def}}{=} \sum_{t_i < t} \frac{d_i}{Y_i} \tag{5.6}$$

Theorem 5.9 (Variance of Nelson-Aalen estimator).

The variance of this estimator is approximately:

$$\begin{split} \hat{Var}\left(\hat{H}_{NA}(t)\right) &= \sum_{t_i < t} \frac{(d_i/Y_i)(1 - d_i/Y_i)}{Y_i} \\ &\approx \sum_{t_i < t} \frac{d_i}{Y_i^2} \end{split} \tag{5.7}$$

Since $S(t) = \exp\{-H(t)\}$, the Nelson-Aalen cumulative hazard estimate can be converted into an alternate estimate of the survival function:

$$\begin{split} \hat{S}_{NA}(t) &= \exp\left\{-\hat{H}_{NA}(t)\right\} \\ &= \exp\left\{-\sum_{t_i < t} \frac{d_i}{Y_i}\right\} \\ &= \prod_{t_i < t} \exp\left\{-\frac{d_i}{Y_i}\right\} \end{split}$$

Compare these with the corresponding Kaplan-Meier estimates:

$$\begin{split} \hat{H}_{KM}(t) &= -\sum_{t_i < t} \log \left\{ 1 - \frac{d_i}{Y_i} \right\} \\ \hat{S}_{KM}(t) &= \prod_{t_i < t} \left[1 - \frac{d_i}{Y_i} \right] \end{split}$$

The product limit estimate and the Nelson-Aalen estimate often do not differ by much. The latter is considered more accurate in small samples and also directly estimates the cumulative hazard. The "fleming-harrington" method for survfit() reduces to Nelson-Aalen when the data are unweighted. We can also estimate the cumulative hazard as the negative log of the KM survival function estimate.

5.11.1. Application to bmt dataset

```
na_fit = survfit(
  formula = surv ~ group,
  type = "fleming-harrington",
  data = bmt)

km_fit = survfit(
  formula = surv ~ group,
  type = "kaplan-meier",
  data = bmt)

km_and_na =
  bind_rows(
    .id = "model",
    "Kaplan-Meier" = km_fit |> fortify(surv.connect = TRUE),
    "Nelson-Aalen" = na_fit |> fortify(surv.connect = TRUE)
) |>
  as_tibble()
```

```
km_and_na |>
  ggplot(aes(x = time, y = surv, col = model)) +
  geom_step() +
  facet_grid(. ~ strata) +
```

```
theme_bw() +
ylab("S(t) = P(T>=t)") +
xlab("Survival time (t, days)") +
theme(legend.position = "bottom")
```




Figure 5.13.: Kaplan-Meier and Nelson-Aalen Survival Function Estimates, stratified by disease group

The Kaplan-Meier and Nelson-Aalen survival estimates are very similar for this dataset.

Configuring R

Functions from these packages will be used throughout this document:

```
library(conflicted) # check for conflicting function definitions
# library(printr) # inserts help-file output into markdown output
library(rmarkdown) # Convert R Markdown documents into a variety of formats.
library(pander) # format tables for markdown
library(ggplot2) # graphics
library(ggeasy) # help with graphics
library(ggfortify) # help with graphics
library(dplyr) # manipulate data
library(tibble) # `tibble`s extend `data.frame`s
library(magrittr) # `%>%` and other additional piping tools
library(haven) # import Stata files
library(knitr) # format R output for markdown
library(tidyr) # Tools to help to create tidy data
library(plotly) # interactive graphics
library(dobson) # datasets from Dobson and Barnett 2018
library(parameters) # format model output tables for markdown
library(haven) # import Stata files
```

```
library(latex2exp) # use LaTeX in R code (for figures and tables)
library(fs) # filesystem path manipulations
library(survival) # survival analysis
library(survminer) # survival analysis graphics
library(KMsurv) # datasets from Klein and Moeschberger
library(parameters) # format model output tables for
library(webshot2) # convert interactive content to static for pdf
library(forcats) # functions for categorical variables ("factors")
library(stringr) # functions for dealing with strings
library(lubridate) # functions for dealing with dates and times
```

Here are some R settings I use in this document:

```
rm(list = ls()) # delete any data that's already loaded into R

conflicts_prefer(dplyr::filter)
ggplot2::theme_set(
    ggplot2::theme_bw() +
        # ggplot2::labs(col = "") +
        ggplot2::theme(
        legend.position = "bottom",
        text = ggplot2::element_text(size = 12, family = "serif")))

knitr::opts_chunk$set(message = FALSE)
options('digits' = 4)

panderOptions("big.mark", ",")
pander::panderOptions("table.emphasize.rownames", FALSE)
pander::panderOptions("table.split.table", Inf)
conflicts_prefer(dplyr::filter) # use the `filter()` function from dplyr() by defaul legend_text_size = 9
```

6.1. Introduction

Recall: the exponential distribution has constant hazard:

$$f(t) = \lambda e^{-\lambda t}$$

$$S(t) = e^{-\lambda t}$$

$$h(t) = \lambda$$

Let's make two generalizations. First, we let the hazard depend on some covariates x_1, x_2, \ldots, x_p ; we will indicate this dependence by extending our notation for hazard:

$$h(t|\tilde{x}) \stackrel{\mathrm{def}}{=} \mathrm{p}(T=t|T \geq t, \tilde{X}=\tilde{x})$$

Definition 6.1 (baseline hazard).

The baseline hazard, base hazard, or reference hazard, denoted $h_0(t)$ or $\lambda_0(t)$, is the hazard function for the subpopulation of individuals whose covariates are all equal to their reference levels:

$$h_0(t) \stackrel{\mathrm{def}}{=} h(t|\tilde{X} = \tilde{0})$$

Similarly:

Definition 6.2 (baseline cumulative hazard).

The baseline cumulative hazard, base cumulative hazard, or reference cumulative hazard, denoted $H_0(t)$ or $\Lambda_0(t)$, is the cumulative hazard function for the subpopulation of individuals whose covariates are all equal to their reference levels:

$$H_0(t) \stackrel{\mathrm{def}}{=} H(t|\tilde{X} = \tilde{0})$$

Also:

Definition 6.3 (Baseline survival function). The **baseline survival function** is the survival function for an individual whose covariates are all equal to their default values.

$$S_0(t) \stackrel{\text{def}}{=} S(t|\tilde{X} = \tilde{0})$$

As the second generalization, we let the base hazard, cumulative hazard, and survival functions depend on t, but not on the covariates (for now). We can do this using either parametric or semi-parametric approaches.

6.2. Cox's Proportional Hazards Model

The generalization is that the hazard function is:

$$\begin{split} h(t|x) &= h_0(t)\theta(x) \\ \theta(x) &= \exp\left\{\eta(x)\right\} \\ \eta(x) &= \tilde{x}^\top \tilde{\beta} \\ &\stackrel{\text{def}}{=} \beta_1 x_1 + \dots + \beta_p x_p \end{split}$$

The relationship between h(t|x) and $\eta(x)$ is typically modeled using a log link, as in a generalized linear model; that is:

$$\log \{h(t|x)\} = \log \{h_0(t)\} + \eta(x)$$

This model is **semi-parametric**, because the linear predictor depends on estimated parameters but the base hazard function is unspecified. There is no constant term in $\eta(x)$, because it is absorbed in the base hazard.

Alternatively, we could define $\beta_0(t) = \log \{h_0(t)\}$, and then:

$$\eta(x,t) = \beta_0(t) + \beta_1 x_1 + \dots + \beta_p x_p$$

For two different individuals with covariate patterns x_1 and x_2 , the ratio of the hazard functions (a.k.a. **hazard ratio**, a.k.a. **relative hazard**) is:

$$\begin{split} \frac{h(t|x_1)}{h(t|x_2)} &= \frac{h_0(t)\theta(x_1)}{h_0(t)\theta(x_2)} \\ &= \frac{\theta(x_1)}{\theta(x_2)} \end{split}$$

Under the proportional hazards model, this ratio (a.k.a. proportion) does not depend on t. This property is a structural limitation of the model; it is called the **proportional hazards assumption**.

Definition 6.4 (proportional hazards). A conditional probability distribution p(T|X) has **proportional hazards** if the hazard ratio $h(t|x_1)/h(t|x_2)$ does not depend on t. Mathematically, it can be written as:

$$\frac{h(t|x_1)}{h(t|x_2)} = \theta(x_1, x_2)$$

As we saw above, Cox's proportional hazards model has this property, with $\theta(x_1,x_2)=rac{\theta(x_1)}{\theta(x_2)}.$

Note

We are using two similar notations, $\theta(x_1, x_2)$ and $\theta(x)$. We can link these notations if we define $\theta(x) \stackrel{\text{def}}{=} \theta(x, 0)$ and $\theta(0) = 1$.

The proportional hazards model also has additional notable properties:

$$\begin{split} \frac{h(t|x_1)}{h(t|x_2)} &= \frac{\theta(x_1)}{\theta(x_2)} \\ &= \frac{\exp\left\{\eta(x_1)\right\}}{\exp\left\{\eta(x_2)\right\}} \\ &= \exp\left\{\eta(x_1) - \eta(x_2)\right\} \\ &= \exp\left\{x_1'\beta - x_2'\beta\right\} \\ &= \exp\left\{(x_1 - x_2)'\beta\right\} \end{split}$$

Hence on the log scale, we have:

$$\begin{split} \log\left\{\frac{h(t|x_1)}{h(t|x_2)}\right\} &= \eta(x_1) - \eta(x_2) \\ &= x_1'\beta - x_2'\beta \\ &= (x_1 - x_2)'\beta \end{split}$$

If only one covariate x_j is changing, then:

$$\begin{split} \log \left\{ \frac{h(t|x_1)}{h(t|x_2)} \right\} &= (x_{1j} - x_{2j}) \cdot \beta_j \\ &\propto (x_{1j} - x_{2j}) \end{split}$$

That is, under Cox's model $h(t|x) = h_0(t) \exp\{x'\beta\}$, the log of the hazard ratio is proportional to the difference in x_j , with the proportionality coefficient equal to β_j .

Further,

$$\log \{h(t|x)\} = \log \{h_0(t)\} + x'\beta$$

That is, the covariate effects are additive on the log-hazard scale; hazard functions for different covariate patterns should be vertical shifts of each other.

See also:

 $https://en.wikipedia.org/wiki/Proportional_hazards_model\#Why_it_is_called_\%22proportional\%22$

6.2.1. Additional properties of the proportional hazards model

If $h(t|x) = h_0(t)\theta(x)$, then:

Theorem 6.1 (Cumulative hazards are also proportional to $H_0(t)$).

$$\begin{split} H(t|x) &\stackrel{def}{=} \int_{u=0}^{t} h(u) du \\ &= \int_{u=0}^{t} h_0(u) \theta(x) du \\ &= \theta(x) \int_{u=0}^{t} h_0(u) du \\ &= \theta(x) H_0(t) \end{split}$$

where $H_0(t) \stackrel{\text{def}}{=} H(t|0) = \int_{u=0}^t h_0(u) du$.

Theorem 6.2 (The logarithms of cumulative hazard should be parallel).

$$\log\{H(t|\tilde{x})\} = \log\{H_0(t)\} + \tilde{x}'\tilde{\beta}$$

Theorem 6.3 (Survival functions are exponential multiples of $S_0(t)$).

$$\begin{split} S(t|x) &= \exp\left\{-H(t|x)\right\} \\ &= \exp\left\{-\theta(x) \cdot H_0(t)\right\} \\ &= \left(\exp\left\{-H_0(t)\right\}\right)^{\theta(x)} \\ &= \left[S_0(t)\right]^{\theta(x)} \end{split}$$

6.2.2. Testing the proportional hazards assumption

The Nelson-Aalen estimate of the cumulative hazard is usually used for estimates of the hazard and often the cumulative hazard.

If the hazards of the three groups are proportional, that means that the ratio of the hazards is constant over t. We can test this using the ratios of the estimated cumulative hazards, which also would be proportional, as shown above.

```
library(KMsurv)
library(survival)
library(dplyr)
data(bmt)
bmt =
```

```
bmt |>
  as_tibble() |>
  mutate(
    group =
      group |>
      factor(
        labels = c("ALL","Low Risk AML","High Risk AML")))
nafit = survfit(
  formula = Surv(t2,d3) ~ group,
 type = "fleming-harrington",
  data = bmt)
bmt_curves = tibble(timevec = 1:1000)
sf1 <- with(nafit[1], stepfun(time,c(1,surv)))</pre>
sf2 <- with(nafit[2], stepfun(time,c(1,surv)))</pre>
sf3 <- with(nafit[3], stepfun(time,c(1,surv)))</pre>
bmt_curves =
  bmt_curves |>
 mutate(
    cumhaz1 = -log(sf1(timevec)),
    cumhaz2 = -log(sf2(timevec)),
    cumhaz3 = -log(sf3(timevec)))
library(ggplot2)
bmt_rel_hazard_plot =
  bmt_curves |>
  ggplot(
```

```
bmt_rel_hazard_plot =
bmt_curves |>
ggplot(
   aes(
    x = timevec,
    y = cumhaz1/cumhaz2)
) +
```

```
geom_line(aes(col = "ALL/Low Risk AML")) +
ylab("Hazard Ratio") +
xlab("Time") +
ylim(0,6) +
geom_line(aes(y = cumhaz3/cumhaz1, col = "High Risk AML/ALL")) +
geom_line(aes(y = cumhaz3/cumhaz2, col = "High Risk AML/Low Risk AML")) +
theme_bw() +
labs(colour = "Comparison") +
theme(legend.position="bottom")
print(bmt_rel_hazard_plot)
```



Figure 6.1.: Hazard Ratios by Disease Group for bmt data

We can zoom in on 30-300 days to take a closer look:

```
bmt_rel_hazard_plot + xlim(c(30,300))
```



Figure 6.2.: Hazard Ratios by Disease Group (30-300 Days)

The cumulative hazard curves should also be proportional

```
library(ggfortify)
plot_cuhaz_bmt =
  bmt |>
  survfit(formula = Surv(t2, d3) ~ group) |>
  autoplot(fun = "cumhaz",
```

```
mark.time = TRUE) +
ylab("Cumulative hazard")

plot_cuhaz_bmt |> print()
```



Figure 6.3.: Disease-Free Cumulative Hazard by Disease Group

```
plot_cuhaz_bmt +
  scale_y_log10() +
  scale_x_log10()
```



Figure 6.4.: Disease-Free Cumulative Hazard by Disease Group (log-scale)

6.2.3. Smoothed hazard functions

The Nelson-Aalen estimate of the cumulative hazard is usually used for estimates of the hazard. Since the hazard is the derivative of the cumulative hazard, we need a smooth estimate of the cumulative hazard, which is provided by smoothing the step-function cumulative hazard.

The R package muhaz handles this for us. What we are looking for is whether the hazard function is more or less the same shape, increasing, decreasing, constant, etc. Are the hazards "proportional"?

```
library(muhaz)
muhaz(bmt$t2,bmt$d3,bmt$group=="High Risk AML") |> plot(lwd=2,col=3)
```

```
muhaz(bmt$t2,bmt$d3,bmt$group=="ALL") |> lines(lwd=2,col=1)
muhaz(bmt$t2,bmt$d3,bmt$group=="Low Risk AML") |> lines(lwd=2,col=2)
legend("topright",c("ALL","Low Risk AML","High Risk AML"),col=1:3,lwd=2)
```



Figure 6.5.: Smoothed Hazard Rate Estimates by Disease Group

Group 3 was plotted first because it has the highest hazard.

Except for an initial blip in the high risk AML group, the hazards look roughly proportional. They are all strongly decreasing.

6.2.4. Fitting the Proportional Hazards Model

How do we fit a proportional hazards regression model? We need to estimate the coefficients of the covariates, and we need to estimate the base

hazard $h_0(t)$. For the covariates, supposing for simplicity that there are no tied event times, let the event times for the whole data set be t_1, t_2, \dots, t_D . Let the risk set at time t_i be $R(t_i)$ and

$$\begin{split} \eta(x) &= \beta_1 x_1 + \dots + \beta_p x_p \\ \theta(x) &= e^{\eta(x)} \\ h(t|X=x) &= h_0(t) e^{\eta(x)} = \theta(x) h_0(t) \end{split}$$

Conditional on a single failure at time t, the probability that the event is due to subject $f \in R(t)$ is approximately

$$\begin{split} \Pr(f \text{ fails} | 1 \text{ failure at } t) &= \frac{h_0(t)e^{\eta(x_f)}}{\sum_{k \in R(t)} h_0(t)e^{\eta(x_f)}} \\ &= \frac{\theta(x_f)}{\sum_{k \in R(t)} \theta(x_k)} \end{split}$$

The logic behind this has several steps. We first fix (ex post) the failure times and note that in this discrete context, the probability p_j that a subject j in the risk set fails at time t is just the hazard of that subject at that time.

If all of the p_i are small, the chance that exactly one subject fails is

$$\sum_{k \in R(t)} p_k \left[\prod_{m \in R(t), m \neq k} (1 - p_m) \right] \approx \sum_{k \in R(t)} p_k$$

If subject i is the one who experiences the event of interest at time t_i , then the **partial likelihood** is

$$\mathcal{L}^*(\boldsymbol{\beta}|T) = \prod_i \frac{\theta(\boldsymbol{x}_i)}{\sum_{k \in R(t_i)} \theta(\boldsymbol{x}_k)}$$

and we can numerically maximize this with respect to the coefficients β that specify $\eta(x) = x'\beta$. When there are tied event times adjustments need to be made, but the likelihood is still similar. Note that we don't need to know the base hazard to solve for the coefficients.

Once we have coefficient estimates $\hat{\beta} = (\hat{\beta}_1, \dots, \hat{\beta}_p)$, this also defines $\hat{\eta}(x)$ and $\hat{\theta}(x)$ and then the estimated base cumulative hazard function is

$$\hat{H}(t) = \sum_{t_i < t} \frac{d_i}{\sum_{k \in R(t_i)} \theta(x_k)}$$

which reduces to the Nelson-Aalen estimate when there are no covariates. There are numerous other estimates that have been proposed as well.

6.3. Cox Model for the bmt data

6.3.1. Fit the model

```
bmt.cox <- coxph(Surv(t2, d3) ~ group, data = bmt)
summary(bmt.cox)
#> Call:
#> coxph(formula = Surv(t2, d3) ~ group, data = bmt)
#>
#> n= 137, number of events= 83
#>
#> coef exp(coef) se(coef) z Pr(>|z|)
#> groupLow Risk AML -0.574 0.563 0.287 -2.00 0.046;
```

```
#> groupHigh Risk AML 0.383
                                 1.467
                                          0.267 1.43
                                                         0.152
#> ---
#> Signif. codes: 0 '*** 0.001 '** 0.01 '* 0.05 '.' 0.1 ' ' 1
#>
#>
                      exp(coef) exp(-coef) lower .95 upper .95
                          0.563
#> groupLow Risk AML
                                     1.776
                                               0.321
                                                         0.989
#> groupHigh Risk AML
                          1.467
                                     0.682
                                               0.869
                                                         2.478
#>
\# Concordance= 0.625 (se = 0.03)
#> Likelihood ratio test= 13.4 on 2 df,
                                           p=0.001
                                         p=0.001
#> Wald test
                       = 13 on 2 df,
#> Score (logrank) test = 13.8 on 2 df,
                                          p=0.001
```

The table provides hypothesis tests comparing groups 2 and 3 to group 1. Group 3 has the highest hazard, so the most significant comparison is not directly shown.

The coefficient 0.3834 is on the log-hazard-ratio scale, as in log-risk-ratio. The next column gives the hazard ratio 1.4673, and a hypothesis (Wald) test.

The (not shown) group 3 vs. group 2 log hazard ratio is 0.3834 + 0.5742 = 0.9576. The hazard ratio is then $\exp(0.9576)$ or 2.605.

Inference on all coefficients and combinations can be constructed using coef(bmt.cox) and vcov(bmt.cox) as with logistic and poisson regression.

Concordance is agreement of first failure between pairs of subjects and higher predicted risk between those subjects, omitting non-informative pairs.

The Rsquare value is Cox and Snell's pseudo R-squared and is not very useful.

6.3.2. Tests for nested models

summary() prints three tests for whether the model with the group covariate is better than the one without

- Likelihood ratio test (chi-squared)
- Wald test (also chi-squared), obtained by adding the squares of the z-scores
- Score = log-rank test, as with comparison of survival functions.

The likelihood ratio test is probably best in smaller samples, followed by the Wald test.

6.3.3. Survival Curves from the Cox Model

We can take a look at the resulting group-specific curves:

```
#| fig-cap: "Survival Functions for Three Groups by KM and Cox Model"
km_fit = survfit(Surv(t2, d3) ~ group, data = as.data.frame(bmt))

cox_fit = survfit(
   bmt.cox,
   newdata =
        data.frame(
        group = unique(bmt$group),
        row.names = unique(bmt$group)))

library(survminer)

list(KM = km_fit, Cox = cox_fit) |>
   survminer::ggsurvplot(
        # facet.by = "group",
```

```
legend = "bottom",
legend.title = "",
combine = TRUE,
fun = 'pct',
size = .5,
ggtheme = theme_bw(),
conf.int = FALSE,
censor = FALSE) |>
suppressWarnings() # ggsurvplot() throws some warnings that aren't too worrying
```



When we use survfit() with a Cox model, we have to specify the covariate levels we are interested in; the argument newdata should include a data.frame with the same named columns as the predictors in the Cox model and one or more levels of each.

From ?survfit.coxph:

If the newdata argument is missing, a curve is produced for a single "pseudo" subject with covariate values equal to the means component of the fit. The resulting curve(s) almost never make sense, but the default remains due to an unwarranted attachment to the option shown by some users and by other packages. Two particularly egregious examples are factor variables and interactions. Suppose one were studying interspecies transmission of a virus, and the data set has a factor variable with levels ("pig", "chicken") and about equal numbers of observations for each. The "mean" covariate level will be 0.5 – is this a flying pig?

6.3.4. Examining survfit

```
survfit(Surv(t2, d3) ~ group, data = bmt)
#> Call: survfit(formula = Surv(t2, d3) ~ group, data = bmt)
#>
#>
                         n events median 0.95LCL 0.95UCL
#> group=ALL
                        38
                               24
                                      418
                                              194
                                                        NA
#> group=Low Risk AML
                               25
                                     2204
                                              704
                        54
                                                        NA
#> group=High Risk AML 45
                               34
                                      183
                                              115
                                                       456
```

```
survfit(Surv(t2, d3) ~ group, data = bmt) |> summary()
#> Call: survfit(formula = Surv(t2, d3) ~ group, data = bmt)
#>
#>
                    group=ALL
#>
    time n.risk n.event survival std.err lower 95% CI upper 95% CI
#>
       1
             38
                       1
                            0.974
                                   0.0260
                                                  0.924
                                                                1.000
      55
             37
                       1
                            0.947
                                    0.0362
                                                   0.879
                                                                1.000
#>
      74
#>
             36
                            0.921 0.0437
                                                  0.839
                                                                1.000
```

| #> | 86 | 35 | 1 | 0.895 | 0.0498 | 0.802 | 0.998 |
|----|------|--------|---------|------------|---------|--------------|--------------|
| #> | 104 | 34 | 1 | 0.868 | 0.0548 | 0.767 | 0.983 |
| #> | 107 | 33 | 1 | 0.842 | 0.0592 | 0.734 | 0.966 |
| #> | 109 | 32 | 1 | 0.816 | 0.0629 | 0.701 | 0.949 |
| #> | 110 | 31 | 1 | 0.789 | 0.0661 | 0.670 | 0.930 |
| #> | 122 | 30 | 2 | 0.737 | 0.0714 | 0.609 | 0.891 |
| #> | 129 | 28 | 1 | 0.711 | 0.0736 | 0.580 | 0.870 |
| #> | 172 | 27 | 1 | 0.684 | 0.0754 | 0.551 | 0.849 |
| #> | 192 | 26 | 1 | 0.658 | 0.0770 | 0.523 | 0.827 |
| #> | 194 | 25 | 1 | 0.632 | 0.0783 | 0.495 | 0.805 |
| #> | 230 | 23 | 1 | 0.604 | 0.0795 | 0.467 | 0.782 |
| #> | 276 | 22 | 1 | 0.577 | 0.0805 | 0.439 | 0.758 |
| #> | 332 | 21 | 1 | 0.549 | 0.0812 | 0.411 | 0.734 |
| #> | 383 | 20 | 1 | 0.522 | 0.0817 | 0.384 | 0.709 |
| #> | 418 | 19 | 1 | 0.494 | 0.0819 | 0.357 | 0.684 |
| #> | 466 | 18 | 1 | 0.467 | 0.0818 | 0.331 | 0.658 |
| #> | 487 | 17 | 1 | 0.439 | 0.0815 | 0.305 | 0.632 |
| #> | 526 | 16 | 1 | 0.412 | 0.0809 | 0.280 | 0.605 |
| #> | 609 | 14 | 1 | 0.382 | 0.0803 | 0.254 | 0.577 |
| #> | 662 | 13 | 1 | 0.353 | 0.0793 | 0.227 | 0.548 |
| #> | | | | | | | |
| #> | | | grou | p=Low Risl | k AML | | |
| #> | time | n.risk | n.event | survival | std.err | lower 95% CI | upper 95% CI |
| #> | 10 | 54 | 1 | 0.981 | 0.0183 | 0.946 | 1.000 |
| #> | 35 | 53 | 1 | 0.963 | 0.0257 | 0.914 | 1.000 |
| #> | 48 | 52 | 1 | 0.944 | 0.0312 | 0.885 | 1.000 |
| #> | 53 | 51 | 1 | 0.926 | 0.0356 | 0.859 | 0.998 |
| #> | 79 | 50 | 1 | 0.907 | 0.0394 | 0.833 | 0.988 |
| #> | 80 | 49 | 1 | 0.889 | 0.0428 | 0.809 | 0.977 |
| #> | 105 | 48 | 1 | 0.870 | 0.0457 | 0.785 | 0.965 |
| #> | 211 | 47 | 1 | 0.852 | 0.0483 | 0.762 | 0.952 |
| #> | 219 | 46 | 1 | 0.833 | 0.0507 | 0.740 | 0.939 |
| #> | 248 | 45 | 1 | 0.815 | 0.0529 | 0.718 | 0.925 |
| | | | | | | | |

| #> | 272 | 44 | 1 | 0.796 | 0.0548 | 0.696 | 0.911 |
|----------------------------------|--|--|----------------------------|--|--|---|---|
| #> | 288 | 43 | 1 | 0.778 | 0.0566 | 0.674 | 0.897 |
| #> | 381 | 42 | 1 | 0.759 | 0.0582 | 0.653 | 0.882 |
| #> | 390 | 41 | 1 | 0.741 | 0.0596 | 0.633 | 0.867 |
| #> | 414 | 40 | 1 | 0.722 | 0.0610 | 0.612 | 0.852 |
| #> | 421 | 39 | 1 | 0.704 | 0.0621 | 0.592 | 0.837 |
| #> | 481 | 38 | 1 | 0.685 | 0.0632 | 0.572 | 0.821 |
| #> | 486 | 37 | 1 | 0.667 | 0.0642 | 0.552 | 0.805 |
| #> | 606 | 36 | 1 | 0.648 | 0.0650 | 0.533 | 0.789 |
| #> | 641 | 35 | 1 | 0.630 | 0.0657 | 0.513 | 0.773 |
| #> | 704 | 34 | 1 | 0.611 | 0.0663 | 0.494 | 0.756 |
| #> | 748 | 33 | 1 | 0.593 | 0.0669 | 0.475 | 0.739 |
| #> | 1063 | 26 | 1 | 0.570 | 0.0681 | 0.451 | 0.720 |
| #> | 1074 | 25 | 1 | 0.547 | 0.0691 | 0.427 | 0.701 |
| #> | 2204 | 6 | 1 | 0.456 | 0.1012 | 0.295 | 0.704 |
| #> | | | | | | | |
| #> | | | | p=High Ris | | | |
| #> | | | | | | lower 95% CI | |
| #> | 2 | 45 | 1 | 0.978 | 0.0220 | 0.936 | 1.000 |
| #> | 16 | 44 | 1 | 0.956 | 0.0307 | 0.897 | 1.000 |
| #> | 32 | 43 | 1 | 0.933 | 0.0372 | 0.863 | 1.000 |
| #> | 47 | 42 | 2 | 0.889 | 0.0468 | 0.802 | 0.986 |
| #> | 48 | 40 | 1 | 0.867 | 0.0507 | 0.773 | 0.972 |
| #> | 63 | 39 | 1 | 0.844 | 0.0540 | 0.745 | 0.957 |
| #> | | | | | | | |
| | 64 | 38 | 1 | 0.822 | 0.0570 | 0.718 | 0.942 |
| #> | 74 | 37 | 1 | 0.822 0.800 | 0.0570 0.0596 | 0.718 0.691 | 0.926 |
| #> #> | 74 76 | 37 36 | 1 1 | 0.822 0.800 0.778 | 0.0570 0.0596 0.0620 | 0.718 0.691 0.665 | 0.926 0.909 |
| #> #> #> | 74 76 80 | 37 36 35 | 1 1 1 | 0.822 0.800 0.778 0.756 | 0.0570 0.0596 0.0620 0.0641 | 0.718 0.691 0.665 0.640 | 0.926 0.909 0.892 |
| #> #> #> #> | 74 76 80 84 | 37 36 35 34 | 1 1 1 1 | 0.822 0.800 0.778 0.756 0.733 | 0.0570 0.0596 0.0620 0.0641 0.0659 | 0.718 0.691 0.665 0.640 0.615 | 0.926 0.909 0.892 0.875 |
| #> #> #> #> | 74 76 80 84 93 | 37 36 35 34 33 | 1 1 1 1 | 0.822 0.800 0.778 0.756 0.733 0.711 | 0.0570 0.0596 0.0620 0.0641 0.0659 0.0676 | 0.718 0.691 0.665 0.640 0.615 0.590 | 0.926 0.909 0.892 0.875 0.857 |
| #> #> #> #> #> | 74 76 80 84 93 100 | 37 36 35 34 33 | 1 1 1 1 1 | 0.822 0.800 0.778 0.756 0.733 0.711 0.689 | 0.0570 0.0596 0.0620 0.0641 0.0659 0.0676 0.0690 | 0.718 0.691 0.665 0.640 0.615 0.590 | 0.926 0.909 0.892 0.875 0.857 |
| #> #> #> #> #> #> | 74 76 80 84 93 100 105 | 37 36 35 34 33 32 31 | 1 1 1 1 1 1 | 0.822 0.800 0.778 0.756 0.733 0.711 0.689 0.667 | 0.0570 0.0596 0.0620 0.0641 0.0659 0.0676 0.0690 0.0703 | 0.718 0.691 0.665 0.640 0.615 0.590 0.566 | 0.926 0.909 0.892 0.875 0.857 0.838 0.820 |
| #> #> #> #> #> | 74 76 80 84 93 100 | 37 36 35 34 33 | 1 1 1 1 1 | 0.822 0.800 0.778 0.756 0.733 0.711 0.689 | 0.0570 0.0596 0.0620 0.0641 0.0659 0.0676 0.0690 | 0.718 0.691 0.665 0.640 0.615 0.590 | 0.926 0.909 0.892 0.875 0.857 0.838 |

```
#>
     115
              29
                             0.622
                                     0.0723
                                                    0.496
                                                                   0.781
     120
                                     0.0730
                                                     0.473
                                                                   0.762
#>
              28
                        1
                             0.600
#>
     157
              27
                        1
                             0.578
                                     0.0736
                                                     0.450
                                                                   0.742
     162
                        1
                             0.556
                                     0.0741
                                                     0.428
                                                                   0.721
#>
              26
                                                                   0.701
#>
     164
              25
                        1
                             0.533
                                     0.0744
                                                    0.406
                                     0.0745
                                                     0.384
                                                                   0.680
#>
     168
              24
                        1
                             0.511
#>
     183
              23
                        1
                             0.489
                                     0.0745
                                                    0.363
                                                                   0.659
#>
     242
              22
                        1
                             0.467
                                     0.0744
                                                    0.341
                                                                   0.638
     268
                                     0.0741
                                                                   0.616
#>
              21
                        1
                             0.444
                                                    0.321
#>
     273
              20
                        1
                             0.422
                                     0.0736
                                                    0.300
                                                                   0.594
                             0.400
                                     0.0730
#>
     318
              19
                        1
                                                    0.280
                                                                   0.572
#>
     363
              18
                        1
                             0.378
                                     0.0723
                                                    0.260
                                                                   0.550
#>
     390
              17
                             0.356
                                     0.0714
                                                     0.240
                                                                   0.527
                        1
                                                                   0.504
#>
     422
              16
                        1
                             0.333
                                     0.0703
                                                    0.221
#>
     456
              15
                        1
                             0.311
                                     0.0690
                                                    0.201
                                                                   0.481
#>
     467
              14
                        1
                             0.289
                                     0.0676
                                                    0.183
                                                                   0.457
#>
     625
              13
                        1
                             0.267
                                     0.0659
                                                    0.164
                                                                   0.433
#>
     677
              12
                             0.244 0.0641
                                                    0.146
                                                                   0.409
```

```
survfit(bmt.cox)
#> Call: survfit(formula = bmt.cox)
#>
#>
          n events median 0.95LCL 0.95UCL
#> [1,] 137
                83
                       422
                               268
                                        NA
survfit(bmt.cox, newdata = tibble(group = unique(bmt$group)))
#> Call: survfit(formula = bmt.cox, newdata = tibble(group = unique(bmt$group)))
#>
#>
       n events median 0.95LCL 0.95UCL
#> 1 137
             83
                    422
                            268
                                     NA
#> 2 137
             83
                    NA
                            625
                                     NA
#> 3 137
             83
                    268
                            162
                                    467
```

```
bmt.cox |>
  survfit(newdata = tibble(group = unique(bmt$group))) |>
  summary()
#> Call: survfit(formula = bmt.cox, newdata = tibble(group = unique(bmt$group)))
#>
    time n.risk n.event survival1 survival2 survival3
#>
                        1
                               0.993
                                          0.996
#>
       1
             137
                                                     0.989
#>
       2
             136
                        1
                               0.985
                                          0.992
                                                     0.978
      10
             135
                        1
#>
                               0.978
                                          0.987
                                                     0.968
#>
      16
             134
                        1
                               0.970
                                          0.983
                                                     0.957
                        1
#>
      32
             133
                               0.963
                                          0.979
                                                     0.946
             132
#>
      35
                        1
                               0.955
                                          0.975
                                                     0.935
#>
      47
             131
                        2
                               0.941
                                          0.966
                                                     0.914
#>
      48
             129
                        2
                               0.926
                                          0.957
                                                     0.893
#>
      53
             127
                        1
                               0.918
                                          0.953
                                                     0.882
#>
      55
             126
                        1
                               0.911
                                          0.949
                                                     0.872
#>
      63
             125
                        1
                               0.903
                                          0.944
                                                     0.861
#>
      64
             124
                        1
                               0.896
                                          0.940
                                                     0.851
                        2
             123
#>
      74
                               0.881
                                          0.931
                                                     0.830
#>
      76
             121
                        1
                               0.873
                                          0.926
                                                     0.819
#>
      79
             120
                        1
                               0.865
                                          0.922
                                                     0.809
                        2
#>
      80
             119
                               0.850
                                          0.913
                                                     0.788
#>
      84
             117
                        1
                               0.843
                                          0.908
                                                     0.778
                        1
#>
      86
             116
                               0.835
                                          0.903
                                                     0.768
             115
                        1
                               0.827
                                          0.899
                                                     0.757
#>
      93
                        1
#>
     100
             114
                               0.820
                                          0.894
                                                     0.747
#>
     104
             113
                        1
                               0.812
                                          0.889
                                                     0.737
#>
     105
             112
                        2
                               0.797
                                          0.880
                                                     0.717
#>
     107
             110
                        1
                               0.789
                                          0.875
                                                     0.707
#>
     109
             109
                        1
                               0.782
                                          0.870
                                                     0.697
#>
     110
             108
                        1
                               0.774
                                          0.866
                                                     0.687
                        1
#>
     113
             107
                               0.766
                                          0.861
                                                     0.677
             106
                        1
#>
     115
                               0.759
                                          0.856
                                                     0.667
```

| #> | 120 | 105 | 1 | 0.751 | 0.851 | 0.657 | |
|----|-----|-----|---|-------|-------|-------|--|
| #> | 122 | 104 | 2 | 0.735 | 0.841 | 0.637 | |
| #> | 129 | 102 | 1 | 0.727 | 0.836 | 0.627 | |
| #> | 157 | 101 | 1 | 0.720 | 0.831 | 0.617 | |
| #> | 162 | 100 | 1 | 0.712 | 0.826 | 0.607 | |
| #> | 164 | 99 | 1 | 0.704 | 0.821 | 0.598 | |
| #> | 168 | 98 | 1 | 0.696 | 0.815 | 0.588 | |
| #> | 172 | 97 | 1 | 0.688 | 0.810 | 0.578 | |
| #> | 183 | 96 | 1 | 0.680 | 0.805 | 0.568 | |
| #> | 192 | 95 | 1 | 0.672 | 0.800 | 0.558 | |
| #> | 194 | 94 | 1 | 0.664 | 0.794 | 0.549 | |
| #> | 211 | 93 | 1 | 0.656 | 0.789 | 0.539 | |
| #> | 219 | 92 | 1 | 0.648 | 0.783 | 0.530 | |
| #> | 230 | 90 | 1 | 0.640 | 0.778 | 0.520 | |
| #> | 242 | 89 | 1 | 0.632 | 0.773 | 0.511 | |
| #> | 248 | 88 | 1 | 0.624 | 0.767 | 0.501 | |
| #> | 268 | 87 | 1 | 0.616 | 0.761 | 0.492 | |
| #> | 272 | 86 | 1 | 0.608 | 0.756 | 0.482 | |
| #> | 273 | 85 | 1 | 0.600 | 0.750 | 0.473 | |
| #> | 276 | 84 | 1 | 0.592 | 0.745 | 0.464 | |
| #> | 288 | 83 | 1 | 0.584 | 0.739 | 0.454 | |
| #> | 318 | 82 | 1 | 0.576 | 0.733 | 0.445 | |
| #> | 332 | 81 | 1 | 0.568 | 0.727 | 0.436 | |
| #> | 363 | 80 | 1 | 0.560 | 0.722 | 0.427 | |
| #> | 381 | 79 | 1 | 0.552 | 0.716 | 0.418 | |
| #> | 383 | 78 | 1 | 0.544 | 0.710 | 0.409 | |
| #> | 390 | 77 | 2 | 0.528 | 0.698 | 0.392 | |
| #> | 414 | 75 | 1 | 0.520 | 0.692 | 0.383 | |
| #> | 418 | 74 | 1 | 0.512 | 0.686 | 0.374 | |
| #> | 421 | 73 | 1 | 0.504 | 0.680 | 0.366 | |
| #> | 422 | 72 | 1 | 0.496 | 0.674 | 0.357 | |
| #> | 456 | 71 | 1 | 0.488 | 0.667 | 0.349 | |
| #> | 466 | 70 | 1 | 0.480 | 0.661 | 0.340 | |
| | | | | | | | |

| #> | 467 | 69 | 1 | 0.472 | 0.655 | 0.332 | |
|----|------|----|---|-------|-------|-------|--|
| #> | 481 | 68 | 1 | 0.464 | 0.649 | 0.324 | |
| #> | 486 | 67 | 1 | 0.455 | 0.642 | 0.315 | |
| #> | 487 | 66 | 1 | 0.447 | 0.636 | 0.307 | |
| #> | 526 | 65 | 1 | 0.439 | 0.629 | 0.299 | |
| #> | 606 | 63 | 1 | 0.431 | 0.623 | 0.291 | |
| #> | 609 | 62 | 1 | 0.423 | 0.616 | 0.283 | |
| #> | 625 | 61 | 1 | 0.415 | 0.609 | 0.275 | |
| #> | 641 | 60 | 1 | 0.407 | 0.603 | 0.267 | |
| #> | 662 | 59 | 1 | 0.399 | 0.596 | 0.260 | |
| #> | 677 | 58 | 1 | 0.391 | 0.589 | 0.252 | |
| #> | 704 | 57 | 1 | 0.383 | 0.582 | 0.244 | |
| #> | 748 | 56 | 1 | 0.374 | 0.575 | 0.237 | |
| #> | 1063 | 47 | 1 | 0.365 | 0.567 | 0.228 | |
| #> | 1074 | 46 | 1 | 0.356 | 0.559 | 0.220 | |
| #> | 2204 | 9 | 1 | 0.313 | 0.520 | 0.182 | |
| | | | | | | | |

6.4. Adjustment for Ties (optional)

At each time t_i at which more than one of the subjects has an event, let d_i be the number of events at that time, D_i the set of subjects with events at that time, and let s_i be a covariate vector for an artificial subject obtained by adding up the covariate values for the subjects with an event at time t_i . Let

$$\bar{\eta}_i = \beta_1 s_{i1} + \dots + \beta_p s_{ip}$$

and $\bar{\theta}_i = \exp{\{\bar{\eta}_i\}}$.

Let s_i be a covariate vector for an artificial subject obtained by adding up the covariate values for the subjects with an event at time t_i . Note that

$$\begin{split} \bar{\eta}_i &= \sum_{j \in D_i} \beta_1 x_{j1} + \dots + \beta_p x_{jp} \\ &= \beta_1 s_{i1} + \dots + \beta_p s_{ip} \\ \bar{\theta}_i &= \exp\left\{\bar{\eta}_i\right\} \\ &= \prod_{j \in D_i} \theta_i \end{split}$$

6.4.0.1. Breslow's method for ties

Breslow's method estimates the partial likelihood as

$$\begin{split} L(\beta|T) &= \prod_i \frac{\bar{\theta}_i}{[\sum_{k \in R(t_i)} \theta_k]^{d_i}} \\ &= \prod_i \prod_{j \in D_i} \frac{\theta_j}{\sum_{k \in R(t_i)} \theta_k} \end{split}$$

This method is equivalent to treating each event as distinct and using the non-ties formula. It works best when the number of ties is small. It is the default in many statistical packages, including PROC PHREG in SAS.

6.4.0.2. Efron's method for ties

The other common method is Efron's, which is the default in R.

$$L(\beta|T) = \prod_i \frac{\bar{\theta}_i}{\prod_{j=1}^{d_i} [\sum_{k \in R(t_i)} \theta_k - \frac{j-1}{d_i} \sum_{k \in D_i} \theta_k]}$$

This is closer to the exact discrete partial likelihood when there are many ties.

The third option in R (and an option also in SAS as discrete) is the "exact" method, which is the same one used for matched logistic regression.

6.4.0.3. Example: Breslow's method

Suppose as an example we have a time t where there are 20 individuals at risk and three failures. Let the three individuals have risk parameters $\theta_1, \theta_2, \theta_3$ and let the sum of the risk parameters of the remaining 17 individuals be θ_R . Then the factor in the partial likelihood at time t using Breslow's method is

$$\left(\frac{\theta_1}{\theta_R + \theta_1 + \theta_2 + \theta_3}\right) \left(\frac{\theta_2}{\theta_R + \theta_1 + \theta_2 + \theta_3}\right) \left(\frac{\theta_3}{\theta_R + \theta_1 + \theta_2 + \theta_3}\right)$$

If on the other hand, they had died in the order 1,2, 3, then the contribution to the partial likelihood would be:

$$\left(\frac{\theta_1}{\theta_R+\theta_1+\theta_2+\theta_3}\right)\left(\frac{\theta_2}{\theta_R+\theta_2+\theta_3}\right)\left(\frac{\theta_3}{\theta_R+\theta_3}\right)$$

as the risk set got smaller with each failure. The exact method roughly averages the results for the six possible orderings of the failures.

6.4.0.4. Example: Efron's method

But we don't know the order they failed in, so instead of reducing the denominator by one risk coefficient each time, we reduce it by the same fraction. This is Efron's method.

$$\left(\frac{\theta_1}{\theta_R+\theta_1+\theta_2+\theta_3}\right)\left(\frac{\theta_2}{\theta_R+2(\theta_1+\theta_2+\theta_3)/3}\right)\left(\frac{\theta_3}{\theta_R+(\theta_1+\theta_2+\theta_3)/3}\right)$$

6.5. Building Cox Proportional Hazards models

6.5.1. hodg Lymphoma Data Set from KMsurv

6.5.1.1. Participants

43 bone marrow transplant patients at Ohio State University (Avalos 1993)

6.5.1.2. Variables

- dtype: Disease type (Hodgkin's or non-Hodgkins lymphoma)
- gtype: Bone marrow graft type:
- allogeneic: from HLA-matched sibling
- autologous: from self (prior to chemo)
- time: time to study exit
- delta: study exit reason (death/relapse vs censored)
- wtime: waiting time to transplant (in months)
- score: Karnofsky score:
- 80–100: Able to carry on normal activity and to work; no special care needed.
- 50–70: Unable to work; able to live at home and care for most personal needs; varying amount of assistance needed.
- 10–60: Unable to care for self; requires equivalent of institutional or hospital care; disease may be progressing rapidly.

6.5.1.3. Data

```
data(hodg, package = "KMsurv")
hodg2 = hodg |>
```

```
as_tibble() |>
 mutate(
    # We add factor labels to the categorical variables:
    gtype = gtype |>
      case_match(
        1 ~ "Allogenic",
        2 ~ "Autologous"),
    dtype = dtype |>
      case_match(
        1 ~ "Non-Hodgkins",
        2 ~ "Hodgkins") |>
      factor() |>
      relevel(ref = "Non-Hodgkins"),
    delta = delta |>
      case_match(
        1 ~ "dead",
        0 ~ "alive"),
    surv = Surv(
      time = time,
      event = delta == "dead")
  )
hodg2 |> print()
#> # A tibble: 43 x 7
#>
      gtype
                dtype
                             time delta score wtime
                                                       surv
#>
      <chr>
                <fct>
                             <int> <chr> <int> <int> <Surv>
#>
   1 Allogenic Non-Hodgkins
                               28 dead
                                            90
                                                  24
                                                        28
                                                  7
   2 Allogenic Non-Hodgkins
                             32 dead
                                            30
                                                        32
#> 3 Allogenic Non-Hodgkins
                               49 dead
                                            40
                                                  8
                                                        49
#> 4 Allogenic Non-Hodgkins
                               84 dead
                                            60
                                                  10
                                                       84
#> 5 Allogenic Non-Hodgkins
                               357 dead
                                            70
                                                  42
                                                       357
#> 6 Allogenic Non-Hodgkins
                               933 alive
                                           90
                                                  9
                                                       933+
#> 7 Allogenic Non-Hodgkins 1078 alive
                                           100
                                                  16 1078+
#> 8 Allogenic Non-Hodgkins
                             1183 alive
                                            90
                                                  16
                                                      1183+
```

```
#> 9 Allogenic Non-Hodgkins 1560 alive 80 20 1560+
#> 10 Allogenic Non-Hodgkins 2114 alive 80 27 2114+
#> # i 33 more rows
```

6.5.2. Proportional hazards model

```
hodg.cox1 = coxph(
 formula = surv ~ gtype * dtype + score + wtime,
 data = hodg2)
summary(hodg.cox1)
#> Call:
#> coxph(formula = surv ~ gtype * dtype + score + wtime, data = hodg2)
#>
  n= 43, number of events= 26
#>
#>
#>
                            coef exp(coef) se(coef) z Pr(>|z|)
#> gtypeAutologous
                           0.6394 1.8953 0.5937 1.08 0.2815
#> dtypeHodgkins
                           2.7603 15.8050 0.9474 2.91
                                                      0.0036 **
#> score
                          -0.0166 0.9836 0.0102 -1.62 0.1046
#> wtime
#> Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
#>
#>
                          exp(coef) exp(-coef) lower .95 upper .95
#> gtypeAutologous
                                    0.5276 0.5920
                                                      6.068
                            1.8953
#> dtypeHodgkins
                            15.8050
                                     0.0633
                                             2.4682
                                                     101.207
#> score
                            0.9517
                                    1.0507 0.9288
                                                      0.975
#> wtime
                            0.9836
                                    1.0167 0.9641
                                                      1.003
#> gtypeAutologous:dtypeHodgkins
                            0.0934
                                    10.7074 0.0123
                                                      0.711
```

6.6. Diagnostic graphs for proportional hazards assumption

6.6.1. Analysis plan

- **survival function** for the four combinations of disease type and graft type.
- observed (nonparametric) vs. expected (semiparametric) survival functions.
- complementary log-log survival for the four groups.

6.6.2. Kaplan-Meier survival functions

```
km_model = survfit(
  formula = surv ~ dtype + gtype,
  data = hodg2)

km_model |>
  autoplot(conf.int = FALSE) +
  theme_bw() +
  theme(
   legend.position="bottom",
   legend.title = element_blank(),
```

```
legend.text = element_text(size = legend_text_size)
) +
guides(col=guide_legend(ncol=2)) +
ylab('Survival probability, S(t)') +
xlab("Time since transplant (days)")
```




Figure 6.6.: Kaplan-Meier Survival Curves for HOD/NHL and Allo/Auto Grafts

6.6.3. Observed and expected survival curves

```
# we need to create a tibble of covariate patterns;
# we will set score and wtime to mean values for disease and graft types:
```

```
means = hodg2 |>
  summarize(
    .by = c(dtype, gtype),
   score = mean(score),
    wtime = mean(wtime)) |>
  arrange(dtype, gtype) |>
  mutate(strata = paste(dtype, gtype, sep = ",")) |>
  as.data.frame()
# survfit.coxph() will use the rownames of its `newdata`
# argument to label its output:
rownames(means) = means$strata
cox_model =
 hodg.cox1 |>
  survfit(
    data = hodg2, # ggsurvplot() will need this
    newdata = means)
# I couldn't find a good function to reformat `cox_model` for ggplot,
# so I made my own:
stack_surv_ph = function(cox_model)
  cox_model$surv |>
    as_tibble() |>
    mutate(time = cox_model$time) |>
    pivot_longer(
     cols = -time,
      names_to = "strata",
     values_to = "surv") |>
    mutate(
      cumhaz = -log(surv),
      model = "Cox PH")
```

```
km_and_cph =
  km_model |>
  fortify(surv.connect = TRUE) |>
  mutate(
    strata = trimws(strata),
    model = "Kaplan-Meier",
    cumhaz = -log(surv)) |>
  bind_rows(stack_surv_ph(cox_model))
```

```
km_and_cph |>
  ggplot(aes(x = time, y = surv, col = model)) +
  geom_step() +
  facet_wrap(~strata) +
  theme_bw() +
  ylab("S(t) = P(T>=t)") +
  xlab("Survival time (t, days)") +
  theme(legend.position = "bottom")
```



Figure 6.7.: Observed and expected survival curves for bmt data

6.6.4. Cumulative hazard (log-scale) curves

Also known as "complementary log-log (clog-log) survival curves".

```
na_model = survfit(
  formula = surv ~ dtype + gtype,
  data = hodg2,
  type = "fleming")

na_model |>
  survminer::ggsurvplot(
  legend = "bottom",
  legend.title = "",
```

```
ylab = "log(Cumulative Hazard)",
xlab = "Time since transplant (days, log-scale)",
fun = 'cloglog',
size = .5,
ggtheme = theme_bw(),
conf.int = FALSE,
censor = TRUE) |>
magrittr::extract2("plot") +
guides(
col =
    guide_legend(
    ncol = 2,
    label.theme =
    element_text(
    size = legend_text_size)))
```



Figure 6.8.: Complementary log-log survival curves - Nelson-Aalen estimates

Let's compare these empirical (i.e., non-parametric) curves with the fitted curves from our coxph() model:

```
cox_model |>
  survminer::ggsurvplot(
   facet_by = "",
   legend = "bottom",
   legend.title = "",
   ylab = "log(Cumulative Hazard)",
   xlab = "Time since transplant (days, log-scale)",
   fun = 'cloglog',
   size = .5,
   ggtheme = theme_bw(),
```

```
censor = FALSE, # doesn't make sense for cox model
  conf.int = FALSE) |>
magrittr::extract2("plot") +
guides(
  col =
    guide_legend(
    ncol = 2,
    label.theme =
    element_text(
        size = legend_text_size)))
```



Figure 6.9.: Complementary log-log survival curves - PH estimates

Now let's overlay these cumulative hazard curves:

```
na_and_cph =
 na_model |>
  fortify(fun = "cumhaz") |>
  # `fortify.survfit()` doesn't name cumhaz correctly:
  rename(cumhaz = surv) |>
  mutate(
    surv = exp(-cumhaz),
    strata = trimws(strata)) |>
 mutate(model = "Nelson-Aalen") |>
  bind_rows(stack_surv_ph(cox_model))
na_and_cph |>
  ggplot(
    aes(
     x = time,
     y = cumhaz,
      col = model)) +
  geom_step() +
  facet_wrap(~strata) +
  theme_bw() +
  scale_y_continuous(
   trans = "log10",
    name = "Cumulative hazard H(t) (log-scale)") +
  scale_x_continuous(
   trans = "log10",
   name = "Survival time (t, days, log-scale)") +
  theme(legend.position = "bottom")
```



Figure 6.10.: Observed and expected cumulative hazard curves for bmt data (cloglog format)

6.7. Predictions and Residuals

6.7.1. Review: Predictions in Linear Regression

• In linear regression, we have a linear predictor for each data point i

$$\begin{split} &\eta_i = \beta_0 + \beta_1 x_{1i} + \dots + \beta_p x_{pi} \\ &\hat{y}_i = \hat{\eta}_i = \hat{\beta}_0 + \hat{\beta}_1 x_{1i} + \dots + \hat{\beta}_p x_{pi} \\ &y_i \sim N(\eta_i, \sigma^2) \end{split}$$

• \hat{y}_i estimates the conditional mean of y_i given the covariate values \tilde{x}_i . This together with the prediction error says that we are predicting the distribution of values of y.

6.7.2. Review: Residuals in Linear Regression

- The usual residual is $r_i = y_i \hat{y}_i$, the difference between the actual value of y and a prediction of its mean.
- The residuals are also the quantities the sum of whose squares is being minimized by the least squares/MLE estimation.

6.7.3. Predictions and Residuals in survival models

- In survival analysis, the equivalent of y_i is the event time t_i , which is unknown for the censored observations.
- The expected event time can be tricky to calculate:

$$\hat{\mathbf{E}}[T|X=x] = \int_{t=0}^{\infty} \hat{S}(t)dt$$

6.7.4. Wide prediction intervals

The nature of time-to-event data results in very wide prediction intervals:

- Suppose a cancer patient is predicted to have a mean lifetime of 5 years after diagnosis and suppose the distribution is exponential.
- If we want a 95% interval for survival, the lower end is at the 0.025 percentage point of the exponential which is qexp(.025, rate = 1/5) = 0.1266 years, or 1/40 of the mean lifetime.

- The upper end is at the 0.975 point which is qexp(.975, rate = 1/5) = 18.4444 years, or 3.7 times the mean lifetime.
- Saying that the survival time is somewhere between 6 weeks and 18 years does not seem very useful, but it may be the best we can do.
- For survival analysis, something is like a residual if it is small when the model is accurate or if the accumulation of them is in some way minimized by the estimation algorithm, but there is no exact equivalence to linear regression residuals.
- And if there is, they are mostly quite large!

6.7.5. Types of Residuals in Time-to-Event Models

- It is often hard to make a decision from graph appearances, though the process can reveal much.
- Some diagnostic tests are based on residuals as with other regression methods:
 - Schoenfeld residuals (via cox.zph) for proportionality
 - Cox-Snell residuals for goodness of fit (Section 6.8)
 - martingale residuals for non-linearity
 - **dfbeta** for influence.

6.7.6. Schoenfeld residuals

- There is a Schoenfeld residual for each subject i with an event (not censored) and for each predictor x_k .
- At the event time t for that subject, there is a risk set R, and each subject j in the risk set has a risk coefficient θ_j and also a value x_{jk} of the predictor.
- The Schoenfeld residual is the difference between x_{ik} and the risk-weighted average of all the x_{ik} over the risk set.

$$r_{ik}^S = x_{ik} - \frac{\sum_{k \in R} x_{jk} \theta_k}{\sum_{k \in R} \theta_k}$$

This residual measures how typical the individual subject is with respect to the covariate at the time of the event. Since subjects should fail more or less uniformly according to risk, the Schoenfeld residuals should be approximately level over time, not increasing or decreasing.

We can test this with the correlation with time on some scale, which could be the time itself, the log time, or the rank in the set of failure times.

The default is to use the KM curve as a transform, which is similar to the rank but deals better with censoring.

The cox.zph() function implements a score test proposed in Grambsch and Therneau (1994).


```
hodg.zph = cox.zph(hodg.cox1)
print(hodg.zph)
#>
                chisq df
                             р
#> gtype
               0.5400 1 0.462
#> dtype
               1.8012 1 0.180
               3.8805 1 0.049
#> score
#> wtime
             0.0173 1 0.895
#> gtype:dtype 4.0474 1 0.044
#> GLOBAL
            13.7573 5 0.017
```

6.7.6.1. gtype

```
ggcoxzph(hodg.zph, var = "gtype")
```

Global Schoenfeld Test p: 0.01723

Schoenfeld Individual Test p: 0.4624




6.7.6.2. dtype

ggcoxzph(hodg.zph, var = "dtype")

Global Schoenfeld Test p: 0.01723

Schoenfeld Individual Test p: 0.1796




6.7.6.3. score

ggcoxzph(hodg.zph, var = "score")

Global Schoenfeld Test p: 0.01723

Schoenfeld Individual Test p: 0.0489




6.7.6.4. wtime

```
ggcoxzph(hodg.zph, var = "wtime")
```

Global Schoenfeld Test p: 0.01723

Schoenfeld Individual Test p: 0.8954



6.7.6.5. gtype:dtype

```
ggcoxzph(hodg.zph, var = "gtype:dtype")
```

Global Schoenfeld Test p: 0.01723



6.7.6.6. Conclusions

- From the correlation test, the Karnofsky score and the interaction with graft type disease type induce modest but statistically significant non-proportionality.
- The sample size here is relatively small (26 events in 43 subjects). If the sample size is large, very small amounts of non-proportionality can induce a significant result.
- As time goes on, autologous grafts are over-represented at their own event times, but those from HOD patients become less represented.
- Both the statistical tests and the plots are useful.

6.8. Goodness of Fit using the Cox-Snell Residuals

(references: Klein and Moeschberger (2003), §11.2, and Dobson and Barnett (2018), §10.6)

Suppose that an individual has a survival time T which has survival function S(t), meaning that $\Pr(T > t) = S(t)$. Then S(T) has a uniform distribution on (0,1):

$$\begin{split} \Pr(S(T_i) \leq u) &= \Pr(T_i > S_i^{-1}(u)) \\ &= S_i(S_i^{-1}(u)) \\ &= u \end{split}$$

Also, if U has a uniform distribution on (0,1), then what is the distribution of $-\ln(U)$?

$$\begin{split} \Pr(-\ln(U) < x) &= \Pr(U > \exp\left\{-x\right\}) \\ &= 1 - e^{-x} \end{split}$$

which is the CDF of an exponential distribution with parameter $\lambda = 1$.

Definition 6.5 (Cox-Snell generalized residuals).

The Cox-Snell generalized residuals are defined as:

$$r_i^{CS} \stackrel{\text{def}}{=} \hat{H}(t_i | \tilde{x}_i)$$

If the estimate \hat{S}_i is accurate, r_i^{CS} should have an exponential distribution with constant hazard $\lambda=1$, which means that these values should look like a censored sample from this exponential distribution.

```
hodg2 = hodg2 |>
  mutate(cs = predict(hodg.cox1, type = "expected"))

surv.csr = survfit(
  data = hodg2,
  formula = Surv(time = cs, event = delta == "dead") ~ 1,
  type = "fleming-harrington")

autoplot(surv.csr, fun = "cumhaz") +
  geom_abline(aes(intercept = 0, slope = 1), col = "red") +
  theme_bw()
```



Figure 6.11.: Cumulative Hazard of Cox-Snell Residuals

The line with slope 1 and intercept 0 fits the curve relatively well, so we don't see lack of fit using this procedure.

6.9. Martingale Residuals

The martingale residuals are a slight modification of the Cox-Snell residuals. If the censoring indicator is δ_i , then

$$r_i^M = \delta_i - r_i^{CS}$$

These residuals can be interpreted as an estimate of the excess number of events seen in the data but not predicted by the model. We will use these to examine the functional forms of continuous covariates.

6.9.1. Using Martingale Residuals

Martingale residuals can be used to examine the functional form of a numeric variable.

- We fit the model without that variable and compute the martingale residuals.
- We then plot these martingale residuals against the values of the variable.
- We can see curvature, or a possible suggestion that the variable can be discretized.

Let's use this to examine the score and wtime variables in the wtime data set.

Karnofsky score

```
hodg2 = hodg2 |>
mutate(
    mres =
    hodg.cox1 |>
    update(. ~ . - score) |>
    residuals(type="martingale"))

hodg2 |>
    ggplot(aes(x = score, y = mres)) +
    geom_point() +
    geom_smooth(method = "loess", aes(col = "loess")) +
    geom_smooth(method = 'lm', aes(col = "lm")) +
    theme_classic() +
    xlab("Karnofsky Score") +
    ylab("Martingale Residuals") +
    guides(col=guide_legend(title = ""))
```




Figure 6.12.: Martingale Residuals vs. Karnofsky Score

The line is almost straight. It could be some modest transformation of the Karnofsky score would help, but it might not make much difference.

Waiting time

```
hodg2$mres =
hodg.cox1 |>
update(. ~ . - wtime) |>
residuals(type="martingale")
```

```
hodg2 |>
  ggplot(aes(x = wtime, y = mres)) +
  geom_point() +
  geom_smooth(method = "loess", aes(col = "loess")) +
  geom_smooth(method = 'lm', aes(col = "lm")) +
  theme_classic() +
  xlab("Waiting Time") +
  ylab("Martingale Residuals") +
  guides(col=guide_legend(title = ""))
```




Figure 6.13.: Martingale Residuals vs. Waiting Time

The line could suggest a step function. To see where the drop is, we can look at the largest waiting times and the associated martingale residual.

The martingale residuals are all negative for wtime >83 and positive for the next smallest value. A reasonable cut-point is 80 days.

Updating the model

Let's reformulate the model with dichotomized wtime.

```
hodg2 =
hodg2 |>
mutate(
    wt2 = cut(
        wtime,c(0, 80, 200),
        labels=c("short","long")))

hodg.cox2 =
    coxph(
    formula =
        Surv(time, event = delta == "dead") ~
        gtype*dtype + score + wt2,
        data = hodg2)
```

```
hodg.cox1 |> drop1(test="Chisq")

#> # A tibble: 4 x 4

#> Df AIC LRT `Pr(>Chi)`

#> <dbl> <dbl> <dbl> <dbl>
#> 1 NA 152. NA NA

#> 2 1 168. 17.2 0.0000330

#> 3 1 154. 3.28 0.0702

#> 4 1 156. 5.44 0.0197
```

```
hodg.cox2 |> drop1(test="Chisq")
#> # A tibble: 4 x 4
```

```
#>
             AIC
                    LRT
                         `Pr(>Chi)`
#>
     <dbl> <dbl> <dbl>
                               <dbl>
#> 1
            149. NA
        NA
                        NA
            169. 21.6
                         0.00000335
         1
            154.
                  6.61
                         0.0102
#> 4
         1
            152. 4.97
                         0.0258
```

The new model has better (lower) AIC.

6.10. Checking for Outliers and Influential Observations

We will check for outliers using the deviance residuals. The martingale residuals show excess events or the opposite, but highly skewed, with the maximum possible value being 1, but the smallest value can be very large negative. Martingale residuals can detect unexpectedly long-lived patients, but patients who die unexpectedly early show up only in the deviance residual. Influence will be examined using dfbeta in a similar way to linear regression, logistic regression, or Poisson regression.

6.10.1. Deviance Residuals

$$\begin{split} r_i^D &= \mathrm{sign}(r_i^M) \sqrt{-2 \left[r_i^M + \delta_i \ln(\delta_i - r_i^M)\right]} \\ r_i^D &= \mathrm{sign}(r_i^M) \sqrt{-2 \left[r_i^M + \delta_i \ln(r_i^{CS})\right]} \end{split}$$

Roughly centered on 0 with approximate standard deviation 1.

6.10.2.

```
hodg.mart = residuals(hodg.cox2,type="martingale")
hodg.dev = residuals(hodg.cox2,type="deviance")
hodg.dfb = residuals(hodg.cox2,type="dfbeta")
hodg.preds = predict(hodg.cox2) #linear predictor

plot(hodg.preds,
    hodg.mart,
    xlab="Linear Predictor",
    ylab="Martingale Residual")
```



Figure 6.14.: Martingale Residuals vs. Linear Predictor

The smallest three martingale residuals in order are observations 1, 29,

and 18.

plot(hodg.preds,hodg.dev,xlab="Linear Predictor",ylab="Deviance Residual")



Figure 6.15.: Deviance Residuals vs. Linear Predictor

The two largest deviance residuals are observations 1 and 29. Worth examining.

6.10.3. dfbeta

- dfbeta is the approximate change in the coefficient vector if that observation were dropped
- dfbetas is the approximate change in the coefficients, scaled by the standard error for the coefficients.

6.10.3.1. Graft type

```
plot(hodg.dfb[,1],xlab="Observation Order",ylab="dfbeta for Graft Type")
```



Figure 6.16.: dfbeta Values by Observation Order for Graft Type

The smallest dfbeta for graft type is observation 1.

6.10.3.2. Disease type

```
plot(hodg.dfb[,2],
     xlab="Observation Order",
     ylab="dfbeta for Disease Type")
```



Figure 6.17.: dfbeta Values by Observation Order for Disease Type

The smallest two dfbeta values for disease type are observations 1 and 16.

6.10.3.3. Karnofsky score



Figure 6.18.: dfbeta Values by Observation Order for Karnofsky Score

The two highest dfbeta values for score are observations 1 and 18. The next three are observations 17, 29, and 19. The smallest value is observation 2.

6.10.3.4. Waiting time (dichotomized)

```
plot(
  hodg.dfb[,4],
  xlab="Observation Order",
  ylab="dfbeta for `Waiting Time < 80`")</pre>
```



Figure 6.19.: dfbeta Values by Observation Order for Waiting Time (dichotomized)

The two large values of dfbeta for dichotomized waiting time are observations 15 and 16. This may have to do with the discretization of waiting time.

6.10.3.5. Interaction: graft type and disease type

```
plot(hodg.dfb[,5],
     xlab="Observation Order",
     ylab="dfbeta for dtype:gtype")
```



Figure 6.20.: dfbeta Values by Observation Order for dtype:gtype

The two largest values are observations 1 and 16. The smallest value is observation 35.

Table 6.1.: Observations to Examine by Residuals and Influence

| Diagnostic | Observations to Examine |
|----------------------------|-------------------------|
| Martingale Residuals | 1, 29, 18 |
| Deviance Residuals | 1, 29 |
| Graft Type Influence | 1 |
| Disease Type Influence | 1, 16 |
| Karnofsky Score Influence | 1, 18 (17, 29, 19) |
| Waiting Time Influence | 15, 16 |
| Graft by Disease Influence | 1, 16, 35 |

The most important observations to examine seem to be 1, 15, 16, 18, and 29.

6.10.4.

```
with(hodg, summary(time[delta==1]))
     Min. 1st Qu. Median
                             Mean 3rd Qu.
                                             Max.
#>
       2.0 41.2
                   62.5
                             97.6 83.2
                                            524.0
with(hodg,summary(wtime))
     Min. 1st Qu. Median
                             Mean 3rd Qu.
                                             Max.
       5.0 16.0
                    24.0
                             37.7 55.5
                                            171.0
with(hodg,summary(score))
     Min. 1st Qu. Median
#>
                             Mean 3rd Qu.
                                             Max.
     20.0 60.0
                     80.0
                             76.3
                                     90.0
                                            100.0
hodg.cox2
#> Call:
#> coxph(formula = Surv(time, event = delta == "dead") ~ gtype *
       dtype + score + wt2, data = hodg2)
#>
#>
#>
                                 coef exp(coef) se(coef) z
                                                    0.59 1 0.263
#> gtypeAutologous
                                 0.67
                                           1.94
#> dtypeHodgkins
                                 2.33
                                          10.25
                                                    0.73 3 0.002
#> score
                                -0.06
                                           0.95
                                                    0.01 -4 8e-06
#> wt2long
                                -2.06
                                           0.13
                                                    1.05 - 20.050
#> gtypeAutologous:dtypeHodgkins -2.07
                                           0.13
                                                    0.93 -2 0.026
#>
#> Likelihood ratio test=35 on 5 df, p=1e-06
\# n= 43, number of events= 26
```

```
hodg2[c(1,15,16,18,29),] |>
  select(gtype, dtype, time, delta, score, wtime) |>
  mutate(
    comment =
      c(
        "early death, good score, low risk",
        "high risk grp, long wait, poor score",
        "high risk grp, short wait, poor score",
        "early death, good score, med risk grp",
        "early death, good score, med risk grp"
      ))
#> # A tibble: 5 x 7
     gtype
                dtype
                              time delta score wtime comment
                              <int> <chr> <int> <int> <chr>
#>
     <chr>
                <fct>
#> 1 Allogenic Non-Hodgkins
                                 28 dead
                                             90
                                                   24 early death, good score, low ~
#> 2 Allogenic Hodgkins
                                 77 dead
                                             60
                                                  102 high risk grp, long wait, poo~
#> 3 Allogenic Hodgkins
                                                   71 high risk grp, short wait, po~
                                 79 dead
                                             70
#> 4 Autologous Non-Hodgkins
                                 53 dead
                                             90
                                                   17 early death, good score, med ~
#> 5 Autologous Hodgkins
                                                   73 early death, good score, med ~
                                 30 dead
                                             90
```

6.10.5. Action Items

- Unusual points may need checking, particularly if the data are not completely cleaned. In this case, observations 15 and 16 may show some trouble with the dichotomization of waiting time, but it still may be useful.
- The two largest residuals seem to be due to unexpectedly early deaths, but unfortunately this can occur.
- If hazards don't look proportional, then we may need to use strata, between which the base hazards are permitted to be different. For this problem, the natural strata are the two diseases, because they could need to be managed differently anyway.

• A main point that we want to be sure of is the relative risk difference by disease type and graft type.

```
hodg.cox2 |>
  predict(
    reference = "zero",
    newdata = means |>
        mutate(
        wt2 = "short",
        score = 0),
    type = "lp") |>
  data.frame('linear predictor' = _) |>
  pander()
```

Table 6.2.: Linear Risk Predictors for Lymphoma

| | linear.predictor |
|-------------------------|------------------|
| Non-Hodgkins, Allogenic | 0 |
| Non- | 0.6651 |
| Hodgkins, Autologous | |
| Hodgkins, Allogenic | 2.327 |
| Hodgkins, Autologous | 0.9256 |

For Non-Hodgkin's, the allogenic graft is better. For Hodgkin's, the autologous graft is much better.

6.11. Stratified survival models

6.11.1. Revisiting the leukemia dataset (anderson)

We will analyze remission survival times on 42 leukemia patients, half on new treatment, half on standard treatment.

This is the same data as the drug6mp data from KMsurv, but with two other variables and without the pairing. This version comes from Kleinbaum and Klein (2012) (e.g., p281):

```
anderson =
 paste0(
    "http://web1.sph.emory.edu/dkleinb/allDatasets/",
    "surv2datasets/anderson.dta") |>
 haven::read_dta() |>
 mutate(
   status = status |>
      case_match(
       1 ~ "relapse",
       0 ~ "censored"
      ),
    sex = sex |>
      case_match(
       0 ~ "female",
        1 ~ "male"
      ) |>
      factor() |>
     relevel(ref = "female"),
   rx = rx >
      case_match(
       0 ~ "new",
        1 ~ "standard"
      factor() |> relevel(ref = "standard"),
    surv = Surv(
      time = survt,
```

```
event = (status == "relapse"))
)
print(anderson)
```

6.11.2. Cox semi-parametric proportional hazards model

```
anderson.cox1 = coxph(
 formula = surv ~ rx + sex + logwbc,
 data = anderson)
summary(anderson.cox1)
#> Call:
#> coxph(formula = surv ~ rx + sex + logwbc, data = anderson)
#>
  n= 42, number of events= 30
#>
#>
         coef exp(coef) se(coef) z Pr(>|z|)
#> rxnew -1.504 0.222 0.462 -3.26 0.0011 **
#> sexmale 0.315
                  1.370
                          0.455 0.69 0.4887
#> logwbc 1.682
                  #> ---
#> Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
       exp(coef) exp(-coef) lower .95 upper .95
#> rxnew
          0.222
                   4.498
                             0.090 0.549
#> sexmale
           1.370
                      0.730
                              0.562
                                      3.338
#> logwbc
            5.376
                      0.186
                              2.779 10.398
#>
#> Concordance= 0.851 (se = 0.041 )
#> Likelihood ratio test= 47.2 on 3 df, p=3e-10
```

6.11.2.1. Test the proportional hazards assumption

```
cox.zph(anderson.cox1)

#> chisq df p

#> rx 0.036 1 0.85

#> sex 5.420 1 0.02

#> logwbc 0.142 1 0.71

#> GLOBAL 5.879 3 0.12
```

6.11.2.2. Graph the K-M survival curves

```
anderson_km_model = survfit(
  formula = surv ~ sex,
  data = anderson)

anderson_km_model |>
  autoplot(conf.int = FALSE) +
  theme_bw() +
  theme(legend.position="bottom")
```



The survival curves cross, which indicates a problem in the proportionality assumption by sex.

6.11.3. Graph the Nelson-Aalen cumulative hazard

We can also look at the log-hazard ("cloglog survival") plots:

```
anderson_na_model = survfit(
  formula = surv ~ sex,
  data = anderson,
  type = "fleming")

anderson_na_model |>
  autoplot(
   fun = "cumhaz",
   conf.int = FALSE) +
```

```
theme_classic() +
theme(legend.position="bottom") +
ylab("log(Cumulative Hazard)") +
scale_y_continuous(
   trans = "log10",
   name = "Cumulative hazard (H(t), log scale)") +
scale_x_continuous(
   breaks = c(1,2,5,10,20,50),
   trans = "log"
)
```



Figure 6.21.: Cumulative hazard (cloglog scale) for anderson data

This can be fixed by using strata or possibly by other model alterations.

6.11.4. The Stratified Cox Model

- In a stratified Cox model, each stratum, defined by one or more factors, has its own base survival function $h_0(t)$.
- But the coefficients for each variable not used in the strata definitions are assumed to be the same across strata.
- To check if this assumption is reasonable one can include interactions with strata and see if they are significant (this may generate a warning and NA lines but these can be ignored).
- Since the sex variable shows possible non-proportionality, we try stratifying on sex.

```
anderson.coxph.strat =
  coxph(
   formula =
     surv ~ rx + logwbc + strata(sex),
   data = anderson)
summary(anderson.coxph.strat)
#> Call:
#> coxph(formula = surv ~ rx + logwbc + strata(sex), data = anderson)
#>
#>
    n= 42, number of events= 30
#>
#>
            coef exp(coef) se(coef)
                                        z Pr(>|z|)
#> rxnew -0.998
                     0.369
                              0.474 - 2.11
                                             0.035 *
                     4.279
                              0.344 4.22 2.4e-05 ***
#> logwbc 1.454
#> ---
#> Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
#>
          exp(coef) exp(-coef) lower .95 upper .95
#> rxnew
              0.369
                         2.713
                                   0.146
                                             0.932
              4.279
                         0.234
#> logwbc
                                   2.180
                                             8.398
```

Let's compare this to a model fit only on the subset of males:

```
anderson.coxph.male =
 coxph(
   formula = surv ~ rx + logwbc,
   subset = sex == "male",
   data = anderson)
summary(anderson.coxph.male)
#> Call:
#> coxph(formula = surv ~ rx + logwbc, data = anderson, subset = sex ==
     "male")
#>
#>
#> n= 20, number of events= 14
#>
           coef exp(coef) se(coef) z Pr(>|z|)
#> rxnew -1.978 0.138 0.739 -2.68 0.0075 **
#> logwbc 1.743
                   5.713
                            0.536 3.25
                                        0.0011 **
#> Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
#>
         exp(coef) exp(-coef) lower .95 upper .95
#> rxnew
             0.138
                       7.227
                                0.0325
             5.713
                        0.175
                                1.9991
                                         16.328
#> logwbc
#>
\# Concordance = 0.905 (se = 0.043)
```

```
#> Likelihood ratio test= 29.2 on 2 df, p=5e-07
\#> Wald test = 15.3 on 2 df, p=5e-04
#> Score (logrank) test = 26.4 on 2 df, p=2e-06
anderson.coxph.female =
 coxph(
   formula =
    surv ~ rx + logwbc,
   subset = sex == "female",
   data = anderson)
summary(anderson.coxph.female)
#> Call:
#> coxph(formula = surv ~ rx + logwbc, data = anderson, subset = sex ==
#>
     "female")
#>
#> n= 22, number of events= 16
#>
#>
         coef exp(coef) se(coef) z Pr(>|z|)
#> rxnew -0.311 0.733 0.564 -0.55 0.581
                 3.341 0.503 2.40 0.017 *
#> logwbc 1.206
#> ---
#> Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
#>
       exp(coef) exp(-coef) lower .95 upper .95
#> rxnew 0.733 1.365 0.243 2.21
#> logwbc
            3.341
                      0.299
                             1.245
                                        8.96
#>
```

#> Concordance= 0.692 (se = 0.085)

#> Likelihood ratio test= 6.65 on 2 df, p=0.04 #> Wald test = 6.36 on 2 df, p=0.04 #> Score (logrank) test = 6.74 on 2 df, p=0.03

The coefficients of treatment look different. Are they statistically different?

```
anderson.coxph.strat.intxn =
  coxph(
   formula = surv ~ strata(sex) * (rx + logwbc),
   data = anderson)
anderson.coxph.strat.intxn |> summary()
#> Call:
#> coxph(formula = surv ~ strata(sex) * (rx + logwbc), data = anderson)
#>
    n= 42, number of events= 30
#>
#>
                           coef exp(coef) se(coef)
#>
                                                     z Pr(>|z|)
#> rxnew
                         -0.311
                                   0.733
                                             0.564 - 0.55
                                                            0.581
#> logwbc
                          1.206
                                    3.341
                                             0.503 2.40
                                                            0.017 *
#> strata(sex)male:rxnew -1.667
                                          0.930 -1.79
                                    0.189
                                                            0.073 .
#> strata(sex)male:logwbc 0.537
                                 1.710
                                          0.735 0.73
                                                            0.465
#> ---
#> Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
#>
#>
                         exp(coef) exp(-coef) lower .95 upper .95
#> rxnew
                             0.733
                                        1.365
                                                 0.2427
                                                             2.21
#> logwbc
                             3.341
                                        0.299
                                                 1.2452
                                                             8.96
#> strata(sex)male:rxnew
                             0.189
                                        5.294
                                                 0.0305
                                                             1.17
#> strata(sex)male:logwbc
                             1.710
                                    0.585 0.4048
                                                             7.23
#>
#> Concordance= 0.797 (se = 0.058 )
#> Likelihood ratio test= 35.8 on 4 df,
                                          p = 3e - 07
#> Wald test
                       = 21.7 on 4 df,
                                          p = 2e - 04
#> Score (logrank) test = 33.1 on 4 df,
                                          p=1e-06
```

We don't have enough evidence to tell the difference between these two models.

6.11.5. Conclusions

- We chose to use a stratified model because of the apparent nonproportionality of the hazard for the sex variable.
- When we fit interactions with the strata variable, we did not get an improved model (via the likelihood ratio test).
- So we use the stratifed model with coefficients that are the same across strata.

6.11.6. Another Modeling Approach

- We used an additive model without interactions and saw that we might need to stratify by sex.
- Instead, we could try to improve the model's functional form maybe the interaction of treatment and sex is real, and after fitting that we might not need separate hazard functions.
- Either approach may work.

```
anderson.coxph.intxn =
 coxph(
   formula = surv ~ (rx + logwbc) * sex,
   data = anderson)
anderson.coxph.intxn |> summary()
#> Call:
#> coxph(formula = surv ~ (rx + logwbc) * sex, data = anderson)
#>
   n= 42, number of events= 30
#>
#>
#>
                  coef exp(coef) se(coef) z Pr(>|z|)
#> rxnew
               #> logwbc
               1.0637
                         2.8971 0.4726 2.25
                                              0.024 *
               -2.8052 0.0605 2.0323 -1.38 0.167
#> sexmale
#> rxnew:sexmale -2.1782 0.1132 0.9109 -2.39 0.017 *
#> logwbc:sexmale 1.2303
                        3.4223 0.6301 1.95
                                              0.051 .
#> ---
#> Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
#>
#>
               exp(coef) exp(-coef) lower .95 upper .95
#> rxnew
                 0.6874
                            1.455
                                  0.23185
                                              2.038
#> logwbc
                 2.8971
                            0.345 1.14730
                                              7.315
                          16.531 0.00113
#> sexmale
                 0.0605
                                              3.248
#> rxnew:sexmale 0.1132
                           8.830 0.01899
                                             0.675
                                           11.766
#> logwbc:sexmale 3.4223
                           0.292 0.99539
#> Concordance= 0.861 (se = 0.036 )
#> Likelihood ratio test= 57 on 5 df, p=5e-11
\#> Wald test = 35.6 on 5 df, p=1e-06
\# Score (logrank) test = 57.1 on 5 df, p=5e-11
```

```
cox.zph(anderson.coxph.intxn)
#>
              chisq df
#> rx
              0.136
                    1 0.71
#> logwbc
                    1 0.20
             1.652
#> sex
              1.266
                    1 0.26
#> rx:sex
             0.149 1 0.70
#> logwbc:sex 0.102
                    1 0.75
#> GLOBAL
             3.747 5 0.59
```

6.12. Time-varying covariates

(adapted from Klein and Moeschberger (2003), §9.2)

6.12.1. Motivating example: back to the leukemia dataset

```
# load the data:
data(bmt, package = 'KMsurv')
bmt |> as_tibble() |> print(n = 5)
#> # A tibble: 137 x 22
    group
            t1
                  t2
                       d1
                            d2
                                  d3
                                             da
                                                  tc
                                       ta
                                                        dc
                                                             tp
                                                                  dp
                                                                        z1
    #> 1
        1
          2081 2081
                        0
                             0
                                   0
                                       67
                                              1
                                                 121
                                                        1
                                                             13
                                                                   1
                                                                        26
#> 2
        1
          1602
               1602
                        0
                             0
                                   0
                                     1602
                                              0
                                                 139
                                                        1
                                                             18
                                                                   1
                                                                        21
#> 3
        1
         1496
               1496
                        0
                             0
                                   0
                                     1496
                                              0
                                                 307
                                                        1
                                                             12
                                                                   1
                                                                        26
#> 4
       1 1462
               1462
                             0
                                   0
                                       70
                                             1
                                                  95
                                                        1
                                                             13
                                                                        17
#> 5
       1 1433
               1433
                             0
                                     1433
                                              0
                                                 236
                                                        1
                                                             12
                                                                        32
#> # i 132 more rows
#> # i 9 more variables: z2 <int>, z3 <int>, z4 <int>, z5 <int>, z6 <int>,
      z7 <int>, z8 <int>, z9 <int>, z10 <int>
```

This dataset comes from the Copelan et al. (1991) study of allogenic bone marrow transplant therapy for acute myeloid leukemia (AML) and acute lymphoblastic leukemia (ALL).

Outcomes (endpoints)

• The main endpoint is disease-free survival (t2 and d3) for the three risk groups, "ALL", "AML Low Risk", and "AML High Risk".

Possible intermediate events

- graft vs. host disease (**GVHD**), an immunological rejection response to the transplant (bad)
- acute (AGVHD)
- chronic (CGVHD)
- platelet recovery, a return of platelet count to normal levels (good)

One or the other, both in either order, or neither may occur.

Covariates

- We are interested in possibly using the covariates z1-z10 to adjust for other factors.
- In addition, the time-varying covariates for acute GVHD, chronic GVHD, and platelet recovery may be useful.

6.12.1.1. Preprocessing

We reformat the data before analysis:

```
# reformat the data:
bmt1 =
 bmt |>
 as_tibble() |>
 mutate(
   id = 1:n(), # will be used to connect multiple records for the same individual
   group = group |>
     case_match(
       1 ~ "ALL",
       2 ~ "Low Risk AML",
        3 ~ "High Risk AML") |>
     factor(levels = c("ALL", "Low Risk AML", "High Risk AML")),
    `patient age` = z1,
    'donor age' = z2,
    \epsilon = z3 >
      case_match(
       0 ~ "Female",
       1 ~ "Male"),
    donor sex = z4 >
      case_match(
       0 ~ "Female",
        1 ~ "Male"),
    `Patient CMV Status` = z5 |>
      case_match(
       0 ~ "CMV Negative",
        1 ~ "CMV Positive"),
```

```
`Donor CMV Status` = z6 |>
      case_match(
       0 ~ "CMV Negative",
       1 ~ "CMV Positive"),
    `Waiting Time to Transplant` = z7,
   FAB = z8 \mid >
     case_match(
       1 ~ "Grade 4 Or 5 (AML only)",
       0 ~ "Other") |>
     factor() |>
     relevel(ref = "Other"),
   hospital = z9 |> # `z9` is hospital
      case_match(
       1 ~ "Ohio State University",
       2 ~ "Alferd",
       3 ~ "St. Vincent",
        4 ~ "Hahnemann") |>
      factor() |>
     relevel(ref = "Ohio State University"),
   MTX = (z10 == 1) # a prophylatic treatment for GVHD
 ) |>
  select(-(z1:z10)) # don't need these anymore
bmt1 |>
 select(group, id:MTX) |>
 print(n = 10)
#> # A tibble: 137 x 12
#> group id `patient age` `donor age` `patient sex` `donor sex`
```

| #> | <1 | fct> | <int></int> | <int></int> | <int></int> | <chr></chr> | <chr></chr> |
|----|-------|------|-------------|-------------|-------------|-------------|-------------|
| #> | 1 AI | LL | 1 | 26 | 33 | Male | Female |
| #> | 2 AI | LL | 2 | 21 | 37 | Male | Male |
| #> | 3 AI | LL | 3 | 26 | 35 | Male | Male |
| #> | 4 AI | LL | 4 | 17 | 21 | Female | Male |
| #> | 5 AI | LL | 5 | 32 | 36 | Male | Male |
| #> | 6 AI | LL | 6 | 22 | 31 | Male | Male |
| #> | 7 AI | LL | 7 | 20 | 17 | Male | Female |
| #> | 8 AI | LL | 8 | 22 | 24 | Male | Female |
| #> | 9 AI | LL | 9 | 18 | 21 | Female | Male |
| #> | 10 AI | LL | 10 | 24 | 40 | Male | Male |
| #> | # i ' | 127 | more roug | | | | |

#> # i 6 more variables: `Patient CMV Status` <chr>, `Donor CMV Status` <chr>,
#> # `Waiting Time to Transplant` <int>, FAB <fct>, hospital <fct>, MTX <lgl>

6.12.2. Time-Dependent Covariates

- A time-dependent covariate ("TDC") is a covariate whose value changes during the course of the study.
- For variables like age that change in a linear manner with time, we can just use the value at the start.
- But it may be plausible that when and if GVHD occurs, the risk of relapse or death increases, and when and if platelet recovery occurs, the risk decreases.

6.12.3. Analysis in R

- We form a variable precovery which is = 0 before platelet recovery and is = 1 after platelet recovery, if it occurs.
- For each subject where platelet recovery occurs, we set up multiple records (lines in the data frame); for example one from t = 0 to the

time of platelet recovery, and one from that time to relapse, recovery, or death.

- We do the same for acute GVHD and chronic GVHD.
- For each record, the covariates are constant.

```
bmt2 = bmt1 |>
    #set up new long-format data set:
    tmerge(bmt1, id = id, tstop = t2) |>

# the following three steps can be in any order,
# and will still produce the same result:
#add aghvd as tdc:
    tmerge(bmt1, id = id, agvhd = tdc(ta)) |>
#add cghvd as tdc:
    tmerge(bmt1, id = id, cgvhd = tdc(tc)) |>
#add platelet recovery as tdc:
    tmerge(bmt1, id = id, precovery = tdc(tp))

bmt2 = bmt2 |>
    as_tibble() |>
    mutate(status = as.numeric((tstop == t2) & d3))
# status only = 1 if at end of t2 and not censored
```

Let's see how we've rearranged the first row of the data:

```
bmt1 |>
 dplyr::filter(id == 1) |>
 dplyr::select(id, t1, d1, t2, d2, d3, ta, da, tc, dc, tp, dp)
#> # A tibble: 1 x 12
     id
         t1
                 t2
                     d2
                         d3
                                 da
                             ta
                                     tc
                                         dc
                                             tp
                                                 dp
   #> 1 1 2081 0 2081 0 0
                             67
                               1 121
```

The event times for this individual are:

- t = 0 time of transplant
- tp = 13 platelet recovery
- ta = 67 acute GVHD onset
- tc = 121 chronic GVHD onset
- t2 = 2081 end of study, patient not relapsed or dead

After converting the data to long-format, we have:

```
bmt2 |>
  select(
    id,
    tstart,
    tstop,
    agvhd,
    cgvhd,
    precovery,
    status
  ) |>
  dplyr::filter(id == 1)
#> # A tibble: 4 x 7
        id tstart tstop agvhd cgvhd precovery status
            <dbl> <int> <int> <int>
                                           <int>
#> 1
         1
                 0
                       13
                              0
                                     0
                                                0
                                                        0
#> 2
         1
                13
                       67
                              0
                                     0
                                                1
                                                        0
#> 3
         1
                67
                     121
                              1
                                     0
                                                1
                                                        0
                    2081
                                     1
                                                        0
#> 4
         1
               121
                              1
```

Note that status could have been 1 on the last row, indicating that relapse or death occurred; since it is false, the participant must have exited the study without experiencing relapse or death (i.e., they were censored).

6.12.4. Event sequences

Let:

- A = acute GVHD
- C = chronic GVHD
- P = platelet recovery

Each of the eight possible combinations of A or not-A, with C or not-C, with P or not-P occurs in this data set.

- A always occurs before C, and P always occurs before C, if both occur.
- Thus there are ten event sequences in the data set: None, A, C, P, AC, AP, PA, PC, APC, and PAC.
- In general, there could be as many as 1+3+(3)(2)+6=16 sequences, but our domain knowledge tells us that some are missing: CA, CP, CAP, CPA, PCA, PC, PAC
- Different subjects could have 1, 2, 3, or 4 intervals, depending on which of acute GVHD, chronic GVHD, and/or platelet recovery occurred.
- The final interval for any subject has status = 1 if the subject relapsed or died at that time; otherwise status = 0.
- Any earlier intervals have status = 0.
- Even though there might be multiple lines per ID in the dataset, there is never more than one event, so no alterations need be made in the estimation procedures or in the interpretation of the output.
- The function tmerge in the survival package eases the process of constructing the new long-format dataset.

6.12.5. Model with Time-Fixed Covariates

```
bmt1 =
 bmt1 |>
 mutate(surv = Surv(t2,d3))
bmt_coxph_TF = coxph(
 formula = surv ~ group + `patient age`*`donor age` + FAB,
 data = bmt1)
summary(bmt_coxph_TF)
#> Call:
#> coxph(formula = surv ~ group + `patient age` * `donor age` +
#>
     FAB, data = bmt1)
#>
#>
   n= 137, number of events= 83
#>
#>
                                 coef exp(coef) se(coef) z Pr(>|z|)
#> groupLow Risk AML
                           -1.090648 0.335999 0.354279 -3.08 0.00208 **
#> groupHigh Risk AML
                           -0.403905 0.667707
                                               0.362777 -1.11 0.26555
#> `patient age`
                           -0.081639  0.921605  0.036107  -2.26  0.02376 *
#> `donor age`
                           #> FABGrade 4 Or 5 (AML only) 0.837416 2.310388 0.278464 3.01 0.00264 **
#> `patient age`: `donor age` 0.003159 1.003164 0.000951 3.32 0.00089 ***
#> ---
#> Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
#>
#>
                            exp(coef) exp(-coef) lower .95 upper .95
#> groupLow Risk AML
                               0.336
                                          2.976
                                                  0.168
                                                             0.673
#> groupHigh Risk AML
                               0.668
                                          1.498
                                                   0.328
                                                             1.360
#> `patient age`
                               0.922
                                          1.085
                                                  0.859
                                                             0.989
#> `donor age`
                               0.919
                                          1.088
                                                  0.866
                                                             0.975
#> FABGrade 4 Or 5 (AML only)
                               2.310
                                                  1.339
                                                             3.988
                                          0.433
#> `patient age`: `donor age`
                               1.003
                                          0.997
                                                  1.001
                                                             1.005
#>
```

```
#> Concordance= 0.665 (se = 0.033 )
#> Likelihood ratio test= 32.8 on 6 df, p=1e-05
\#> Wald test = 33 on 6 df, p=1e-05
#> Score (logrank) test = 35.8 on 6 df, p=3e-06
drop1(bmt_coxph_TF, test = "Chisq")
#> # A tibble: 4 x 4
      Df AIC LRT `Pr(>Chi)`
    <dbl> <dbl> <dbl> <dbl>
#>
#> 1 NA 726. NA
                    NA
#> 2
      2 734. 12.5 0.00192
#> 3
      1 733. 9.22 0.00240
#> 4 1 733. 9.51 0.00204
```

```
bmt1$mres =
  bmt_coxph_TF |>
  update(. ~ . - `donor age`) |>
  residuals(type="martingale")

bmt1 |>
  ggplot(aes(x = `donor age`, y = mres)) +
  geom_point() +
  geom_smooth(method = "loess", aes(col = "loess")) +
  geom_smooth(method = 'lm', aes(col = "lm")) +
  theme_classic() +
  xlab("Donor age") +
  ylab("Martingale Residuals") +
  guides(col=guide_legend(title = ""))
```



Figure 6.22.: Martingale residuals for Donor age

A more complex functional form for donor age seems warranted; left as an exercise for the reader.

Now we will add the time-varying covariates:

```
# add counting process formulation of Surv():
bmt2 =
  bmt2 |>
  mutate(
    surv =
    Surv(
       time = tstart,
       time2 = tstop,
```

```
event = status,
type = "counting"))
```

Let's see how the data looks for patient 15:

```
bmt1 |> dplyr::filter(id == 15) |> dplyr::select(tp, dp, tc,dc, ta, da, FAB, surv, t
#> # A tibble: 1 x 13
           dp tc
                              da FAB
      tp
                                       surv
                                              t1
                                                   d1
                                                        t2
                     dc
                          ta
   21 1 220 1
                        418 0 Other
                                        418
                                            418
                                                   1
                                                       418
bmt2 |> dplyr::filter(id == 15) |> dplyr::select(id, agvhd, cgvhd, precovery, surv)
#> # A tibble: 3 x 5
#>
      id agvhd cgvhd precovery
                               surv
   <int> <int> <int>
                     <int>
                              <Surv>
                         0 ( 0, 21+]
#> 1
      15
            0
                0
                         1 (21,220+]
#> 2
      15
            0
                 0
#> 3
      15
                         1 (220,418]
```

6.12.6. Model with Time-Dependent Covariates

```
bmt_coxph_TV = coxph(
  formula =
    surv ~
    group + `patient age`*`donor age` + FAB + agvhd + cgvhd + precovery,
  data = bmt2)

summary(bmt_coxph_TV)

#> Call:
#> coxph(formula = surv ~ group + `patient age` * `donor age` +
#> FAB + agvhd + cgvhd + precovery, data = bmt2)
#>
```

```
n= 341, number of events= 83
#>
#>
#>
                                  coef exp(coef)
                                                  se(coef)
                                                               z Pr(>|z|)
#> groupLow Risk AML
                             -1.038514 0.353980
                                                  0.358220 - 2.90
                                                                   0.0037 **
#> groupHigh Risk AML
                             -0.380481 0.683533
                                                  0.374867 - 1.01
                                                                   0.3101
#> `patient age`
                             -0.073351 0.929275
                                                  0.035956 - 2.04
                                                                   0.0413 *
#> `donor age`
                             -0.076406 0.926440 0.030196 -2.53
                                                                   0.0114 *
#> FABGrade 4 Or 5 (AML only) 0.805700 2.238263 0.284273 2.83
                                                                   0.0046 **
#> agvhd
                              0.150565 1.162491 0.306848 0.49
                                                                   0.6237
#> cgvhd
                             -0.116136  0.890354  0.289046  -0.40
                                                                   0.6878
                             -0.941123 0.390190 0.347861 -2.71
                                                                   0.0068 **
#> precovery
#> `patient age`: `donor age` 0.002895 1.002899 0.000944 3.07
                                                                   0.0022 **
#> Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
#>
                             exp(coef) exp(-coef) lower .95 upper .95
#>
#> groupLow Risk AML
                                 0.354
                                            2.825
                                                     0.175
                                                                0.714
#> groupHigh Risk AML
                                 0.684
                                            1.463
                                                      0.328
                                                                1.425
#> `patient age`
                                 0.929
                                            1.076
                                                     0.866
                                                                0.997
#> `donor age`
                                 0.926
                                                                0.983
                                            1.079
                                                     0.873
#> FABGrade 4 Or 5 (AML only)
                                 2.238
                                            0.447
                                                    1.282
                                                                3.907
#> agvhd
                                 1.162
                                            0.860
                                                     0.637
                                                                2.121
#> cgvhd
                                 0.890
                                                      0.505
                                                                1.569
                                            1.123
#> precovery
                                 0.390
                                            2.563
                                                      0.197
                                                                0.772
#> `patient age`: `donor age`
                                 1.003
                                            0.997
                                                      1.001
                                                                1.005
#>
#> Concordance= 0.702 (se = 0.028 )
#> Likelihood ratio test= 40.3 on 9 df,
                                          p = 7e - 06
#> Wald test
                       = 42.4 on 9 df,
                                          p = 3e - 06
#> Score (logrank) test = 47.2 on 9 df,
                                          p=4e-07
```

Platelet recovery is highly significant.

Neither acute GVHD (agvhd) nor chronic GVHD (cgvhd) has a statisti-

cally significant effect here, nor are they significant in models with the other one removed.

```
update(bmt_coxph_TV, .~.-agvhd) |> summary()
#> Call:
#> coxph(formula = surv ~ group + `patient age` + `donor age` +
      FAB + cgvhd + precovery + `patient age`: `donor age`, data = bmt2)
#>
#>
    n= 341, number of events= 83
#>
#>
                                  coef exp(coef) se(coef)
                                                              z Pr(>|z|)
#> groupLow Risk AML
                             -1.049870 0.349983 0.356727 -2.94
                                                                   0.0032 **
#> groupHigh Risk AML
                             -0.417049 0.658988 0.365348 -1.14
                                                                   0.2537
#> `patient age`
                             -0.070749 0.931696 0.035477 -1.99
                                                                   0.0461 *
#> `donor age`
                             -0.075693 0.927101 0.030075 -2.52
                                                                  0.0118 *
#> FABGrade 4 Or 5 (AML only) 0.807035 2.241253 0.283437 2.85
                                                                   0.0044 **
#> cgvhd
                             -0.095393 0.909015 0.285979 -0.33
                                                                  0.7387
#> precovery
                             -0.983653 0.373942 0.338170 -2.91
                                                                   0.0036 **
#> `patient age`: `donor age` 0.002859 1.002863 0.000936 3.05
                                                                  0.0023 **
#> ---
#> Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
#>
#>
                             exp(coef) exp(-coef) lower .95 upper .95
#> groupLow Risk AML
                                 0.350
                                            2.857
                                                     0.174
                                                                0.704
#> groupHigh Risk AML
                                                     0.322
                                 0.659
                                            1.517
                                                               1.349
#> `patient age`
                                 0.932
                                            1.073
                                                     0.869
                                                               0.999
#> `donor age`
                                 0.927
                                            1.079
                                                    0.874
                                                               0.983
#> FABGrade 4 Or 5 (AML only)
                                 2.241
                                            0.446
                                                    1.286
                                                               3.906
#> cgvhd
                                 0.909
                                            1.100
                                                    0.519
                                                                1.592
#> precovery
                                 0.374
                                            2.674
                                                    0.193
                                                               0.726
#> `patient age`: `donor age`
                                 1.003
                                            0.997
                                                     1.001
                                                               1.005
#> Concordance= 0.701 (se = 0.027 )
```

```
#> Likelihood ratio test= 40 on 8 df, p=3e-06
\#> Wald test = 42.4 on 8 df, p=1e-06
\# Score (logrank) test = 47.2 on 8 df, p=1e-07
update(bmt_coxph_TV, .~.-cgvhd) |> summary()
#> Call:
#> coxph(formula = surv ~ group + `patient age` + `donor age` +
      FAB + agvhd + precovery + `patient age`: `donor age`, data = bmt2)
#>
   n= 341, number of events= 83
#>
#>
#>
                                 coef exp(coef) se(coef) z Pr(>|z|)
#> groupLow Risk AML
                            -1.019638 0.360725 0.355311 -2.87
                                                                0.0041 **
                            -0.381356 0.682935 0.374568 -1.02
#> groupHigh Risk AML
                                                                0.3086
#> `patient age`
                            -0.073189 0.929426 0.035890 -2.04
                                                                0.0414 *
#> `donor age`
                            -0.076753 0.926118 0.030121 -2.55
                                                                0.0108 *
#> FABGrade 4 Or 5 (AML only) 0.811716 2.251769 0.284012 2.86
                                                                0.0043 **
#> agvhd
                            0.131621 1.140676 0.302623 0.43
                                                                0.6636
#> precovery
                            -0.946697 0.388021 0.347265 -2.73
                                                                0.0064 **
#> `patient age`: `donor age`     0.002904     1.002908     0.000943     3.08
                                                                0.0021 **
#> ---
#> Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
#>
#>
                            exp(coef) exp(-coef) lower .95 upper .95
#> groupLow Risk AML
                                0.361
                                          2.772 0.180
                                                             0.724
#> groupHigh Risk AML
                                0.683
                                          1.464
                                                  0.328
                                                             1.423
                                0.929
#> `patient age`
                                          1.076
                                                  0.866
                                                             0.997
#> `donor age`
                                0.926
                                          1.080
                                                  0.873
                                                             0.982
#> FABGrade 4 Or 5 (AML only)
                                2.252
                                          0.444
                                                  1.291
                                                             3.929
#> agvhd
                                1.141
                                          0.877
                                                  0.630
                                                             2.064
#> precovery
                                0.388
                                          2.577
                                                  0.196
                                                             0.766
#> `patient age`: `donor age`
                                                           1.005
                               1.003
                                          0.997
                                                  1.001
#> Concordance= 0.701 (se = 0.027 )
```

```
#> Likelihood ratio test= 40.1 on 8 df, p=3e-06
#> Wald test = 42.1 on 8 df, p=1e-06
#> Score (logrank) test = 47.1 on 8 df, p=1e-07
```

Let's drop them both:

```
bmt_coxph_TV2 = update(bmt_coxph_TV, . ~ . - agvhd -cgvhd)
bmt_coxph_TV2 |> summary()
#> Call:
#> coxph(formula = surv ~ group + `patient age` + `donor age` +
      FAB + precovery + `patient age`: `donor age`, data = bmt2)
#>
   n= 341, number of events= 83
#>
#>
#>
                                 coef exp(coef) se(coef)
                                                             z Pr(>|z|)
#> groupLow Risk AML
                            -1.032520 0.356108 0.353202 -2.92
                                                                 0.0035 **
#> groupHigh Risk AML
                            -0.413888 0.661075 0.365209 -1.13
                                                                 0.2571
#> `patient age`
                            -0.070965 0.931495 0.035453 -2.00
                                                                 0.0453 *
#> `donor age`
                            -0.076052 0.926768 0.030007 -2.53
                                                                 0.0113 *
#> FABGrade 4 Or 5 (AML only) 0.811926 2.252242 0.283231 2.87
                                                                 0.0041 **
                            -0.983505 0.373998 0.337997 -2.91
                                                                 0.0036 **
#> precovery
#> `patient age`: `donor age` 0.002872 1.002876 0.000936 3.07
                                                                 0.0021 **
#> ---
#> Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
#>
                            exp(coef) exp(-coef) lower .95 upper .95
#> groupLow Risk AML
                                0.356
                                           2.808
                                                    0.178
                                                              0.712
#> groupHigh Risk AML
                                0.661
                                           1.513
                                                    0.323
                                                              1.352
#> `patient age`
                                0.931
                                           1.074
                                                   0.869
                                                              0.999
#> `donor age`
                                           1.079
                                0.927
                                                   0.874
                                                              0.983
#> FABGrade 4 Or 5 (AML only)
                                2.252
                                           0.444
                                                   1.293
                                                              3.924
#> precovery
                                0.374
                                           2.674
                                                   0.193
                                                              0.725
#> `patient age`: `donor age`
                                1.003
                                           0.997
                                                   1.001
                                                              1.005
```

6.13. Recurrent Events

(Adapted from Kleinbaum and Klein (2012), Ch 8)

- Sometimes an appropriate analysis requires consideration of recurrent events.
- A patient with arthritis may have more than one flareup. The same is true of many recurring-remitting diseases.
- In this case, we have more than one line in the data frame, but each line may have an event.
- We have to use a "robust" variance estimator to account for correlation of time-to-events within a patient.

6.13.1. Bladder Cancer Data Set

The bladder cancer dataset from Kleinbaum and Klein (2012) contains recurrent event outcome information for eighty-six cancer patients followed for the recurrence of bladder cancer tumor after transurethral surgical excision (Byar and Green 1980). The exposure of interest is the effect of the drug treatment of thiotepa. Control variables are the initial number and initial size of tumors. The data layout is suitable for a counting processes approach.

This drug is still a possible choice for some patients. Another therapeutic choice is Bacillus Calmette-Guerin (BCG), a live bacterium related to cow tuberculosis.

6.13.1.1. Data dictionary

Table 6.3.: Variables in the bladder dataset

| Variable | Definition |
|----------|---|
| id | Patient unique ID |
| status | for each time interval: $1 = \text{recurred}$, $0 = \text{censored}$ |
| interval | 1 = first recurrence, etc. |
| intime | 'tstop - tstart (all times in months) |
| tstart | start of interval |
| tstop | end of interval |
| tx | treatment code, $1 = \text{thiotepa}$ |
| num | number of initial tumors |
| size | size of initial tumors (cm) |

- There are 85 patients and 190 lines in the dataset, meaning that many patients have more than one line.
- Patient 1 with 0 observation time was removed.
- Of the 85 patients, 47 had at least one recurrence and 38 had none.
- 18 patients had exactly one recurrence.
- There were up to 4 recurrences in a patient.
- Of the 190 intervals, 112 terminated with a recurrence and 78 were censored.

6.13.1.2. Different intervals for the same patient are correlated.

- Is the effective sample size 47 or 112? This might narrow confidence intervals by as much as a factor of $\sqrt{112/47} = 1.54$
- What happens if I have 5 treatment and 5 control values and want to do a t-test and I then duplicate the 10 values as if the sample size was 20? This falsely narrows confidence intervals by a factor of $\sqrt{2} = 1.41$.

```
bladder =
  paste0(
    "http://web1.sph.emory.edu/dkleinb/allDatasets",
    "/surv2datasets/bladder.dta") |>
  read_dta() |>
  as_tibble()
bladder = bladder[-1,] #remove subject with 0 observation time
print(bladder)
bladder =
  bladder |>
  mutate(
    surv =
     Surv(
        time = start,
        time2 = stop,
        event = event,
        type = "counting"))
bladder.cox1 = coxph(
 formula = surv~tx+num+size,
  data = bladder)
#results with biased variance-covariance matrix:
summary(bladder.cox1)
#> Call:
#> coxph(formula = surv ~ tx + num + size, data = bladder)
#>
#>
    n= 190, number of events= 112
#>
```

coef exp(coef) se(coef) z Pr(>|z|)

#>

```
-0.4116 0.6626
                         0.1999 -2.06 0.03947 *
#> tx
                 1.1778
#> num 0.1637
                         0.0478 3.43 0.00061 ***
#> size -0.0411
                 #> Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
#>
#>
       exp(coef) exp(-coef) lower .95 upper .95
          0.663
                    1.509
                              0.448
                                        0.98
#> tx
           1.178
                     0.849
                              1.073
#> num
                                        1.29
#> size
           0.960
                     1.042
                              0.836
                                        1.10
#>
#> Concordance= 0.624 (se = 0.032 )
#> Likelihood ratio test= 14.7 on 3 df,
                                       p=0.002
#> Wald test
                      = 15.9 \text{ on } 3 \text{ df},
                                       p=0.001
#> Score (logrank) test = 16.2 on 3 df,
                                       p=0.001
```

Note

The likelihood ratio and score tests assume independence of observations within a cluster. The Wald and robust score tests do not.

6.13.1.3. adding cluster = id

If we add cluster= id to the call to coxph, the coefficient estimates don't change, but we get an additional column in the summary() output: robust se:

```
bladder.cox2 = coxph(
  formula = surv ~ tx + num + size,
  cluster = id,
  data = bladder)
```

```
#unbiased though this reduces power:
summary(bladder.cox2)
#> Call:
#> coxph(formula = surv ~ tx + num + size, data = bladder, cluster = id)
#>
#>
    n= 190, number of events= 112
#>
#>
          coef exp(coef) se(coef) robust se
                                                z Pr(>|z|)
       -0.4116
                0.6626
                           0.1999
                                     0.2488 -1.65
                                                    0.0980 .
#> tx
#> num 0.1637
                  1.1778
                           0.0478
                                     0.0584 2.80
                                                  0.0051 **
#> size -0.0411 0.9598
                          0.0703
                                     0.0742 - 0.55
                                                    0.5799
#> Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
#>
       exp(coef) exp(-coef) lower .95 upper .95
           0.663
                      1.509
                                0.407
                                           1.08
#> tx
            1.178
#> num
                      0.849
                                1.050
                                           1.32
#> size
           0.960
                      1.042
                                0.830
                                          1.11
#> Concordance= 0.624 (se = 0.031)
#> Likelihood ratio test= 14.7 on 3 df,
                                         p=0.002
#> Wald test
                       = 11.2 on 3 df,
                                          p=0.01
#> Score (logrank) test = 16.2 on 3 df,
                                         p=0.001,
                                                     Robust = 10.8 p=0.01
#>
     (Note: the likelihood ratio and score tests assume independence of
#>
        observations within a cluster, the Wald and robust score tests do not).
```

robust se is larger than se, and accounts for the repeated observations from the same individuals:

```
round(bladder.cox2$naive.var, 4)
#> [,1] [,2] [,3]
```

```
#> [1,] 0.0400 -0.0014 0.0000
#> [2,] -0.0014 0.0023 0.0007
#> [3,] 0.0000 0.0007 0.0049
round(bladder.cox2$var, 4)

#> [,1] [,2] [,3]
#> [1,] 0.0619 -0.0026 -0.0004
#> [2,] -0.0026 0.0034 0.0013
#> [3,] -0.0004 0.0013 0.0055
```

These are the ratios of correct confidence intervals to naive ones:

```
with(bladder.cox2, diag(var)/diag(naive.var)) |> sqrt()
#> [1] 1.244 1.223 1.056
```

We might try dropping the non-significant size variable:

```
#remove non-significant size variable:
bladder.cox3 = bladder.cox2 |> update(. ~ . - size)
summary(bladder.cox3)
#> Call:
#> coxph(formula = surv ~ tx + num, data = bladder, cluster = id)
#>
#>
   n= 190, number of events= 112
#>
         coef exp(coef) se(coef) robust se
                                            z Pr(>|z|)
#> tx -0.4117 0.6625 0.2003
                                  0.2515 -1.64 0.1017
#> num 0.1700
                1.1853
                         0.0465
                                  0.0564 3.02 0.0026 **
#> Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
#>
      exp(coef) exp(-coef) lower .95 upper .95
#> tx 0.663 1.509 0.405 1.08
```

```
#> num
          1.185
                     0.844
                               1.061
                                          1.32
#>
#> Concordance= 0.623 (se = 0.031 )
#> Likelihood ratio test= 14.3 on 2 df,
                                          p=8e-04
                       = 10.2 on 2 df,
#> Wald test
                                          p=0.006
#> Score (logrank) test = 15.8 on 2 df,
                                          p=4e-04,
                                                     Robust = 10.6 p=0.005
#>
     (Note: the likelihood ratio and score tests assume independence of
       observations within a cluster, the Wald and robust score tests do not).
```

Ways to check PH assumption:

- cloglog
- schoenfeld residuals
- interaction with time

6.14. Age as the time scale

See Canchola et al. (2003).

Configuring R

Functions from these packages will be used throughout this document:

```
library(conflicted) # check for conflicting function definitions
# library(printr) # inserts help-file output into markdown output
library(rmarkdown) # Convert R Markdown documents into a variety of formats.
library(pander) # format tables for markdown
library(ggplot2) # graphics
library(ggeasy) # help with graphics
library(ggfortify) # help with graphics
library(dplyr) # manipulate data
library(tibble) # `tibble`s extend `data.frame`s
library(magrittr) # `%>%` and other additional piping tools
library(haven) # import Stata files
library(knitr) # format R output for markdown
library(tidyr) # Tools to help to create tidy data
library(plotly) # interactive graphics
library(dobson) # datasets from Dobson and Barnett 2018
library(parameters) # format model output tables for markdown
library(haven) # import Stata files
```

```
library(latex2exp) # use LaTeX in R code (for figures and tables)
library(fs) # filesystem path manipulations
library(survival) # survival analysis
library(survminer) # survival analysis graphics
library(KMsurv) # datasets from Klein and Moeschberger
library(parameters) # format model output tables for
library(webshot2) # convert interactive content to static for pdf
library(forcats) # functions for categorical variables ("factors")
library(stringr) # functions for dealing with strings
library(lubridate) # functions for dealing with dates and times
```

Here are some R settings I use in this document:

```
rm(list = ls()) # delete any data that's already loaded into R

conflicts_prefer(dplyr::filter)
ggplot2::theme_set(
    ggplot2::theme_bw() +
        # ggplot2::labs(col = "") +
        ggplot2::theme(
        legend.position = "bottom",
        text = ggplot2::element_text(size = 12, family = "serif")))

knitr::opts_chunk$set(message = FALSE)
options('digits' = 4)

panderOptions("big.mark", ",")
pander::panderOptions("table.emphasize.rownames", FALSE)
pander::panderOptions("table.split.table", Inf)
conflicts_prefer(dplyr::filter) # use the `filter()` function from dplyr() by defaul
legend_text_size = 9
```

7.1. Parametric Survival Models

7.1.1. Exponential Distribution

• The exponential distribution is the basic distribution for survival analysis.

$$\begin{split} f(t) &= \lambda e^{-\lambda t} \\ \log \left\{ f(t) \right\} &= \log \left\{ \lambda \right\} - \lambda t \\ F(t) &= 1 - e^{-\lambda t} \\ S(t) &= e^{-\lambda t} \\ H(t) &= \log \left\{ S(t) \right\} \\ &= -\lambda t \\ h(t) &= \lambda \\ \mathrm{E}(T) &= \lambda^{-1} \end{split}$$

7.1.2. Weibull Distribution


Using the Kalbfleisch and Prentice (2002) notation:

$$\begin{split} f(t) &= \lambda p (\lambda t)^{p-1} e^{-(\lambda t)^p} \\ F(t) &= 1 - e^{-(\lambda t)^p} \\ S(t) &= e^{-(\lambda t)^p} \\ h(t) &= \lambda p (\lambda t)^{p-1} \\ H(t) &= (\lambda t)^p \\ \log \left\{ H(t) \right\} &= p \log \left\{ \lambda t \right\} \\ &= p \log \left\{ \lambda \right\} + p \log \left\{ t \right\} \\ \mathrm{E}(T) &= \lambda^{-1} \cdot \Gamma \left(1 + \frac{1}{p} \right) \end{split}$$

Note

Recall from calculus:

- $\Gamma(t) \stackrel{\text{def}}{=} \int_{u=0}^{\infty} u^{t-1} e^{-u} du$
- $\Gamma(t) = (t-1)!$ for integers $t \in \mathbb{Z}$
- It is implemented by the gamma() function in R.



Here are some Weibull density functions, with $\lambda=1$ and p varying:

```
library(ggplot2)
lambda = 1
ggplot() +
  geom_function(
   aes(col = "0.25"),
  fun = \((x)\) dweibull(x, shape = 0.25, scale = 1/lambda)) +
```

```
geom_function(
  aes(col = "0.5"),
  fun = \(x) dweibull(x, shape = 0.5, scale = 1/lambda)) +
geom_function(
  aes(col = "1"),
  fun = \(x) dweibull(x, shape = 1, scale = 1/lambda)) +
geom_function(
  aes(col = "1.5"),
  fun = \(x) dweibull(x, shape = 1.5, scale = 1/lambda)) +
geom_function(
  aes(col = "2"),
  fun = \(x) dweibull(x, shape = 2, scale = 1/lambda)) +
geom_function(
  aes(col = "5"),
  fun = (x) dweibull(x, shape = 5, scale = 1/lambda)) +
theme_bw() +
xlim(0, 2.5) +
ylab("f(t)") +
theme(axis.title.y = element_text(angle=0)) +
theme(legend.position="bottom") +
guides(
  col =
    guide_legend(
      title = "p",
      label.theme =
        element text(
          size = 12)))
```



Figure 7.1.: Density functions for Weibull distribution

7.1.2.1. Properties of Weibull hazard functions

Theorem 7.1. If T has a Weibull distribution, then:

- When p = 1, the Weibull distribution simplifies to the exponential distribution
- When p > 1, the hazard is increasing: h'(t) > 0
- When p < 1, the hazard is decreasing: h'(t) < 0
- $\log\{H(t)\}$ is a straight line relative to $\log\{t\}$: $\log\{H(t)\} = p\log\{\lambda\} + p\log\{t\}$

Exercise 7.1. Prove Theorem 7.1.

The Weibull distribution provides more flexibility than the exponential. Figure 7.2 shows some Weibull hazard functions, with $\lambda=1$ and p varying:

```
library(ggplot2)
library(eha)
lambda = 1
ggplot() +
  geom_function(
    aes(col = "0.25"),
    fun = \(x) hweibull(x, shape = 0.25, scale = 1/lambda)) +
  geom_function(
    aes(col = "0.5"),
    fun = \(x) hweibull(x, shape = 0.5, scale = 1/lambda)) +
  geom_function(
    aes(col = "1"),
    fun = (x) hweibull(x, shape = 1, scale = 1/lambda)) +
  geom_function(
    aes(col = "1.5"),
    fun = \(x) hweibull(x, shape = 1.5, scale = 1/lambda)) +
  geom_function(
    aes(col = "2"),
    fun = \(x) hweibull(x, shape = 2, scale = 1/lambda)) +
  theme_bw() +
  xlim(0, 2.5) +
  ylab("h(t)") +
  theme(axis.title.y = element_text(angle=0)) +
  theme(legend.position="bottom") +
  guides(
    col =
```

```
guide_legend(
  title = "p",
  label.theme =
    element_text(
      size = 12)))
```



Figure 7.2.: Hazard functions for Weibull distribution

```
library(ggplot2)
lambda = 1

ggplot() +
  geom_function(
```

```
aes(col = "0.25"),
  fun = \(x) pweibull(lower = FALSE, x, shape = 0.25, scale = 1/lambda)) +
geom_function(
  aes(col = "0.5"),
  fun = \(x) pweibull(lower = FALSE, x, shape = 0.5, scale = 1/lambda)) +
geom_function(
  aes(col = "1"),
  fun = (x) pweibull(lower = FALSE, x, shape = 1, scale = 1/lambda)) +
geom_function(
  aes(col = "1.5"),
  fun = (x) pweibull(lower = FALSE, x, shape = 1.5, scale = 1/lambda)) +
geom_function(
  aes(col = "2"),
  fun = (x) pweibull(lower = FALSE, x, shape = 2, scale = 1/lambda)) +
theme_bw() +
xlim(0, 2.5) +
ylab("S(t)") +
theme(axis.title.y = element_text(angle=0)) +
theme(legend.position="bottom") +
guides(
  col =
    guide_legend(
     title = "p",
      label.theme =
        element_text(
          size = 12)))
```



Figure 7.3.: Survival functions for Weibull distribution

7.1.3. Exponential Regression

For each subject i, define a linear predictor:

$$\begin{split} \eta(x) &= \beta_0 + (\beta_1 x_1 + \dots + \beta_p x_p) \\ h(t|x) &= \exp \left\{ \eta(x) \right\} \\ h_0 &\stackrel{\text{def}}{=} h(t|0) \\ &= \exp \left\{ \eta(0) \right\} \\ &= \exp \left\{ \beta_0 + (\beta_1 \cdot 0 + \dots + \beta_p \cdot 0) \right\} \\ &= \exp \left\{ \beta_0 + 0 \right\} \\ &= \exp \left\{ \beta_0 \right\} \end{split}$$

We let the linear predictor have a constant term, and when there are no additional predictors the hazard is $\lambda = \exp{\{\beta_0\}}$. This has a log link as in a generalized linear model. Since the hazard does not depend on t, the hazards are (trivially) proportional.

7.1.4. Accelerated Failure Time

Previously, we assumed the hazards were proportional; that is, the covariates multiplied the baseline hazard function:

$$\begin{split} h(T=t|X=x) &\stackrel{\text{def}}{=} p(T=t|X=x, T \geq t) \\ &= h(t|X=0) \cdot \exp\left\{\eta(x)\right\} \\ &= h(t|X=0) \cdot \theta(x) \\ &= h_0(t) \cdot \theta(x) \end{split}$$

and correspondingly,

$$\begin{split} H(t|x) &= \theta(x) H_0(t) \\ S(t|x) &= \exp\left\{-H(t|x)\right\} \\ &= \exp\left\{-\theta(x) \cdot H_0(t)\right\} \\ &= \left(\exp\left\{-H_0(t)\right\}\right)^{\theta(x)} \\ &= \left(S_0(t)\right)^{\theta(x)} \end{split}$$

An alternative modeling assumption would be

$$S(t|X=x) = S_0(t \cdot \theta(x))$$

where $\theta(x)=\exp\{\eta(x)\}$, $\eta(x)=\beta_1x_1+\cdots+\beta_px_p$, and $S_0(t)=P(T\geq t|X=0)$ is the base survival function.

Then

$$\begin{split} E(T|X=x) &= \int_{t=0}^{\infty} S(t|x) dt \\ &= \int_{t=0}^{\infty} S_0(t \cdot \theta(x)) dt \\ &= \int_{u=0}^{\infty} S_0(u) du \cdot \theta(x)^{-1} \\ &= \theta(x)^{-1} \cdot \int_{u=0}^{\infty} S_0(u) du \\ &= \theta(x)^{-1} \cdot \mathrm{E}(T|X=0) \end{split}$$

So the mean of T given X=x is the baseline mean divided by $\theta(x)=\exp{\{\eta(x)\}}.$

This modeling strategy is called an accelerated failure time model, because covariates cause uniform acceleration (or slowing) of failure times.

Additionally:

$$H(t|x) = H_0(\theta(x) \cdot t)$$

$$h(t|x) = \theta(x) \cdot h_0(\theta(x) \cdot t)$$

If the base distribution is exponential with parameter λ then

$$\begin{split} S(t|x) &= \exp\left\{-\lambda \cdot t\theta(x)\right\} \\ &= \left[\exp\left\{-\lambda t\right\}\right]^{\theta(x)} \end{split}$$

which is an exponential model with base hazard multiplied by $\theta(x)$, which is also the proportional hazards model.

In terms of the log survival time $Y = \log \{T\}$ the model can be written as

$$Y = \alpha - \eta + W$$
$$\alpha = -\log \{\lambda\}$$

where W has the extreme value distribution. The estimated parameter λ is the intercept and the other coefficients are those of η , which will be the opposite sign of those for coxph.

For a Weibull distribution, the hazard function and the survival function are

$$h(t) = \lambda p(\lambda t)^{p-1}$$

$$S(t) = e^{-(\lambda t)^p}$$

We can construct a proportional hazards model by using a linear predictor η_i without constant term and letting $\theta_i = e^{\eta_i}$ we have

$$h(t) = \lambda p(\lambda t)^{p-1} \theta_i$$

A distribution with $h(t)=\lambda p(\lambda t)^{p-1}\theta_i$ is a Weibull distribution with parameters $\lambda^*=\lambda\theta_i^{1/p}$ and p so the survival function is

$$S^*(t) = e^{-(\lambda^* t)^p}$$

$$= e^{-(\lambda \theta^{1/p} t)^p}$$

$$= S(t\theta^{1/p})$$

so this is also an accelerated failure time model.

In terms of the log survival time $Y = \log\{T\}$ the model can be written as

$$Y = \alpha - \sigma \eta + \sigma W$$
$$\alpha = -\log \{\lambda\}$$
$$\sigma = 1/p$$

where W has the extreme value distribution. The estimated parameter λ is the intercept and the other coefficients are those of η , which will be the opposite sign of those for coxph.

These AFT models are log-linear, meaning that the linear predictor has a log link. The exponential and the Weibull are the only log-linear models that are simultaneously proportional hazards models. Other parametric distributions can be used for survival regression either as a proportional hazards model or as an accelerated failure time model.

7.1.5. Dataset: Leukemia treatments

Remission survival times on 42 leukemia patients, half on new treatment, half on standard treatment.

This is the same data as the drug6mp data from KMsurv, but with two other variables and without the pairing.

```
library(haven)
library(survival)
anderson =
  paste0(
    "http://web1.sph.emory.edu/dkleinb/allDatasets",
    "/surv2datasets/anderson.dta") |>
  read_dta() |>
  mutate(
    status = status |>
        case_match(
        1 ~ "relapse",
```

```
0 ~ "censored"
),
sex = sex |>
    case_match(
    0 ~ "female",
    1 ~ "male"
),

rx = rx |>
    case_match(
    0 ~ "new",
    1 ~ "standard"
),

surv = Surv(time = survt, event = (status == "relapse"))
)
print(anderson)
```

7.1.5.1. Cox semi-parametric model

```
anderson.cox0 = coxph(
  formula = surv ~ rx,
  data = anderson)
summary(anderson.cox0)
#> Call:
#> coxph(formula = surv ~ rx, data = anderson)
#>
#> n= 42, number of events= 30
#>
#> coef exp(coef) se(coef) z Pr(>|z|)
```

7.1.5.2. Weibull parametric model

```
anderson.weib <- survreg(</pre>
formula = surv ~ rx,
 data = anderson,
 dist = "weibull")
summary(anderson.weib)
#>
#> Call:
#> survreg(formula = surv ~ rx, data = anderson, dist = "weibull")
       Value Std. Error z p
#> (Intercept) 3.516 0.252 13.96 < 2e-16
#> rxstandard -1.267
                        0.311 -4.08 4.5e-05
#> Log(scale) -0.312 0.147 -2.12 0.034
#> Scale= 0.732
#>
#> Weibull distribution
#> Loglik(model) = -106.6 Loglik(intercept only) = -116.4
```

```
#> Chisq= 19.65 on 1 degrees of freedom, p= 9.3e-06
#> Number of Newton-Raphson Iterations: 5
#> n= 42
```


7.1.5.3. Exponential parametric model

```
anderson.exp <- survreg(</pre>
 formula = surv ~ rx,
 data = anderson,
 dist = "exp")
summary(anderson.exp)
#>
#> Call:
#> survreg(formula = surv ~ rx, data = anderson, dist = "exp")
              Value Std. Error z p
#> (Intercept) 3.686 0.333 11.06 < 2e-16
#> rxstandard -1.527 0.398 -3.83 0.00013
#> Scale fixed at 1
#>
#> Exponential distribution
#> Loglik(model)= -108.5 Loglik(intercept only)= -116.8
\# Chisq= 16.49 on 1 degrees of freedom, p= 4.9e-05
#> Number of Newton-Raphson Iterations: 4
\#> n= 42
```

7.1.5.4. Diagnostic - complementary log-log survival plot

```
library(survminer)
survfit(
```

```
formula = surv ~ rx,
data = anderson) |>
ggsurvplot(fun = "cloglog")
```



If the cloglog plot is linear, then a Weibull model may be ok.

7.2. Combining left-truncation and interval-censoring

 $From \ [https://stat.ethz.ch/pipermail/r-help/2015-August/431733.html]: \\$

coxph does left truncation but not left (or interval) censoring survreg does interval censoring but not left truncation (or time dependent covariates).

• Terry Therneau, August 31, 2015

- Agresti, Alan. 2012. Categorical Data Analysis. Vol. 792. John Wiley & Sons. https://www.wiley.com/en-us/Categorical+Data+Analysis% 2C+3rd+Edition-p-9780470463635.
- ———. 2015. Foundations of Linear and Generalized Linear Models. John Wiley & Sons. https://www.wiley.com/en-us/Foundations+of+Linear+and+Generalized+Linear+Models-p-9781118730034.
- ——. 2018. An Introduction to Categorical Data Analysis. John Wiley & Sons. https://www.wiley.com/en-us/An+Introduction+to+Categorical+Data+Analysis%2C+3rd+Edition-p-9781119405283.
- Anderson, Edgar. 1935. "The Irises of the Gaspe Peninsula." Bulletin of American Iris Society 59: 2–5.
- Bache, Stefan Milton, and Hadley Wickham. 2022. Magrittr: A Forward-Pipe Operator for r. https://CRAN.R-project.org/package=magrittr.
- Banerjee, Sudipto, and Anindya Roy. 2014. Linear Algebra and Matrix Analysis for Statistics. Vol. 181. Crc Press Boca Raton. https://www.routledge.com/Linear-Algebra-and-Matrix-Analysis-for-Statistics/Banerjee-Roy/p/book/9781420095388.
- Canchola, Alison J, Susan L Stewart, Leslie Bernstein, Dee W West, Ronald K Ross, Dennis Deapen, Richard Pinder, et al. 2003. "Cox Regression Using Different Time-Scales." Western Users of SAS Software. https://www.lexjansen.com/wuss/2003/DataAnalysis/i-cox_time_scales.pdf.
- Casella, George, and Roger Berger. 2002. Statistical Inference. 2nd ed. Cengage Learning. https://www.cengage.com/c/statistical-inference-2e-casella-berger/9780534243128/.

- Chang, Winston. 2024. R Graphics Cookbook: Practical Recipes for Visualizing Data. O'Reilly Media. https://r-graphics.org/.
- Chatterjee, Samprit, and Ali S Hadi. 2015. Regression Analysis by Example. John Wiley & Sons. https://www.wiley.com/en-us/Regression+Analysis+by+Example%2C+4th+Edition-p-9780470055458.
- Clayton, David, and Michael Hills. 2013. Statistical Models in Epidemiology. Oxford University Press. https://global.oup.com/academic/product/statistical-models-in-epidemiology-9780199671182.
- Copelan, Edward A, James C Biggs, James M Thompson, Pamela Crilley, Jeff Szer, John P Klein, Neena Kapoor, Belinda R Avalos, Isabel Cunningham, and Kerry Atkinson. 1991. "Treatment for Acute Myelocytic Leukemia with Allogeneic Bone Marrow Transplantation Following Preparation with BuCy2." https://doi.org/10.1182/blood.V78.3.838.838.
- Dalgaard, Peter. 2008. *Introductory Statistics with r.* New York, NY: Springer New York. https://link.springer.com/book/10.1007/978-0-387-79054-1.
- Dobson, Annette J, and Adrian G Barnett. 2018. An Introduction to Generalized Linear Models. 4th ed. CRC press. https://doi.org/10.1201/9781315182780.
- Dunn, Peter K, Gordon K Smyth, et al. 2018. Generalized Linear Models with Examples in r. Vol. 53. Springer. https://link.springer.com/book/10.1007/978-1-4419-0118-7.
- Efron, Bradley, and David V Hinkley. 1978. "Assessing the Accuracy of the Maximum Likelihood Estimator: Observed Versus Expected Fisher Information." *Biometrika* 65 (3): 457–83.
- Faraway, Julian J. 2016. Extending the Linear Model with r: Generalized Linear, Mixed Effects and Nonparametric Regression Models. 2nd ed. Chapman; Hall/CRC. https://doi.org/10.1201/9781315382722.
- Fay, Colin, Sébastien Rochette, Vincent Guyader, and Cervan Girard. 2021. Engineering Production-Grade Shiny Apps. Chapman; Hall/CRC. https://engineering-shiny.org/.
- Fieller, Nick. 2016. Basics of Matrix Algebra for Statistics with r. Chapman; Hall/CRC. https://doi.org/10.1201/9781315370200.

- Fox, John. 2015. Applied Regression Analysis and Generalized Linear Models. Sage publications.
- Grambsch, Patricia M, and Terry M Therneau. 1994. "Proportional Hazards Tests and Diagnostics Based on Weighted Residuals." *Biometrika* 81 (3): 515–26. https://doi.org/10.1093/biomet/81.3.515.
- Harrell, Frank E. 2015. Regression Modeling Strategies: With Applications to Linear Models, Logistic Regression, and Survival Analysis. 2nd ed. Springer. https://doi.org/10.1007/978-3-319-19425-7.
- Hosmer, David W, Stanley Lemeshow, and Rodney X Sturdivant. 2013. Applied Logistic Regression. John Wiley & Sons. https://onlinelibrary.wiley.com/doi/book/10.1002/9781118548387.
- James, Gareth, Daniela Witten, Trevor Hastie, Robert Tibshirani, et al. 2013. An Introduction to Statistical Learning. Vol. 112. Springer. https://www.statlearning.com/.
- Jewell, Nicholas P. 2003. Statistics for Epidemiology. Oxford, UK: Chapman; Hall/CRC. https://www.routledge.com/Statistics-for-Epidemiology/Jewell/p/book/9781584884330.
- Kalbfleisch, John D, and Ross L Prentice. 2011. The Statistical Analysis of Failure Time Data. John Wiley & Sons.
- Khuri, André I. 2003. Advanced Calculus with Applications in Statistics. John Wiley & Sons. https://www.wiley.com/en-us/Advanced+Calculus+with+Applications+in+Statistics%2C+2nd+Edition-p-9780471391043.
- Klein, John P, and Melvin L Moeschberger. 2003. Survival Analysis: Techniques for Censored and Truncated Data. Vol. 1230. Springer. https://link.springer.com/book/10.1007/b97377.
- Kleinbaum, David G, and Mitchel Klein. 2010. Logistic Regression. 3rd ed. Springer. https://link.springer.com/book/10.1007/978-1-4419-1742-3.
- ——. 2012. Survival Analysis a Self-Learning Text. 3rd ed. Springer. https://link.springer.com/book/10.1007/978-1-4419-6646-9.
- Kleinbaum, David G, Lawrence L Kupper, Azhar Nizam, K Muller, and ES Rosenberg. 2014. Applied Regression Analysis and Other Multivariable Methods. 5th ed. Cengage Learning. https://www.

- cengage.com/c/applied-regression-analysis-and-other-multivariable-methods-5e-kleinbaum/9781285051086/.
- Kleinman, Ken, and Nicholas J Horton. 2009. SAS and r: Data Management, Statistical Analysis, and Graphics. Chapman; Hall/CRC. https://www.routledge.com/SAS-and-R-Data-Management-Statistical-Analysis-and-Graphics-Second-Edition/Kleinman-Horton/p/book/9781466584495.
- Kuhn, Max, and Julia Silge. 2022. *Tidy Modeling with r.* "O'Reilly Media, Inc.". https://www.tmwr.org/.
- Kutner, Michael H, Christopher J Nachtsheim, John Neter, and William Li. 2005. Applied Linear Statistical Models. McGraw-Hill.
- Lawrance, Rachael, Evgeny Degtyarev, Philip Griffiths, Peter Trask, Helen Lau, Denise D'Alessio, Ingolf Griebsch, Gudrun Wallenstein, Kim Cocks, and Kaspar Rufibach. 2020. "What Is an Estimand, and How Does It Relate to Quantifying the Effect of Treatment on Patient-Reported Quality of Life Outcomes in Clinical Trials?" Journal of Patient-Reported Outcomes 4 (1): 1–8. https://link.springer.com/article/10.1186/s41687-020-00218-5.
- McCullagh, Peter, and J. A. Nelder. 1989. Generalized Linear Models. 2nd ed. Routledge. https://www.utstat.toronto.edu/~brunner/oldclass/2201s11/readings/glmbook.pdf.
- McLachlan, Geoffrey J, and Thriyambakam Krishnan. 2007. The EM Algorithm and Extensions. 2nd ed. John Wiley & Sons. https://doi.org/10.1002/9780470191613.
- Moore, Dirk F. 2016. Applied Survival Analysis Using r. Vol. 473. Springer. https://doi.org/10.1007/978-3-319-31245-3.
- Nahhas, Ramzi W. 2023. An Introduction to r for Research. https://bookdown.org/rwnahhas/IntroToR/.
- Pebesma, Edzer, and Roger Bivand. 2023. Spatial Data Science: With Applications in R. Boca Raton: Chapman; Hall/CRC. https://doi.org/10.1201/9780429459016.
- Pohl, Moritz, Lukas Baumann, Rouven Behnisch, Marietta Kirchner, Jo-

- hannes Krisam, and Anja Sander. 2021. "Estimands—A Basic Element for Clinical Trials." *Deutsches Ärzteblatt International* 118 (51-52): 883–88. https://doi.org/10.3238/arztebl.m2021.0373.
- Polin, Richard A, William W Fox, and Steven H Abman. 2011. Fetal and Neonatal Physiology. 4th ed. Elsevier health sciences.
- Rosenman, Ray H, Richard J Brand, C David Jenkins, Meyer Friedman, Reuben Straus, and Moses Wurm. 1975. "Coronary Heart Disease in the Western Collaborative Group Study: Final Follow-up Experience of 8 1/2 Years." *JAMA* 233 (8): 872–77. https://doi.org/10.1001/jama.1975.03260080034016.
- Searle, Shayle R, and Andre I Khuri. 2017. Matrix Algebra Useful for Statistics. John Wiley & Sons.
- Seber, George AF, and Alan J Lee. 2012. Linear Regression Analysis. 2nd ed. John Wiley & Sons. https://www.wiley.com/en-us/Linear+Regression+Analysis%2C+2nd+Edition-p-9781118274422.
- Selvin, Steve. 2001. Epidemiologic Analysis: A Case-Oriented Approach. Oxford University Press.
- Van Buuren, Stef. 2018. Flexible Imputation of Missing Data. CRC press. https://stefvanbuuren.name/fimd/.
- Venables, Bill. 2023. codingMatrices: Alternative Factor Coding Matrices for Linear Model Formulae. https://CRAN.R-project.org/package=codingMatrices.
- Vittinghoff, Eric, David V Glidden, Stephen C Shiboski, and Charles E McCulloch. 2012. Regression Methods in Biostatistics: Linear, Logistic, Survival, and Repeated Measures Models. 2nd ed. Springer. https://doi.org/10.1007/978-1-4614-1353-0.
- Wickham, Hadley. 2019. Advanced r. Chapman; Hall/CRC. https://adv-r.hadley.nz/index.html.
- ——. 2021. Mastering Shiny. "O'Reilly Media, Inc.". https://mastering-shiny.org/.
- Wickham, Hadley, Mara Averick, Jennifer Bryan, Winston Chang, Lucy D'Agostino McGowan, Romain François, Garrett Grolemund, et al. 2019. "Welcome to the tidyverse." Journal of Open Source Software 4 (43): 1686. https://doi.org/10.21105/joss.01686.

- Wickham, Hadley, and Jennifer Bryan. 2023. *R Packages*. "O'Reilly Media, Inc.". https://r-pkgs.org/.
- Wickham, Hadley, Mine Çetinkaya-Rundel, and Garrett Grolemund. 2023. R for Data Science. "O'Reilly Media, Inc.". https://r4ds.hadley.nz/.
- Woodward, Mark. 2013. Epidemiology: Study Design and Data Analysis. CRC press. https://www.routledge.com/Epidemiology-Study-Design-and-Data-Analysis-Third-Edition/Woodward/p/book/9781439839706.
- Zeileis, Achim, Christian Kleiber, and Simon Jackman. 2008. "Regression Models for Count Data in R." *Journal of Statistical Software* 27 (8). https://www.jstatsoft.org/v27/i08/.

These lecture notes use:

- algebra
- precalculus
- univariate calculus
- linear algebra
- vector calculus

Some key results are listed here.

A.1. Elementary Algebra

Mastery of Elementary Algebra¹ (a.k.a. "College Algebra") is a prerequisite for calculus, which is a prerequisite for Epi 202 and Epi 203, which are prerequisites for this course (Epi 204). Nevertheless, each year, some Epi 204 students are still uncomfortable with algebraic manipulations of mathematical formulas. Therefore, I include this section as a quick reference.

A.1.1. Equalities

Theorem A.1 (Equalities are transitive). If a = b and b = c, then a = c

¹https://en.wikipedia.org/wiki/Elementary_algebra

Theorem A.2 (Substituting equivalent expressions). If a=b, then for any function f(x), f(a)=f(b)

A.1.2. Inequalities

Theorem A.3. If a < b, then a + c < b + c

Theorem A.4 (negating both sides of an inequality). If a < b, then: -a > -b

Theorem A.5. If a < b and $c \ge 0$, then ca < cb.

Theorem A.6.

-a = (-1) * a

A.1.3. Sums

Theorem A.7 (adding zero changes nothing).

$$a + 0 = a$$

Theorem A.8 (Sums are symmetric).

$$a + b = b + a$$

Theorem A.9 (Sums are associative).

When summing three or more terms, the order in which you sum them does not matter:

$$(a+b) + c = a + (b+c)$$

A.1.4. Products

Theorem A.10 (Multiplying by 1 changes nothing).

$$a \times 1 = a$$

Theorem A.11 (Products are symmetric).

$$a \times b = b \times a$$

Theorem A.12 (Products are associative).

$$(a \times b) \times c = a \times (b \times c)$$

A.1.5. Sums and products together

Theorem A.13 (Multiplication is distributive).

$$a(b+c) = ab + ac$$

A.1.6. Quotients

Definition A.1 (Quotients, fractions, rates).

A quotient, fraction, or rate is a division of one quantity by another:

 $\frac{a}{h}$

In epidemiology, rates typically have a quantity involving time or population in the denominator.

c.f. https://en.wikipedia.org/wiki/Rate_(mathematics)

Definition A.2 (Ratios). A **ratio** is a quotient in which the numerator and denominator are measured using the same unit scales.

c.f. https://en.wikipedia.org/wiki/Ratio

Additional reference: https://en.wikipedia.org/wiki/Elementary_algebra

A.2. Exponentials and Logarithms

Theorem A.14 (logarithm of a product is the sum of the logs of the factors).

$$log\{a \cdot b\} = log\{a\} + log\{b\}$$

Corollary A.1 (logarithm of a quotient).

The logarithm of a quotient is equal to the difference of the logs of the factors:

$$\log\left\{\frac{a}{b}\right\} = \log\left\{a\right\} - \log\left\{b\right\}$$

Theorem A.15 (logarithm of an exponential function).

$$\log\left\{a^b\right\} = b \cdot \log\left\{a\right\}$$

Theorem A.16 (exponential of a sum).

The exponential of a sum is equal to the product of the exponentials of the addends:

$$exp\{a+b\} = exp\{a\} \cdot exp\{b\}$$

Corollary A.2 (exponential of a difference).

The exponential of a difference is equal to the quotient of the exponentials of the addends:

$$\exp\left\{a-b\right\} = \frac{\exp\left\{a\right\}}{\exp\left\{b\right\}}$$

Theorem A.17 (exponential of a product).

$$a^{bc} = \left(a^b\right)^c = \left(a^c\right)^b$$

Corollary A.3 (natural exponential of a product).

$$exp \{ab\} = (exp \{a\})^b = (exp \{b\})^a$$

Theorem A.18 ($\exp\{\}$ and $\log\{\}$ are mutual inverses).

$$exp\{log\{a\}\} = log\{exp\{a\}\} = a$$

A.3. Derivatives

Theorem A.19 (Derivatives of polynomials).

$$\frac{\partial}{\partial x}x^q = qx^{q-1}$$

Theorem A.20 (derivative of natural logarithm).

$$\log'\{x\} = \frac{1}{x} = x^{-1}$$

Theorem A.21 (derivative of exponential).

$$exp'\{x\} = exp\{x\}$$

Theorem A.22 (Product rule).

$$(ab)' = ab' + ba'$$

Theorem A.23 (Quotient rule).

$$(a/b)' = 1/b - (a/b^2)b'$$

Theorem A.24 (Chain rule).

$$\frac{\partial a}{\partial c} = \frac{\partial a}{\partial b} \frac{\partial b}{\partial c}$$

Corollary A.4 (Chain rule for logarithms).

$$\frac{\partial}{\partial x} \log \{f(x)\} = \frac{f'(x)}{f(x)}$$

Proof. Apply Theorem A.24 and Theorem A.20.

A.4. Vector Calculus

(adapted from Fieller (2016), §7.2)

Let \tilde{x} and $\tilde{\beta}$ be vectors of length p, or in other words, matrices of length $p \times 1$:

$$\tilde{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_p \end{bmatrix}$$

$$\tilde{\beta} = \begin{bmatrix} \beta_1 \\ \beta_2 \\ \vdots \\ \beta_p \end{bmatrix}$$

Definition A.3 (Transpose). The transpose of a row vector is the column vector with the same sequence of entries:

$$\tilde{x}' \equiv \tilde{x}^{\top} \equiv [x_1, x_2, ..., x_p]$$

Example A.1 (Dot product as matrix multiplication).

$$\begin{split} \tilde{x} \cdot \tilde{\beta} &= \tilde{x}^{\intercal} \tilde{\beta} \\ &= [x_1, x_2, ..., x_p] \begin{bmatrix} \beta_1 \\ \beta_2 \\ \vdots \\ \beta_p \end{bmatrix} \\ &= x_1 \beta_1 + x_2 \beta_2 + ... + x_p \beta_p \end{split}$$

Theorem A.25 (Transpose of a sum).

$$(\tilde{x} + \tilde{y})^\top = \tilde{x}^\top + \tilde{y}^\top$$

Definition A.4 (Vector derivative of a vector-to-scalar function). If $f(\tilde{\beta})$ is a function that takes a vector $\tilde{\beta}$ as input and outputs a scalar, such as $f(\tilde{\beta}) = x'\tilde{\beta}$, then:

$$\frac{\partial}{\partial \tilde{\beta}} f(\tilde{\beta}) = \begin{bmatrix} \frac{\partial}{\partial \beta_1} f(\tilde{\beta}) \\ \frac{\partial}{\partial \beta_2} f(\tilde{\beta}) \\ \vdots \\ \frac{\partial}{\partial \beta_p} f(\tilde{\beta}) \end{bmatrix}$$

Theorem A.26 (Derivative of a linear combination).

$$\frac{\partial}{\partial \tilde{\beta}} \tilde{x}^{\top} \tilde{\beta} = x$$

This looks a lot like non-vector calculus, except that you have to transpose the coefficient.

Proof.

$$\begin{split} \frac{\partial}{\partial \beta}(x^{\intercal}\beta) &= \begin{bmatrix} \frac{\partial}{\partial \beta_{1}}(x_{1}\beta_{1} + x_{2}\beta_{2} + \ldots + x_{p}\beta_{p}) \\ \frac{\partial}{\partial \beta_{2}}(x_{1}\beta_{1} + x_{2}\beta_{2} + \ldots + x_{p}\beta_{p}) \\ \vdots \\ \frac{\partial}{\partial \beta_{p}}(x_{1}\beta_{1} + x_{2}\beta_{2} + \ldots + x_{p}\beta_{p}) \end{bmatrix} \\ &= \begin{bmatrix} x_{1} \\ x_{2} \\ \vdots \\ x_{p} \end{bmatrix} \\ &= \tilde{x} \end{split}$$

Definition A.5 (Quadratic form). A $\mathbf{quadratic}$ form is a mathematical expression with the structure

 $\tilde{x}^{\top}\mathbf{S}\tilde{x}$

where \tilde{x} is a vector and \mathbf{S} is a matrix with compatible dimensions for vector-matrix multiplication.

Quadratic forms occur frequently in regression models. They are the matrix-vector generalizations of the scalar quadratic form $cx^2 = xcx$.

460

Theorem A.27 (Derivative of a quadratic form). If S is a $p \times p$ matrix that is constant with respect to β , then:

$$\frac{\partial}{\partial \beta} \beta' S \beta = 2S \beta$$

This is like taking the derivative of cx^2 with respect to x in non-vector calculus.

Corollary A.5 (Derivative of a simple quadratic form).

$$\frac{\partial}{\partial \tilde{\beta}} \tilde{\beta}' \tilde{\beta} = 2 \tilde{\beta}$$

This is like taking the derivative of x^2 .

A.5. Additional resources

A.5.1. Calculus

• Khuri (2003)

A.5.2. Linear Algebra and Vector Calculus

- Fieller (2016)
- Banerjee and Roy (2014)
- Searle and Khuri (2017)

A.5.3. Numerical Analysis

Methods for Biostatistics" $(2023 \text{ Fall})^3$

 $[\]frac{^2 https://hua-zhou.github.io/}{^3 https://ucla-biostat-216.github.io/2023fall/schedule/schedule.html}$

Configuring R

Functions from these packages will be used throughout this document:

```
library(conflicted) # check for conflicting function definitions
# library(printr) # inserts help-file output into markdown output
library(rmarkdown) # Convert R Markdown documents into a variety of formats.
library(pander) # format tables for markdown
library(ggplot2) # graphics
library(ggeasy) # help with graphics
library(ggfortify) # help with graphics
library(dplyr) # manipulate data
library(tibble) # `tibble`s extend `data.frame`s
library(magrittr) # `%>%` and other additional piping tools
library(haven) # import Stata files
library(knitr) # format R output for markdown
library(tidyr) # Tools to help to create tidy data
library(plotly) # interactive graphics
library(dobson) # datasets from Dobson and Barnett 2018
library(parameters) # format model output tables for markdown
library(haven) # import Stata files
library(latex2exp) # use LaTeX in R code (for figures and tables)
library(fs) # filesystem path manipulations
```

```
library(survival) # survival analysis
library(survminer) # survival analysis graphics
library(KMsurv) # datasets from Klein and Moeschberger
library(parameters) # format model output tables for
library(webshot2) # convert interactive content to static for pdf
library(forcats) # functions for categorical variables ("factors")
library(stringr) # functions for dealing with strings
library(lubridate) # functions for dealing with dates and times
```

Here are some R settings I use in this document:

```
rm(list = ls()) # delete any data that's already loaded into R

conflicts_prefer(dplyr::filter)
ggplot2::theme_set(
    ggplot2::theme_bw() +
        # ggplot2::labs(col = "") +
        ggplot2::theme(
        legend.position = "bottom",
        text = ggplot2::element_text(size = 12, family = "serif")))

knitr::opts_chunk$set(message = FALSE)
options('digits' = 4)

panderOptions("big.mark", ",")
pander::panderOptions("table.emphasize.rownames", FALSE)
pander::panderOptions("table.split.table", Inf)
conflicts_prefer(dplyr::filter) # use the `filter()` function from dplyr() by defaul legend_text_size = 9
```

Most of the content in this chapter should be review from UC Davis Epi 202.

B.1. Statistical events

Theorem B.1. If A and B are statistical events and $A \subseteq B$, then p(A, B) = p(A).

Proof. Left to the reader.

B.2. Random variables

B.2.1. Binary variables

Definition B.1 (binary variable). A **binary variable** is a random variable which has only two possible values in its range.

Exercise B.1 (Examples of binary variables). What are some examples of binary variables in the health sciences?

Solution. Examples of binary outcomes include:

- exposure (exposed vs unexposed)
- disease (diseased vs healthy)

- recovery (recovered vs unrecovered)
- relapse (relapse vs remission)
- return to hospital (returned vs not)
- vital status (dead vs alive)

B.2.2. Count variables

Definition B.2 (Count variable). A **count variable** is a random variable whose possible values are some subset of the non-negative integers; that is, a random variable X such that:

$$\mathcal{R}(X)\in\mathbb{N}$$

Exercise B.2. What are some examples of count variables?

Solution.

- Number of fish in a pond
- Number of cyclones per season
- Seconds of tooth-brushing per session (if rounded)¹
- Infections per person-year
- Visits to ER per person-month
- Car accidents per 1000 miles driven

¹https://pubmed.ncbi.nlm.nih.gov/35587489/

B.2.2.1. Exposure magnitude

Definition B.3 (Exposure magnitude). For many count outcomes, there is some sense of **exposure magnitude**, **population size**, or **duration of observation**.

Exercise B.3. What are some examples of exposure magnitudes?

Solution.

Table B.1.: Examples of exposure units

| outcome | exposure units | | | |
|---------------------------------------|---|--|--|--|
| disease incidence | number of individuals exposed; time at risk | | | |
| car accidents | miles driven | | | |
| worksite accidents population size | person-hours worked size of habitat | | | |

Exposure units are similar to the number of trials in a binomial distribution, but in non-binomial count outcomes, there can be more than one event per unit of exposure.

We can use t to represent continuous-valued exposures/observation durations, and n to represent discrete-valued exposures.

Definition B.4 (Event rate).

For a count outcome Y with exposure magnitude t, the **event rate** (denoted λ) is defined as the mean of Y divided by the exposure magnitude. That is:

$$\mu \stackrel{\text{def}}{=} \mathbb{E}[Y|T=t]$$

$$\lambda \stackrel{\text{def}}{=} \frac{\mu}{t} \tag{B.1}$$

Event rate is somewhat analogous to odds in binary outcome models; it typically serves as an intermediate transformation between the mean of the outcome and the linear component of the model. However, in contrast with the odds function, the transformation $\lambda = \mu/t$ is not considered part of the Poisson model's link function, and it treats the exposure magnitude covariate differently from the other covariates.

Theorem B.2 (Transformation function from event rate to mean). For a count variable with mean μ , event rate λ , and exposure magnitude t:

$$\therefore \mu = \lambda \cdot t \tag{B.2}$$

Solution. Start from definition of event rate and use algebra to solve for μ .

| Equation | B.2 | is | analogous | to | the | inverse-odds | function | for | binary | vari- |
|----------|-----|----|-----------|----|-----|--------------|----------|-----|--------|-------|
| ables. | | | | | | | | | | |
| | | | | | | | | | | |
| | | | | | | | | | | |

Theorem B.3. When the exposure magnitude is 0, there is no opportunity for events to occur:

$$\mathbb{E}[Y|T=0]=0$$

Proof.

$$\mathbb{E}[Y|T=0] = \lambda \cdot 0 = 0$$

B.2.2.2. Probability distributions for count outcomes

- Poisson distribution
- Negative binomial distribution

469

B.3. Key probability distributions

B.3.1. The Bernoulli distribution

Definition B.5 (Bernoulli distribution). The **Bernoulli distribution** family for a random variable X is defined as:

$$\begin{split} \Pr(X = x) &= \mathbb{1}_{x \in \{0,1\}} \pi^x (1 - \pi)^{1 - x} \\ &= \left\{ \begin{matrix} \pi, x = 1 \\ 1 - \pi, x = 0 \end{matrix} \right. \end{split}$$

B.3.2. The Poisson distribution



(a) Siméon Denis Poisson

Figure B.2.: "Les Poissons"

Definition B.6 (Poisson distribution).

$$\mathcal{R}(Y) = \{0, 1, 2, ...\} = \mathbb{N}$$

$$P(Y = y) = \frac{\mu^y e^{-\mu}}{y!}, y \in \mathbb{N}$$
(B.3)

(see Figure B.3)

$$P(Y \le y) = e^{-\mu} \sum_{j=0}^{\lfloor y \rfloor} \frac{\mu^j}{j!}$$
 (B.4)

(see Figure B.4)

```
library(dplyr)
pois_dists = tibble(
    mu = c(0.5, 1, 2, 5, 10, 20)) |>
    reframe(
        .by = mu,
        x = 0:30
) |>
    mutate(
        `P(X = x)` = dpois(x, lambda = mu),
        `P(X <= x)` = ppois(x, lambda = mu),
        mu = factor(mu)
)</pre>
```

```
library(ggplot2)
library(latex2exp)
plot0 = pois_dists |>
  ggplot(
   aes(
     x = x,
     y = P(X = x),
     fill = mu,
      col = mu)) +
  theme(legend.position = "bottom") +
  labs(
   fill = latex2exp::TeX("$\\mu$"),
   col = latex2exp::TeX("$\\mu$"),
    y = latex2exp::TeX("$\Pr_{\mu}(X = x)$"))
plot1 = plot0 +
 geom_col(position = "identity", alpha = .5) +
  facet_wrap(~mu)
 \# geom_point(alpha = 0.75) +
  \# geom_line(alpha = 0.75)
print(plot1)
```



Figure B.3.: Poisson PMFs, by mean parameter μ

```
plot2 =
  plot0 +
  geom_step(alpha = 0.75) +
  aes(y = `P(X <= x)`) +
  labs(y = latex2exp::TeX("$\\Pr_{\mu}(X \\leq x)$"))
print(plot2)</pre>
```



Figure B.4.: Poisson CDFs

Exercise B.4 (Poisson distribution functions). Let $X \sim \text{Pois}(\mu = 3.75)$.

Compute:

- $P(X = 4|\mu = 3.75)$
- $P(X \le 7 | \mu = 3.75)$ $P(X > 5 | \mu = 3.75)$

Solution.

•
$$P(X = 4) = 0.1938$$

•
$$P(X \le 7) = 0.9624$$

•
$$P(X > 5) = 0.1771$$

Theorem B.4 (Properties of the Poisson distribution). If $X \sim Pois(\mu)$, then:

•
$$\mathbb{E}[Y] = \mu$$

•
$$Var(Y) = \mu$$

Exercise B.5. Prove Theorem B.4.

Solution.

$$\begin{split} & \operatorname{E}[X] = \sum_{x=0}^{\infty} x \cdot P(X = x) \\ & = 0 \cdot P(X = 0) + \sum_{x=1}^{\infty} x \cdot P(X = x) \\ & = 0 + \sum_{x=1}^{\infty} x \cdot P(X = x) \\ & = \sum_{x=1}^{\infty} x \cdot \frac{\lambda^x e^{-\lambda}}{x!} \\ & = \sum_{x=1}^{\infty} x \cdot \frac{\lambda^x e^{-\lambda}}{x \cdot (x-1)!} & \text{[definition of factorial ("!") function]} \\ & = \sum_{x=1}^{\infty} \frac{\lambda^x e^{-\lambda}}{(x-1)!} & \\ & = \sum_{x=1}^{\infty} \frac{(\lambda \cdot \lambda^{x-1}) e^{-\lambda}}{(x-1)!} & \\ & = \lambda \cdot \sum_{x=1}^{\infty} \frac{(\lambda^y) e^{-\lambda}}{(y)!} & \text{[substituting } y \stackrel{\text{def}}{=} x - 1] \\ & = \lambda \cdot 1 & \text{[because PDFs sum to 1]} \\ & = \lambda \end{split}$$

B.3.2.1. Accounting for exposure

If the exposures/observation durations, denoted T = t or N = n, vary between observations, we model:

$$\mu = \lambda \cdot t$$

 λ is interpreted as the "expected event rate per unit of exposure"; that is,

$$\lambda = \frac{\mathbb{E}[Y|T=t]}{t}$$

! Important

The exposure magnitude, T, is *similar* to a covariate in linear or logistic regression. However, there is an important difference: in count regression, **there is no intercept corresponding to** $\mathbb{E}[Y|T=0]$. In other words, this model assumes that if there is no exposure, there can't be any events.

Theorem B.5. If $\mu = \lambda \cdot t$, then:

$$log\{\mu\} = log\{\lambda\} + log\{t\}$$

Definition B.7 (Offset). When the linear component of a model involves a term without an unknown coefficient, that term is called an **offset**.

B.3.3. The Negative-Binomial distribution

Definition B.8 (Negative binomial distribution).

$$P(Y = y) = \frac{\mu^y}{y!} \cdot \frac{\Gamma(\rho + y)}{\Gamma(\rho) \cdot (\rho + \mu)^y} \cdot \left(1 + \frac{\mu}{\rho}\right)^{-\rho}$$

where ρ is an overdispersion parameter and $\Gamma(x) = (x-1)!$ for integers x.

You don't need to memorize or understand this expression.

As $\rho \to \infty$, the second term converges to 1 and the third term converges to exp $\{-\mu\}$, which brings us back to the Poisson distribution.

Theorem B.6. If $Y \sim NegBin(\mu, \rho)$, then:

- $\begin{array}{ll} \bullet & \mathbb{E}[Y] = \mu \\ \bullet & Var(Y) = \mu + \frac{\mu^2}{\rho} > \mu \end{array}$

B.3.4. Weibull Distribution

$$p(t) = \alpha \lambda x^{\alpha - 1} e^{-\lambda x^{\alpha}}$$

$$h(t) = \alpha \lambda x^{\alpha - 1}$$

$$S(t) = e^{-\lambda x^{\alpha}}$$

$$E(T) = \Gamma(1+1/\alpha) \cdot \lambda^{-1/\alpha}$$

When $\alpha = 1$ this is the exponential. When $\alpha > 1$ the hazard is increasing and when $\alpha < 1$ the hazard is decreasing. This provides more flexibility than the exponential.

We will see more of this distribution later.

B.4. Characteristics of probability distributions

B.4.1. Probability density function

Definition B.9 (probability density). If X is a continuous random variable, then the **probability density** of X at value x, denoted f(x), $f_X(x)$, p(x), $p_X(x)$, or p(X=x), is defined as the limit of the probability (mass) that X is in an interval around x, divided by the width of that interval, as that width reduces to 0.

$$f(x) \stackrel{\text{def}}{=} \lim_{\delta \to 0} \frac{P(X \in [x, x + \delta])}{\delta}$$

See also https://en.wikipedia.org/wiki/Probability_density_function#Formal_definition

Theorem B.7 (Density function is derivative of CDF). The density function f(t) or p(T = t) for a random variable T at value t is equal to the derivative of the cumulative probability function $F(t) \stackrel{def}{=} P(T \le t)$; that is:

$$f(t) \stackrel{\textit{def}}{=} \frac{\partial}{\partial t} F(t)$$

Theorem B.8 (Density functions integrate to 1). For any density function f(x),

$$\int_{x \in \mathcal{R}(X)} f(x) dx = 1$$

B.4.2. Hazard function

Definition B.10 (Hazard function, hazard rate, hazard rate function).

The hazard function, hazard rate, hazard rate function, for a random variable T at value t, typically denoted as h(t) or $\lambda(t)$, is the conditional density of T at t, given $T \ge t$. That is:

$$h(t) \stackrel{\text{def}}{=} p(T = t | T \ge t)$$

If T represents the time at which an event occurs, then h(t) is the probability that the event occurs at time t, given that it has not occurred prior to time t.

B.4.3. Expectation

Definition B.11 (Expectation, expected value, population mean). The **expectation**, **expected value**, or **population mean** of a *continuous* random variable X, denoted $\mathbb{E}[X]$, $\mu(X)$, or μ_X , is the weighted mean of X's possible values, weighted by the probability density function of those values:

$$\mathbb{E}\left[X\right] = \int_{x \in \mathcal{R}(X)} x \cdot \mathbf{p}(X = x) dx$$

The expectation, expected value, or population mean of a discrete random variable X, denoted $\mathbb{E}[X]$, $\mu(X)$, or μ_X , is the mean of X's possible values, weighted by the probability mass function of those values:

$$\mathbb{E}\left[X\right] = \sum_{x \in \mathcal{R}(X)} x \cdot \mathrm{P}(X = x)$$

(c.f. https://en.wikipedia.org/wiki/Expected_value)

Theorem B.9 (Expectation of the Bernoulli distribution). The expectation of a Bernoulli random variable with parameter π is:

$$\mathbb{E}\left[X\right] =\pi$$

Proof.

$$\begin{split} \mathbb{E}\left[X\right] &= \sum_{x \in \mathcal{R}(X)} x \cdot \mathbf{P}(X = x) \\ &= \sum_{x \in \{0,1\}} x \cdot \mathbf{P}(X = x) \\ &= (0 \cdot \mathbf{P}(X = 0)) + (1 \cdot \mathbf{P}(X = 1)) \\ &= (0 \cdot (1 - \pi)) + (1 \cdot \pi) \\ &= 0 + \pi \\ &= \pi \end{split}$$

B.4.4. Variance and related characteristics

Definition B.12 (Variance). The variance of a random variable X is the expectation of the squared difference between X and $\mathbb{E}[X]$; that is:

$$\mathrm{Var}\left(X\right) \stackrel{\mathrm{def}}{=} \mathbb{E}\left[(X - \mathbb{E}\left[X\right])^2\right]$$

Theorem B.10 (Simplified expression for variance).

$$Var(X) = \mathbb{E}\left[X^2\right] - \left(\mathbb{E}\left[X\right]\right)^2$$

Proof. By linearity of expectation, we have:

$$\begin{split} \operatorname{Var}\left(X\right) &\stackrel{\text{def}}{=} \mathbb{E}\left[(X - \mathbb{E}\left[X\right])^2\right] \\ &= \mathbb{E}\left[X^2 - 2X\mathbb{E}\left[X\right] + \left(\mathbb{E}\left[X\right]\right)^2\right] \\ &= \mathbb{E}\left[X^2\right] - \mathbb{E}\left[2X\mathbb{E}\left[X\right]\right] + \mathbb{E}\left[\left(\mathbb{E}\left[X\right]\right)^2\right] \\ &= \mathbb{E}\left[X^2\right] - 2\mathbb{E}\left[X\right]\mathbb{E}\left[X\right] + \left(\mathbb{E}\left[X\right]\right)^2 \\ &= \mathbb{E}\left[X^2\right] - \left(\mathbb{E}\left[X\right]\right)^2 \end{split}$$

Definition B.13 (Precision). The **precision** of a random variable X, often denoted $\tau(X)$, τ_X , or shorthanded as τ , is the inverse of that random variable's variance; that is:

$$\tau(X) \stackrel{\text{def}}{=} (\text{Var}(X))^{-1}$$

Definition B.14 (Standard deviation). The standard deviation of a random variable X is the square-root of the variance of X:

$$\mathrm{SD}\left(X\right)\overset{\mathrm{def}}{=}\sqrt{\mathrm{Var}\left(X\right)}$$

Definition B.15 (Covariance). For any two one-dimensional random vari-

$$\operatorname{Cov}\left(X,Y\right) \stackrel{\operatorname{def}}{=} \mathbb{E}\left[(X - \mathbb{E}\left[X\right])(Y - \mathbb{E}\left[Y\right])\right]$$

Theorem B.11.

ables, X, Y:

$$Cov(X, Y) = \mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y]$$

Proof. Left to the reader.

Lemma B.1 (The covariance of a variable with itself is its variance). For any random variable X:

$$Cov(X, X) = Var(X)$$

Proof.

$$\begin{aligned} \operatorname{Cov}\left(X,X\right) &= E[XX] - E[X]E[X] \\ &= E[X^2] - (E[X])^2 \\ &= \operatorname{Var}\left(X\right) \end{aligned}$$

Definition B.16 (Variance/covariance of a $p \times 1$ random vector). For a $p \times 1$ dimensional random vector X,

$$\begin{aligned} \operatorname{Var}(X) &\stackrel{\operatorname{def}}{=} \operatorname{Cov}(X) \\ &\stackrel{\operatorname{def}}{=} E[(X - E[X])^\top \left(X - E[X]\right)] \end{aligned}$$

Theorem B.12 (Alternate expression for variance of a random vector).

$$Var(X) = E[X^{\top}X] - E[X]^{\top}E[X]$$

Proof.

$$\begin{aligned} \operatorname{Var}\left(X\right) &= E[\left(X^{\top} - E[X]^{\top}\right) \left(X - E[X]\right)] \\ &= E[X^{\top}X - E[X]^{\top}X - X^{\top}E[X] + E[X]^{\top}E[X]] \\ &= E[X^{\top}X] - E[X]^{\top}E[X] - E[X]^{\top}E[X] + E[X]^{\top}E[X] \\ &= E[X^{\top}X] - 2E[X]^{\top}E[X] + E[X]^{\top}E[X] \\ &= E[X^{\top}X] - E[X]^{\top}E[X] \end{aligned}$$

Theorem B.13 (Variance of a linear combination). For any set of random variables $X_1, ..., X_n$ and corresponding constants $a_1, ..., a_n$:

$$Var\left(\sum_{i=1}^{n}a_{i}X_{i}\right)=\sum_{i=1}^{n}\sum_{j=1}^{n}a_{i}a_{j}Cov\left(X_{i},X_{j}\right)$$

Proof. Left to the reader...

Lemma B.2. For any two random variables X and Y and scalars a and b:

$$Var(aX + bY) = a^{2} Var(X) + b^{2} Var(Y) + 2(a \cdot b) Cov(X, Y)$$

Proof. Apply Theorem B.13 with n = 2, $X_1 = X$, and $X_2 = Y$.

Or, see https://statproofbook.github.io/P/var-lincomb.html

Definition B.17 (homoskedastic, heteroskedastic). A random variable Y is **homoskedastic** (with respect to covariates X) if the variance of Y does not vary with X:

$$Var(Y|X=x) = \sigma^2, \forall x$$

Otherwise it is **heteroskedastic**.

Definition B.18 (Statistical independence). A set of random variables X_1, \dots, X_n are statistically independent if their joint probability is equal to the product of their marginal probabilities:

$$\Pr(X_1=x_1,\dots,X_n=x_n)=\prod_{i=1}^n\Pr(X_i=x_i)$$

Tip

The symbol for independence, $\$, is essentially just $\$ upside-down. So the symbol can remind you of its definition (Definition B.18).

Definition B.19 (Conditional independence). A set of random variables Y_1, \ldots, Y_n are **conditionally statistically independent** given a set of covariates X_1, \ldots, X_n if the joint probability of the Y_i s given the X_i s is equal to the product of their marginal probabilities:

$$\Pr(Y_1=y_1,\ldots,Y_n=y_n|X_1=x_1,\ldots,X_n=x_n)=\prod_{i=1}^n\Pr(Y_i=y_i|X_i=x_i)$$

Definition B.20 (Identically distributed). A set of random variables X_1, \ldots, X_n are **identically distributed** if they have the same range $\mathcal{R}(X)$ and if their marginal distributions $P(X_1 = x_1), \ldots, P(X_n = x_n)$ are all equal to some shared distribution P(X = x):

$$\forall i \in \left\{1:n\right\}, \forall x \in \mathcal{R}(X): \mathrm{P}(X_i = x) = \mathrm{P}(X = x)$$

Definition B.21 (Conditionally identically distributed). A set of random variables Y_1, \ldots, Y_n are **conditionally identically distributed** given a set of covariates X_1, \ldots, X_n if Y_1, \ldots, Y_n have the same range $\mathcal{R}(X)$ and if the distributions $P(Y_i = y_i | X_i = x_i)$ are all equal to the same distribution P(Y = y | X = x):

$$\mathrm{P}(Y_i = y | X_i = x) = \mathrm{P}(Y = y | X = x)$$

Definition B.22 (Independent and identically distributed). A set of random variables X_1, \ldots, X_n are **independent and identically distributed** (shorthand: " X_i iid") if they are statistically independent and identically distributed.

Definition B.23 (Conditionally independent and identically distributed). A set of random variables Y_1,\ldots,Y_n are **conditionally independent and identically distributed** (shorthand: " $Y_i|X_i$ ciid" or just " $Y_i|X_i$ iid") given a set of covariates X_1,\ldots,X_n if Y_1,\ldots,Y_n are conditionally independent given X_1,\ldots,X_n and Y_1,\ldots,Y_n are identically distributed given X_1,\ldots,X_n .

C.1. Probabilistic models

Definition C.1 (Scientific models). **Scientific models** are attempts to describe *physical conditions or changes* that occur in the world and universe around us.

Example C.1 (Scientific models in epidemiology). Epidemiologists typically study *biological conditions and changes*, such as the spread of infectious diseases through populations, or the effects of environmental factors on individuals.

C.1.1. Statistical analysis of scientific models

When we perform statistical analyses, we use data to help us choose between models - specifically, to determine which models best explain that data.

However, physical processes do not produce data on their own. Data is only produced when scientists implement an *observation process* (i.e., a *scientific study*), which is distinct from the underlying *physical process*. In

some cases, the observation process and the physical process interact with each other. This phenomenon is called the "observer effect".

In order to learn about the physical processes we are ultimately interested in, we often need to make special considerations for the observation process that produced the data which we are analyzing. In particular, if some of the planned observations in the study design were not completed, we will likely need to account for the incompleteness of the resulting data set in our analysis. If we are not sure why some observations are incomplete, we may need to model the observation process in addition to the physical process we were originally interested in. For example, if some participants in a study dropped out part-way through the study, we may need investigate why those participants dropped out, as opposed to other participants who completed the study.

These kinds of *missing data* issues are outside of the scope of this course; see Van Buuren (2018) for more details.

C.2. Estimands, estimates, and estimators

C.2.1. Estimands

Definition C.2 (Estimand). An **estimand** is an unknown quantity whose value we want to know (Pohl et al. 2021; Lawrance et al. 2020).

Example C.2 (Mean height of students). If we are trying to determine the mean height of students at our school, then the *population mean* is our **estimand**.

In statistical contexts, most estimands are parameters of probabilistic models, or functions of model parameters.

¹https://en.wikipedia.org/wiki/Observer effect

Notation for estimands

Model paramaters and other estimands are often symbolized using lower-case Greek letters: $\alpha, \beta, \gamma, \delta$, etc.

C.2.2. Estimates

Definition C.3 (Estimate/estimated value). In statistics, an **estimate** or **estimated value** is an informed guess of an estimand's value, based on observed data.

Example C.3 (Mean height of students). Suppose we measure the heights of 50 random students from our school, and the sample mean was 175cm. We might use 175cm as an *estimate* of the population mean.

C.2.3. Estimators

Definition C.4 (Estimator). An **estimator** is a function $\hat{\theta}(x_1,...x_n)$ that transforms data $x_1,...x_n$ into an estimate.

i Estimators are random variables

When estimators are applied to random variables, the estimators are also random variables.

Notation for estimators

Estimators are often symbolized by placing a $\hat{\theta}$ ("hat") symbol on top of the corresponding estimand; for example, $\hat{\theta}$. Usually, their dependence on the data is implicit:

$$\hat{\theta} \stackrel{\text{def}}{=} \hat{\theta}(x_1,...x_n)$$

Example C.4 (Mean height of students). If we want to estimate the mean height of students at our university, which we will represent as μ , we might measure the heights of n=50 randomly sampled students as random variables $X_1, ..., X_n$. Then we could use the function

$$\hat{\mu}(X_1,...,X_n) = \frac{1}{n} \sum_{i=1}^n X_i \stackrel{\text{def}}{=} \bar{X}$$

as an estimator to produce an estimate $\hat{\mu} = \bar{x}$ of μ .

Another estimator would be just the height of the first student sampled:

$$\hat{\mu}^{(2)}(X_1,...,X_n) = X_1$$

A third possible estimator would be the mean of all sampled students' heights, except for the two most extreme; that is, if we re-order the observations $X_{(1)} = \min_{i \in 1:n} X_i$, $X_{(2)} = \min_{i \in \{1:n\} - \arg X_{(1)}} X_i$, ..., $X_{(n)} = \max_{i \in 1:n} X_i$, then we could define the estimator:

$$\hat{\mu}^{(3)}(X_1,...,X_n) = \frac{1}{n} \sum_{i=2}^{n-1} X_{(i)}$$

Which of these estimators is best? It depends on how we evaluate them (see Section C.3 below).

C.2.4. Contrasting estimands, estimates, and estimators

It's helpful to keep in mind the mathematical type of each estimation concept:

- estimands are numbers (or vector of numbers)
- estimates are also numbers (or vectors)
- estimators are functions of random variables, so they are also random variables

C.3. Accuracy of estimators

C.3.1. Accuracy

To determine which estimator is best, we need to define *best*. "Accuracy" is usually most important; easy computation is usually secondary.

Definition C.5 (Accuracy). The **accuracy** of an estimator for a given estimand does not have a consensus formal definition, but all of the usual candidates are related to the distributions of the *errors* made by the resulting estimates.

C.3.2. Error

Definition C.6 (Error). The **error** of an estimate $\hat{\theta}$ of a true value θ , often denoted $\varepsilon(\hat{\theta})$, or more completely $\varepsilon(\hat{\theta}, \theta)$, is the difference between the estimate and its estimand θ ; that is:

$$\varepsilon(\hat{\theta}) \stackrel{\text{def}}{=} \hat{\theta} - \theta$$

Some frequently-used measures of accuracy include:

C.3.3. Mean squared error

Definition C.7 (Mean squared error). The **mean squared error** of an estimator $\hat{\theta}$, denoted MSE $(\hat{\theta})$, is the expectation of the square of the error²:

$$\mathrm{MSE}\left(\hat{\theta}\right) \stackrel{\mathrm{def}}{=} \mathbb{E}\left[(\varepsilon(\hat{\theta}))^2\right]$$

C.3.4. Mean absolute error

Definition C.8 (Mean absolute error). The **mean absolute error** of an estimator is the expectation of the absolute value of the error:

$$\operatorname{MAE}\left(\hat{\theta}\right) \stackrel{\operatorname{def}}{=} \mathbb{E}\left[\left|\varepsilon(\hat{\theta})\right|\right]$$

C.3.5. Bias

Definition C.9 (Bias). The **bias** of an estimator $\hat{\theta}$ for an estimand θ is the expected value of the error:

$$\operatorname{Bias}\left(\hat{\theta}\right) \stackrel{\operatorname{def}}{=} \mathbb{E}\left[\varepsilon(\hat{\theta})\right] \tag{C.1}$$

Theorem C.1 (Bias equals Expectation minus Truth).

$$Bias\left(\widehat{\theta}\right) =\mathbb{E}\left[\widehat{\theta}\right] -\theta$$

 $^{^{2}}$ depending on whether it is applied to a matrix or a function

Proof.

$$\begin{aligned} \operatorname{Bias}\left(\hat{\theta}\right) & \stackrel{\operatorname{def}}{=} \mathbb{E}\left[\varepsilon(\hat{\theta})\right] \\ &= \mathbb{E}\left[\hat{\theta} - \theta\right] \\ &= \mathbb{E}\left[\hat{\theta}\right] - \mathbb{E}\left[\theta\right] \\ &= \mathbb{E}\left[\hat{\theta}\right] - \theta \end{aligned}$$

The third equality is by the linearity of expectation.

Theorem C.2 (Mean Squared Error equals Bias Squared plus Variance). For any one-dimensional estimator $\hat{\theta}$:

$$MSE(\hat{\theta}) = \left(Bias(\hat{\theta})\right)^2 + Var(\hat{\theta})$$
 (C.2)

Proof. Let's start by expanding each term of the right-hand side:

$$\begin{split} \left(\mathrm{Bias} \left(\hat{\theta} \right) \right)^2 &= \left(\mathbb{E} \left[\hat{\theta} \right] - \theta \right)^2 \\ &= \left(\mathbb{E} \left[\hat{\theta} \right] \right)^2 - 2 \mathbb{E} \left[\hat{\theta} \right] \theta + \theta^2 \end{split}$$

$$\mathrm{Var} \left(\hat{\theta} \right) &= \mathbb{E} \left[\hat{\theta}^2 \right] - \left(\mathbb{E} \left[\hat{\theta} \right] \right)^2 \end{split}$$

Now, add them together and simplify:

$$\begin{split} \left(\mathrm{Bias} \left(\hat{\theta} \right) \right)^2 + \mathrm{Var} \left(\hat{\theta} \right) &= \left(\mathbb{E} \left[\hat{\theta} \right] \right)^2 - 2 \mathbb{E} \left[\hat{\theta} \right] \theta + \theta^2 + \mathbb{E} \left[\hat{\theta}^2 \right] - \left(\mathbb{E} \left[\hat{\theta} \right] \right)^2 \\ &= \mathbb{E} \left[\hat{\theta}^2 \right] - 2 \mathbb{E} \left[\hat{\theta} \right] \theta + \theta^2 \end{split}$$

Now let's expand the left-hand side to reach the same expression:

$$\begin{split} \text{MSE} \left(\hat{\theta} \right) &= \mathbb{E} \left[(\mathbf{e}^{(\hat{\theta})})^2 \right] \\ &= \mathbb{E} \left[(\hat{\theta} - \theta)^2 \right] \\ &= \mathbb{E} \left[\hat{\theta}^2 - 2 \hat{\theta} \theta - \theta^2 \right] \\ &= \mathbb{E} \left[\hat{\theta}^2 \right] - \mathbb{E} \left[2 \hat{\theta} \theta \right] + \mathbb{E} \left[\theta^2 \right] \\ &= \mathbb{E} \left[\hat{\theta}^2 \right] - 2 \mathbb{E} \left[\hat{\theta} \right] \theta + \theta^2 \end{split}$$

$$MSE(\hat{\theta}) = (Bias(\hat{\theta}))^{2} + Var(\hat{\theta})$$

C.3.5.1. Unbiased estimators

Definition C.10 (unbiased estimator). An estimator $\hat{\theta}$ is **unbiased** if Bias $(\hat{\theta}) = 0$.

Theorem C.3 (properties of unbiased estimators). If $\hat{\theta}$ is unbiased, then:

$$\mathbb{E}\left[\hat{\theta}\right] = \theta \tag{C.3}$$

$$MSE(\hat{\theta}) = Var(\hat{\theta})$$
 (C.4)

Proof. If $\hat{\theta}$ is unbiased, then:

Equation C.3:

$$\begin{aligned} \operatorname{Bias}\left(\hat{\theta}\right) &= 0 \\ \mathbb{E}\left[\hat{\theta}\right] - \theta &= 0 \\ \mathbb{E}\left[\hat{\theta}\right] &= \theta \end{aligned}$$

Equation C.4:

$$\begin{aligned} \operatorname{MSE}\left(\hat{\theta}\right) &\stackrel{\text{def}}{=} \mathbb{E}\left[\left(\varepsilon(\hat{\theta})\right)^{2}\right] \\ &= \mathbb{E}\left[\left(\hat{\theta} - \theta\right)^{2}\right] \\ &= \mathbb{E}\left[\left(\hat{\theta} - \mathbb{E}\left[\hat{\theta}\right]\right)^{2}\right] \\ &\stackrel{\text{def}}{=} \operatorname{Var}\left(\hat{\theta}\right) \end{aligned}$$

(Alternative proof of Equation C.4) We could have started from Theorem C.2 instead:

$$\begin{aligned} \text{MSE} \left(\hat{\theta} \right) &= \left(\text{Bias} \left(\hat{\theta} \right) \right)^2 + \text{Var} \left(\hat{\theta} \right) \\ &= \left(0 \right)^2 + \text{Var} \left(\hat{\theta} \right) \\ &= 0 + \text{Var} \left(\hat{\theta} \right) \\ &= \text{Var} \left(\hat{\theta} \right) \end{aligned}$$

C.3.6. Standard error

Definition C.11 (Standard error). The **standard error** of an estimator $\hat{\theta}$ is just the **standard deviation** of $\hat{\theta}$; that is:

$$\operatorname{SE}\left(\hat{\theta}\right) \stackrel{\mathrm{def}}{=} \operatorname{SD}\left(\hat{\theta}\right)$$

"Standard error" is a confusing concept in a few ways. First of all, it isn't even defined as a characteristic of the error, $\varepsilon(\hat{\theta})$! Moreover, it is just a synonym for standard deviation, so it seems like a redundant concept. However, standard errors help us construct p-values and confidence intervals, so they come up a lot - often enough to give them their own name.

We can relate standard error to actual error, by rearranging the result from Theorem C.2:

$$\begin{aligned} \operatorname{Var}\left(\hat{\theta}\right) &= \operatorname{Var}\left(\hat{\theta} - \theta\right) \\ &= \operatorname{Var}\left(\varepsilon(\hat{\theta})\right) \end{aligned}$$

So the variance of the estimator is equal to the variance of the error, and the standard error is equal to the standard deviation of the error:

$$\operatorname{SE}\left(\hat{\theta}\right) = \operatorname{SD}\left(\varepsilon(\hat{\theta})\right)$$

Corollary C.1 (Standard error squared equals MSE minus squared bias). standard error is what is left over of MSE after bias is removed:

$$\left(SE\left(\hat{\theta}\right)\right)^{2} = MSE\left(\hat{\theta}\right) - \left(Bias\left(\hat{\theta}\right)\right)^{2}$$

C. Estimation

Proof.

$$\begin{aligned} \operatorname{MSE}\left(\widehat{\theta}\right) &= \left(\operatorname{Bias}\left(\widehat{\theta}\right)\right)^2 + \operatorname{Var}\left(\widehat{\theta}\right) \\ & \operatorname{...}\operatorname{Var}\left(\widehat{\theta}\right) &= \operatorname{MSE}\left(\widehat{\theta}\right) - \left(\operatorname{Bias}\left(\widehat{\theta}\right)\right)^2 \\ & \operatorname{...}\left(\operatorname{SE}\left(\widehat{\theta}\right)\right)^2 &= \operatorname{MSE}\left(\widehat{\theta}\right) - \left(\operatorname{Bias}\left(\widehat{\theta}\right)\right)^2 \end{aligned}$$

Corollary C.2 (For unbiased estimators, SE = RMSE). If $\mathbb{E}\left[\varepsilon\left(\hat{\theta}\right)\right]=0$, then:

$$SE\left(\hat{\theta}\right) = \sqrt{MSE\left(\hat{\theta}\right)}$$

(this result is equivalent to Equation C.4)

499

D. Inference

D.1. Interpretation of Negative Findings

See Vittinghoff et al. (2012) §3.7 (p64).

These notes are derived primarily from Dobson and Barnett (2018) (mostly chapters 1-5).

Some material was also taken from McLachlan and Krishnan (2007) and Casella and Berger (2002).

Configuring R

Functions from these packages will be used throughout this document:

```
library(conflicted) # check for conflicting function definitions
# library(printr) # inserts help-file output into markdown output
library(rmarkdown) # Convert R Markdown documents into a variety of formats.
library(pander) # format tables for markdown
library(ggplot2) # graphics
library(ggeasy) # help with graphics
library(ggfortify) # help with graphics
```

```
library(dplyr) # manipulate data
library(tibble) # `tibble`s extend `data.frame`s
library(magrittr) # `%>%` and other additional piping tools
library(haven) # import Stata files
library(knitr) # format R output for markdown
library(tidyr) # Tools to help to create tidy data
library(plotly) # interactive graphics
library(dobson) # datasets from Dobson and Barnett 2018
library(parameters) # format model output tables for markdown
library(haven) # import Stata files
library(latex2exp) # use LaTeX in R code (for figures and tables)
library(fs) # filesystem path manipulations
library(survival) # survival analysis
library(survminer) # survival analysis graphics
library(KMsurv) # datasets from Klein and Moeschberger
library(parameters) # format model output tables for
library(webshot2) # convert interactive content to static for pdf
library(forcats) # functions for categorical variables ("factors")
library(stringr) # functions for dealing with strings
library(lubridate) # functions for dealing with dates and times
```

Here are some R settings I use in this document:

```
knitr::opts_chunk$set(message = FALSE)
options('digits' = 4)

panderOptions("big.mark", ",")
pander::panderOptions("table.emphasize.rownames", FALSE)
pander::panderOptions("table.split.table", Inf)
conflicts_prefer(dplyr::filter) # use the `filter()` function from dplyr() by defaul
legend_text_size = 9
```

E.1. Overview of maximum likelihood estimation

E.1.1. The likelihood function

Definition E.1 (Likelihood of a single observation). Let X be a random variable and let x be X's observed data value. Let $p_{\Theta}(X = x)$ be a probability model for the distribution of X, with parameter vector Θ .

Then the **likelihood** of parameter value θ , for model $p_{\Theta}(X = x)$ and data X = x, is simply the probability of the event X = x given $\Theta = \theta$:

$$\mathcal{L}(\theta) \stackrel{\mathrm{def}}{=} \mathrm{P}_{\theta}(X = x)$$

Definition E.2 (Likelihood of a dataset). Let \tilde{x} be a dataset with corresponding random variable \tilde{X} . Let $p_{\Theta}(\tilde{X})$ be a probability model for the distribution of \tilde{X} with unknown parameter vector Θ .

Then the **likelihood** of parameter value θ , for model $p_{\Theta}(X)$ and data $\tilde{X} = \tilde{x}$, is the *joint probability* of $\tilde{X} = \tilde{x}$ given $\Theta = \theta$:

$$\begin{split} \mathcal{L}(\theta) &\stackrel{\text{def}}{=} p_{\theta}(\tilde{X} = \tilde{x}) \\ &= p_{\theta}(X_1 = x_1, ..., X_n = x_n) \end{split}$$

Notation for the likelihood function

The likelihood function can be written as:

- $\begin{array}{ccc} & \mathcal{L}(\widetilde{x};\theta) \\ \bullet & \mathcal{L}(\widetilde{x};\theta) \\ \bullet & \mathcal{L}(\theta;\widetilde{x}) \\ \bullet & \mathcal{L}_{\widetilde{x}}(\theta) \\ \bullet & \mathcal{L}_{\theta}(\widetilde{x}) \\ \bullet & \mathcal{L}(\widetilde{x}|\theta) \end{array}$

All of these notations mean the same thing.

The likelihood is a function that takes θ (and implicitly, \tilde{X}) as inputs and outputs a single number, the joint probability of \tilde{x} for model $p_{\Theta}(\tilde{X} = \tilde{x})$ with $\Theta = \theta$.

Theorem E.1 (Likelihood of an independent sample). For mutually in $dependent\ data\ X_1,...,X_n\colon$

$$\mathcal{L}(\tilde{x}|\theta) = \prod_{i=1}^{n} p(X_i = x_i|\theta) \tag{E.1}$$

Proof.

$$\begin{split} \mathcal{L}(\tilde{x}|\theta) &\stackrel{\text{def}}{=} \mathrm{p}(X_1 = x_1, ..., X_n = x_n|\theta) \\ &= \prod_{i=1}^n \mathrm{p}(X_i = x_i|\theta) \end{split}$$

The second equality is by the definition of statistical independence. \Box

Definition E.3 (Likelihood components). Given an iid dataset \tilde{x} , the likelihood component or likelihood factor of observation $X_i = x_i$ is the marginal likelihood of $X_i = x_i$:

$$\mathcal{L}_i(\theta) = \mathrm{P}(X_i = x_i)$$

Theorem E.2. For iid data $\tilde{x} \stackrel{def}{=} x_1, \dots, x_n$, the likelihood of the dataset is equal to the product of the observation-specific likelihood factors:

$$\mathcal{L}(\theta) = \prod_{i=1}^n \mathcal{L}_i(\theta)$$

E.1.2. The maximum likelihood estimate

Definition E.4 (Maximum likelihood estimate). The **maximum likelihood estimate** of a parameter vector Θ , denoted $\hat{\theta}_{ML}$, is the value of Θ that maximizes the likelihood:

$$\hat{\theta}_{\mathrm{ML}} \stackrel{\mathrm{def}}{=} \arg \max_{\Theta} \mathcal{L}(\Theta) \tag{E.2}$$

E.1.3. Finding the maximum of a function

Recall from calculus: the maxima of a continuous function f(x) over a range of input values $\mathcal{R}(x)$ can be found either:

- at the edges of the range of input values, OR:
- where the function is flat (i.e. where the gradient function f'(x) = 0) AND the second derivative is negative definite (f''(x) < 0).

E.1.4. Directly maximizing the likelihood function for iid data

To find the maximizer(s) of the likelihood function, we need to solve $\mathcal{L}'(\theta) = 0$ for θ . However, even for mutually independent data, we quickly run into a problem:

$$\begin{split} \mathcal{L}'(\theta) &= \frac{\partial}{\partial \theta} \mathcal{L}(\theta) \\ &= \frac{\partial}{\partial \theta} \prod_{i=1}^n p(X_i = x_i | \theta) \end{split} \tag{E.3}$$

The derivative of the likelihood of independent data is the derivative of a product. We will have to perform a massive application of the product rule to evaluate this derivative.

E.1.5. The log-likelihood function

It is typically easier to work with the log of the likelihood function:

Definition E.5 (Log-likelihood). The **log-likelihood** of parameter value θ , for model $p_{\Theta}(\tilde{X})$ and data $\tilde{X} = \tilde{x}$, is the natural logarithm of the likelihood¹:

 $^{^{1}} https://en.wikipedia.org/wiki/Does_exactly_what_it_says_on_the_tin$

$$\ell(\theta) \stackrel{\text{def}}{=} \log \left\{ \mathcal{L}(\theta) \right\}$$

Definition E.6 (Log-likelihood components). Given a dataset $\tilde{X} = \tilde{x}$, the **log-likelihood component of observation** $X_i = x_i$ is the natural logarithm of the likelihood component:

$$\ell_i(\theta) \stackrel{\text{def}}{=} \log \left\{ \mathcal{L}_i(\theta) \right\}$$

Theorem E.3. The likelihood and log-likelihood have the same maximizer:

$$\arg\max_{\theta} \mathcal{L}(\theta) = \arg\max_{\theta} \ell(\theta)$$

Proof. Left to the reader.

reader.

Theorem E.4 (Log-likelihood of an iid sample). For iid data $X_1,...,X_n$ with shared distribution p(X=x):

$$\ell(x|\theta) = \sum_{i=1}^{n} \log \left\{ p(X = x_i|\theta) \right\}$$
 (E.4)

Proof.

$$\begin{split} \ell(x|\theta) &\stackrel{\text{def}}{=} \log \left\{ \mathcal{L}(\tilde{x}|\theta) \right\} \\ &= \log \left\{ \prod_{i=1}^n \mathrm{p}(X_i = x_i|\theta) \right\} \\ &= \sum_{i=1}^n \log \left\{ p(X = x_i|\theta) \right\} \end{split}$$

For iid data, we will have a much easier time taking the derivative of the log-likelihood:

Theorem E.5 (Derivative of the log-likelihood function for iid data). For $iid\ data$:

$$\ell'(\theta) = \sum_{i=1}^{n} \frac{\partial}{\partial \theta} log \{ p(X = x_i | \theta) \}$$
 (E.5)

Proof.

$$\begin{split} \ell'(\theta) &= \frac{\partial}{\partial \theta} \ell(\theta) \\ &= \frac{\partial}{\partial \theta} \sum_{i=1}^n \log \left\{ \mathbf{p}(X = x_i | \theta) \right\} \\ &= \sum_{i=1}^n \frac{\partial}{\partial \theta} \log \left\{ \mathbf{p}(X = x_i | \theta) \right\} \end{split}$$

E.1.6. The score function

The first derivative² of the log-likelihood, $\ell'(\theta)$, is important enough to have its own name: the *score function*.

Definition E.7 (Score function). The **score function** of a statistical model $p(\tilde{X} = \tilde{x})$ is the gradient (i.e., first derivative) of the log-likelihood of that model:

$$\ell'(\theta) \stackrel{\text{def}}{=} \frac{\partial}{\partial \theta} \ell(\theta)$$

We often skip writing the arguments x and/or θ), so $\ell' \stackrel{\text{def}}{=} \ell'(\tilde{x}; \theta) \stackrel{\text{def}}{=} \ell'(\theta)$. Some statisticians use U or S instead of ℓ' . I prefer ℓ' . Why use up extra letters?

E.1.7. Asymptotic distribution of the maximum likelihood estimate

We learned how to quantify our uncertainty about these maximum likelihood estimates; with sufficient sample size, $\hat{\theta}_{\rm ML}$ has the approximate distribution:

$$\hat{\theta}_{ML} \dot{\sim} N(\theta, \mathcal{I}(\theta)^{-1})$$

Recall:

- $\mathcal{I}(\theta) \stackrel{\text{def}}{=} \mathbb{E}\left[I(\tilde{X}; \theta)\right]$
- $I(\tilde{X}, \theta) \stackrel{\text{def}}{=} -\ell''(\tilde{X}; \theta)$

²a.k.a. the gradient³

 $^{^4}$ depending on whether it is applied to a matrix or a function

We can estimate $\mathcal{I}(\theta)$ using either $\mathcal{I}(\hat{\theta}_{\mathrm{ML}})$ or $I(\tilde{x}; \hat{\theta}_{\mathrm{ML}})$.

So we can estimate the standard error of $\hat{\theta}_k$ as:

$$\widehat{\mathrm{SE}}\left(\widehat{\boldsymbol{\theta}}_{k}\right) = \sqrt{\left[\left(\widehat{\mathcal{I}}\left(\widehat{\boldsymbol{\theta}}_{\mathrm{ML}}\right)\right)^{-1}\right]_{kk}}$$

E.1.8. The (Fisher) (expected) information matrix

The variance of $\ell'(x,\theta)$, $Cov\{\ell'(x,\theta)\}$, is also very important; we call it the "expected information matrix", "Fisher information matrix", or just "information matrix", and we represent it using the symbol $\mathcal{I}(I)$ (\frakturI in Unicode, \mathfrak{I} in LaTeX).

$$\begin{split} \mathcal{I} & \stackrel{\text{def}}{=} \mathcal{I}(\theta) \\ & \stackrel{\text{def}}{=} \operatorname{Cov}\left(\ell'|\theta\right) \\ & = \mathbb{E}[\ell'{\ell'}^\top] - \mathbb{E}[\ell'] \ \mathbb{E}[\ell']^\top \end{split}$$

The elements of \Im are:

$$\begin{split} \mathfrak{I}_{ij} &\stackrel{\text{def}}{=} \operatorname{Cov}\left({\ell'}_i, {\ell'}_j\right) \\ &= \mathbb{E}[\ell'_i \ell'_j] - \mathbb{E}[{\ell'}_i] \mathbb{E}[{\ell'}_j] \end{split}$$

Here,

$$\mathbb{E}\left[\ell'\right] \stackrel{\text{def}}{=} \int_{x \in \mathcal{R}(x)} \ell'(x, \theta) p(X = x | \theta) dx$$

$$= \int_{x \in \mathcal{R}(X)} \left(\frac{\partial}{\partial \theta} \log \left\{ p(X = x | \theta) \right\} \right) p(X = x | \theta) dx$$

$$= \int_{x \in \mathcal{R}(X)} \frac{\frac{\partial}{\partial \theta} p(X = x | \theta)}{p(X = x | \theta)} p(X = x | \theta) dx$$

$$= \int_{x \in \mathcal{R}(X)} \frac{\partial}{\partial \theta} p(X = x | \theta) dx$$

And similarly

$$\mathbb{E}\left[\ell'\ell'^{\top}\right] \stackrel{\mathrm{def}}{=} \int_{x \in R(x)} \ell'(x,\theta) \ell'(x,\theta)^{\top} \ \mathbf{p}\left(X = x | \theta\right) \ dx$$

Note that $\mathbb{E}\left[\ell'\right]$ and $\mathbb{E}\left[\ell'{\ell'}^{\top}\right]$ are functions of θ but not of x; the expectation operator removed x.

Also note that for most of the distributions you are familiar with (including Gaussian, binomial, Poisson, exponential):

$$\mathbb{E}\left[\ell'\right] = 0$$

So

$$\mathcal{I}\left(\theta\right) = \mathbb{E}\left[\ell'{\ell'}^{\top}\right]$$

Moreover, for those distributions (called the "exponential family"), we have:

$$\Im = -\mathbb{E}\left[\ell''\right] = \mathbb{E}\left[-\ell''\right]$$

(see Dobson and Barnett (2018), §3.17), where

$$\ell'' \stackrel{\mathrm{def}}{=} \frac{\partial}{\partial \theta} \ell^{'(x,\theta)^\top} = \frac{\partial}{\partial \theta} \frac{\partial}{\partial \theta^\top} \ell(x,\theta)$$

is the $p \times p$ matrix whose elements are:

$$\ell_{ij}'' \stackrel{\text{def}}{=} \frac{\partial}{\partial \theta_i} \frac{\partial}{\partial \theta_j} \log \left\{ p \left(X = x \mid \theta \right) \right\}$$

 ℓ'' is called the "Hessian" of the log-likelihood function.

Sometimes, we use $I(\theta;x) \stackrel{\text{def}}{=} -\ell''$ (note the standard-font "I" here). $I(\theta;x)$ is the observed information, precision, or concentration matrix (Negative Hessian).

Key point

The asymptotics of MLEs gives us $\hat{\theta}_{ML} \sim N(\theta, \Im^{-1}(\theta))$, approximately, for large sample sizes.

We can estimate $\mathcal{I}^{-1}(\theta)$ by working out $\mathbb{E}\left[-\ell''\right]$ or $\mathbb{E}\left[\ell'\ell'^{\top}\right]$ and plugging in $\hat{\theta}_{\mathrm{ML}}$, but sometimes we instead use $I(\hat{\theta}_{\mathrm{ML}}, \tilde{x})$ for convenience; there are some cases where it's provably better according to some criteria (Efron and Hinkley (1978)).

E.1.9. Iterative maximization

(c.f., Dobson and Barnett (2018), Chapter 4)

⁵named after mathematician Otto Hesse⁶

Later, when we are trying to find MLEs for likelihoods which we can't easily differentiate, we will "hill-climb" using the Newton-Raphson algorithm:

$$\begin{split} \hat{\theta}^* &\leftarrow \hat{\theta}^* + \left(I\left(\tilde{y}; \hat{\theta}^*\right) \right)^{-1} \ell'\left(\tilde{y}; \hat{\theta}^*\right) \\ &= \hat{\theta}^* - \left(\ell''\left(\tilde{y}; \hat{\theta}^*\right) \right)^{-1} \ell'\left(\tilde{y}; \hat{\theta}^*\right) \end{split}$$

The reasoning for this algorithm is that we can approximate the score function using the first-order Taylor polynomial⁷:

$$\begin{split} \ell'(\theta) &\approx \ell'^*(\theta) \\ &\stackrel{\text{def}}{=} \ell'(\hat{\theta}^*) + \ell''(\hat{\theta}^*)(\theta - \hat{\theta}^*) \end{split}$$

The approximate score function, $\ell'^*(\theta)$, is a linear function of θ , so it is easy to solve the corresponding approximate score equation, $\ell'^*(\theta) = 0$, for θ :

$$\theta = \hat{\theta}^* - \ell'(\hat{\theta}^*) \cdot \left(\ell''(\hat{\theta}^*)\right)^{-1}$$

For computational simplicity, we will sometimes use $\mathfrak{I}^{-1}(\theta)$ in place of $I\left(\hat{\theta},y\right)$; doing so is called "Fisher scoring" or the "method of scoring". Note that this is the opposite of the substitution that we are making for estimating the variance of the MLE; this time we should technically use the observed information but we use the expected information instead.

⁷https://en.wikipedia.org/wiki/Taylor%27s_theorem

There's also an "empirical information matrix" (see McLachlan and Krishnan (2007)):

$$I_e(\theta,y) \stackrel{\mathrm{def}}{=} \sum_{i=1}^n {\ell_i'} \ {\ell_i'}^\top - \frac{1}{n} {\ell'} {\ell'}^\top$$

where ℓ_i is the log-likelihood of the ith observation. Note that $\ell' = \sum_{i=1}^n \ell_i'$.

 $\frac{1}{n}I_e(\theta,y)$ is the sample equivalent of

$$\mathfrak{I} \stackrel{\text{def}}{=} \mathfrak{I}(\theta) \stackrel{\text{def}}{=} Cov\left(\ell'|\theta\right) = E[\ell'{\ell'}^\top] - E[\ell'] \ E[\ell']^\top$$

$$\left\{ \Im_{jk} \stackrel{\mathrm{def}}{=} Cov\left({\ell'}_j, {\ell'}_k\right) = E[{\ell'_j}{\ell'_k}] - E[{\ell'}_j]E[{\ell'}_k] \right\}$$

 $I_e(\theta,y)$ is sometimes computationally easier to compute for Newton-Raphson-type maximization algorithms.

 $c.f.\ https://en.wikipedia.org/wiki/Newton\%27s_method_in_optimization$

E.1.10. Quantifying (un)certainty of MLEs

E.1.10.1. Confidence intervals for MLEs

An asymptotic approximation of a 95% confidence interval for θ_k is

$$\widehat{\boldsymbol{\theta}}_{\mathrm{ML}} \pm \boldsymbol{z}_{0.975} \times \widehat{\mathrm{SE}} \left(\widehat{\boldsymbol{\theta}}_{k} \right)$$

where z_{β} the β quantile of the standard Gaussian distribution.

E.1.10.2. p-values and hypothesis tests for MLEs

(to add)

E.1.10.3. Likelihood ratio tests for MLEs

log(likelihood ratio) tests (c.f. Dobson and Barnett 2018, sec. 5.7):

$$-2\ell_0 \sim \chi^2(p-q)$$

See also https://online.stat.psu.edu/stat504/book/export/html/657

E.1.10.4. Prediction intervals for MLEs

$$\overline{X} \in [\hat{\mu} \pm z_{1-\alpha/2} \frac{\sigma}{m}]$$

Where m is the sample size of the new data to be predicted (typically 1, except for binary outcomes, where it needs to be bigger for prediction intervals to make sense)

E.2. Example: Maximum likelihood for Tropical Cyclones in Australia

(Adapted from Dobson and Barnett (2018) §1.6.5)

E.2.1. Data

The cyclones dataset in the dobson package (Table E.1) records the number of tropical cyclones in Northeastern Australia during 13 November-to-April cyclone seasons (more details in Dobson and Barnett (2018) §1.6.5 and help(cyclones, package = "dobson")). Figure E.1 graphs the number of cyclones (y-axis) by season (x-axis). Let's use Y_i to represent these counts, where i is an indexing variable for the seasons and Y_i is the number of cyclones in season i.

E.2.2. Exploratory analysis

Suppose we want to learn about how many cyclones to expect per season.

```
library(dobson)
library(dplyr)
data(cyclones)
library(pander)
pander(cyclones |> relocate(season, .before = everything()))
```

Table E.1.: Number of tropical cyclones during a season from November to April in Northeastern Australia

| season | years | number |
|--------|---------|--------|
| 1 | 1956/7 | 6 |
| 2 | 1957/8 | 5 |
| 3 | 1958/9 | 4 |
| 4 | 1959/60 | 6 |
| 5 | 1960/1 | 6 |
| 6 | 1961/2 | 3 |
| 7 | 1962/3 | 12 |
| 8 | 1963/4 | 7 |

Table E.1.: Number of tropical cyclones during a season from November to April in Northeastern Australia

| season | years | number |
|--------|--------|--------|
| 9 | 1964/5 | 4 |
| 10 | 1965/6 | 2 |
| 11 | 1966/7 | 6 |
| 12 | 1967/8 | 7 |
| 13 | 1968/9 | 4 |

```
library(ggplot2)
library(dplyr)
cyclones |>
  mutate(years = years |> factor(levels = years)) |>
  ggplot(aes(x = years, y = number, group = 1)) +
  geom_point() +
  geom_line() +
  xlab("Season") +
  ylab("Number of cyclones") +
  expand_limits(y = 0) +
  theme(axis.text.x = element_text(vjust = .5, angle = 45))
```




Figure E.1.: Number of tropical cyclones per season in northeastern Australia, 1956-1969

There's no obvious correlation between adjacent seasons, so let's assume that each season is independent of the others.

Let's also assume that they are identically distributed; let's denote this distribution as P(Y=y) (note that there's no index i in this expression, since we are assuming the Y_i s are identically distributed). We can visualize the distribution using a bar plot (Figure E.2). Table E.2 provides summary statistics.

```
cyclones |>
  ggplot() +
  geom_histogram(aes(x = number)) +
  expand_limits(x = 0) +
```

```
xlab("Number of cyclones") +
ylab("Count (number of seasons)")
```



Figure E.2.: Bar plot of cyclones per season

```
n = nrow(cyclones)
sumx = cyclones |> pull(number) |> sum()
xbar = cyclones |> pull(number) |> mean()
cyclones |> table1::table1(x = ~ number)
```

E.2.3. Model

We want to estimate P(Y = y); that is, P(Y = y) is our estimand.

Table E.2.: Summary statistics for cyclones data

| | 0 11 |
|-------------------|-------------------|
| | Overall |
| | (N=13) |
| number | |
| Mean (SD) | 5.54(2.47) |
| Median [Min, Max] | 6.00 [2.00, 12.0] |

We could estimate P(Y = y) for each value of y in $0 : \infty$ separately ("nonparametrically") using the fraction of our data with $Y_i = y$, but then we would be estimating an infinitely large set of parameters, and we would have low precision. We will probably do better with a parametric model.

Exercise E.1. What parametric probability distribution family might we use to model this empirical distribution?

Solution. Let's use the Poisson. The Poisson distribution is appropriate for this data , because the data are counts that could theoretically take any integer value (discrete) in the range $0:\infty$. Visually, the plot of our data closely resembles a Poisson or binomial distribution. Since cyclones do not have an "upper limit" on the number of events we could potentially observe in one season, the Poisson distribution is more appropriate than the binomial.

Exercise E.2. Write down the Poisson distribution's probability mass function.

Solution.

$$P(Y=y) = \frac{\lambda^y e^{-\lambda}}{y!} \tag{E.6}$$

E.2.4. Estimating the model parameters using maximum likelihood

Now, we can estimate the parameter λ for this distribution using maximum likelihood estimation.

What is the likelihood?

Exercise E.3. Write down the likelihood (probability mass function or probability density function) of a single observation x, according to your model.

Solution.

$$\begin{split} \mathcal{L}(\lambda;x) &= p(X=x|\Lambda=\lambda) \\ &= \frac{\lambda^x e^{-\lambda}}{x!} \end{split}$$

Exercise E.4. Write down the vector of parameters in your model.

Solution. There is only one parameter, λ :

$$\theta = (\lambda)$$

Exercise E.5. Write down the population mean and variance of a single observation from your chosen probability model, as a function of the parameters (extra credit - derive them).

Solution.

• Population mean: $E[X] = \lambda$

• Population variance: $Var(X) = \lambda$

Exercise E.6. Write down the likelihood of the full dataset.

Solution.

$$\begin{split} \mathcal{L}(\lambda; \tilde{x}) &= \mathbf{P}(\tilde{X} = \tilde{x}) \\ &= \mathbf{P}(X_1 = x_1, X_2 = x_2, ..., X_{13} = x_{13}) \\ &= \prod_{i=1}^{13} \mathbf{P}(X_i = x_i) \\ &= \prod_{i=1}^{13} \frac{\lambda^{x_i} e^{-\lambda}}{x_i!} \end{split}$$

Exercise E.7. Graph the likelihood as a function of λ .

Solution.

```
lik = function(lambda, y = cyclones$number, n = length(y))
{
lambda^sum(y) * exp(-n*lambda) / prod(factorial(y))
}

library(ggplot2)
lik_plot =
ggplot() +
geom_function(fun = lik, n = 1001) +
xlim(min(cyclones$number), max(cyclones$number)) +
ylab("likelihood") +
xlab('lambda')

print(lik_plot)
```



Figure E.3.: Likelihood of Dobson cyclone data

Exercise E.8. Write down the log-likelihood of the full dataset.

Solution.

$$\begin{split} \ell(\lambda; \tilde{x}) &= \log \left\{ \mathcal{L}(\lambda; \tilde{x}) \right\} \\ &= \log \left\{ \prod_{i=1}^n \frac{\lambda^{x_i} \mathrm{e}^{-\lambda}}{x_i!} \right\} \\ &= \sum_{i=1}^n \log \left\{ \frac{\lambda^{x_i} \mathrm{e}^{-\lambda}}{x_i!} \right\} \\ &= \sum_{i=1}^n \log \left\{ \lambda^{x_i} \right\} + \log \left\{ \mathrm{e}^{-\lambda} \right\} - \log \left\{ x_i! \right\} \\ &= \sum_{i=1}^n x_i \log \left\{ \lambda \right\} - \lambda - \log \left\{ x_i! \right\} \\ &= \sum_{i=1}^n x_i \log \left\{ \lambda \right\} - \sum_{i=1}^n \lambda - \sum_{i=1}^n \log \left\{ x_i! \right\} \\ &= \sum_{i=1}^n x_i \log \left\{ \lambda \right\} - n\lambda - \sum_{i=1}^n \log \left\{ x_i! \right\} \end{split}$$

Exercise E.9. Graph the log-likelihood as a function of λ .

Solution.

```
loglik = function(lambda, y = cyclones$number, n = length(y))
{
sum(y) * log(lambda) - n*lambda - sum(log(factorial(y)))
}

ll_plot = ggplot() +
geom_function(fun = loglik, n = 1001) +
xlim(min(cyclones$number), max(cyclones$number)) +
ylab("log-likelihood") +
xlab('lambda')
ll_plot
```




Figure E.4.: log-likelihood of Dobson cyclone data

E.2.4.1. The score function

Exercise E.10. Derive the score function for the dataset.

Solution. The score function is the first derivative of the log-likelihood:

$$\begin{split} \ell'(\lambda;\tilde{x}) &= \frac{\partial}{\partial \lambda} \sum_{i=1}^n x_i \log \left\{ \lambda \right\} - n\lambda - \sum_{i=1}^n \log \left\{ x_i ! \right\} \\ &= \frac{\partial}{\partial \lambda} \sum_{i=1}^n x_i \log \left\{ \lambda \right\} - \frac{\partial}{\partial \lambda} n\lambda - \frac{\partial}{\partial \lambda} \sum_{i=1}^n \log \left\{ x_i ! \right\} \\ &= \sum_{i=1}^n x_i \frac{\partial}{\partial \lambda} \log \left\{ \lambda \right\} - n \frac{\partial}{\partial \lambda} \lambda - \sum_{i=1}^n \frac{\partial}{\partial \lambda} \log \left\{ x_i ! \right\} \\ &= \sum_{i=1}^n x_i \frac{1}{\lambda} - n - 0 \\ &= \frac{1}{\lambda} \sum_{i=1}^n x_i - n \\ &= \left(\frac{1}{\lambda} n \bar{x} \right) - n \\ &= \left(\frac{1}{\lambda} 72 \right) - 13 \end{split}$$

Exercise E.11. Graph the score function.

Solution.

```
score = function(lambda, y = cyclones$number, n = length(y))
{
   (sum(y) / lambda) - n
}

ggplot() +
geom_function(fun = score, n = 1001) +
xlim(min(cyclones$number), max(cyclones$number)) +

ylab("l'(lambda)") +
```

```
xlab('lambda') +
geom_hline(yintercept = 0, col = 'red')
```



Figure E.5.: score function of Dobson cyclone data

E.2.4.2. The Hessian matrix

Exercise E.12. Derive the Hessian matrix.

Solution. The Hessian function for an iid sample is the 2nd derivative(s) of the log-likelihood:

$$\begin{split} \ell''(\lambda;\tilde{x}) &= \frac{\partial}{\partial \lambda} \left(\frac{1}{\lambda} \sum_{i=1}^{n} x_i - n \right) \\ &= \frac{\partial}{\partial \lambda} \frac{1}{\lambda} \sum_{i=1}^{n} x_i - \frac{\partial}{\partial \lambda} n \\ &= -\frac{1}{\lambda^2} \sum_{i=1}^{n} x_i \\ &= -\frac{1}{\lambda^2} n \bar{x} \\ &= -\frac{1}{\lambda^2} \cdot 72 \end{split}$$

Exercise E.13. Graph the Hessian.

Solution.

```
hessian = function(lambda, y = cyclones$number, n = length(y))
{
    -sum(y)/(lambda^2)
}

ggplot() +
geom_function(fun = hessian, n = 1001) +
xlim(min(cyclones$number), max(cyclones$number)) +

ylab("l''(lambda)") +
xlab('lambda') +
geom_hline(yintercept = 0, col = 'red')
```



Figure E.6.: Hessian function of Dobson cyclone data

Exercise E.14. Write the score equation (estimating equation).

Solution.

$$\ell'(\lambda;\tilde{x})=0$$

Exercise E.15. Solve the estimating equation for λ :

Solution.

$$0 = \frac{1}{\lambda} \sum_{i=1}^{n} x_i - n$$

$$n = \frac{1}{\lambda} \sum_{i=1}^{n} x_i$$

$$n\lambda = \sum_{i=1}^{n} x_i$$

$$\lambda = \frac{1}{n} \sum_{i=1}^{n} x_i$$

$$= \bar{x}$$

Let's call this solution of the estimating equation $\tilde{\lambda}$ for now:

$$\tilde{\lambda} \stackrel{\text{def}}{=} \bar{x}$$

Exercise E.16. Confirm that the Hessian $\ell''(\lambda; \tilde{x})$ is negative when evaluated at $\tilde{\lambda}$.

Solution.

$$\begin{split} \ell''(\tilde{\lambda};\tilde{x}) &= -\frac{1}{\tilde{\lambda}^2} n \bar{x} \\ &= -\frac{1}{\bar{x}^2} n \bar{x} \\ &= -\frac{n}{\bar{x}} \\ &< 0 \end{split}$$

Exercise E.17. Find the MLE of λ .

Solution. Since $\ell''(\tilde{\lambda}; \tilde{x}) < 0$, $\tilde{\lambda}$ is at least a local maximizer of the likelihood function $\mathcal{L}(\lambda)$. Since there is only one solution to the estimating equation

and the Hessian is negative definite everywhere, $\tilde{\lambda}$ must also be the global maximizer of $\mathcal{L}(\lambda; \tilde{x})$:

$$\hat{\lambda}_{\mathrm{ML}} = \bar{x} = 5.5385$$

 ${\bf Exercise} \ {\bf E.18.} \ {\bf Graph \ the \ log-likelihood \ with \ the \ MLE \ superimposed.}$

Solution.

```
library(dplyr)

mle_data = tibble(x = mle, y = loglik(mle))

ll_plot + geom_point(data = mle_data, aes(x = x, y = y), col = 'red')
```




Figure E.7.: log-likelihood of Dobson cyclone data with MLE

E.2.4.3. Information matrices

```
obs_inf = function(...) -hessian(...)
ggplot() +
geom_function(fun = obs_inf, n = 1001) +
xlim(min(cyclones$number), max(cyclones$number)) +
ylab("I(lambda)") +
xlab('lambda') +
geom_hline(yintercept = 0, col = 'red')
```



Figure E.8.: Observed information function of Dobson cyclone data

Example E.1 (Finding the MLE using the Newton-Raphson algorithm).

We found that the MLE was $\hat{\lambda} = \bar{x}$, by solving the score equation $\ell'(\lambda) = 0$ for λ .

What if we hadn't been able to solve the score equation?

Then we could start with some initial guess $\hat{\lambda}^*$, such as $\hat{\lambda}^* = 3$, and use the Newton-Raphson algorithm.

```
# specify initial guess:
cur_lambda_est = 3
```

In Exercise E.10, we found that the score function was:

$$\ell'(\lambda;\tilde{x}) = \left(\frac{72}{\lambda}\right) - n$$

In Exercise E.12, we found that the Hessian was:

$$\ell''(\lambda;\tilde{x}) = -\frac{72}{\lambda^2}$$

So we can approximate the the score function using the first-order Taylor polynomial⁸:

$$\begin{split} \ell'(\lambda) &\approx \ell'^*(\lambda) \\ &\stackrel{\text{def}}{=} \ell'(\hat{\lambda}^*) + \ell''(\hat{\lambda}^*)(\lambda - \hat{\lambda}^*) \\ &= \left(\frac{72}{\hat{\lambda}^*} - n\right) + \left(-\frac{72}{\left(\hat{\lambda}^*\right)^2}\right)(\lambda - \hat{\lambda}^*) \end{split}$$

Figure E.9 compares the true score function and the approximate score function at $\hat{\lambda}^* = 3$.

⁸https://en.wikipedia.org/wiki/Taylor%27s_theorem

```
approx_score = function(lambda, lhat, ...)
  score(lambda = lhat, ...) +
  hessian(lambda = lhat, ...) * (lambda - lhat)
}
point_size = 5
plot1 = ggplot() +
geom_function(
 fun = score,
  aes(col = "true score function"),
  n = 1001) +
geom_function(
  fun = approx_score,
  aes(col = "approximate score function"),
  n = 1001,
  args = list(lhat = cur_lambda_est)) +
geom_point(
 size = point_size,
  aes(x = cur_lambda_est, y = score(lambda = cur_lambda_est),
      col = "current estimate")
) +
geom_point(
size = point_size,
aes(
x = xbar,
y = 0,
col = "true MLE"
) +
xlim(min(cyclones$number), max(cyclones$number)) +
ylab("1'(lambda)") +
```

```
xlab('lambda') +
geom_hline(yintercept = 0)
print(plot1)
```




Figure E.9.: score function of Dobson cyclone data and approximate score function

This is equivalent to estimating the log-likelihood with a second-order Taylor polynomial:

$$\ell^*(\lambda) = \ell(\hat{\lambda}^*) + (\lambda - \hat{\lambda}^*)\ell'(\hat{\lambda}^*) + \frac{1}{2}\ell''(\hat{\lambda}^*)(\lambda - \hat{\lambda}^*)^2$$

```
approx_loglik = function(lambda, lhat, ...)
loglik(lambda = lhat, ...) +
score(lambda = lhat, ...) * (lambda - lhat) +
1/2 * hessian(lambda = lhat, ...) * (lambda - lhat)^2
}
plot_loglik = ggplot() +
geom_function(
 fun = loglik,
  aes(col = "true log-likelihood"),
 n = 1001) +
geom_function(
 fun = approx_loglik,
  aes(col = "approximate log-likelihood"),
  n = 1001,
  args = list(lhat = cur_lambda_est)) +
geom_point(
  size = point_size,
  aes(x = cur_lambda_est, y = loglik(lambda = cur_lambda_est),
      col = "current estimate")
) +
geom_point(
size = point_size,
aes(
x = xbar,
y = loglik(xbar),
col = "true MLE"
)
) +
xlim(min(cyclones$number) - 1, max(cyclones$number)) +
ylab("1'(lambda)") +
xlab('lambda')
```

print(plot_loglik)




Figure E.10.: log-likelihood of Dobson cyclone data and approximate log-likelihood function

The approximate score function, $\ell'^*(\lambda)$, is a linear function of λ , so it is easy to solve the corresponding approximate score equation, $\ell'^*(\lambda) = 0$, for λ :

$$\begin{split} \lambda &= \hat{\lambda}^* - \ell'(\hat{\lambda}^*) \cdot \left(\ell''(\hat{\lambda}^*)\right)^{-1} \\ &= 4.375 \end{split}$$

```
new_lambda_est <-
   cur_lambda_est -
   score(cur_lambda_est) * hessian(cur_lambda_est)^-1</pre>
```

```
plot2 = plot1 +
  geom_point(size = point_size,
             aes(x = new_lambda_est,
                 y = 0,
                 col = "new estimate")) +
  geom_segment(
    arrow = grid::arrow(),
   linewidth = 2,
    alpha = .7,
    aes(
      x = cur_lambda_est,
     y = approx_score(lhat = cur_lambda_est,
                       lambda = cur_lambda_est),
     xend = new_lambda_est,
     yend = 0,
      col = "update"
    )
  )
print(plot2)
```




Figure E.11.: score function of Dobson cyclone data and approximate score function $\,$

So we update $\hat{\lambda}^* \leftarrow 4.375$ and repeat our estimation process:

```
plot2 +
geom_function(
  fun = approx_score,
  aes(col = "new approximate score function"),
  n = 1001,
  args = list(lhat = new_lambda_est)) +
geom_point(
  size = point_size,
  aes(x = new_lambda_est, y = score(lambda = new_lambda_est),
```




Figure E.12.: score function of Dobson cyclone data and approximate score function

We repeat this process until the likelihood converges:

Compare with Exercise E.17

Table E.3.: Convergence of Newton-Raphson Algorithm for finding MLE of cyclone data

```
library(tibble)
cur_lambda_est = 3 # restarting
diff_loglik = Inf
tolerance = 10^{-4}
max_iter = 100
NR_info = tibble(
 iteration = 0,
 lambda = cur_lambda_est |> num(digits = 4),
 likelihood = lik(cur_lambda_est),
  `log(likelihood)` = loglik(cur_lambda_est) |> num(digits = 4),
 score = score(cur_lambda_est),
 hessian = hessian(cur_lambda_est)
)
for (cur_iter in 1:max_iter)
 new lambda est <-
   cur_lambda_est - score(cur_lambda_est) * hessian(cur_lambda_est) ^ -1
 diff_loglik = loglik(new_lambda_est) - loglik(cur_lambda_est)
 new_NR_info = tibble(
   iteration = cur_iter,
   lambda = new_lambda_est,
   likelihood = lik(new_lambda_est),
   `log(likelihood)` = loglik(new_lambda_est),
   score = score(new_lambda_est),
   hessian = hessian(new_lambda_est),
   `diff(loglik)` = diff_loglik
  )
 NR_info = NR_info |> bind_rows(new_NR_info)
  if (abs(diff_loglik) < tolerance)</pre>
   break
}
NR_info
#> # A tibble: 6 x 7
#> iteration lambda likelihood `log(likelihood)` score hessian `diff(loglik)`
       <dbl> <num:.> <dbl> <num:.4!> <dbl>
#>
                                                            <dbl>
#> 1
          0 3.0000 4.00e-18
                                        -40.0610 1.1 e+ 1
                                                            -8
                                                                      NA
```

-30.7708 3.46e+ 0

-3.76

#> 2

1 4.3750 4.33e-14

<dbl>

9.29e+ 0

```
11_plot +
  geom_segment(
    data = NR_info,
    arrow = grid::arrow(),
    # linewidth = 2,
    alpha = .7,
    aes(
        x = lambda,
        xend = lead(lambda),
        y = `log(likelihood)`,
        yend = lead(`log(likelihood)`),
        col = factor(iteration)
    )
)
```



Figure E.13.: Newton-Raphson algorithm for finding MLE of model E.6 for cyclone data

E.3. Maximum likelihood inference for univariate Gaussian models

Suppose $X_1,...,X_n \sim_{\mathrm{iid}} N(\mu,\sigma^2)$. Let $X=(X_1,...,X_n)^\top$ be these random variables in vector format. Let x_i and x denote the corresponding observed data. Then $\theta=(\mu,\sigma^2)$ is the vector of true parameters, and $\Theta=(\mathrm{M},\Sigma^2)$ is the vector of parameters as a random vector.

Then the log-likelihood is:

$$\begin{split} \ell & \propto -\frac{n}{2} \log \left\{ \sigma^2 \right\} - \frac{1}{2} \sum_{i=1}^n \frac{(x_i - \mu)^2}{\sigma^2} \\ & = -\frac{n}{2} \log \left\{ \sigma^2 \right\} - \frac{1}{2\sigma^2} \sum_{i=1}^n x_i^2 - 2x_i \mu + \mu^2 \end{split}$$

E.3.1. The score function

$$\ell'(x,\theta) \stackrel{\mathrm{def}}{=} \frac{\partial}{\partial \theta} \ell(x,\theta) = \left(\begin{array}{c} \frac{\partial}{\partial \mu} \ell(\theta;x) \\ \frac{\partial}{\partial \sigma^2} \ell(\theta;x) \end{array} \right) = \left(\begin{array}{c} \ell'_{\mu}(\theta;x) \\ \ell'_{\sigma^2}(\theta;x) \end{array} \right)$$

 $\ell'(x,\theta)$ is the function we set equal to 0 and solve to find the MLE:

$$\hat{\theta}_{ML} = \{\theta : \ell'(x, \theta) = 0\}$$

E.3.2. MLE of μ

$$\begin{split} \frac{d\ell}{d\mu} &= -\frac{1}{2} \sum_{i=1}^{n} \frac{-2(x_i - \mu)}{\sigma^2} \\ &= \frac{1}{\sigma^2} \left[\left(\sum_{i=1}^{n} x_i \right) - n\mu \right] \end{split}$$

If $\frac{d\ell}{d\mu} = 0$, then $\mu = \overline{x} \stackrel{\text{def}}{=} \frac{1}{n} \sum_{i=1}^{n} x_i$.

$$\frac{d^2\ell}{(d\mu)^2} = \frac{-n}{\sigma^2} < 0$$

So $\hat{\mu}_{ML} = \overline{x}$.

E.3.3. MLE of σ^2

• Reparametrizing the Gaussian distribution

When solving for $\hat{\sigma}_{ML}$, you can treat σ^2 as an atomic variable (don't differentiate with respect to σ or things get messy). In fact, you can replace σ^2 with $1/\tau$ and differentiate with respect to τ instead, and the process might be even easier.

$$\begin{split} \frac{d\ell}{d\sigma^2} &= \frac{\partial}{\partial \sigma^2} \left(-\frac{n}{2} \log \left\{ \sigma^2 \right\} - \frac{1}{2} \sum_{i=1}^n \frac{\left(x_i - \mu \right)^2}{\sigma^2} \right) \\ &= -\frac{n}{2} \left(\sigma^2 \right)^{-1} + \frac{1}{2} \left(\sigma^2 \right)^{-2} \sum_{i=1}^n \left(x_i - \mu \right)^2 \end{split}$$

If $\frac{d\ell}{d\sigma^2} = 0$, then:

$$\begin{split} \frac{n}{2} \left(\sigma^2\right)^{-1} &= \frac{1}{2} \left(\sigma^2\right)^{-2} \sum_{i=1}^n \left(x_i - \mu\right)^2 \\ \sigma^2 &= \frac{1}{n} \sum_{i=1}^n \left(x_i - \mu\right)^2 \end{split}$$

We plug in $\hat{\mu}_{ML} = \overline{x}$ to maximize globally (a technique called profiling):

$$\sigma_{ML}^2 = \frac{1}{n} \sum_{i=1}^{n} \left(x_i - \overline{x} \right)^2$$

Now:

$$\begin{split} \frac{d^2\ell}{\left(d\sigma^2\right)^2} &= \frac{\partial}{\partial\sigma^2} \left\{ -\frac{n}{2} \left(\sigma^2\right)^{-1} + \frac{1}{2} \left(\sigma^2\right)^{-2} \sum_{i=1}^n \left(x_i - \mu\right)^2 \right\} \\ &= \left\{ -\frac{n}{2} \frac{\partial}{\partial\sigma^2} \left(\sigma^2\right)^{-1} + \frac{1}{2} \frac{\partial}{\partial\sigma^2} \left(\sigma^2\right)^{-2} \sum_{i=1}^n \left(x_i - \mu\right)^2 \right\} \\ &= \left\{ \frac{n}{2} \left(\sigma^2\right)^{-2} - \left(\sigma^2\right)^{-3} \sum_{i=1}^n \left(x_i - \mu\right)^2 \right\} \\ &= \left(\sigma^2\right)^{-2} \left\{ \frac{n}{2} - \left(\sigma^2\right)^{-1} \sum_{i=1}^n \left(x_i - \mu\right)^2 \right\} \end{split}$$

Evaluated at $\mu = \overline{x}, \sigma^2 = \frac{1}{n} \sum_{i=1}^n \left(x_i - \overline{x}\right)^2$, we have:

$$\begin{split} \frac{d^2\ell}{\left(d\sigma^2\right)^2} &= \left(\hat{\sigma}^2\right)^{-2} \left\{ \frac{n}{2} - \left(\hat{\sigma}^2\right)^{-1} \sum_{i=1}^n \left(x_i - \overline{x}\right)^2 \right\} \\ &= \left(\hat{\sigma}^2\right)^{-2} \left\{ \frac{n}{2} - \left(\hat{\sigma}^2\right)^{-1} n \hat{\sigma}^2 \right\} \\ &= \left(\hat{\sigma}^2\right)^{-2} \left\{ \frac{n}{2} - n \right\} \\ &= \left(\hat{\sigma}^2\right)^{-2} n \left\{ \frac{1}{2} - 1 \right\} \\ &= \left(\hat{\sigma}^2\right)^{-2} n \left(-\frac{1}{2} \right) < 0 \end{split}$$

Finally, we have:

$$\begin{split} \frac{d^2\ell}{d\mu\;d\sigma^2} &= \frac{\partial}{\partial\mu} \left\{ -\frac{n}{2} \left(\sigma^2\right)^{-1} + \frac{1}{2} \left(\sigma^2\right)^{-2} \sum_{i=1}^n \left(x_i - \mu\right)^2 \right\} \\ &= \frac{1}{2} \left(\sigma^2\right)^{-2} \frac{\partial}{\partial\mu} \sum_{i=1}^n \left(x_i - \mu\right)^2 \\ &= \frac{1}{2} \left(\sigma^2\right)^{-2} \sum_{i=1}^n -2(x_i - \mu) \\ &= -\left(\sigma^2\right)^{-2} \sum_{i=1}^n \left(x_i - \mu\right) \end{split}$$

Evaluated at $\mu=\hat{\mu}=\overline{x},\sigma^2=\hat{\sigma}^2=\frac{1}{n}\sum_{i=1}^n\left(x_i-\overline{x}\right)^2$, we have:

$$\frac{d^2\ell}{d\mu\;d\sigma^2} = -\left(\hat{\sigma}^2\right)^{-2}(n\overline{x}-n\overline{x}) = 0$$

E.3.4. Covariance matrix

$$I = \begin{bmatrix} \frac{n}{\sigma^2} & 0\\ 0 & (\hat{\sigma}^2)^{-2} n\left(-\frac{1}{2}\right) \end{bmatrix} = \begin{bmatrix} a & 0\\ 0 & d \end{bmatrix}$$

So:

$$I^{-1} = \frac{1}{ad} \begin{bmatrix} d & 0 \\ 0 & a \end{bmatrix} = \begin{bmatrix} \frac{1}{a} & 0 \\ 0 & \frac{1}{d} \end{bmatrix}$$

$$I^{-1} = \begin{bmatrix} \frac{\widehat{\sigma}^2}{n} & 0 \\ 0 & \frac{2(\widehat{\sigma}^2)^2}{n} \end{bmatrix}$$

See Casella and Berger (2002) p322, example 7.2.12.

To prove it's a maximum, we need:

- $\ell' = 0$
- At least one diagonal element of ℓ'' is negative.
- Determinant of ℓ'' is positive.

E.4. Example: hormone therapy study

Now, we're going to analyze some real-world data using a Gaussian model, and then we're going to do a simulation to examine the properties of maximum likelihood estimation for that Gaussian model.

Here we look at the "heart and estrogen/progestin study" (HERS), a clinical trial of hormone therapy for prevention of recurrent heart attacks and death among 2,763 post-menopausal women with existing coronary heart disease (CHD) (Hulley et al. 1998).

We are going to model the distribution of fasting glucose among nondiabetics who don't exercise.

```
# load the data directly from a UCSF website
hers = haven::read_dta(
  paste0( # I'm breaking up the url into two chunks for readability
    "https://regression.ucsf.edu/sites/g/files",
    "/tkssra6706/f/wysiwyg/home/data/hersdata.dta"
  )
)
```

Table E.4.: HERS dataset

```
hers |> head()
#> # A tibble: 6 x 37
#>
     ΗТ
                     age raceth nonwhite smoking drinkany exercise physact globrat
     <dbl+lbl>
                   <dbl> <dbl+1> <dbl+1> <dbl+1> <dbl+1> <dbl+1> <dbl+1> <dbl+1>
#> 1 0 [placebo]
                      70 2 [Afr~ 1 [yes]
                                           0 [no] 0 [no]
                                                            0 [no]
                                                                      5 [muc~ 3 [goo~
#> 2 0 [placebo]
                      62 2 [Afr~ 1 [yes]
                                                            0 [no]
                                                                      1 [muc~ 3 [goo~
                                           0 [no]
                                                   0 [no]
#> 3 1 [hormone t~
                      69 1 [Whi~ 0 [no]
                                           0 [no]
                                                   0 [no]
                                                            0 [no]
                                                                      3 [abo~ 3 [goo~
                                                                      1 [muc~ 3 [goo~
#> 4 0 [placebo]
                      64 1 [Whi~ 0 [no]
                                           1 [yes] 1 [yes]
                                                            0 [no]
#> 5 0 [placebo]
                      65 1 [Whi~ 0 [no]
                                           0 [no]
                                                   0 [no]
                                                            0 [no]
                                                                      2 [som~ 3 [goo~
#> 6 1 [hormone t~
                      68 2 [Afr~ 1 [yes] 0 [no] 1 [yes]
                                                            0 [no]
                                                                      3 [abo~ 3 [goo~
#> # i 28 more variables: poorfair <dbl+lbl>, medcond <dbl>, htnmeds <dbl+lbl>,
       statins <dbl+lbl>, diabetes <dbl+lbl>, dmpills <dbl+lbl>,
#> #
#> #
       insulin <dbl+lbl>, weight <dbl>, BMI <dbl>, waist <dbl>, WHR <dbl>,
       glucose <dbl>, weight1 <dbl>, BMI1 <dbl>, waist1 <dbl>, WHR1 <dbl>,
#> #
       glucose1 <dbl>, tchol <dbl>, LDL <dbl>, HDL <dbl>, TG <dbl>, tchol1 <dbl>,
#> #
       LDL1 <dbl>, HDL1 <dbl>, TG1 <dbl>, SBP <dbl>, DBP <dbl>, age10 <dbl>
#> #
```

```
n.obs = 100 \text{ # we're going to take a small subset of the data to look at;}
# if we took the whole data set, the likelihood function would be hard to
# graph nicely
library(dplyr)
data1 =
  hers |>
  filter(
    diabetes == 0,
    exercise == 0) >
  head(n.obs)
glucose_data =
  data1 |>
  pull(glucose)
library(ggplot2)
library(ggeasy)
plot1 =
  data1 |>
  ggplot(aes(x = glucose)) +
  geom_histogram(aes(x = glucose, after_stat(density))) +
  theme_classic() +
  easy_labs()
print(plot1)
```



Looks somewhat plausibly Gaussian. Good enough for this example!

E.4.1. Find the MLEs


```
mu_hat = mean(glucose_data)
sigma_sq_hat = mean((glucose_data - mean(glucose_data))^2)
```

Our MLEs are:

- $\hat{\mu} = 98.66$
- $\hat{\sigma}^2 = 104.7444$

Here's the estimated distribution, superimposed on our histogram:

```
plot1 +
  geom_function(
   fun = function(x) dnorm(x, mean = mu_hat, sd = sqrt(sigma_sq_hat)),
   col = "red"
)
```



Looks like a somewhat decent fit? We could probably do better, but that's for another time.

E.4.2. Construct the likelihood and log-likelihood functions

it's often computationally more effective to construct the log-likelihood first and then exponentiate it to get the likelihood

```
loglik = function(
   mu, # I'm assigning default values, which the function will use
    # unless we tell it otherwise
    sigma = sd(x), # note that you can define some defaults based on other arguments
   x = glucose_data,
   n = length(x)
)
{
 normalizing_constants = -n/2 * \log((sigma^2) * 2 * pi)
  likelihood_kernel = - 1/(2 * sigma^2) *
      # I have to do this part in a somewhat complicated way
      # so that we can pass in vectors of possible values of mu
      # and get the likelihood for each value;
      # for the binomial case it's easier
      sum(x^2) - 2 * sum(x) * mu + n * mu^2
 answer = normalizing_constants + likelihood_kernel
 return(answer)
}
# `...` means pass any inputs to lik() along to loglik()
lik = function(...) exp(loglik(...))
```

E.4.3. Graph the Likelihood as a function of μ

```
(fixing \sigma^2 at \hat{\sigma}^2 = 104.7444)
```

```
ggplot() +
  geom_function(fun = function(x) lik(mu = x, sigma = sigma_sq_hat)) +
  xlim(mean(glucose_data) + c(-1,1) * sd(glucose_data)) +
  xlab("possible values of mu") +
  ylab("likelihood") +
  geom_vline(xintercept = mean(glucose_data), col = "red")
```



E.4.4. Graph the Log-likelihood as a function of μ

```
(fixing \sigma^2 at \hat{\sigma}^2 = 104.7444)
```

```
ggplot() +
  geom_function(fun = function(x) loglik(mu = x, sigma = sigma_sq_hat)) +
  xlim(mean(glucose_data) + c(-1,1) * sd(glucose_data)) +
```

```
xlab("possible values of mu") +
ylab('log(likelihood)') +
geom_vline(xintercept = mean(glucose_data), col = "red")
```



E.4.5. Likelihood and log-likelihood for σ , conditional on $\mu = \hat{\mu}$:

```
ggplot() +
  geom_function(fun = function(x) lik(sigma = x, mu = mean(glucose_data))) +
  xlim(sd(glucose_data) * c(.9,1.1)) +
  geom_vline(
     xintercept = sd(glucose_data) * sqrt(n.obs - 1)/sqrt(n.obs),
     col = "red") +
```

```
xlab("possible values for sigma") +
ylab('Likelihood')
```



```
ggplot() +
  geom_function(
    fun = function(x) loglik(sigma = x, mu = mean(glucose_data))
) +
  xlim(sd(glucose_data) * c(0.9, 1.1)) +
  geom_vline(
    xintercept =
      sd(glucose_data) * sqrt(n.obs - 1) / sqrt(n.obs),
    col = "red") +
  xlab("possible values for sigma") +
  ylab("log(likelihood)")
```



E.4.6. Standard errors by sample size:

```
se.mu.hat = function(n, sigma = sd(glucose_data)) sigma/sqrt(n)
ggplot() +
  geom_function(fun = se.mu.hat) +
  scale_x_continuous(trans = "log10", limits = c(10, 10^5), name = "Sample size") +
  ylab("Standard error of mu (mg/dl)") +
  theme_classic()
```



E.4.7. Simulations

E.4.7.1. Create simulation framework

Here's a function that performs a single simulation of a Gaussian modeling analysis:

```
do_one_sim = function(
    n = 100,
    mu = mean(glucose_data),
    mu0 = mean(glucose_data) * 0.9,
    sigma2 = var(glucose_data),
    return_data = FALSE # if this is set to true, we will create a list() containing
    # the analytic results and the vector of simulated data
)
```

```
{
  # generate data
  x = rnorm(n = 100, mean = mu, sd = sqrt(sigma2))
  # analyze data
  mu_hat = mean(x)
  sigmahat = sd(x)
  se_hat = sigmahat/sqrt(n)
  confint = mu_hat + c(-1, 1) * se_hat * qt(.975, df = n - 1)
  tstat = abs(mu_hat - mu0) / se_hat
  pval = pt(df = n - 1, q = tstat, lower = FALSE) * 2
  confint_covers = between(mu, confint[1], confint[2])
  test_rejects = pval < 0.05</pre>
  # if you want spaces, hyphens, or characters in your column names, use "", '', or
  to_return = tibble(
    "mu-hat" = mu_hat,
    "sigma-hat" = sigmahat,
    "se_hat" = se_hat,
    "confint_left" = confint[1],
    "confint_right" = confint[2],
    "tstat" = tstat,
    "pval" = pval,
    "confint covers true mu" = confint_covers,
    "test rejects null hypothesis" = test_rejects
  if(return_data)
   return(
      list(
        data = x,
```

```
results = to_return))
} else
{
   return(to_return)
}
```

Let's see what this function outputs for us:

```
do_one_sim()
#> # A tibble: 1 x 9
    `mu-hat` `sigma-hat` se_hat confint_left confint_right tstat
                                                                    pval
#>
       <dbl>
               <dbl> <dbl>
                                      <dbl>
                                                    <dbl> <dbl>
                                                                   <dbl>
#> 1
        99.3
                    9.92 0.992
                                       97.3
                                                     101. 10.6 6.74e-18
#> # i 2 more variables: `confint covers true mu` <lgl>,
    `test rejects null hypothesis` <lgl>
```

Looks good!

Now let's check it against the t.test() function from the stats package:

```
set.seed(1)
mu = mean(glucose_data)
mu0 = 80
sim.output = do_one_sim(mu0 = mu0, return_data = TRUE)
our_results =
   sim.output$results |>
   mutate(source = "`do_one_sim()`")

results_t.test = t.test(sim.output$data, mu = mu0)
```

```
results2 =
 tibble(
   source = "`stats::t.test()`",
    "mu-hat" = results_t.test$estimate,
   "sigma-hat" = results_t.test$stderr*sqrt(length(sim.output$data)),
    "se_hat" = results_t.test$stderr,
   confint_left = results_t.test$conf.int[1],
   confint_right = results_t.test$conf.int[2],
   tstat = results_t.test$statistic,
   pval = results_t.test$p.value,
    "confint covers true mu" = between(mu, confint_left, confint_right),
     `test rejects null hypothesis` = pval < 0.05
  )
comparison =
 bind_rows(
   our_results,
   results2
 ) |>
 relocate(
   "source",
    .before = everything()
  )
comparison
#> # A tibble: 2 x 10
    source
            `mu-hat` `sigma-hat` se_hat confint_left confint_right tstat
                                                                               pval
    <chr>
                 <dbl>
                            <dbl> <dbl>
                                                 <dbl>
                                                               <dbl> <dbl>
                                                                               <dbl>
#> 1 `do_one~
                              9.24 0.924
                                                                102. 21.4 6.23e-39
                  99.8
                                                  97.9
                              9.24 0.924
#> 2 `stats:~
                 99.8
                                                  97.9
                                                                102. 21.4 6.23e-39
#> # i 2 more variables: `confint covers true mu` <lgl>,
#> # `test rejects null hypothesis` <lgl>
```

Looks like we got it right!

E.4.7.2. Run 1000 simulations

Here's a function that calls the previous function n_sims times and summarizes the results:

```
do_n_sims = function(
   n_sims = 1000,
    ... # this symbol means "allow additional arguments to be passed on to the `do_s
)
{
  sim_results = NULL # we're going to create a "tibble" of results,
  # row by row (slightly different from the hint on the homework)
  for (i in 1:n_sims)
  {
    set.seed(i)
    current_results =
      do_one_sim(...) |> # here's where the simulation actually gets run
      mutate(
        sim_number = i
      ) |>
      relocate(sim_number, .before = everything())
    sim_results =
      sim_results |>
      bind_rows(current_results)
```

```
}
  return(sim_results)
sim_results = do_n_sims(
 n_sims = 100,
 mu = mean(glucose_data),
  sigma2 = var(glucose_data),
  n = 100 # this is the number of samples per simulated data set
sim_results |> head(10)
#> # A tibble: 10 x 10
#>
      sim_number `mu-hat` `sigma-hat` se_hat confint_left confint_right tstat
#>
           <int>
                    <dbl>
                                 <dbl> <dbl>
                                                     <dbl>
                                                                    <dbl> <dbl>
#>
   1
               1
                     99.8
                                  9.24 0.924
                                                      97.9
                                                                    102. 11.9
#>
   2
               2
                     98.3
                                11.9
                                        1.19
                                                      96.0
                                                                    101.
                                                                           8.00
   3
               3
                     98.8
                                  8.81 0.881
                                                      97.0
                                                                    101.
                                                                          11.3
#>
   4
               4
                                                                          11.6
#>
                     99.7
                                  9.40 0.940
                                                      97.8
                                                                    102.
   5
               5
                     99.0
                                  9.72 0.972
                                                      97.1
                                                                    101.
                                                                          10.5
#>
   6
               6
                     98.6
                                10.6
                                        1.06
                                                      96.4
                                                                    101.
                                                                           9.18
               7
                                                                    102. 11.5
#>
   7
                    100.
                                  9.86 0.986
                                                      98.1
#>
   8
               8
                     97.7
                                 11.1
                                                      95.5
                                                                     99.9 8.03
                                        1.11
#>
   9
               9
                     98.1
                                  9.86 0.986
                                                      96.2
                                                                    100.
                                                                           9.45
                     97.3
#> 10
              10
                                  9.68 0.968
                                                      95.3
                                                                     99.2 8.74
#> # i 3 more variables: pval <dbl>, `confint covers true mu` <lgl>,
       `test rejects null hypothesis` <lgl>
```

The simulation results are in! Now we have to analyze them.

E.4.7.3. Analyze simulation results

To do that, we write another function:

```
summarize_sim = function(
    sim_results,
    mu = mean(glucose_data),
    sigma2 = var(glucose_data),
    n = 100)
  # calculate the true standard error based on the data-generating parameters:
  `se(mu-hat)` = sqrt(sigma2/n)
  sim_results |>
    summarize(
      `bias[mu-hat]` = mean(`mu-hat`) - mu,
      `SE(mu-hat)` = sd(`mu-hat`),
      `bias[SE-hat]` = mean(se_hat) - `se(mu-hat)`,
      `SE(SE-hat)` = sd(se_hat),
      coverage = mean(`confint covers true mu`),
      power = mean(`test rejects null hypothesis`)
}
```

Let's try it out:

```
sim_summary = summarize_sim(
   sim_results,
   mu = mean(glucose_data),
```

```
# this function needs to know the true parameter values in order to assess bias
  sigma2 = var(glucose_data),
 n = 100
sim_summary
#> # A tibble: 1 x 6
     `bias[mu-hat]` `SE(mu-hat)` `bias[SE-hat]` `SE(SE-hat)` coverage power
#>
              <dbl>
                           <dbl>
                                           <dbl>
                                                        <dbl>
                                                                  <dbl> <dbl>
#> 1
            -0.0334
                           0.999
                                        -0.00826
                                                       0.0752
                                                                  0.98
```

From this simulation, we observe that our estimate of μ , $\hat{\mu}$, has minimal bias, and so does our estimate of $SE(\hat{\mu})$, $\hat{SE}(\hat{\mu})$.

The confidence intervals captured the true value even more often than they were supposed to, and the hypothesis test always rejected the null hypothesis.

I wonder what would happen with a different sample size, a different true μ value, or a different σ^2 value...

F. Common Mistakes

Configuring R

Functions from these packages will be used throughout this document:

```
library(conflicted) # check for conflicting function definitions
# library(printr) # inserts help-file output into markdown output
library(rmarkdown) # Convert R Markdown documents into a variety of formats.
library(pander) # format tables for markdown
library(ggplot2) # graphics
library(ggeasy) # help with graphics
library(ggfortify) # help with graphics
library(dplyr) # manipulate data
library(tibble) # `tibble`s extend `data.frame`s
library(magrittr) # `%>%` and other additional piping tools
library(haven) # import Stata files
library(knitr) # format R output for markdown
library(tidyr) # Tools to help to create tidy data
library(plotly) # interactive graphics
library(dobson) # datasets from Dobson and Barnett 2018
library(parameters) # format model output tables for markdown
library(haven) # import Stata files
```

F. Common Mistakes

```
library(latex2exp) # use LaTeX in R code (for figures and tables)
library(fs) # filesystem path manipulations
library(survival) # survival analysis
library(survminer) # survival analysis graphics
library(KMsurv) # datasets from Klein and Moeschberger
library(parameters) # format model output tables for
library(webshot2) # convert interactive content to static for pdf
library(forcats) # functions for categorical variables ("factors")
library(stringr) # functions for dealing with strings
library(lubridate) # functions for dealing with dates and times
```

Here are some R settings I use in this document:

```
rm(list = ls()) # delete any data that's already loaded into R


conflicts_prefer(dplyr::filter)
ggplot2::theme_set(
    ggplot2::theme_bw() +
        # ggplot2::labs(col = "") +
        ggplot2::theme(
        legend.position = "bottom",
        text = ggplot2::element_text(size = 12, family = "serif")))

knitr::opts_chunk$set(message = FALSE)
options('digits' = 4)

panderOptions("big.mark", ",")
pander::panderOptions("table.emphasize.rownames", FALSE)
pander::panderOptions("table.split.table", Inf)
conflicts_prefer(dplyr::filter) # use the `filter()` function from dplyr() by defaul
legend_text_size = 9
```

F.1. Parameters versus random variables

The parameters of a probability distribution shouldn't involve the random variables being modeled:



Solution.

$$\hat{\lambda}_{ML} \to_D N(\lambda, \lambda/n)$$

Expectations are means, not sums, despite the similarity of Σ and E. Really, we should use μ instead of E.

F.2. Quarto

F.2.1.

Make sure not to put a div ::: on the next line after a slide break ---:

--::: notes

There needs to be an empty line between them:

F. Common Mistakes

::: notes

F.2.2. library(printr) currently breaks df-print: paged

See https://github.com/yihui/printr/issues/41

F.3. LaTeX

Double superscript issues: $\label{lem:https://www.overleaf.com/learn/latex/Errors/Double_superscript}$

G. Notation

Table G.1.: Notation used in this book

| symbol | meaning | LaTeX |
|------------------|--|--------------|
| | not | \neg |
| \forall | all | \forall |
| ∃ | some | \exists |
| U | union, "or" | \cup |
| \cap | intersection, "and" | \cap |
| | given, conditional | \mid, |
| | on | |
| \sum | sum | \sum |
| $\overline{\Pi}$ | $\operatorname{product}$ | \prod |
| μ | mean | \mu |
| E | expectation | \mathbb{E} |
| <i>'</i> | transpose or | 1 |
| | $derivative^1$ | |
| | independent | |
| . . | therefore, thus | \therefore |
| η | linear component of a GLM^2 | \eta |

 $^{^1} depending on whether it is applied to a matrix or a function <math display="inline">^2 https://en.wikipedia.org/wiki/Generalized_linear_model#:~:text=The%20linear% 20predictor%20is%20the,data%20through%20the%20link%20function$

G.1. Information matrices

There is no consistency in the notation for observed and expected information.

book observed information expected information

G.2. The percent sign

The percent sign "%" is just a shorthand for " $\times \frac{1}{100}$ ". The word "percent" comes from the Latin "per centum"; "centum" means 100 in Latin, so "percent" means "per hundred" (c.f., https://en.wikipedia.org/wiki/Percentage)

So, contrary to what you may have learned previously, 10% = 0.1 is a true and correct equality.

Proof.

$$10\% = 10 \times \frac{1}{100}$$
$$= \frac{10}{100}$$
$$= 0.1$$

H. Statistical computing in R

There are an overwhelming number of great resources for learning R; here are some recommendations:

- Introduction to modern R: Wickham, Çetinkaya-Rundel, and Grolemund (2023)
- Advanced R programming: Wickham (2019)
- Examples of graphics: Chang (2024)
- Building R packages: Wickham and Bryan (2023)
- Translations from SAS: Kleinman and Horton (2009)
- UC Davis courses:
 - ENV 224, a.k.a. ECL 224¹
 - BIS $015L^2$

H.1. Functions

- Read this ASAP: https://r4ds.hadley.nz/functions.html
- Use this as a reference: https://adv-r.hadley.nz/functions.html

H.1.1. Methods versus functions

See https://adv-r.hadley.nz/oo.html#oop-systems

¹https://ucd-cepb.github.io/R-DAVIS/

²https://jmledford3115.github.io/datascibiol/

H.1.2. Debugging R and C code

See https://www.maths.ed.ac.uk/ \sim swood34/RCdebug/RCdebug.html

H.2. data.frames and tibbles

H.2.1. Displaying tibbles

See vignette("digits", package = "tibble")

H.3. The tidyverse

The tidyverse is an opinionated collection of R packages designed for data science. All packages share an underlying design philosophy, grammar, and data structures.

• https://www.tidyverse.org/

These packages are being actively developed by Hadley Wickham³ and his colleagues at posit⁴⁵.

Details:

- Wickham et al. (2019)
- Wickham, Çetinkaya-Rundel, and Grolemund (2023)
- Kuhn and Silge (2022)

³https://hadley.nz/

⁴https://posit.co/

⁵the company formerly known as RStudio⁶

H.4. Piping

See Wickham, Cetinkaya-Rundel, and Grolemund (2023)⁷ for details.

There are currently (2024) two commonly-used pipe operators in R:

- %>%: the "magrittr pipe", from the magrittr⁸ package (Bache and Wickham (2022); re-exported⁹ by $\tt dplyr^{10}$ and others).
- |>: the "native pipe", from base R (4.1.0)

H.4.1. Which pipe should I use?

Wickham, Çetinkaya-Rundel, and Grolemund (2023) recommends the native pipe¹¹:

For simple cases, |> and %>% behave identically. So why do we recommend the base pipe? Firstly, because it's part of base R, it's always available for you to use, even when you're not using the tidyverse. Secondly, |> is quite a bit simpler than %>%: in the time between the invention of %>% in 2014 and the inclusion of |> in R 4.1.0 in 2021, we gained a better understanding of the pipe. This allowed the base implementation to jettison infrequently used and less important features.

⁷https://r4ds.hadley.nz/data-transform.html#sec-the-pipe

 $^{^{8} \}rm https://cran.r-project.org/web/packages/magrittr/index.html$

⁹https://r-pkgs.org/dependencies-in-practice.html#re-exporting

¹⁰https://cran.r-project.org/web/packages/dplyr/index.html

 $^{^{11}}$ https://r4ds.hadley.nz/data-transform.html#sec-the-pipe:~:text=So%20why% 20do%20we%20recommend%20the%20base%20pipe%3F

H.4.2. Why doesn't ggplot2 use piping?

Here's tidyverse creator Hadley Wickham's answer (from 2018):

I think it's worth unpacking this question into a few smaller pieces:

- Should ggplot2 use the pipe? IMO, yes.
- Could ggplot2 support both the pipe and plus? No
- Would it be worth it to create a ggplot3 that uses the pipe? No.

https://forum.posit.co/t/why-cant-ggplot2-use/4372/7

H.5. Quarto

Quarto is a system for writing documents with embedded R code and/or results:

- Read this ASAP: https://r4ds.hadley.nz/communicate
- Then use this for reference: https://quarto.org/docs/reference/

H.6. Packages

This book espouses our philosophy of package development: anything that can be automated, should be automated. Do as little as possible by hand. Do as much as possible with functions. The goal is to spend your time thinking about what you want your package to do rather than thinking about the minutiae of package structure.

• $https://r-pkgs.org/introduction.html\#:\sim:text=This\%20book\%20espouses,of\%20package\%20struckers.$

H. Statistical computing in R

- Read this ASAP: https://r-pkgs.org/whole-game.html
- Use the rest of Wickham and Bryan (2023) as a reference

H.7. Git

94% of respondents to a 2022 Stack Overflow survey reported using git for version control link 12

More details¹³

- https://happygitwithr.com/
- https://usethis.r-lib.org/articles/pr-functions.html

H.8. Spatial data science

• Pebesma and Bivand (2023)

H.9. Shiny apps

- Read this first: Wickham (2021)
- Use this as a reference: Fay et al. (2021)

 $^{^{12} \}rm https://survey.stackoverflow.co/2022/\#section-version-control-version-control-systems$

 $^{^{13} \}rm https://r-pkgs.org/software-development-practices.html \#sec-sw-dev-practices-git-github$

H.10. Other resources for learning R

I. Contributing to rme

Contributions to these notes are very much appreciated; anything from one-character typo corrections to new chapters or rewrites. The GitHub repository for this project¹ provides a Pull Request system for submitting contributions. See https://happygitwithr.com/pr-extend for an explanation of the pull request system and the available R utility functions for working with pull requests.

I.1. Style guide

- Every abstract concept (definition or theorem) should have at least one concrete example immediately following it.
- More structure (headers, labels) is better.
- Make each conceptual chunk as compact as possible:
 - Decompose long derivations into smaller lemmas.
 - When manipulating part of a larger expression, isolate that part in a lemma.

 $^{^{1}}$ https://github.com/d-morrison/rme

I.2. Fixing typos

This book is written using Quarto². You can fix typos, spelling mistakes, or grammatical errors directly using the GitHub web interface by making changes in the corresponding *source* file. This generally means you'll need to edit a .qmd file.

I.3. Bigger changes

If you want to make a bigger change, it's a good idea to first file an issue and make sure someone from the development team agrees that it's needed.

I.3.1. Pull request³ process

- Fork the package and clone onto your computer. If you haven't done this before, we recommend using usethis::create_from_github("d-morrison/rme", fork = TRUE).
- Install all development dependencies with devtools::install_dev_deps(). Make sure you can build the book by running quarto render in a Terminal.
- Create a Git branch for your pull request (PR). We recommend using usethis::pr_init("brief-description-of-change"). Details at https://usethis.r-lib.org/articles/pr-functions.html
- Make your changes, commit to git, and then create a PR by running usethis::pr_push(), and following the prompts in your browser.
 The title of your PR should briefly describe the change. The body of your PR should contain Fixes #issue-number.

²https://quarto.org/docs/books/

³https://usethis.r-lib.org/articles/pr-functions.html#whats-a-pull-request

I. Contributing to rme

• Add a bullet to the top of NEWS.md (i.e. just below the first header). Follow the style described in https://style.tidyverse.org/news.html.

I.3.2. Code style

• New code should follow the tidyverse style guide⁴. You can use the styler⁵ package to apply these styles, but please don't restyle code that has nothing to do with your PR.

I.4. Code of Conduct

Please note that the rme project is released with a Contributor Code of Conduct⁶. By contributing to this project you agree to abide by its terms.

I.5. Additional references

For a detailed discussion on contributing to this and other projects, please see the Tidyverse development contributing guide⁷ and the Tidyverse code review principles⁸. This project is not part of the tidyverse, but we have borrowed their development processes.

⁴https://style.tidyverse.org

⁵https://CRAN.R-project.org/package=styler

⁶CODE OF CONDUCT.md

⁷https://rstd.io/tidy-contrib

⁸https://code-review.tidyverse.org/

Index

```
estimand, 490
estimate, 491
estimated value, 491
estimator, 491
expectation, 480
expected value, 480
```