

Supporting Information for: Caching in: A seed dispersal mutualism signals the decline of a songbird-pine system

Author List: Ana Miller-ter Kuile^{1,2}, Andreas P. Wion³, Kyle C. Rodman⁴, Jamie S. Sanderlin², Kiona Ogle¹

Author Affiliations:

¹ Northern Arizona University School of Informatics, Computing, and Cyber Systems, Flagstaff, Arizona, USA

² USDA Forest Service Rocky Mountain Research Station, Flagstaff, Arizona, USA

³ Forest Stewards Guild, Santa Fe, New Mexico, USA

⁴ Ecological Restoration Institute, Northern Arizona University, Flagstaff, Arizona, USA

Corresponding author: Ana Miller-ter Kuile, ana.miller-ter-kuile@nau.edu, Northern Arizona University School of Informatics, Computing, and Cyber Systems, 1295 Knoles Drive, Flagstaff, Arizona 86011

16 **SI Materials and Methods**

17 **Model Implementation, Convergence, and Diagnostics.** We implemented the above model in
18 the Bayesian software JAGS ¹ (version 4.3.0) using R ² (version 4.5.1) and the jagsUI wrapper
19 package ³ (version 1.6.2). JAGS uses Markov chain Monte Carlo (MCMC) methods to sample
20 from the joint posterior distribution of the model parameters. We initially produced 4000
21 samples for three parallel MCMC sequences to determine the number of iterations (samples)
22 needed to sufficiently sample from the posterior based on the Raftery diagnostic ⁴, and we re-ran
23 the model to obtain a sufficient number of iterations. We qualitatively assessed convergence of
24 the MCMC sequences by inspecting history and autocorrelation plots generated with the
25 mcmcplots package ⁵ (version 0.4.3). We quantitatively evaluated convergence using the
26 raftery.diag function in the coda package ⁶ (version 0.19.4.1); the MCMC sequences were
27 deemed to have converged if the Gelman-Rubin statistic, $\hat{R}$, for all root nodes in the model was
28 less than 1.1 ⁷. We assessed model goodness-of-fit by comparing observed eBird count data to
29 predicted count data based on simulating “replicated” count data from the Binomial distribution
30 in Equation (1). We tested for spatial dependence in the residuals (observed – predicted counts)
31 within each year using spline correlograms produced by the R package ncf ⁸ (version 1.3-2).

32 **Out of Sample Model Validation.** To evaluate predictions outside of the data used to fit the
33 model, we first removed the in-sample (“training”) data (22,295 checklists) from the available
34 eBird checklist data (833,834 checklists). We performed the same set of steps for spatial and
35 temporal stratification on the remaining checklists (811,539 checklists) for an additional 20,380
36 checklists that represented “out-of-sample” (“validation”) data. We evaluated out-of-sample
37 model fit using ~1000 posterior samples for covariate effects and intercepts from the in-sample

dataset and model. We then examined model replicative ability (RMSE) of the in-sample and out-of-sample datasets.

Variable Importance. We determined the relative contribution to model fit (R^2) of cones by running a model with no biological covariates (“null”) and a model excluding cone (seed) availability and its interactions (“climate and habitat” model) and tracking R^2 for each model. Based on R^2 improvement from the sequential addition of covariates (e.g., null vs. climate and habitat; climate and habitat vs. full model), we determined the relative contribution to the full model R^2 from adding the cone availability covariate in addition to habitat and climate variables (Figure 3).⁹


Software Used. We prepared data using the here ¹⁰ (version 1.0.1), tidyverse ¹¹ (version 2.0.0), sf ¹² (version 1.0.21), terra ¹³ (version 1.8.54), readxl ¹⁴ (version 1.4.5), exactextractr ¹⁵ (version 0.10.0), spatialEco ¹⁶ (version 2.0-2), ngeo ¹⁷ (version 0.4.8), auk ¹⁸ (version 0.8.2), lubridate ¹⁹ (version 1.9.4), prism ²⁰ (version 0.0.6), FNN ²¹ (version 1.1.4.1), and data.table ²² (version 1.17.6) packages.

SI Results

Model Performance and Goodness-of-fit. All stochastic model parameters converged with an $\hat{R} \leq 1.1$ (Fig. S3). Regressions of observed data versus predicted data (based on the posterior mean of 1000 samples of replicated data) yielded a mean R^2 of 0.17 (s.e. = 0.005). In general, the model over-estimated small pinyon jay counts but performed well for most other values (Fig. S4). The RMSE for our in-sample dataset had an average value of 3.25 (s.e. = 0.001) and the out-of-sample dataset had a value of 5.10 (s.e. = 0.0002; Fig. S5). Thus, our model predicted our in-sample data slightly better than the out-of-sample data, but neither RMSE values are large

considering the scale of the count data (counts per checklist ranged from 0 to 350), suggesting good model fit and predictive capacity. We observed no evidence for strong or consistent spatial autocorrelation in model residuals (Fig. S6 SI).

Covariates Influencing Detection. Pinyon jay detection probability was most influenced by the number of observers, such that fewer birds were observed on checklists involving more observers (observer number effect: -1.01 [-1.10, -1.04]). Checklists where observers walked at faster speeds had more detections (speed effect: 0.43 [0.40, 0.46]). Speed incorporated the distance (“effort”) and duration variables from the original checklists, and this effect is likely influenced by checklists where observers went further in a traveling checklist. Fewer birds were observed at later start times (time effect: -0.19 [-0.25, -0.14]) and there was a weak effect in which longer surveys had fewer detections (survey length effect: -0.03 [-0.07, -0.003]).



73
74 **Fig. S1.**Posterior estimates (median and 95% CI) for the seasonal importance weights for climate
75 (precipitation and maximum temperature; w terms, see equation 5 in the main text). The dashed
76 horizontal line indicates the ‘null’ weight in which all time periods would have equal weights.
77 Any weights that extend clearly above this line suggest significant importance of the
78 corresponding climate variable during that season to pinyon jay abundance. Seasons are based on
79 pinyon jay biology: Br (breeding; February - May), FI (fledging; May - June), Su (summer
80 localized foraging; July), and SW (summer and winter dispersed foraging or ‘irruption’ period;
81 August - January).

82









Fig. S2. Validation results supporting the use of space-for-time substitution of replicate checklists within sampling polygons. To use replicate eBird surveys within a year at a location in models such as N-mixture or occupancy models, checklists need to represent sites that display the full range of values for covariates of interest. In general, checklists in our dataset meet these criteria. Data represented here are values for all raster grid cells for all years (~1 million grid cells, depending on covariate). Thus, for the two covariates for which we are missing the upper tails, (a) cone availability and (e) pinyon basal area, these represent extreme outliers (<1% of total values in the dataset; ~20-200 raster grid cells, depending on covariate).



95

96 **Fig. S3.** Convergence statistics for stochastic parameters in the N-mixture for representing
 97 biological (α and α_0) and detection (γ and γ_0) covariate effects, deviance (an overall index of
 98 model fit), and the cone and climate importance weights (w_1 , w_2 , and w_3). All quantities
 99 converged with potential scale reduction ($\hat{R}$) values ≤ 1.1 (dashed line).




Fig. S4. As an evaluation of goodness-of-fit, we examined a) the relationship between observed pinyon jay counts and counts predicted from the model (median and 95% Bayesian Credible Interval) and b) the R^2 of this relationship from ~ 1000 posterior samples. In a), the black dashed line represents the 1:1 line; the pink line corresponds to the best-fit line between log-log predicted and observed bird count values. In b) the dashed vertical line represents the mean R^2 .












Fig. S5. Root mean squared error (RMSE) to evaluate the predictive ability of the model applied to the in-sample (“test”) and out-of-sample datasets. Values for RMSE are on the scale of the data (counts of pinyon jays per square kilometer). The RMSE values are relatively small compared to the range of possible counts in a checklist (0 to 350 birds per hectare in our test dataset).




114




115



9




10




118

11



12



119




Fig. S6. Evaluation of spatial autocorrelation (Pearson correlation on the y-axis) for residuals in each year (year indices are indicated above each figure panel, where 1 = 2010, 2 = 2011, ..., 13 = 2022). Distance values are in kilometers. There was no consistent evidence of spatial autocorrelation of residuals.

References

1. Plummer, M. JAGS: A program for analysis of Bayesian graphical models using Gibbs sampling. *Proceedings of the 3rd international workshop on distributed statistical computing* **124**, (2003).
2. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing (2020).
3. Kellner, K. jagsUI: a wrapper around ‘rjags’ to streamline JAGS analyses. (2021).
4. Raftery, A. E. & Lewis, S. M. The number of iterations, convergence diagnostics and generic Metropolis algorithms. in *Practical Markov Chain Monte Carlo* (Chapman and Hall, London, U.K., 1995).

- 137 5. Curtis, S. M. mcmcplots: Create Plots from MCMC Output. 0.4.3
138 <https://doi.org/10.32614/CRAN.package.mcmcplots> (2010).
- 139 6. Plummer, M. *et al.* coda: Output Analysis and Diagnostics for MCMC. 0.19-4.1
140 <https://doi.org/10.32614/CRAN.package.coda> (1999).
- 141 7. Gelman, A. & Rubin, D. B. Inference from Iterative Simulation Using Multiple Sequences.
142 *Statist. Sci.* **7**, (1992).
- 143 8. Bjornstad, O. N. ncf: Spatial Covariance Functions. The R Foundation
144 <https://doi.org/10.32614/cran.package.ncf> (2008).
- 145 9. Liu, Y., Schwalm, C. R., Samuels-Crow, K. E. & Ogle, K. Ecological memory of daily
146 carbon exchange across the globe and its importance in drylands. *Ecology Letters* **22**, 1806–
147 1816 (2019).
- 148 10. Muller, K. here: A simpler way to find your files. (2020).
- 149 11. Wickham, H. *et al.* Welcome to the Tidyverse. *JOSS* **4**, 1686 (2019).
- 150 12. Pebesma, E. & Bivand, R. *Spatial Data Science: With Applications in R.* (Chapman and
151 Hall/CRC, New York, 2023). doi:10.1201/9780429459016.
- 152 13. Hijmans, R. J. terra: Spatial Data Analysis. The R Foundation
153 <https://doi.org/10.32614/cran.package.terra> (2020).
- 154 14. Wickham, H. & Bryan, J. readxl: Read Excel Files. 1.4.3
155 <https://doi.org/10.32614/CRAN.package.readxl> (2015).
- 156 15. Daniel Baston. exactextractr: Fast Extraction from Raster Datasets using Polygons. The R
157 Foundation <https://doi.org/10.32614/cran.package.exactextractr> (2019).
- 158 16. Evans, J. S. & Murphy, M. A. spatialEco. (2023).

- 159 17. Dorman, M. ngeo: k-Nearest Neighbor Join for Spatial Data. The R Foundation
160 <https://doi.org/10.32614/cran.package.ngeo> (2018).
- 161 18. Strimas-Mackey, M., Miller, E. & Hochachka, W. auk: eBird Data Extraction and Processing
162 in R. The R Foundation <https://doi.org/10.32614/cran.package.auk> (2017).
- 163 19. Grolemund, G. & Wickham, H. Dates and Times Made Easy with **lubridate**. *J. Stat. Soft.* **40**,
164 (2011).
- 165 20. Hart, E. & Bell, K. prism: Access data from the Oregon State Prism climate project. Zenodo
166 <https://doi.org/10.5281/ZENODO.33663> (2015).
- 167 21. Beygelzimer, A. *et al.* FNN: Fast Nearest Neighbor Search Algorithms and Applications.
168 The R Foundation <https://doi.org/10.32614/cran.package.fnn> (2010).
- 169 22. Barrett, T. *et al.* data.table: Extension of `data.frame`. The R Foundation
170 <https://doi.org/10.32614/cran.package.data.table> (2006).