20.02.19

FP-62-NMR

We test the different operations on the p20 contol unit and adjust the 90° and 180° pulse with the help of an oscilloscope whiche we use to diplay the change of the Magnitization.

After adjusting the Pulses we start with measuring T_2 with Spin-Echo methode for Ga 500 and 600.
Then we proceed withe measuring T_2 with the Carr Purcell methode.
And at last we measure T_1 with the Spin-Echo-Methode again.

Table 1: Diffrent Relaxation Times

	$T_{2,SE}$ [ms]	$T_{2,CP}$ [ms]	$T_{1,SE}$ [ms]
Ga 500	62,4	66,1	68,1
Ga 600	78,6	~~86,4~~ 84,0	99,2

We continue with the chemical ~~shift~~ identification. After takeing all the Frequency profiles, we ~~try~~ match the peaks with the different parts of the ~~chemi~~ molecules. First identify that for each "+"-probe the most left peak is from TMS and use the relatis distance of the other peaks to identify the responsible part of the molecule. ~~We identified~~

Table 2: Identifying the probes with relative distance

	A+	B+	C+	D+	E+
Δ(TMS)-(1st)	2.2	2.1	2.0	3.9	2.6
Δ(TMS)-(3rd)	3.9	6.9	11.6	6.3	7.5
Δ(TMS)-(4th)	6.3				
→ Molecule	$FCH_2-CO-CH_3$	CH_3COOH	FCH_2-CN		
					

$\Delta T = 1.29\,\mu s$

$f_{HF} \sim 200\,MHz\,@\,B_0 = f_L$

$B_0 = \frac{f_L \cdot 2\pi}{\gamma}$

$B_1 = \frac{\alpha}{\gamma \cdot \Delta T}$

$\Delta E_{Res} = \sigma_{Halbwidth} \cdot h$

$\Delta E_{Dip} = \Delta f_{peak\,FWHM} \cdot h$

The difference is here calculated via

$$\delta = \frac{\omega_{res} - \omega_i(\text{i-th-peak})}{\omega_L (\approx 19.8\ \text{MHz} \cdot 2\pi)} \quad (1)$$

It should be mentioned that we didn't expect to se 2 peaks for FCH_2-CN or 3 peaks for $FCH_2-CO-CH_3$ because the resonce Freqeuncy for Fluor is a lot higher than the one we are using here.

But if you consides the Spin-Spin Interation From Fluor and the proton (hydrogen) we can have two different energy states of $\uparrow\uparrow$ and $\uparrow\downarrow$. Hence we can see two peaks for FCH_2- around 4 and 6.5.

We can also calculate the strength of the B-Fields with

$$B_0 = \frac{\omega_L}{\gamma} \quad (2)$$

$$B_1 = \frac{\alpha}{\Delta T \cdot \gamma} \quad (3)$$

with B_0 as static magnetic field used and B_1 the induced magnetic Field in the coil. $\gamma = 2.6 \cdot 10^8 \frac{1}{s \cdot T}$ is the gyromagnetic ratio of the nucleus for protons in this cas $f_L \approx f_{HF} = 19.8\ MHz$ is taken From the LabView program and ΔT for a $90°$-Pulse is identified with an oscilo

$\rightarrow \Delta T_{90°} = 1.29 \cdot 10^{-6}\ s$

$$\rightarrow B_0 = \frac{19.8 \cdot 2\pi \cdot 10^6}{2.6 \cdot 10^8} \cdot T = 0,48T$$

$$B_1 = \frac{\pi/2}{1.29 \cdot 10^{-6} \cdot 2.6 \cdot 10^8} T = 4.3\ mT$$

Lastly we can also deduce our Energy Resolution by looking at th width of a peak.

$$\Delta E_{res} = f_{FWHM} \cdot h \quad (4)$$

and calculate the Energy diffence for the different Fluor states.

$$\Delta E_{dipol} = f_{\Delta F} \cdot h \quad (5)$$

For $f_{\Delta F}$ we take from A+ the 3rd and 4th peak and calculate the Δf and we get $f_{\Delta F} = 48\,Hz$, $f_{FWHM} = 19.9\,Hz$ is taken from the 2nd Peak in A+ with the 2 coursors. and $h = 4.136\,eVs$.

$$\rightarrow \quad \Delta E_{res} \approx 8 \cdot 10^{-14}\,eV$$

$$\Delta E_{dipol} \approx 2 \cdot 10^{-13}\,eV$$

Next we start with the 1-d MRI and 2-d MRI. We use a Bruker minispec mq7.5 to take the images. We compare the way oil sinks through sand with the diffusion from a gas and we can conclude that it can be described with such model.

At last we take pictures with it from different objects.

T2_meas_freq.vi
C:\F61\sw\T2_meas_freq.vi
Last modified on 26.11.2018 at 15:49
Printed on 20.02.2019 at 10:28

Measurement of $T_{2,SE}$, Ga 500

Front Panel



T2_meas_freq.vi
C:\F61\sw\T2_meas_freq.vi
Last modified on 26.11.2018 at 15:49
Printed on 20.02.2019 at 10:44

Retaken -> check next one
(T2,SE Ga 600)

Front Panel



T2_meas_freq.vi
C:\F61\sw\T2_meas_freq.vi
Last modified on 26.11.2018 at 15:49
Printed on 20.02.2019 at 10:57

Measurment of $T_{2,SE}$, Ga 600

Front Panel



CP_meas_freq.vi
C:\F61\sw\CP_meas_freq.vi
Last modified on 16.10.2018 at 15:12
Printed on 20.02.2019 at 11:38

Measurment of $T_{2,CP}$ @ Ga500



CP_meas_freq.vi
C:\F61\sw\CP_meas_freq.vi
Last modified on 16.10.2018 at 15:12
Printed on 20.02.2019 at 11:21

Retaken because of low working frequency
Measurment of $T_{2,CP}$, Ca 600



CP_meas_freq.vi
C:\F61\sw\CP_meas_freq.vi
Last modified on 16.10.2018 at 15:12
Printed on 20.02.2019 at 11:26

Measurment of $T_{2,CP}$, ba 600



working frequency → 1000 Hz

T1_meas_freq.vi
C:\F61\sw\T1_meas_freq.vi
Last modified on 26.11.2018 at 16:30
Printed on 20.02.2019 at 12:10

Measurement of T_1 for Cu 500



We changed the amplitude from positive to negative for $\tau = 30$ ms

T1_meas_freq.vi

C:\F61\sw\T1_meas_freq.vi

Last modified on 26.11.2018 at 16:30

Printed on 20.02.2019 at 12:43

Measurement of $T_{1,SE}$, Ga 600



changed sign for τ = ~~33~~ 40 ms

chem_shift.vi
C:\F61\sw\chem_shift.vi
Last modified on 10.01.2019 at 14:36
Printed on 20.02.2019 at 15:36

A+ . 2



2: ~~1.8~~ 2.2

3: 3.9

4: 6.3

$FCH_2-CO-CH_3$

Dipole interaction with F-Atom:

$\Delta E = 48\ Hz \cdot h = 1.99 \cdot 10^{-13}\ eV$

↑

peak 4 - peak 3

Energy resolution:

we read the WDFH of a peak → $\Delta E \approx 8 \cdot 10^{-14}\ eV$

chem_shift.vi
C:\F61\sw\chem_shift.vi
Last modified on 10.01.2019 at 14:36
Printed on 20.02.2019 at 14:18

A+ Rebaken -> One Peak not fitted see A+.2



2 : ~~4.7~~ 1.7

3 : 4.1

$FCH_2-CO-CH_3$

chem_shift.vi
C:\F61\sw\chem_shift.vi
Last modified on 10.01.2019 at 14:36
Printed on 20.02.2019 at 14:04

B+



2: 7.1
3: 6.9



chem_shift.vi
C:\F61\sw\chem_shift.vi
Last modified on 10.01.2019 at 14:36
Printed on 20.02.2019 at 14:01

B



chem_shift.vi
C:\F61\sw\chem_shift.vi
Last modified on 10.01.2019 at 14:36
Printed on 20.02.2019 at 14:11

C+



2: 2.0
3: 11.6 CH_3COOH

chem_shift.vi
C:\F61\sw\chem_shift.vi
Last modified on 10.01.2019 at 14:36
Printed on 20.02.2019 at 14:05



chem_shift.vi
C:\F61\sw\chem_shift.vi
Last modified on 10.01.2019 at 14:36
Printed on 20.02.2019 at 14:13

D+



2: 3.9

3: 6.3

FCH_2-CN

chem_shift.vi
C:\F61\sw\chem_shift.vi
Last modified on 10.01.2019 at 14:36
Printed on 20.02.2019 at 14:12



chem_shift.vi
C:\F61\sw\chem_shift.vi
Last modified on 10.01.2019 at 14:36
Printed on 20.02.2019 at 14:17

E+



2: 2.6

3: 7.5



chem_shift.vi

C:\F61\sw\chem_shift.vi

Last modified on 10.01.2019 at 14:36

Printed on 20.02.2019 at 14:14



many_profiles.vi
C:\F62\sw\day3\many_profiles.vi
Last modified on 19.02.2019 at 10:50
Printed on 20.02.2019 at 16:15



many_profiles.vi
C:\F62\sw\day3\many_profiles.vi
Last modified on 19.02.2019 at 10:50
Printed on 20.02.2019 at 16:20



many_profiles.vi
C:\F62\sw\day3\many_profiles.vi
Last modified on 19.02.2019 at 10:50
Printed on 20.02.2019 at 16:57



nmr_2d.vi
C:\F62\sw\day3\nmr_2d.vi
Last modified on 25.01.2019 at 13:47
Printed on 21.02.2019 at 09:37

Pic. 3.1 15mm oil horizontal



nmr_2d.vi
C:\F62\sw\day3\nmr_2d.vi
Last modified on 25.01.2019 at 13:47
Printed on 21.02.2019 at 09:43

Pic 3.2 15 mm oil vertical back to front



capillar effect + distortion on the right side

nmr_2d.vi
C:\F62\sw\day3\nmr_2d.vi
Last modified on 25.01.2019 at 13:47
Printed on 21.02.2019 at 09:52

Pic 3.3 15 mm oil vertical left to right



nmr_2d.vi
C:\F62\sw\day3\nmr_2d.vi
Last modified on 25.01.2019 at 13:47
Printed on 21.02.2019 at 10:12

Pic 3.4 Celery (field of view 22,5 Avgs)