

Fourier Transform Theorems

- Addition Theorem
- Shift Theorem
- Convolution Theorem
- Similarity Theorem
- Rayleigh's Theorem
- Differentiation Theorem

Addition Theorem

$$\mathcal{F} \{f + g\} = F + G$$

Proof:

$$\begin{aligned}\mathcal{F} \{f + g\}(s) &= \int_{-\infty}^{\infty} [f(t) + g(t)] e^{-j2\pi st} dt \\&= \int_{-\infty}^{\infty} f(t) e^{-j2\pi st} dt + \int_{-\infty}^{\infty} g(t) e^{-j2\pi st} dt \\&= F(s) + G(s)\end{aligned}$$

Shift Theorem

$$\mathcal{F} \{f(t - t_0)\}(s) = e^{-j2\pi st_0} F(s)$$

Proof:

$$\mathcal{F} \{f(t - t_0)\}(s) = \int_{-\infty}^{\infty} f(t - t_0) e^{-j2\pi st} dt$$

Multiplying the r.h.s. by $e^{j2\pi st_0} e^{-j2\pi st_0} = 1$ yields:

$$\begin{aligned} & \mathcal{F} \{f(t - t_0)\}(s) \\ &= \int_{-\infty}^{\infty} f(t - t_0) e^{-j2\pi st} e^{j2\pi st_0} e^{-j2\pi st_0} dt \\ &= e^{-j2\pi st_0} \int_{-\infty}^{\infty} f(t - t_0) e^{-j2\pi s(t - t_0)} dt. \end{aligned}$$

Substituting $u = t - t_0$ and $du = dt$ yields:

$$\begin{aligned} \mathcal{F} \{f(t - t_0)\}(s) &= e^{-j2\pi st_0} \int_{-\infty}^{\infty} f(u) e^{-j2\pi su} du \\ &= e^{-j2\pi st_0} F(s). \end{aligned}$$

Shift Theorem Example

$$\begin{aligned}& \mathcal{F} \{ \sin(2\pi(t + 1/4)) \} (s) \\&= e^{\frac{j2\pi s}{4}} \mathcal{F} \{ \sin(2\pi t) \} \\&= e^{\frac{j\pi s}{2}} \cdot \frac{j}{2} [\delta(s + 1) - \delta(s - 1)] \\&= \frac{j}{2} \left[e^{\frac{j\pi s}{2}} \delta(s + 1) - e^{\frac{j\pi s}{2}} \delta(s - 1) \right] \\&= \frac{j}{2} \left[e^{\frac{j\pi(-1)}{2}} \delta(s + 1) - e^{\frac{j\pi(+1)}{2}} \delta(s - 1) \right] \\&= \frac{j}{2} [-j\delta(s + 1) - j\delta(s - 1)] \\&= \frac{1}{2} [\delta(s + 1) + \delta(s - 1)] \\&= \mathcal{F} \{ \cos(2\pi s) \}\end{aligned}$$

Shift Theorem (variation)

$$\mathcal{F}^{-1}\{F(s - s_0)\}(t) = e^{j2\pi s_0 t} f(t)$$

Proof:

$$\mathcal{F}^{-1}\{F(s - s_0)\}(t) = \int_{-\infty}^{\infty} F(s - s_0) e^{j2\pi s t} ds$$

Multiplying the r.h.s. by $e^{j2\pi s_0 t} e^{-j2\pi s_0 t} = 1$ yields:

$$\begin{aligned} & \mathcal{F}^{-1}\{F(s - s_0)\}(t) \\ &= \int_{-\infty}^{\infty} F(s - s_0) e^{j2\pi s t} e^{j2\pi s_0 t} e^{-j2\pi s_0 t} ds \\ &= e^{j2\pi s_0 t} \int_{-\infty}^{\infty} F(s - s_0) e^{j2\pi(s - s_0)t} ds. \end{aligned}$$

Substituting $u = s - s_0$ and $du = ds$ yields:

$$\begin{aligned} \mathcal{F}^{-1}\{F(s - s_0)\}(t) &= e^{j2\pi s_0 t} \int_{-\infty}^{\infty} F(u) e^{j2\pi u t} du \\ &= e^{j2\pi s_0 t} f(t). \end{aligned}$$

Convolution Theorem

$$\mathcal{F} \{f * g\} = F \cdot G$$

Proof:

$$\begin{aligned} & \mathcal{F} \{f * g\}(s) \\ &= \int_{-\infty}^{\infty} \left[\int_{-\infty}^{\infty} f(u)g(t-u)du \right] e^{-j2\pi st} dt \end{aligned}$$

Changing the order of integration:

$$\begin{aligned} & \mathcal{F} \{f * g\}(s) \\ &= \int_{-\infty}^{\infty} f(u) \left[\int_{-\infty}^{\infty} g(t-u)e^{-j2\pi st} dt \right] du \end{aligned}$$

By the Shift Theorem, we recognize that

$$\left[\int_{-\infty}^{\infty} g(t-u)e^{-j2\pi st} dt \right] = e^{-j2\pi su} G(s)$$

so that

$$\begin{aligned} \mathcal{F} \{f * g\}(s) &= \int_{-\infty}^{\infty} f(u)e^{-j2\pi su} G(s) du \\ &= G(s) \int_{-\infty}^{\infty} f(u)e^{-j2\pi su} du \\ &= G(s) \cdot F(s) \end{aligned}$$

Convolution Theorem Example

The pulse, Π , is defined as:

$$\Pi(t) = \begin{cases} 1 & \text{if } |t| \leq \frac{1}{2} \\ 0 & \text{otherwise.} \end{cases}$$

The triangular pulse, Λ , is defined as:

$$\Lambda(t) = \begin{cases} 1 - |t| & \text{if } |t| \leq 1 \\ 0 & \text{otherwise.} \end{cases}$$

It is straightforward to show that $\Lambda = \Pi * \Pi$.

Using this fact, we can compute $\mathcal{F} \{ \Lambda \}$:

$$\begin{aligned} \mathcal{F} \{ \Lambda \}(s) &= \mathcal{F} \{ \Pi * \Pi \}(s) \\ &= \mathcal{F} \{ \Pi \}(s) \cdot \mathcal{F} \{ \Pi \}(s) \\ &= \frac{\sin(\pi s)}{\pi s} \cdot \frac{\sin(\pi s)}{\pi s} \\ &= \frac{\sin^2(\pi s)}{\pi^2 s^2}. \end{aligned}$$

Convolution Theorem (variation)

$$\mathcal{F}^{-1}\{F * G\} = f \cdot g$$

Proof:

$$\begin{aligned} & \mathcal{F}^{-1}\{F * G\}(t) \\ &= \int_{-\infty}^{\infty} \left[\int_{-\infty}^{\infty} F(u)G(s-u)du \right] e^{j2\pi st} ds \end{aligned}$$

Changing the order of integration:

$$\begin{aligned} & \mathcal{F}^{-1}\{F * G\}(t) \\ &= \int_{-\infty}^{\infty} F(u) \left[\int_{-\infty}^{\infty} G(s-u)e^{j2\pi st} ds \right] du \end{aligned}$$

By the Shift Theorem, we recognize that

$$\left[\int_{-\infty}^{\infty} G(s-u)e^{j2\pi st} ds \right] = e^{j2\pi tu} g(t)$$

so that

$$\begin{aligned} \mathcal{F}^{-1}\{F * G\}(t) &= \int_{-\infty}^{\infty} F(u)e^{j2\pi tu} g(t) du \\ &= g(t) \int_{-\infty}^{\infty} F(u)e^{j2\pi tu} du \\ &= g(t) \cdot f(t) \end{aligned}$$

Similarity Theorem

$$\mathcal{F} \{f(at)\}(s) = \frac{1}{|a|} F\left(\frac{s}{a}\right)$$

Proof:

$$\mathcal{F} \{f(at)\}(s) = \int_{-\infty}^{\infty} f(at) e^{-j2\pi st} dt$$

There are two cases.

- $a > 0$. Multiplying the integral by $a/|a| = 1$ and the exponent by $a/a = 1$ yields:

$$\mathcal{F} \{f(at)\}(s) = \frac{1}{|a|} \int_{-\infty}^{\infty} f(at) e^{-j2\pi(s/a)at} a dt$$

We now make the substitution $u = at$ and $du = a dt$:

$$\begin{aligned} \mathcal{F} \{f(at)\}(s) &= \frac{1}{|a|} \int_{-\infty}^{\infty} f(u) e^{-j2\pi(s/a)u} du \\ &= \frac{1}{|a|} F\left(\frac{s}{a}\right) \end{aligned}$$

Similarity Theorem (contd.)

- $a < 0$. Multiplying the integral by $|a|/|a| = 1$ and the exponent by $-|a|/a = 1$ and using the fact that $a = -|a|$ yields:

$$\mathcal{F} \{f(at)\}(s) = \frac{1}{|a|} \int_{t=-\infty}^{t=\infty} f(-|a|t) e^{-j2\pi(s/a)(-|a|t)} |a| dt$$

We now make the substitution $u = -|a|t$ and $du = -|a|dt$:

$$\begin{aligned} \mathcal{F} \{f(at)\}(s) &= -\frac{1}{|a|} \int_{u=\infty}^{u=-\infty} f(u) e^{-j2\pi(s/a)u} du \\ &= \frac{1}{|a|} \int_{-\infty}^{\infty} f(u) e^{-j2\pi(s/a)u} du \\ &= \frac{1}{|a|} F\left(\frac{s}{a}\right) \end{aligned}$$

Similarity Theorem Example

Let's compute, $G(s)$, the Fourier transform of:

$$g(t) = e^{-t^2/9}.$$

We know that the Fourier transform of a Gaussian:

$$f(t) = e^{-\pi t^2}$$

is a Gaussian:

$$F(s) = e^{-\pi s^2}.$$

We also know that :

$$\mathcal{F} \{f(at)\}(s) = \frac{1}{|a|} F\left(\frac{s}{a}\right).$$

We need to write $g(t)$ in the form $f(at)$:

$$g(t) = f(at) = e^{-\pi(at)^2}.$$

Let $a = \frac{1}{3\sqrt{\pi}}$:

$$g(t) = e^{-t^2/9} = e^{-\pi\left(\frac{1}{3\sqrt{\pi}}t\right)^2} = f\left(\frac{1}{3\sqrt{\pi}}t\right).$$

It follows that:

$$G(s) = 3\sqrt{\pi} e^{-\pi(3\sqrt{\pi}s)^2}.$$

Rayleigh's Theorem

$$\int_{-\infty}^{\infty} |f(t)|^2 dt = \int_{-\infty}^{\infty} |F(s)|^2 ds$$

Proof:

$$\int_{-\infty}^{\infty} |f(t)|^2 dt = \int_{-\infty}^{\infty} f^*(t) f(t) dt$$

Substituting $\{\mathcal{F}^{-1}\{F\}\}^*(t)$ for $f^*(t)$:

$$\begin{aligned} &= \int_{-\infty}^{\infty} \left[\int_{-\infty}^{\infty} F(s) e^{j2\pi st} ds \right]^* f(t) dt \\ &= \int_{-\infty}^{\infty} \left[\int_{-\infty}^{\infty} F^*(s) e^{-j2\pi st} ds \right] f(t) dt \end{aligned}$$

Rayleigh's Theorem (contd.)

Changing the order of integration:

$$\begin{aligned} &= \int_{t=-\infty}^{t=\infty} \left[\int_{s=-\infty}^{s=\infty} F^*(s) e^{-j2\pi st} ds \right] f(t) dt \\ &= \int_{s=-\infty}^{s=\infty} F^*(s) \left[\int_{t=-\infty}^{t=\infty} e^{-j2\pi st} f(t) dt \right] ds \\ &= \int_{-\infty}^{\infty} F^*(s) F(s) ds \\ &= \int_{-\infty}^{\infty} |F(s)|^2 ds \end{aligned}$$

Differentiation Theorem

$$\mathcal{F} \{f'\}(s) = j2\pi s F(s)$$

Proof:

$$f = \mathcal{F}^{-1}\{F\}$$

Therefore

$$f(t) = \int_{-\infty}^{\infty} F(s) e^{j2\pi st} ds$$

Differentiating both sides with respect to t :

$$f'(t) = \int_{-\infty}^{\infty} j2\pi s F(s) e^{j2\pi st} ds$$

or

$$f' = \mathcal{F}^{-1}\{j2\pi s F(s)\}$$

Taking the Fourier transform of both sides:

$$\begin{aligned} \mathcal{F} \{f'\}(s) &= \mathcal{F} \{ \mathcal{F}^{-1}\{j2\pi s F(s)\} \} \\ &= j2\pi s F(s) \end{aligned}$$

Differentiation Theorem Example

$$\begin{aligned} & \mathcal{F} \left\{ \frac{d \sin(2\pi t)}{dt} \right\} (s) \\ &= j2\pi s \mathcal{F} \{ \sin(2\pi t) \} (s) \\ &= j2\pi s \cdot \frac{j}{2} [\delta(s+1) - \delta(s-1)] \\ &= -\pi s [\delta(s+1) - \delta(s-1)] \\ &= \pi [s\delta(s-1) - s\delta(s+1)] \\ &= \pi [(+1)\delta(s-1) - (-1)\delta(s+1)] \\ &= \pi [\delta(s-1) + \delta(s+1)] \\ &= \pi \mathcal{F} \{ 2 \cos(2\pi t) \} (s) \\ &= \mathcal{F} \{ 2\pi \cos(2\pi t) \} (s) \end{aligned}$$