

Week 2: Limit of Functions

July 30 – August 3, 2012

1 Limits of functions at a point

2 Limits at infinity

► Lecture 4

1 Limits of functions at a point

2 Limits at infinity

Meaning of limits

A function may or may not be defined at a particular point, but it can have a limiting value at that point.

Meaning of limits

A function may or may not be defined at a particular point, but it can have a limiting value at that point.

Let us consider the function $f(x) = \frac{x^2 - 1}{x - 1}$.

Meaning of limits

A function may or may not be defined at a particular point, but it can have a limiting value at that point.

Let us consider the function $f(x) = \frac{x^2 - 1}{x - 1}$.

x	0.9	0.999	1.01	1.0001
$\frac{x^2 - 1}{x - 1}$	1.9	1.999	2.01	2.0001

Meaning of limits

A function may or may not be defined at a particular point, but it can have a limiting value at that point.

Let us consider the function $f(x) = \frac{x^2 - 1}{x - 1}$.

x	0.9	0.999	1.01	1.0001
$\frac{x^2 - 1}{x - 1}$	1.9	1.999	2.01	2.0001

$f(x)$ is not defined at $x = 1$, but as x gets closer and closer to 1, $f(x)$ gets closer and closer to 2.

Meaning of limits

A function may or may not be defined at a particular point, but it can have a limiting value at that point.

Let us consider the function $f(x) = \frac{x^2 - 1}{x - 1}$.

x	0.9	0.999	1.01	1.0001
$\frac{x^2 - 1}{x - 1}$	1.9	1.999	2.01	2.0001

$f(x)$ is not defined at $x = 1$, but as x gets closer and closer to 1, $f(x)$ gets closer and closer to 2. We say that *the limit of $f(x)$ as x approaches to 1, is 2*.

Definition of limits

Definition

Let a be a real number. Suppose the function $y = f(x)$ is defined for all x near a on both sides. That is, for some real numbers b, c with $b < a < c$, the intervals (b, a) and (a, c) are in the domain of $y = f(x)$.

Definition of limits

Definition

Let a be a real number. Suppose the function $y = f(x)$ is defined for all x near a on both sides. That is, for some real numbers b, c with $b < a < c$, the intervals (b, a) and (a, c) are in the domain of $y = f(x)$. If there is a real number L such that $y = f(x)$ is as close to L as desired by taking x close ENOUGH to a on both sides of a , then L is the **limit** of $f(x)$ as x approaches a , written

$$\lim_{x \rightarrow a} f(x) = L.$$

Definition of limits

Definition

Let a be a real number. Suppose the function $y = f(x)$ is defined for all x near a on both sides. That is, for some real numbers b, c with $b < a < c$, the intervals (b, a) and (a, c) are in the domain of $y = f(x)$. If there is a real number L such that $y = f(x)$ is as close to L as desired by taking x close ENOUGH to a on both sides of a , then L is the **limit** of $f(x)$ as x approaches a , written

$$\lim_{x \rightarrow a} f(x) = L.$$

Example

$$\lim_{x \rightarrow 1} x^2 = 1.$$

Definition of limits

Remark

The limit is not affected by whether $f(a)$ is defined or what $f(a)$ is.

Definition of limits

Remark

The limit is not affected by whether $f(a)$ is defined or what $f(a)$ is.

Example

① Calculate $\lim_{x \rightarrow 1} \frac{x^2 - 1}{x - 1}$.

Definition of limits

Remark

The limit is not affected by whether $f(a)$ is defined or what $f(a)$ is.

Example

- ① Calculate $\lim_{x \rightarrow 1} \frac{x^2 - 1}{x - 1}$.

Solution: The domain is $(-\infty, 1) \cup (1, \infty)$. When $x \neq 1$,

$$\frac{x^2 - 1}{x - 1} = \frac{(x + 1)(x - 1)}{x - 1} = x + 1.$$

Definition of limits

Remark

The limit is not affected by whether $f(a)$ is defined or what $f(a)$ is.

Example

- ① Calculate $\lim_{x \rightarrow 1} \frac{x^2 - 1}{x - 1}$.

Solution: The domain is $(-\infty, 1) \cup (1, \infty)$. When $x \neq 1$,
$$\frac{x^2 - 1}{x - 1} = \frac{(x + 1)(x - 1)}{x - 1} = x + 1. \text{ Thus } \lim_{x \rightarrow 1} (x + 1) = 2.$$

Definition of limits

Remark

The limit is not affected by whether $f(a)$ is defined or what $f(a)$ is.

Example

- ① Calculate $\lim_{x \rightarrow 1} \frac{x^2 - 1}{x - 1}$.

Solution: The domain is $(-\infty, 1) \cup (1, \infty)$. When $x \neq 1$,
$$\frac{x^2 - 1}{x - 1} = \frac{(x + 1)(x - 1)}{x - 1} = x + 1. \text{ Thus } \lim_{x \rightarrow 1} (x + 1) = 2.$$

- ② Calculate $\lim_{x \rightarrow -1} \frac{2x + 2}{x + 1}$.

Definition of limits

Remark

The limit is not affected by whether $f(a)$ is defined or what $f(a)$ is.

Example

- ① Calculate $\lim_{x \rightarrow 1} \frac{x^2 - 1}{x - 1}$.

Solution: The domain is $(-\infty, 1) \cup (1, \infty)$. When $x \neq 1$,

$$\frac{x^2 - 1}{x - 1} = \frac{(x + 1)(x - 1)}{x - 1} = x + 1. \text{ Thus } \lim_{x \rightarrow 1} (x + 1) = 2.$$

- ② Calculate $\lim_{x \rightarrow -1} \frac{2x + 2}{x + 1}$.

Solution: The domain is $(-\infty, -1) \cup (-1, \infty)$. When $x \neq -1$,

$$\frac{2x + 2}{x + 1} = 2.$$

Definition of limits

Remark

The limit is not affected by whether $f(a)$ is defined or what $f(a)$ is.

Example

- ① Calculate $\lim_{x \rightarrow 1} \frac{x^2 - 1}{x - 1}$.

Solution: The domain is $(-\infty, 1) \cup (1, \infty)$. When $x \neq 1$,
$$\frac{x^2 - 1}{x - 1} = \frac{(x + 1)(x - 1)}{x - 1} = x + 1. \text{ Thus } \lim_{x \rightarrow 1} (x + 1) = 2.$$

- ② Calculate $\lim_{x \rightarrow -1} \frac{2x + 2}{x + 1}$.

Solution: The domain is $(-\infty, -1) \cup (-1, \infty)$. When $x \neq -1$,
$$\frac{2x + 2}{x + 1} = 2. \text{ Limit is } 2.$$

Limits of functions at a point

Examples.

③ if $f(x) = \begin{cases} \sin(x) & \text{if } x \neq 0 \\ 5 & \text{if } x = 0 \end{cases}$ then $\lim_{x \rightarrow 0} f(x) = 0.$

Limits of functions at a point

Examples.

③ if $f(x) = \begin{cases} \sin(x) & \text{if } x \neq 0 \\ 5 & \text{if } x = 0 \end{cases}$ then $\lim_{x \rightarrow 0} f(x) = 0$.

④ Sometimes, the limit does not exist.

$\lim_{x \rightarrow 0} \log_2 |x|$ does not exist.

Notation: When the limit $\lim_{x \rightarrow a} f(x)$ doesn't exist because $f(x)$ becomes arbitrarily large and positive (resp. negative) as x approaches a , we write $\lim_{x \rightarrow a} f(x) = \infty$ (resp.

$\lim_{x \rightarrow a} f(x) = -\infty$). Thus,

$$\lim_{x \rightarrow 0} \log_2 |x| = -\infty.$$

Limits of functions at a point

Examples.

③ if $f(x) = \begin{cases} \sin(x) & \text{if } x \neq 0 \\ 5 & \text{if } x = 0 \end{cases}$ then $\lim_{x \rightarrow 0} f(x) = 0$.

④ Sometimes, the limit does not exist.

$\lim_{x \rightarrow 0} \log_2 |x|$ does not exist.

Notation: When the limit $\lim_{x \rightarrow a} f(x)$ doesn't exist because $f(x)$ becomes arbitrarily large and positive (resp. negative) as x approaches a , we write $\lim_{x \rightarrow a} f(x) = \infty$ (resp.

$\lim_{x \rightarrow a} f(x) = -\infty$). Thus,

$$\lim_{x \rightarrow 0} \log_2 |x| = -\infty.$$

⑤ $\lim_{x \rightarrow 0} \frac{1}{x^2} = \infty$.

Definition of limits

Remark

One-sided limits:

- (i) The limit from the left: $\lim_{x \rightarrow a^-} f(x) = L.$
- (ii) The limit from the right: $\lim_{x \rightarrow a^+} f(x) = L.$

Definition of limits

Remark

One-sided limits:

- (i) The limit from the left: $\lim_{x \rightarrow a^-} f(x) = L$.
- (ii) The limit from the right: $\lim_{x \rightarrow a^+} f(x) = L$.

Example

$$\lim_{x \rightarrow 0} \frac{|x|}{x}$$

Solution: The domain is $(-\infty, 0) \cup (0, \infty)$.

Definition of limits

Remark

One-sided limits:

- (i) The limit from the left: $\lim_{x \rightarrow a^-} f(x) = L$.
- (ii) The limit from the right: $\lim_{x \rightarrow a^+} f(x) = L$.

Example

$$\lim_{x \rightarrow 0} \frac{|x|}{x}$$

Solution: The domain is $(-\infty, 0) \cup (0, \infty)$.

$$\frac{|x|}{x} = \begin{cases} +1 & x > 0 \\ -1 & x < 0. \end{cases}$$

Definition of limits

Remark

One-sided limits:

- (i) The limit from the left: $\lim_{x \rightarrow a^-} f(x) = L$.
- (ii) The limit from the right: $\lim_{x \rightarrow a^+} f(x) = L$.

Example

$$\lim_{x \rightarrow 0} \frac{|x|}{x}$$

Solution: The domain is $(-\infty, 0) \cup (0, \infty)$.

$$\frac{|x|}{x} = \begin{cases} +1 & x > 0 \\ -1 & x < 0. \end{cases}$$

$$\lim_{x \rightarrow 0^+} \frac{|x|}{x} = 1$$

Definition of limits

Remark

One-sided limits:

- (i) The limit from the left: $\lim_{x \rightarrow a^-} f(x) = L$.
- (ii) The limit from the right: $\lim_{x \rightarrow a^+} f(x) = L$.

Example

$$\lim_{x \rightarrow 0} \frac{|x|}{x}$$

Solution: The domain is $(-\infty, 0) \cup (0, \infty)$.

$$\frac{|x|}{x} = \begin{cases} +1 & x > 0 \\ -1 & x < 0. \end{cases}$$

$$\lim_{x \rightarrow 0^+} \frac{|x|}{x} = 1 \text{ and } \lim_{x \rightarrow 0^-} \frac{|x|}{x} = -1.$$

Existence of limits

Remark

The limit $\lim_{x \rightarrow a} f(x) = L$ exists if and only if

- (i) $\lim_{x \rightarrow a^-} f(x)$ and $\lim_{x \rightarrow a^+} f(x)$ exist; and

Existence of limits

Remark

The limit $\lim_{x \rightarrow a} f(x) = L$ exists if and only if

- (i) $\lim_{x \rightarrow a^-} f(x)$ and $\lim_{x \rightarrow a^+} f(x)$ exist; and
- (ii) $\lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^+} f(x) = L$.

Existence of limits

Remark

The limit $\lim_{x \rightarrow a} f(x) = L$ exists if and only if

- (i) $\lim_{x \rightarrow a^-} f(x)$ and $\lim_{x \rightarrow a^+} f(x)$ exist; and
- (ii) $\lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^+} f(x) = L$.

Example

$$\lim_{x \rightarrow 0} \frac{|x|}{x}$$

Existence of limits

Remark

The limit $\lim_{x \rightarrow a} f(x) = L$ exists if and only if

- (i) $\lim_{x \rightarrow a^-} f(x)$ and $\lim_{x \rightarrow a^+} f(x)$ exist; and
- (ii) $\lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^+} f(x) = L$.

Example

$$\lim_{x \rightarrow 0} \frac{|x|}{x}$$

As $\lim_{x \rightarrow 0^+} \frac{|x|}{x} = 1$ and $\lim_{x \rightarrow 0^-} \frac{|x|}{x} = -1$, $\lim_{x \rightarrow 0} \frac{|x|}{x}$ doesn't exist.

Existence of limits

Remark

The limit $\lim_{x \rightarrow a} f(x) = L$ exists if and only if

- (i) $\lim_{x \rightarrow a^-} f(x)$ and $\lim_{x \rightarrow a^+} f(x)$ exist; and
- (ii) $\lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^+} f(x) = L$.

Example

$$\lim_{x \rightarrow 0} \frac{|x|}{x}$$

As $\lim_{x \rightarrow 0^+} \frac{|x|}{x} = 1$ and $\lim_{x \rightarrow 0^-} \frac{|x|}{x} = -1$, $\lim_{x \rightarrow 0} \frac{|x|}{x}$ doesn't exist.

Remark

$L \neq \pm\infty$, otherwise, the limit doesn't exist.

Existence of limits

Example

$$\lim_{x \rightarrow 0} \frac{1}{x}$$

Existence of limits

Example

$$\lim_{x \rightarrow 0} \frac{1}{x}$$

Solution: The domain is $(-\infty, 0) \cup (0, \infty)$. As $x \rightarrow 0^+$, $\frac{1}{x}$ gets larger and larger, indicated by writing

$$\lim_{x \rightarrow 0^+} \frac{1}{x} = +\infty.$$

As ∞ is not a real number, we say $\lim_{x \rightarrow 0} \frac{1}{x}$ does not exist.

Existence of limits

Example

$$\lim_{x \rightarrow 0} \frac{1}{x}$$

Solution: The domain is $(-\infty, 0) \cup (0, \infty)$. As $x \rightarrow 0^+$, $\frac{1}{x}$ gets larger and larger, indicated by writing

$$\lim_{x \rightarrow 0^+} \frac{1}{x} = +\infty.$$

As ∞ is not a real number, we say $\lim_{x \rightarrow 0} \frac{1}{x}$ does not exist.

Remark

If $x = k$ is a vertical asymptote of $y = f(x)$, then $\lim_{x \rightarrow k} f(x)$ doesn't exist.

Rules for calculating limits

Basic limit rules. Suppose that $a \in \mathbb{R}$ and that $\lim_{x \rightarrow a} f(x)$ and $\lim_{x \rightarrow a} g(x)$ exist and are finite real numbers. Then

- (i) $\lim_{x \rightarrow a} (f(x) + g(x)) = \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x)$
- (ii) $\lim_{x \rightarrow a} (f(x) - g(x)) = \lim_{x \rightarrow a} f(x) - \lim_{x \rightarrow a} g(x)$
- (iii) $\lim_{x \rightarrow a} f(x)g(x) = \lim_{x \rightarrow a} f(x) \lim_{x \rightarrow a} g(x)$
- (iv) $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)}$, provided $\lim_{x \rightarrow a} g(x) \neq 0$.

A similar set of rules hold for left- and right-hand limits.

Rules for calculating limits

Limit rule for compositions.

If $\lim_{x \rightarrow a} f(x) = L$ and $\lim_{x \rightarrow L} g(x) = g(L)$ then

$$\lim_{x \rightarrow a} g(f(x)) = g\left(\lim_{x \rightarrow a} f(x)\right).$$

An easier way of remembering this rule will be given later.

Recall from last class


Basic limit rules. Suppose that $a \in \mathbb{R}$ and that $\lim_{x \rightarrow a} f(x)$ and $\lim_{x \rightarrow a} g(x)$ exist and are finite real numbers. Then

- (i) $\lim_{x \rightarrow a} (f(x) + g(x)) = \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x)$
- (ii) $\lim_{x \rightarrow a} (f(x) - g(x)) = \lim_{x \rightarrow a} f(x) - \lim_{x \rightarrow a} g(x)$
- (iii) $\lim_{x \rightarrow a} f(x)g(x) = \lim_{x \rightarrow a} f(x) \lim_{x \rightarrow a} g(x)$
- (iv) $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)}$, provided $\lim_{x \rightarrow a} g(x) \neq 0$.

A similar set of rules hold for left- and right-hand limits.

Rules for calculating limits

Example.




Use the limit laws and the graphs of f and g in the above figure to evaluate the following limits, if they exist.

(a) $\lim_{x \rightarrow -2} [f(x) + 5g(x)]$

Rules for calculating limits

Example.




Use the limit laws and the graphs of f and g in the above figure to evaluate the following limits, if they exist.

(a) $\lim_{x \rightarrow -2} [f(x) + 5g(x)]$

(b) $\lim_{x \rightarrow 1} [f(x)g(x)]$

Rules for calculating limits

Example.



Use the limit laws and the graphs of f and g in the above figure to evaluate the following limits, if they exist.

(a) $\lim_{x \rightarrow -2} [f(x) + 5g(x)]$

(b) $\lim_{x \rightarrow 1} [f(x)g(x)]$

(c) $\lim_{x \rightarrow 2} \frac{f(x)}{g(x)}$

1 Limits of functions at a point

2 Limits at infinity

Limits at infinity

There are many situations where it is desirable to know what the eventual state of a system will be.

Example: Suppose that an initially unpolluted lake containing 1 gigalitre (10^9 litres) of water has a river flowing through it at a rate of 1 megalitre (10^6 litres) per day. A factory is built next to the lake and discharges ten thousand (10^4) litres of pollution per day. Environmental authorities want to know what the eventual impact of the pollution will be. By making some simple assumptions, it can be shown that the amount $P(t)$ of pollution (in litres) in the lake after t days of the factory's operation is given by

$$P(t) = \frac{10^9}{101} (1 - e^{-101t/10^5}). \quad (1)$$

To see whether the pollution in the lake eventually stabilizes, and what the level of pollution will be, one studies the behavior of $P(t)$ as $t \rightarrow \infty$.

Calculating limits at infinity

Definition (imprecise).

if f is a function defined on an interval (a, ∞) , or $(-\infty, \infty)$, then

$$\lim_{x \rightarrow \infty} f(x) = L$$

means that the values of $f(x)$ can be made arbitrarily close to L by taking x sufficiently large.

Sometimes we also write

$$f(x) \longrightarrow L \quad \text{as} \quad x \longrightarrow \infty$$

Examples. True or False?

Examples. True or False?

(a) $\lim_{x \rightarrow \infty} x = \infty$

Examples. True or False?

(a) $\lim_{x \rightarrow \infty} x = \infty$

(b) $\lim_{x \rightarrow \infty} x^2 = \infty$

Examples. True or False?

(a) $\lim_{x \rightarrow \infty} x = \infty$

(b) $\lim_{x \rightarrow \infty} x^2 = \infty$

(c) $\lim_{x \rightarrow \infty} x^3 = \infty$

Examples. True or False?

(a) $\lim_{x \rightarrow \infty} x = \infty$

(b) $\lim_{x \rightarrow \infty} x^2 = \infty$

(c) $\lim_{x \rightarrow \infty} x^3 = \infty$

(d) $\lim_{x \rightarrow \infty} x^{-1} = \infty$

Examples. True or False?

(a) $\lim_{x \rightarrow \infty} x = \infty$

(b) $\lim_{x \rightarrow \infty} x^2 = \infty$

(c) $\lim_{x \rightarrow \infty} x^3 = \infty$

(d) $\lim_{x \rightarrow \infty} x^{-1} = \infty$

(e) $\lim_{x \rightarrow \infty} \sin(x) = 0$

Examples. True or False?

(a) $\lim_{x \rightarrow \infty} x = \infty$

(b) $\lim_{x \rightarrow \infty} x^2 = \infty$

(c) $\lim_{x \rightarrow \infty} x^3 = \infty$

(d) $\lim_{x \rightarrow \infty} x^{-1} = \infty$

(e) $\lim_{x \rightarrow \infty} \sin(x) = 0$

(f) $\lim_{x \rightarrow \infty} e^x = \infty$

Examples. True or False?

(a) $\lim_{x \rightarrow \infty} x = \infty$

(b) $\lim_{x \rightarrow \infty} x^2 = \infty$

(c) $\lim_{x \rightarrow \infty} x^3 = \infty$

(d) $\lim_{x \rightarrow \infty} x^{-1} = \infty$

(e) $\lim_{x \rightarrow \infty} \sin(x) = 0$

(f) $\lim_{x \rightarrow \infty} e^x = \infty$

(g) $\lim_{x \rightarrow \infty} \sqrt{x} = 2$

Examples. True or False?

- (a) $\lim_{x \rightarrow \infty} x = \infty$
- (b) $\lim_{x \rightarrow \infty} x^2 = \infty$
- (c) $\lim_{x \rightarrow \infty} x^3 = \infty$
- (d) $\lim_{x \rightarrow \infty} x^{-1} = \infty$
- (e) $\lim_{x \rightarrow \infty} \sin(x) = 0$
- (f) $\lim_{x \rightarrow \infty} e^x = \infty$
- (g) $\lim_{x \rightarrow \infty} \sqrt{x} = 2$
- (h) $\lim_{x \rightarrow \infty} \ln(x) = 100$

We have a similar definition for the limit as x goes to $-\infty$:
Let f be a function defined on some interval $(-\infty, a)$ or $(-\infty, \infty)$.
Then

$$\lim_{x \rightarrow -\infty} f(x) = L$$

means that the values of $f(x)$ can be made arbitrarily close to L by taking x sufficiently large and negative.

We have a similar definition for the limit as x goes to $-\infty$:
Let f be a function defined on some interval $(-\infty, a)$ or $(-\infty, \infty)$.
Then

$$\lim_{x \rightarrow -\infty} f(x) = L$$

means that the values of $f(x)$ can be made arbitrarily close to L by taking x sufficiently large and negative.

Question. What is the value of $\lim_{x \rightarrow -\infty} e^x$?

We have a similar definition for the limit as x goes to $-\infty$:
Let f be a function defined on some interval $(-\infty, a)$ or $(-\infty, \infty)$.
Then

$$\lim_{x \rightarrow -\infty} f(x) = L$$

means that the values of $f(x)$ can be made arbitrarily close to L by taking x sufficiently large and negative.

Question. What is the value of $\lim_{x \rightarrow -\infty} e^x$?

The line $y = L$ is called a *horizontal asymptote* of the curve $y = f(x)$ if either

$$\lim_{x \rightarrow \infty} f(x) = L \quad \text{or} \quad \lim_{x \rightarrow -\infty} f(x) = L.$$

Basic rules. Suppose that $\lim_{x \rightarrow \infty} f(x)$ and $\lim_{x \rightarrow \infty} g(x)$ exist and are finite real numbers. Then

$$(i) \lim_{x \rightarrow \infty} [f(x) + g(x)] = \lim_{x \rightarrow \infty} f(x) + \lim_{x \rightarrow \infty} g(x)$$

Basic rules. Suppose that $\lim_{x \rightarrow \infty} f(x)$ and $\lim_{x \rightarrow \infty} g(x)$ exist and are finite real numbers. Then

- (i) $\lim_{x \rightarrow \infty} [f(x) + g(x)] = \lim_{x \rightarrow \infty} f(x) + \lim_{x \rightarrow \infty} g(x)$
- (ii) $\lim_{x \rightarrow \infty} [f(x) - g(x)] = \lim_{x \rightarrow \infty} f(x) - \lim_{x \rightarrow \infty} g(x)$

Basic rules. Suppose that $\lim_{x \rightarrow \infty} f(x)$ and $\lim_{x \rightarrow \infty} g(x)$ exist and are finite real numbers. Then

- (i) $\lim_{x \rightarrow \infty} [f(x) + g(x)] = \lim_{x \rightarrow \infty} f(x) + \lim_{x \rightarrow \infty} g(x)$
- (ii) $\lim_{x \rightarrow \infty} [f(x) - g(x)] = \lim_{x \rightarrow \infty} f(x) - \lim_{x \rightarrow \infty} g(x)$
- (iii) $\lim_{x \rightarrow \infty} [f(x)g(x)] = [\lim_{x \rightarrow \infty} f(x)][\lim_{x \rightarrow \infty} g(x)]$

Basic rules. Suppose that $\lim_{x \rightarrow \infty} f(x)$ and $\lim_{x \rightarrow \infty} g(x)$ exist and are finite real numbers. Then

- (i) $\lim_{x \rightarrow \infty} [f(x) + g(x)] = \lim_{x \rightarrow \infty} f(x) + \lim_{x \rightarrow \infty} g(x)$
- (ii) $\lim_{x \rightarrow \infty} [f(x) - g(x)] = \lim_{x \rightarrow \infty} f(x) - \lim_{x \rightarrow \infty} g(x)$
- (iii) $\lim_{x \rightarrow \infty} [f(x)g(x)] = [\lim_{x \rightarrow \infty} f(x)][\lim_{x \rightarrow \infty} g(x)]$
- (iv) $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow \infty} f(x)}{\lim_{x \rightarrow \infty} g(x)}$, provided $\lim_{x \rightarrow \infty} g(x) \neq 0$.

Basic rules. Suppose that $\lim_{x \rightarrow \infty} f(x)$ and $\lim_{x \rightarrow \infty} g(x)$ exist and are finite real numbers. Then

- (i) $\lim_{x \rightarrow \infty} [f(x) + g(x)] = \lim_{x \rightarrow \infty} f(x) + \lim_{x \rightarrow \infty} g(x)$
- (ii) $\lim_{x \rightarrow \infty} [f(x) - g(x)] = \lim_{x \rightarrow \infty} f(x) - \lim_{x \rightarrow \infty} g(x)$
- (iii) $\lim_{x \rightarrow \infty} [f(x)g(x)] = [\lim_{x \rightarrow \infty} f(x)][\lim_{x \rightarrow \infty} g(x)]$
- (iv) $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow \infty} f(x)}{\lim_{x \rightarrow \infty} g(x)}$, provided $\lim_{x \rightarrow \infty} g(x) \neq 0$.
- (v) if $\lim_{x \rightarrow \infty} f(x) = \infty$, then $\lim_{x \rightarrow \infty} \frac{1}{f(x)} = 0$.

Basic rules. Suppose that $\lim_{x \rightarrow \infty} f(x)$ and $\lim_{x \rightarrow \infty} g(x)$ exist and are finite real numbers. Then

- (i) $\lim_{x \rightarrow \infty} [f(x) + g(x)] = \lim_{x \rightarrow \infty} f(x) + \lim_{x \rightarrow \infty} g(x)$
- (ii) $\lim_{x \rightarrow \infty} [f(x) - g(x)] = \lim_{x \rightarrow \infty} f(x) - \lim_{x \rightarrow \infty} g(x)$
- (iii) $\lim_{x \rightarrow \infty} [f(x)g(x)] = [\lim_{x \rightarrow \infty} f(x)][\lim_{x \rightarrow \infty} g(x)]$
- (iv) $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow \infty} f(x)}{\lim_{x \rightarrow \infty} g(x)}$, provided $\lim_{x \rightarrow \infty} g(x) \neq 0$.
- (v) if $\lim_{x \rightarrow \infty} f(x) = \infty$, then $\lim_{x \rightarrow \infty} \frac{1}{f(x)} = 0$.

Examples.

Basic rules. Suppose that $\lim_{x \rightarrow \infty} f(x)$ and $\lim_{x \rightarrow \infty} g(x)$ exist and are finite real numbers. Then

- (i) $\lim_{x \rightarrow \infty} [f(x) + g(x)] = \lim_{x \rightarrow \infty} f(x) + \lim_{x \rightarrow \infty} g(x)$
- (ii) $\lim_{x \rightarrow \infty} [f(x) - g(x)] = \lim_{x \rightarrow \infty} f(x) - \lim_{x \rightarrow \infty} g(x)$
- (iii) $\lim_{x \rightarrow \infty} [f(x)g(x)] = [\lim_{x \rightarrow \infty} f(x)][\lim_{x \rightarrow \infty} g(x)]$
- (iv) $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow \infty} f(x)}{\lim_{x \rightarrow \infty} g(x)}$, provided $\lim_{x \rightarrow \infty} g(x) \neq 0$.
- (v) if $\lim_{x \rightarrow \infty} f(x) = \infty$, then $\lim_{x \rightarrow \infty} \frac{1}{f(x)} = 0$.

Examples.

(a) $\lim_{x \rightarrow \infty} \left(\frac{1}{x} + 1 - e^{-2x} \right)$

Basic rules. Suppose that $\lim_{x \rightarrow \infty} f(x)$ and $\lim_{x \rightarrow \infty} g(x)$ exist and are finite real numbers. Then

- (i) $\lim_{x \rightarrow \infty} [f(x) + g(x)] = \lim_{x \rightarrow \infty} f(x) + \lim_{x \rightarrow \infty} g(x)$
- (ii) $\lim_{x \rightarrow \infty} [f(x) - g(x)] = \lim_{x \rightarrow \infty} f(x) - \lim_{x \rightarrow \infty} g(x)$
- (iii) $\lim_{x \rightarrow \infty} [f(x)g(x)] = [\lim_{x \rightarrow \infty} f(x)][\lim_{x \rightarrow \infty} g(x)]$
- (iv) $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow \infty} f(x)}{\lim_{x \rightarrow \infty} g(x)}$, provided $\lim_{x \rightarrow \infty} g(x) \neq 0$.
- (v) if $\lim_{x \rightarrow \infty} f(x) = \infty$, then $\lim_{x \rightarrow \infty} \frac{1}{f(x)} = 0$.

Examples.

- (a) $\lim_{x \rightarrow \infty} \left(\frac{1}{x} + 1 - e^{-2x} \right)$
- (b) $\lim_{x \rightarrow -\infty} \left(\frac{3}{x} + 1 - \cos(x) \right)$

Not all limits can be evaluated using the direct substitution property and the above rules.

Not all limits can be evaluated using the direct substitution property and the above rules.

The squeeze theorem. Suppose that $f(x) \leq g(x) \leq h(x)$ when x is near a (except possibly at a) and

$$\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} h(x) = L.$$

Then

$$\lim_{x \rightarrow a} g(x) = L.$$

Not all limits can be evaluated using the direct substitution property and the above rules.

The squeeze theorem. Suppose that $f(x) \leq g(x) \leq h(x)$ when x is near a (except possibly at a) and

$$\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} h(x) = L.$$

Then

$$\lim_{x \rightarrow a} g(x) = L.$$

(Similar versions of the squeeze theorem exist for left- and right-hand limits.)

Not all limits can be evaluated using the direct substitution property and the above rules.

The squeeze theorem. Suppose that $f(x) \leq g(x) \leq h(x)$ when x is near a (except possibly at a) and

$$\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} h(x) = L.$$

Then

$$\lim_{x \rightarrow a} g(x) = L.$$

(Similar versions of the squeeze theorem exist for left- and right-hand limits.)

Example. Find $\lim_{x \rightarrow 0} x^2 \sin(1/x)$.

The squeeze theorem for limits at infinity. Suppose that f , g and h are all defined on the interval (b, ∞) , where $b \in \mathbb{R}$. If

$$f(x) \leq g(x) \leq h(x) \quad \forall x \in (b, \infty)$$

and

$$\lim_{x \rightarrow \infty} f(x) = \lim_{x \rightarrow \infty} h(x) = L$$

then

$$\lim_{x \rightarrow \infty} g(x) = L.$$

The squeeze theorem for limits at infinity. Suppose that f , g and h are all defined on the interval (b, ∞) , where $b \in \mathbb{R}$. If

$$f(x) \leq g(x) \leq h(x) \quad \forall x \in (b, \infty)$$

and

$$\lim_{x \rightarrow \infty} f(x) = \lim_{x \rightarrow \infty} h(x) = L$$

then

$$\lim_{x \rightarrow \infty} g(x) = L.$$

Example. Use the squeeze theorem to find $\lim_{x \rightarrow \infty} \frac{\sin x}{x}$.

Indeterminate forms.

Limits of the type

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)}$$

where $f(x) \rightarrow \infty$ and $g(x) \rightarrow \infty$ as $x \rightarrow \infty$ are said to be limits of the form $\frac{\infty}{\infty}$.

Indeterminate forms.

Limits of the type

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)}$$

where $f(x) \rightarrow \infty$ and $g(x) \rightarrow \infty$ as $x \rightarrow \infty$ are said to be limits of the form $\frac{\infty}{\infty}$. While the following limits have the form $\frac{\infty}{\infty}$, each displays very different limiting behavior as $x \rightarrow \infty$:

Indeterminate forms.

Limits of the type

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)}$$

where $f(x) \rightarrow \infty$ and $g(x) \rightarrow \infty$ as $x \rightarrow \infty$ are said to be limits of the form $\frac{\infty}{\infty}$. While the following limits have the form $\frac{\infty}{\infty}$, each displays very different limiting behavior as $x \rightarrow \infty$:

- $$\lim_{x \rightarrow \infty} \frac{x}{x}$$

Indeterminate forms.

Limits of the type

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)}$$

where $f(x) \rightarrow \infty$ and $g(x) \rightarrow \infty$ as $x \rightarrow \infty$ are said to be limits of the form $\frac{\infty}{\infty}$. While the following limits have the form $\frac{\infty}{\infty}$, each displays very different limiting behavior as $x \rightarrow \infty$:

- $\lim_{x \rightarrow \infty} \frac{x}{x}$
- $\lim_{x \rightarrow \infty} \frac{x}{x^2}$

Indeterminate forms.

Limits of the type

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)}$$

where $f(x) \rightarrow \infty$ and $g(x) \rightarrow \infty$ as $x \rightarrow \infty$ are said to be limits of the form $\frac{\infty}{\infty}$. While the following limits have the form $\frac{\infty}{\infty}$, each displays very different limiting behavior as $x \rightarrow \infty$:

- $\lim_{x \rightarrow \infty} \frac{x}{x}$
- $\lim_{x \rightarrow \infty} \frac{x}{x^2}$
- $\lim_{x \rightarrow \infty} \frac{x^3}{x}$

Indeterminate forms.

Limits of the type

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)}$$

where $f(x) \rightarrow \infty$ and $g(x) \rightarrow \infty$ as $x \rightarrow \infty$ are said to be limits of the form $\frac{\infty}{\infty}$. While the following limits have the form $\frac{\infty}{\infty}$, each displays very different limiting behavior as $x \rightarrow \infty$:

- $\lim_{x \rightarrow \infty} \frac{x}{x}$
- $\lim_{x \rightarrow \infty} \frac{x}{x^2}$
- $\lim_{x \rightarrow \infty} \frac{x^3}{x}$
- $\lim_{x \rightarrow \infty} \frac{e^x}{x}$

(deal with this more later)