

Week 7: Fundamental Theorem of Calculus

September 3 – September 7, 2012

1 Fundamental Theorem of Calculus

- FTC and applications
- Evaluating integrals

► Lecture 14

Comparison properties of the integral.

Suppose that $a \leq b$.

- ① If $f(x) \geq 0$ for all x in $[a, b]$ then $\int_a^b f(x) dx \geq 0$.

Comparison properties of the integral.

Suppose that $a \leq b$.

① If $f(x) \geq 0$ for all x in $[a, b]$ then $\int_a^b f(x) dx \geq 0$.

② If $f(x) \leq g(x)$ for all x in $[a, b]$ then

$$\int_a^b f(x) dx \leq \int_a^b g(x) dx.$$

Comparison properties of the integral.

Suppose that $a \leq b$.

① If $f(x) \geq 0$ for all x in $[a, b]$ then $\int_a^b f(x) dx \geq 0$.

② If $f(x) \leq g(x)$ for all x in $[a, b]$ then

$$\int_a^b f(x) dx \leq \int_a^b g(x) dx.$$

③ If $m \leq f(x) \leq M$ for all x in $[a, b]$ then

$$m(b-a) \leq \int_a^b f(x) dx \leq M(b-a).$$

Comparison properties of the integral.

Suppose that $a \leq b$.

① If $f(x) \geq 0$ for all x in $[a, b]$ then $\int_a^b f(x) dx \geq 0$.

② If $f(x) \leq g(x)$ for all x in $[a, b]$ then

$$\int_a^b f(x) dx \leq \int_a^b g(x) dx.$$

③ If $m \leq f(x) \leq M$ for all x in $[a, b]$ then

$$m(b-a) \leq \int_a^b f(x) dx \leq M(b-a).$$

④ If $|f|$ is integrable on $[a, b]$ then

$$\left| \int_a^b f(x) dx \right| \leq \int_a^b |f(x)| dx.$$

1 Fundamental Theorem of Calculus

- FTC and applications
- Evaluating integrals

Functions defined by an integral

Before we discuss the Fundamental Theorem of Calculus, we introduce the notion of a function *defined by an integral*.

Functions defined by an integral

Before we discuss the Fundamental Theorem of Calculus, we introduce the notion of a function *defined by an integral*. Suppose we start with a continuous function f defined on the interval $[a, b]$, and we define a new function g on $[a, b]$ via the rule

$$g(x) = \int_a^x f(t) dt.$$

Functions defined by an integral

Before we discuss the Fundamental Theorem of Calculus, we introduce the notion of a function *defined by an integral*. Suppose we start with a continuous function f defined on the interval $[a, b]$, and we define a new function g on $[a, b]$ via the rule

$$g(x) = \int_a^x f(t) dt.$$

For example, $g(a) = \int_a^a f(t) dt = 0$, and $g(b) = \int_a^b f(t) dt$. If f is positive, then $g(x)$ can be interpreted as the area under the graph of f from a to x , the “area so far” function.

Theorem (Fundamental Theorem of Calculus)

Suppose a function f is continuous on $[a, b]$.

- ① *If $g(x) = \int_a^x f(t) dt$ for $x \in [a, b]$, $g'(x) = f(x)$ for all $x \in (a, b)$.*

Theorem (Fundamental Theorem of Calculus)

Suppose a function f is continuous on $[a, b]$.

- 1 If $g(x) = \int_a^x f(t) dt$ for $x \in [a, b]$, $g'(x) = f(x)$ for all $x \in (a, b)$.
- 2 If F is an antiderivative of f , then

$$\int_a^b f(x) dx = F(b) - F(a).$$

Theorem (Fundamental Theorem of Calculus)

Suppose a function f is continuous on $[a, b]$.

- 1 If $g(x) = \int_a^x f(t) dt$ for $x \in [a, b]$, $g'(x) = f(x)$ for all $x \in (a, b)$.
- 2 If F is an antiderivative of f , then

$$\int_a^b f(x) dx = F(b) - F(a).$$

Remark.

Note that part 1 says that if f is integrated, then differentiated, we get the original function f . Part 2 says that if we take a function F , differentiate it, then integrate the result, we get the original function F , but in the form $F(b) - F(a)$. Together, the two parts say that differentiation and integration are *inverse processes*.

Remarks on Part 1.

- (a) Part 1 tells us that every function defined by an integral is differentiable, and $g'(x) = f(x)$.

Remarks on Part 1.

- (a) Part 1 tells us that every function defined by an integral is differentiable, and $g'(x) = f(x)$.
- (b) Also, every continuous function f on (a, b) has an antiderivative g , where g is defined as the integral of f from a to x .

Remarks on Part 1.

- (a) Part 1 tells us that every function defined by an integral is differentiable, and $g'(x) = f(x)$.
- (b) Also, every continuous function f on (a, b) has an antiderivative g , where g is defined as the integral of f from a to x .

Examples. Find the derivative of g , where g is given by

Remarks on Part 1.

- (a) Part 1 tells us that every function defined by an integral is differentiable, and $g'(x) = f(x)$.
- (b) Also, every continuous function f on (a, b) has an antiderivative g , where g is defined as the integral of f from a to x .

Examples. Find the derivative of g , where g is given by

(i) $g(x) = \int_0^x \sqrt{1+t^2} dt.$

Remarks on Part 1.

- (a) Part 1 tells us that every function defined by an integral is differentiable, and $g'(x) = f(x)$.
- (b) Also, every continuous function f on (a, b) has an antiderivative g , where g is defined as the integral of f from a to x .

Examples. Find the derivative of g , where g is given by

- (i) $g(x) = \int_0^x \sqrt{1+t^2} dt.$
- (ii) $g(x) = \int_5^x \sqrt{1+t^2} dt.$

Remarks on Part 1.

- (a) Part 1 tells us that every function defined by an integral is differentiable, and $g'(x) = f(x)$.
- (b) Also, every continuous function f on (a, b) has an antiderivative g , where g is defined as the integral of f from a to x .

Examples. Find the derivative of g , where g is given by

- (i) $g(x) = \int_0^x \sqrt{1+t^2} dt.$
- (ii) $g(x) = \int_5^x \sqrt{1+t^2} dt.$
- (iii) $g(x) = \int_3^{17} \sqrt{1+t^2} dt.$

Remarks on Part 2.

Remarks on Part 2.

- (a) Note that the left hand side of the above equation is found by taking the limit of Riemann sums, which involves knowing the value of f at **all** points between a and b . The right hand side, however, only requires knowing the value of F at **two** points!

Remarks on Part 2.

- (a) Note that the left hand side of the above equation is found by taking the limit of Riemann sums, which involves knowing the value of f at **all** points between a and b . The right hand side, however, only requires knowing the value of F at **two** points!
- (b) A common misconception is that this formula is the *definition* of the definite integral. It is not, the definition of integral is the limit of Riemann sums.

Remarks on Part 2.

- (a) Note that the left hand side of the above equation is found by taking the limit of Riemann sums, which involves knowing the value of f at **all** points between a and b . The right hand side, however, only requires knowing the value of F at **two** points!
- (b) A common misconception is that this formula is the *definition* of the definite integral. It is not, the definition of integral is the limit of Riemann sums.
- (c) We use the notation $F(x)|_a^b = F(b) - F(a)$. Another common notation is $F(x)]_a^b$, which is used in the book.

Remarks on Part 2.

- (a) Note that the left hand side of the above equation is found by taking the limit of Riemann sums, which involves knowing the value of f at **all** points between a and b . The right hand side, however, only requires knowing the value of F at **two** points!
- (b) A common misconception is that this formula is the *definition* of the definite integral. It is not, the definition of integral is the limit of Riemann sums.
- (c) We use the notation $F(x)|_a^b = F(b) - F(a)$. Another common notation is $F(x)]_a^b$, which is used in the book.
- (d) When we apply the FTC, part 2, we may use *any* antiderivative of f (recall that if $F' = f$, the most general form for an antiderivative of f is $F + C$, where C is an arbitrary constant).

Remarks on Part 2.

Remarks on Part 2.

- (e) The FTC, part 2 is quite plausible from a physical perspective. Recall that when we considered the “Distance Problem” several lectures ago, we determined that the area under the velocity curve was equal to the total distance travelled. That is, if velocity and position are given by $v(t)$ and $s(t)$ respectively at time t , then $s'(t) = v(t)$, so s is an antiderivative of v , and $\int_a^b v(t) dt = s(b) - s(a)$.

Before we do some examples, we introduce the notion of an **indefinite integral**. This is just a new notation and name for an antiderivative.

Before we do some examples, we introduce the notion of an **indefinite integral**. This is just a new notation and name for an antiderivative.

Definition

If $F'(x) = f(x)$, we write $\int f(x) dx = F(x) + C$ and call the symbol $\int f(x) dx$ the **indefinite integral** of f , and C is called the **constant of integration**.

Before we do some examples, we introduce the notion of an **indefinite integral**. This is just a new notation and name for an antiderivative.

Definition

If $F'(x) = f(x)$, we write $\int f(x) dx = F(x) + C$ and call the symbol $\int f(x) dx$ the **indefinite integral** of f , and C is called the **constant of integration**.

Note that the definite integral is a *number*, whereas the indefinite integral is a *function*, or a *family of functions*. The Fundamental Theorem of Calculus, part 2 tells us that if f is continuous on $[a, b]$, then $\int_a^b f(x) dx = (\int f(x) dx)|_a^b$. Also note that an indefinite integral can represent either a particular antiderivative of f or an entire family of antiderivatives.

A table of indefinite integrals.

In the following table, c , C , and k are arbitrary constants.

f	$\int f(x) dx$
$cf(x)$	$c(\int f(x) dx) + C$
$f(x) \pm g(x)$	$(\int f(x) dx) \pm (\int g(x) dx) + C$
k	$kx + C$
x^n (where $n \neq -1$)	$\frac{x^{n+1}}{n+1} + C$
$\cos x$	$\sin x + C$
$\sin x$	$-\cos x + C$
$\sec^2 x$	$\tan x + C$
$\sec x \tan x$	$\sec x + C$

Examples. Evaluate the following integrals:

(i) $\int_{\pi}^{2\pi} \cos \theta \, d\theta$

Examples. Evaluate the following integrals:

(i) $\int_{\pi}^{2\pi} \cos \theta \, d\theta$

(ii) $\int_0^1 x(\sqrt[3]{x} + \sqrt[4]{x}) \, dx$

Examples. Evaluate the following integrals:

(i) $\int_{\pi}^{2\pi} \cos \theta \, d\theta$

(ii) $\int_0^1 x(\sqrt[3]{x} + \sqrt[4]{x}) \, dx$

(iii) True or False: $\int_0^{\pi} \sec^2 x \, dx = \tan x \Big|_0^{\pi} = 0$.

Examples. Evaluate the following integrals:

(i) $\int_{\pi}^{2\pi} \cos \theta \, d\theta$

(ii) $\int_0^1 x(\sqrt[3]{x} + \sqrt[4]{x}) \, dx$

(iii) True or False: $\int_0^{\pi} \sec^2 x \, dx = \tan x \Big|_0^{\pi} = 0$.

(iv) Find the indefinite integral: $\int (1 - t)(2 + t^2) \, dt$.

Applications.

Suppose f is continuous on $[a, b]$ and F is an antiderivative of f , that is, $F' = f$. Then the FTC, part 2 can be rewritten as

$$\int_a^b F'(x) dx = F(b) - F(a).$$

Applications.

Suppose f is continuous on $[a, b]$ and F is an antiderivative of f , that is, $F' = f$. Then the FTC, part 2 can be rewritten as

$$\int_a^b F'(x) dx = F(b) - F(a).$$

We know $F'(x)$ is the rate of change of $y = F(x)$ with respect to x , and $F(b) - F(a)$ is the *net* change in y when x changes from a to b . Thus the FTC part 2 says that the integral of a rate of change over an interval is the net change over that interval.

Applications.

Suppose f is continuous on $[a, b]$ and F is an antiderivative of f , that is, $F' = f$. Then the FTC, part 2 can be rewritten as

$$\int_a^b F'(x) dx = F(b) - F(a).$$

We know $F'(x)$ is the rate of change of $y = F(x)$ with respect to x , and $F(b) - F(a)$ is the *net* change in y when x changes from a to b . Thus the FTC part 2 says that the integral of a rate of change over an interval is the net change over that interval.

For instance,

- 1 If $V(t)$ is the volume of water in a tank at time t , then $V'(t)$ is the rate at which water flows into the tank at time t . Hence $\int_{t_1}^{t_2} V'(t) dt = V(t_2) - V(t_1)$ is the net change in the amount of water in the tank between times t_1 and t_2 .

Applications.

- ② If the rate of growth of a population is $\frac{dP}{dt}$, then $\int_{t_1}^{t_2} \frac{dP}{dt} dt = P(t_2) - P(t_1)$, the net change in population during time t_1 to t_2 .

Applications.

- ② If the rate of growth of a population is $\frac{dP}{dt}$, then $\int_{t_1}^{t_2} \frac{dP}{dt} dt = P(t_2) - P(t_1)$, the net change in population during time t_1 to t_2 .
- ③ If an object has position function s and velocity v , then $\int_{t_1}^{t_2} v(t) dt = s(t_2) - s(t_1)$, the net change of position (or *displacement*) of the particle during the period from t_1 to t_2 .

Recall from last class

Theorem (Fundamental Theorem of Calculus)

Suppose a function f is continuous on $[a, b]$.

- ① *If $g(x) = \int_a^x f(t) dt$ for $x \in [a, b]$, $g'(x) = f(x)$ for all $x \in (a, b)$.*

Recall from last class

Theorem (Fundamental Theorem of Calculus)

Suppose a function f is continuous on $[a, b]$.

- 1 If $g(x) = \int_a^x f(t) dt$ for $x \in [a, b]$, $g'(x) = f(x)$ for all $x \in (a, b)$.
- 2 If F is an antiderivative of f , then

$$\int_a^b f(x) dx = F(b) - F(a).$$

Examples

Find the derivative of g , where g is given by

Examples

Find the derivative of g , where g is given by

$$\textcircled{1} \quad g(x) = \int_1^x \sqrt{1+t^3} \, dt.$$

Examples

Find the derivative of g , where g is given by

① $g(x) = \int_1^x \sqrt{1+t^3} dt.$

② $g(x) = \int_1^{x^4} \sec t dt.$

Examples

Find the derivative of g , where g is given by

$$\textcircled{1} \quad g(x) = \int_1^x \sqrt{1+t^3} \, dt.$$

$$\textcircled{2} \quad g(x) = \int_1^{x^4} \sec t \, dt.$$

$$\textcircled{3} \quad g(x) = \int_2^{\frac{1}{x}} \sin^4 t \, dt.$$

Examples

Find the derivative of g , where g is given by

❶ $g(x) = \int_1^x \sqrt{1+t^3} dt.$

❷ $g(x) = \int_1^{x^4} \sec t dt.$

❸ $g(x) = \int_2^{\frac{1}{x}} \sin^4 t dt.$

❹ Evaluate the following integral:

$$\int_0^{\frac{\pi}{4}} \frac{1 + \cos^2 \theta}{\cos^2 \theta} d\theta$$

Examples.

Examples.

- ① If $w'(t)$ is the rate of growth of a child in pounds per year, what does $\int_5^{10} w'(t) dt$ represent?

Examples.

- 1 If $w'(t)$ is the rate of growth of a child in pounds per year, what does $\int_5^{10} w'(t) dt$ represent?
- 2 If oil leaks from a tank at a rate of $r(t)$ gallons per minute at time t , what does $\int_0^{120} r(t) dt$ represent?

Examples.

- 1 If $w'(t)$ is the rate of growth of a child in pounds per year, what does $\int_5^{10} w'(t) dt$ represent?
- 2 If oil leaks from a tank at a rate of $r(t)$ gallons per minute at time t , what does $\int_0^{120} r(t) dt$ represent?
- 3 A honeybee population starts with 100 bees and increases at a rate of $n'(t)$ bees per week. What does $100 + \int_0^{15} n'(t) dt$ represent?

Further Examples.

- (a) If $v(t) = 3t - 5$ gives the velocity of a particle moving along a line, find the displacement and the total distance traveled by the particle during the time interval $0 \leq t \leq 3$.

Further Examples.

- (a) If $v(t) = 3t - 5$ gives the velocity of a particle moving along a line, find the displacement and the total distance traveled by the particle during the time interval $0 \leq t \leq 3$.
- (b) Water flows from the bottom of a storage tank at a rate of $r(t) = 200 - 4t$ liters per minute, where $0 \leq t \leq 50$. Find the amount of water that flows from the tank during the first 10 minutes.

The average value of a function

The average value of finitely many numbers y_1, y_2, \dots, y_n is given by

$$y_{\text{ave}} = \frac{y_1 + y_2 + \dots + y_n}{n}.$$

In general, how do we calculate the average value of a function $y = f(x)$, where $a \leq x \leq b$?

The average value of a function

The average value of finitely many numbers y_1, y_2, \dots, y_n is given by

$$y_{\text{ave}} = \frac{y_1 + y_2 + \dots + y_n}{n}.$$

In general, how do we calculate the average value of a function $y = f(x)$, where $a \leq x \leq b$?

Definition

Suppose that f is continuous on $[a, b]$. Then the *average value* f_{ave} of f on $[a, b]$ is defined by the formula

$$f_{\text{ave}} = \frac{1}{b-a} \int_a^b f(x) dx.$$

The average value is sometimes denoted by \bar{f} .

The average value of a function

Example. Find the average value of the function f over $[-2, 3]$, where

$$f(x) = 1 + x^2.$$

Suppose that $T(t)$ is the temperature (in $^{\circ}\text{C}$) at time t (in hours) and that T_{ave} is the average temperature on the time interval $[0, 24]$. Is there a specific time t_0 in $[0, 24]$ when the temperature $T(t_0)$ is equal to the average temperature T_{ave} ? More generally, given a function f , is there a specific value c for which $f(c) = f_{\text{ave}}$?

Suppose that $T(t)$ is the temperature (in $^{\circ}\text{C}$) at time t (in hours) and that T_{ave} is the average temperature on the time interval $[0, 24]$. Is there a specific time t_0 in $[0, 24]$ when the temperature $T(t_0)$ is equal to the average temperature T_{ave} ? More generally, given a function f , is there a specific value c for which $f(c) = f_{\text{ave}}$?

The answer is yes! This is called the mean value theorem for integrals.

Suppose that $T(t)$ is the temperature (in $^{\circ}\text{C}$) at time t (in hours) and that T_{ave} is the average temperature on the time interval $[0, 24]$. Is there a specific time t_0 in $[0, 24]$ when the temperature $T(t_0)$ is equal to the average temperature T_{ave} ? More generally, given a function f , is there a specific value c for which $f(c) = f_{\text{ave}}$?

The answer is yes! This is called the mean value theorem for integrals.

Theorem (The Mean Value Theorem for integrals)

If f is continuous on $[a, b]$, then there exists a number c in $[a, b]$ such that

$$f(c) = f_{\text{ave}} = \frac{1}{b-a} \int_a^b f(x) dx,$$

that is,

$$\int_a^b f(x) dx = f(c)(b-a).$$

Example. Find all numbers c that satisfy the conclusion of the MVT for integrals when $f(x) = 1 + x^2$ and $[a, b] = [-2, 3]$.

Integration by substitution

To apply the FTC, we need to be able to find antiderivatives. Sometimes this is difficult, for instance, when we try to find the indefinite integral

$$\int 2x\sqrt{1+x^2} dx.$$

Integration by substitution

To apply the FTC, we need to be able to find antiderivatives. Sometimes this is difficult, for instance, when we try to find the indefinite integral

$$\int 2x\sqrt{1+x^2} dx.$$

However, there is a useful method called “integration by substitution” that we can apply. We illustrate it with the above example.

Integration by substitution

To apply the FTC, we need to be able to find antiderivatives. Sometimes this is difficult, for instance, when we try to find the indefinite integral

$$\int 2x\sqrt{1+x^2} dx.$$

However, there is a useful method called “integration by substitution” that we can apply. We illustrate it with the above example. If we let $u = 1 + x^2$, then $du = 2x dx$. Hence we may formally write

$$\begin{aligned}\int 2x\sqrt{1+x^2} dx &= \int \sqrt{1+x^2}(2x dx) \\ &= \int \sqrt{u} du = \frac{2}{3}u^{3/2} + C \\ &= \frac{2}{3}(x^2 + 1)^{3/2} + C\end{aligned}$$

Integration by substitution

The substitution rule

If $u = g(x)$ is a differentiable function whose range is an interval I and f is continuous on I , then

$$\int f(g(x))g'(x) dx = \int f(u) du.$$

That is, *it is permissible to work with dx and du after integral signs as if they are differentials.*

Integration by substitution

The substitution rule

If $u = g(x)$ is a differentiable function whose range is an interval I and f is continuous on I , then

$$\int f(g(x))g'(x) dx = \int f(u) du.$$

That is, *it is permissible to work with dx and du after integral signs as if they are differentials.*

Examples.

(i) Evaluate $\int \sqrt{2x+1} dx$.

Integration by substitution

The substitution rule

If $u = g(x)$ is a differentiable function whose range is an interval I and f is continuous on I , then

$$\int f(g(x))g'(x) dx = \int f(u) du.$$

That is, *it is permissible to work with dx and du after integral signs as if they are differentials.*

Examples.

- (i) Evaluate $\int \sqrt{2x+1} dx$.
- (ii) Evaluate $\int \frac{x}{\sqrt{1-4x^2}} dx$.

Integration by substitution

The substitution rule

If $u = g(x)$ is a differentiable function whose range is an interval I and f is continuous on I , then

$$\int f(g(x))g'(x) dx = \int f(u) du.$$

That is, *it is permissible to work with dx and du after integral signs as if they are differentials.*

Examples.

- (i) Evaluate $\int \sqrt{2x+1} dx$.
- (ii) Evaluate $\int \frac{x}{\sqrt{1-4x^2}} dx$.
- (iii) Evaluate $\int_1^2 \frac{1}{(3-5x)^2} dx$.

Symmetries

Symmetries

Suppose f is integrable on $[-a, a]$, where $a > 0$. Then

- ❶ If f is even ($f(-x) = f(x)$), then

$$\int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx.$$

Symmetries

Symmetries

Suppose f is integrable on $[-a, a]$, where $a > 0$. Then

- ❶ If f is even ($f(-x) = f(x)$), then

$$\int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx.$$

- ❷ If f is odd ($f(-x) = -f(x)$), then

$$\int_{-a}^a f(x) dx = 0.$$

Midterm Review

§3.7, #46 A car braked with a constant deceleration of 16 ft/s^2 , producing skid marks measuring 200 ft before coming to a stop. How fast was the car travelling when the brakes were first applied?

Midterm Review

§3.7, #46 A car braked with a constant deceleration of 16 ft/s^2 , producing skid marks measuring 200 ft before coming to a stop. How fast was the car travelling when the brakes were first applied?

§2.5, #36 Find the derivative of the function

$$y = \sqrt{x + \sqrt{x + \sqrt{x}}}.$$