

## Week 9: The natural logarithm and exponential functions

October 2 – October 5, 2012

## 1 The natural logarithm & exponential

- The natural logarithm
- The exponential function

[▶ Lecture 17](#)[▶ Lecture 18](#)

- 1 The natural logarithm & exponential
  - The natural logarithm
  - The exponential function

# Recall from last class

## Definition

The function  $\ln$  with domain  $(0, \infty)$  is defined by the formula

$$\ln(x) = \int_1^x \frac{1}{t} dt.$$

# Recall from last class

## Definition

The function  $\ln$  with domain  $(0, \infty)$  is defined by the formula

$$\ln(x) = \int_1^x \frac{1}{t} dt.$$

By interpreting  $\ln$  as the area under a curve, we can see that:

# Recall from last class

## Definition

The function  $\ln$  with domain  $(0, \infty)$  is defined by the formula

$$\ln(x) = \int_1^x \frac{1}{t} dt.$$

By interpreting  $\ln$  as the area under a curve, we can see that:

- $\ln x > 0$  if  $x > 1$  and  $\ln 1 = 0$ ,

# Recall from last class

## Definition

The function  $\ln$  with domain  $(0, \infty)$  is defined by the formula

$$\ln(x) = \int_1^x \frac{1}{t} dt.$$

By interpreting  $\ln$  as the area under a curve, we can see that:

- $\ln x > 0$  if  $x > 1$  and  $\ln 1 = 0$ ,
- $\ln x < 0$  if  $0 < x < 1$ .

# Recall from last class

## Definition

The function  $\ln$  with domain  $(0, \infty)$  is defined by the formula

$$\ln(x) = \int_1^x \frac{1}{t} dt.$$

By interpreting  $\ln$  as the area under a curve, we can see that:

- $\ln x > 0$  if  $x > 1$  and  $\ln 1 = 0$ ,
- $\ln x < 0$  if  $0 < x < 1$ .

We have the following (familiar) properties of the logarithm:



# Recall from last class

## Definition

The function  $\ln$  with domain  $(0, \infty)$  is defined by the formula

$$\ln(x) = \int_1^x \frac{1}{t} dt.$$

By interpreting  $\ln$  as the area under a curve, we can see that:

- $\ln x > 0$  if  $x > 1$  and  $\ln 1 = 0$ ,
- $\ln x < 0$  if  $0 < x < 1$ .

We have the following (familiar) properties of the logarithm:

- (i)  $\ln(xy) = \ln(x) + \ln(y)$  for all positive real numbers  $x$  and  $y$ .

# Recall from last class

## Definition

The function  $\ln$  with domain  $(0, \infty)$  is defined by the formula

$$\ln(x) = \int_1^x \frac{1}{t} dt.$$

By interpreting  $\ln$  as the area under a curve, we can see that:

- $\ln x > 0$  if  $x > 1$  and  $\ln 1 = 0$ ,
- $\ln x < 0$  if  $0 < x < 1$ .

We have the following (familiar) properties of the logarithm:

- (i)  $\ln(xy) = \ln(x) + \ln(y)$  for all positive real numbers  $x$  and  $y$ .
- (ii)  $\ln\left(\frac{x}{y}\right) = \ln(x) - \ln(y)$  for all positive real numbers  $x$  and  $y$ .

# Recall from last class

## Definition

The function  $\ln$  with domain  $(0, \infty)$  is defined by the formula

$$\ln(x) = \int_1^x \frac{1}{t} dt.$$

By interpreting  $\ln$  as the area under a curve, we can see that:

- $\ln x > 0$  if  $x > 1$  and  $\ln 1 = 0$ ,
- $\ln x < 0$  if  $0 < x < 1$ .

We have the following (familiar) properties of the logarithm:

- (i)  $\ln(xy) = \ln(x) + \ln(y)$  for all positive real numbers  $x$  and  $y$ .
- (ii)  $\ln\left(\frac{x}{y}\right) = \ln(x) - \ln(y)$  for all positive real numbers  $x$  and  $y$ .
- (iii)  $\ln(x^r) = r \ln(x)$  whenever  $r$  is a rational number and  $x$  is a positive real number.

**Graphing the natural logarithm.** The natural logarithm has the following limits:

**Graphing the natural logarithm.** The natural logarithm has the following limits:

(a)  $\lim_{x \rightarrow \infty} \ln x = \infty.$

**Graphing the natural logarithm.** The natural logarithm has the following limits:

(a)  $\lim_{x \rightarrow \infty} \ln x = \infty.$

(b)  $\lim_{x \rightarrow 0^+} \ln x = -\infty.$

**Graphing the natural logarithm.** The natural logarithm has the following limits:

(a)  $\lim_{x \rightarrow \infty} \ln x = \infty.$

(b)  $\lim_{x \rightarrow 0^+} \ln x = -\infty.$

Also, if  $y = \ln x$  and  $x > 0$ ,

$$\frac{dy}{dx} = \frac{1}{x} > 0 \quad \text{and} \quad \frac{d^2y}{dx^2} = -\frac{1}{x^2} < 0.$$

**Graphing the natural logarithm.** The natural logarithm has the following limits:

(a)  $\lim_{x \rightarrow \infty} \ln x = \infty.$

(b)  $\lim_{x \rightarrow 0^+} \ln x = -\infty.$

Also, if  $y = \ln x$  and  $x > 0$ ,

$$\frac{dy}{dx} = \frac{1}{x} > 0 \quad \text{and} \quad \frac{d^2y}{dx^2} = -\frac{1}{x^2} < 0.$$

Hence  $\ln x$  is increasing and concave downward. Also,  $\ln 1 = 0$  and  $\ln x$  is continuous, so the graph looks like this:



**Graphing the natural logarithm.** The natural logarithm has the following limits:

(a)  $\lim_{x \rightarrow \infty} \ln x = \infty.$


(b)  $\lim_{x \rightarrow 0^+} \ln x = -\infty.$

Also, if  $y = \ln x$  and  $x > 0$ ,

$$\frac{dy}{dx} = \frac{1}{x} > 0 \quad \text{and} \quad \frac{d^2y}{dx^2} = -\frac{1}{x^2} < 0.$$

Hence  $\ln x$  is increasing and concave downward. Also,  $\ln 1 = 0$  and  $\ln x$  is continuous, so the graph looks like this:

Graph for  $\ln(x)$



**Important definition.** By the intermediate value theorem, there is a number where  $\ln x$  equals 1. We denote this number by  $e$  (for “Euler”). It is (in my opinion) perhaps the most important number in mathematics. It is irrational, and **approximately** equal to 2.718.

**Important definition.** By the intermediate value theorem, there is a number where  $\ln x$  equals 1. We denote this number by  $e$  (for “Euler”). It is (in my opinion) perhaps the most important number in mathematics. It is irrational, and **approximately** equal to 2.718.

**Example.** Differentiate the function given by  $y = \ln(\sin x)$ .

**Remarks.**

- In general,  $\frac{d}{dx} \ln(g(x)) = \frac{g'(x)}{g(x)}.$

## Remarks.

- In general,  $\frac{d}{dx} \ln(g(x)) = \frac{g'(x)}{g(x)}$ .
- Note that when we consider function like  $\ln(g(x))$ , we are assuming that this composite function is defined on some interval  $I$  such that the range of  $g$ , when restricted to  $I$ , lies in the domain of definition of  $\ln$ . *The book does not emphasize this, which can be somewhat confusing.*

## Remarks.

- In general,  $\frac{d}{dx} \ln(g(x)) = \frac{g'(x)}{g(x)}$ .
- Note that when we consider function like  $\ln(g(x))$ , we are assuming that this composite function is defined on some interval  $I$  such that the range of  $g$ , when restricted to  $I$ , lies in the domain of definition of  $\ln$ . *The book does not emphasize this, which can be somewhat confusing.*

**Example.** Suppose  $f$  is defined by  $f(x) = \ln|x|$ . What is  $f'(x)$ ?

We record this formula:

$$\frac{d}{dx} \ln |x| = \frac{1}{x}.$$

Hence, we obtain the formula

$$\int \frac{1}{x} dx = \ln |x| + C.$$

We record this formula:

$$\frac{d}{dx} \ln |x| = \frac{1}{x}.$$

Hence, we obtain the formula

$$\int \frac{1}{x} dx = \ln |x| + C.$$

More generally, we can use the chain rule to derive:

$$\frac{d}{dx} \ln |f(x)| = \frac{f'(x)}{f(x)}$$



We record this formula:

$$\frac{d}{dx} \ln |x| = \frac{1}{x}.$$

Hence, we obtain the formula

$$\int \frac{1}{x} dx = \ln |x| + C.$$

More generally, we can use the chain rule to derive:

$$\frac{d}{dx} \ln |f(x)| = \frac{f'(x)}{f(x)}$$

hence,

$$\int \frac{f'(x)}{f(x)} dx = \ln |f(x)| + C$$

provided that  $f$  is differentiable and restricted to an interval  $I$  where it is nonzero.

**Example.** Evaluate  $\int \frac{x}{x^2+1} dx$

**Example.** Evaluate  $\int \frac{x}{x^2+1} dx$

**Example.** Evaluate  $\int \tan x dx$ .

**Example.** Evaluate  $\int \frac{x}{x^2+1} dx$

**Example.** Evaluate  $\int \tan x dx$ . (This example appears in the book on Page 259. Note that although the book doesn't specify the domain of  $x$ , it is assumed to be an interval  $I$  where  $\tan x$  is defined.)

# Logarithmic differentiation

By taking the logarithm, any problem involving powers, products and quotients can be transformed into a problem involving products, sums and differences. This is a powerful strategy if it is easy to manipulate the resulting logarithms, as is the case with differentiation.

# Logarithmic differentiation

By taking the logarithm, any problem involving powers, products and quotients can be transformed into a problem involving products, sums and differences. This is a powerful strategy if it is easy to manipulate the resulting logarithms, as is the case with differentiation.

**Example.** Find  $\frac{dy}{dx}$  if  $y = \left( \frac{(3x^2 + 4)(x + 2)}{x^3 + 5x} \right)^{3/5}$  and  $x > 0$ .

## The exponential function

Since  $\frac{d}{dx} \ln x = \frac{1}{x} > 0$  on  $(0, \infty)$ ,  $\ln x$  is an increasing differentiable function, hence it has a differentiable inverse, which we denote by  $\exp$ . The domain of  $\exp$  is  $\mathbb{R}$  and the range of  $\exp$  is  $(0, \infty)$ .

## The exponential function

Since  $\frac{d}{dx} \ln x = \frac{1}{x} > 0$  on  $(0, \infty)$ ,  $\ln x$  is an increasing differentiable function, hence it has a differentiable inverse, which we denote by  $\exp$ . The domain of  $\exp$  is  $\mathbb{R}$  and the range of  $\exp$  is  $(0, \infty)$ .

### The graph of $\exp$ .

Reflect the graph of  $\ln$  about the line  $y = x$ .



## The exponential function

Since  $\frac{d}{dx} \ln x = \frac{1}{x} > 0$  on  $(0, \infty)$ ,  $\ln x$  is an increasing differentiable function, hence it has a differentiable inverse, which we denote by  $\exp$ . The domain of  $\exp$  is  $\mathbb{R}$  and the range of  $\exp$  is  $(0, \infty)$ .

### The graph of $\exp$ .

Reflect the graph of  $\ln$  about the line  $y = x$ .

Notice that

- $\exp(0) = 1$  since  $\ln(1) = 0$ .

## The exponential function

Since  $\frac{d}{dx} \ln x = \frac{1}{x} > 0$  on  $(0, \infty)$ ,  $\ln x$  is an increasing differentiable function, hence it has a differentiable inverse, which we denote by  $\exp$ . The domain of  $\exp$  is  $\mathbb{R}$  and the range of  $\exp$  is  $(0, \infty)$ .

### The graph of $\exp$ .

Reflect the graph of  $\ln$  about the line  $y = x$ .

Notice that

- $\exp(0) = 1$  since  $\ln(1) = 0$ .
- $\exp(1) = e$  since  $\ln(e) = 1$ .

## The exponential function

Since  $\frac{d}{dx} \ln x = \frac{1}{x} > 0$  on  $(0, \infty)$ ,  $\ln x$  is an increasing differentiable function, hence it has a differentiable inverse, which we denote by  $\exp$ . The domain of  $\exp$  is  $\mathbb{R}$  and the range of  $\exp$  is  $(0, \infty)$ .

### The graph of $\exp$ .

Reflect the graph of  $\ln$  about the line  $y = x$ .

Notice that

- $\exp(0) = 1$  since  $\ln(1) = 0$ .
- $\exp(1) = e$  since  $\ln(e) = 1$ .
- $\exp(\ln(x)) = x$  and  $\ln(\exp(x)) = x$ .

Recall that if  $r$  is a *rational* number, then

$$\ln(e^r) = r \ln(e) = r$$

Recall that if  $r$  is a *rational* number, then

$$\ln(e^r) = r \ln(e) = r$$

Since  $\ln$  and  $\exp$  are inverses, this means that  $\exp(r) = e^r$ .

Recall that if  $r$  is a *rational* number, then

$$\ln(e^r) = r \ln(e) = r$$

Since  $\ln$  and  $\exp$  are inverses, this means that  $\exp(r) = e^r$ .

Now  $e^x$  is currently defined only when  $x$  is *rational*, and by the above, it agrees with  $\exp(x)$ . But  $\exp(x)$  is defined for all **real**  $x$ , not just all rational  $x$ .

Recall that if  $r$  is a *rational* number, then

$$\ln(e^r) = r \ln(e) = r$$

Since  $\ln$  and  $\exp$  are inverses, this means that  $\exp(r) = e^r$ .

Now  $e^x$  is currently defined only when  $x$  is *rational*, and by the above, it agrees with  $\exp(x)$ . But  $\exp(x)$  is defined for all **real**  $x$ , not just all rational  $x$ . So we *define*  $e^x = \exp(x)$  for all irrational  $x$  also! Now  $e^x$  makes sense for all  $x \in \mathbb{R}$ . Therefore,

$$e^x = y \quad \text{and} \quad \ln y = x.$$

Recall that if  $r$  is a *rational* number, then

$$\ln(e^r) = r \ln(e) = r$$

Since  $\ln$  and  $\exp$  are inverses, this means that  $\exp(r) = e^r$ .

Now  $e^x$  is currently defined only when  $x$  is *rational*, and by the above, it agrees with  $\exp(x)$ . But  $\exp(x)$  is defined for all **real**  $x$ , not just all rational  $x$ . So we *define*  $e^x = \exp(x)$  for all irrational  $x$  also! Now  $e^x$  makes sense for all  $x \in \mathbb{R}$ . Therefore,

$$e^x = y \quad \text{and} \quad \ln y = x.$$

Also, the cancellation equations  $\exp(\ln(x)) = x$  and  $\ln(\exp(x)) = x$  become

$$e^{\ln x} = x \quad \text{and} \quad \ln(e^x) = x.$$

The first cancellation formula requires  $x > 0$ , but the second holds for all  $x$ .



**Examples.** Simplify the following

(a)  $\ln \sqrt{e}$

**Examples.** Simplify the following

(a)  $\ln \sqrt{e}$

(b)  $e^{3 \ln 2}$

**Examples.** Simplify the following

(a)  $\ln \sqrt{e}$

(b)  $e^{3 \ln 2}$

(c)  $e^{x + \ln x}$

**Examples.** Simplify the following

(a)  $\ln \sqrt{e}$

(b)  $e^{3 \ln 2}$

(c)  $e^{x + \ln x}$

Solve each equation for  $x$ :

**Examples.** Simplify the following

(a)  $\ln \sqrt{e}$

(b)  $e^{3 \ln 2}$

(c)  $e^{x + \ln x}$

Solve each equation for  $x$ :

(d)  $2^{x-5} = 3$

**Examples.** Simplify the following

(a)  $\ln \sqrt{e}$

(b)  $e^{3 \ln 2}$

(c)  $e^{x + \ln x}$

Solve each equation for  $x$ :

(d)  $2^{x-5} = 3$

(e)  $\ln x + \ln(x - 1) = 1$

Here are some properties of the exponential function which we can see from the graph:

- $e^x > 0$  for all  $x$ .

Here are some properties of the exponential function which we can see from the graph:

- $e^x > 0$  for all  $x$ .
- $\lim_{x \rightarrow -\infty} e^x = 0$ .



Here are some properties of the exponential function which we can see from the graph:

- $e^x > 0$  for all  $x$ .
- $\lim_{x \rightarrow -\infty} e^x = 0$ .
- $\lim_{x \rightarrow \infty} e^x = \infty$ .

Here are some properties of the exponential function which we can see from the graph:

- $e^x > 0$  for all  $x$ .
- $\lim_{x \rightarrow -\infty} e^x = 0$ .
- $\lim_{x \rightarrow \infty} e^x = \infty$ .

We also have the expected nice properties of the exponential function. If  $x, y$  are real and  $r$  is rational, then

Here are some properties of the exponential function which we can see from the graph:

- $e^x > 0$  for all  $x$ .
- $\lim_{x \rightarrow -\infty} e^x = 0$ .
- $\lim_{x \rightarrow \infty} e^x = \infty$ .

We also have the expected nice properties of the exponential function. If  $x, y$  are real and  $r$  is rational, then

①  $e^{x+y} = e^x e^y$

Here are some properties of the exponential function which we can see from the graph:

- $e^x > 0$  for all  $x$ .
- $\lim_{x \rightarrow -\infty} e^x = 0$ .
- $\lim_{x \rightarrow \infty} e^x = \infty$ .

We also have the expected nice properties of the exponential function. If  $x, y$  are real and  $r$  is rational, then

- 1  $e^{x+y} = e^x e^y$
- 2  $e^{x-y} = \frac{e^x}{e^y}$

Here are some properties of the exponential function which we can see from the graph:

- $e^x > 0$  for all  $x$ .
- $\lim_{x \rightarrow -\infty} e^x = 0$ .
- $\lim_{x \rightarrow \infty} e^x = \infty$ .

We also have the expected nice properties of the exponential function. If  $x, y$  are real and  $r$  is rational, then

- 1  $e^{x+y} = e^x e^y$
- 2  $e^{x-y} = \frac{e^x}{e^y}$
- 3  $(e^x)^r = e^{rx}$ .