

## Week 8: Inverse functions, differentiating inverse functions, the natural logarithm

September 24 – September 28, 2012

- 1 Inverse functions
  - Inverse functions and calculus
  
- 2 The natural logarithm

# Recall from last class

## The substitution rule

If  $u = g(x)$  is a differentiable function whose range is an interval  $I$  and  $f$  is continuous on  $I$ , then

$$\int f(g(x))g'(x) dx = \int f(u) du.$$

That is, *it is permissible to work with  $dx$  and  $du$  after integral signs as if they are differentials.*

# Recall from last class

## The substitution rule

If  $u = g(x)$  is a differentiable function whose range is an interval  $I$  and  $f$  is continuous on  $I$ , then

$$\int f(g(x))g'(x) dx = \int f(u) du.$$

That is, *it is permissible to work with  $dx$  and  $du$  after integral signs as if they are differentials.*

### Example.

① Evaluate  $\int_1^2 \frac{1}{(3-5x)^2} dx$ .

- 1 Inverse functions
  - Inverse functions and calculus

- 2 The natural logarithm

# Inverse functions

Recall that we think of functions as machines that take inputs from the domain, do stuff with them, and give outputs. An important question is “for a ‘machine’  $f$ , when does there exist a ‘machine’  $g$  that undoes  $f$ ?”. If such a  $g$  exists, we call it the inverse of  $f$ .

# Inverse functions

Recall that we think of functions as machines that take inputs from the domain, do stuff with them, and give outputs. An important question is “for a ‘machine’  $f$ , when does there exist a ‘machine’  $g$  that undoes  $f$ ?”. If such a  $g$  exists, we call it the inverse of  $f$ .

**Example.** Suppose  $f$  is a function such that  $f(0) = 0$  and  $f(1) = 0$ . Is there a function  $g$  that undoes  $f$ ?

# One-one functions

## Definition

A function  $f$  is said to be **one-to-one** if it doesn't attain the same value twice. That is,

$$\text{if } x_1 \neq x_2, \text{ then } f(x_1) \neq f(x_2)$$

for any  $x_1, x_2 \in \text{Dom}(f)$ .



# One-one functions

## Definition

A function  $f$  is said to be **one-to-one** if it doesn't attain the same value twice. That is,

$$\text{if } x_1 \neq x_2, \text{ then } f(x_1) \neq f(x_2)$$

for any  $x_1, x_2 \in \text{Dom}(f)$ .

In other words, a function is one-to-one if every output corresponds to a unique input.

# One-one functions

## Definition

A function  $f$  is said to be **one-to-one** if it doesn't attain the same value twice. That is,

$$\text{if } x_1 \neq x_2, \text{ then } f(x_1) \neq f(x_2)$$

for any  $x_1, x_2 \in \text{Dom}(f)$ .

In other words, a function is one-to-one if every output corresponds to a unique input.

Graphically,  $f$  is one-to-one if and only if no horizontal line intersects its graph more than once (this is called the *horizontal line test*, compare this with the *vertical line test* to test if  $f$  is a function in the first place). One-to-one functions are sometimes called **injective** functions.

# One-one functions

**True or false?.** The following functions are injective:

- (i)  $f$  given by  $f(x) = x^2$ .

# One-one functions

**True or false?.** The following functions are injective:

(i)  $f$  given by  $f(x) = x^2$ .

(ii)  $f$  given by  $f(x) = x^3$ .

# One-one functions

**True or false?.** The following functions are injective:

- (i)  $f$  given by  $f(x) = x^2$ .
- (ii)  $f$  given by  $f(x) = x^3$ .
- (iii)  $f$  given by  $f(x) = x^2$  where the domain of  $f$  is  $x \geq 0$ .

# One-one functions

**True or false?.** The following functions are injective:

- (i)  $f$  given by  $f(x) = x^2$ .
- (ii)  $f$  given by  $f(x) = x^3$ .
- (iii)  $f$  given by  $f(x) = x^2$  where the domain of  $f$  is  $x \geq 0$ .
- (iv)  $f$  given by  $f(x) = \sqrt{x}$ .

# One-one functions

**True or false?.** The following functions are injective:

- (i)  $f$  given by  $f(x) = x^2$ .
- (ii)  $f$  given by  $f(x) = x^3$ .
- (iii)  $f$  given by  $f(x) = x^2$  where the domain of  $f$  is  $x \geq 0$ .
- (iv)  $f$  given by  $f(x) = \sqrt{x}$ .
- (v)  $f$  given by  $f(x) = \cos(x)$ .

# Inverse functions

One-to-one functions are exactly those that possess inverse functions!



# Inverse functions

One-to-one functions are exactly those that possess inverse functions!

## Definition

Suppose  $f$  is a function with domain  $A$  and range  $B$ . Then its **inverse function**  $f^{-1}$  has domain  $B$  and range  $A$  and is defined by

$$f^{-1}(y) = x \quad \text{if and only if} \quad f(x) = y$$

for any  $y$  in  $B$ .

# Inverse functions

One-to-one functions are exactly those that possess inverse functions!

## Definition

Suppose  $f$  is a function with domain  $A$  and range  $B$ . Then its **inverse function**  $f^{-1}$  has domain  $B$  and range  $A$  and is defined by

$$f^{-1}(y) = x \quad \text{if and only if} \quad f(x) = y$$

for any  $y$  in  $B$ .

**Example.** Suppose  $f$  is defined by  $f(x) = y = x^3$ , so  $f$  maps  $x$  to  $y$ . The inverse function of  $f$  is given by  $f^{-1}(x) = x^{1/3}$  because

$$f^{-1}(y) = f^{-1}(x^3) = (x^3)^{1/3} = x.$$

# Inverse functions

## Remarks.

- (i) If  $f$  is not one-to-one, then  $f^{-1}$  isn't uniquely defined.

# Inverse functions

## Remarks.

- (i) If  $f$  is not one-to-one, then  $f^{-1}$  isn't uniquely defined.
- (ii) (important!) The  $-1$  in  $f^{-1}$  is **NOT** an exponent. That is,  $f^{-1}(x)$  is **NOT** equal to  $\frac{1}{f(x)}$  in general. For example, if  $f(x) = x^3$  then what is  $f^{-1}(8)$ ? What is  $\frac{1}{f(8)}$ ? (However, we *can* write  $(f(x))^{-1} = \frac{1}{f(x)}$ .)

# Inverse functions

## Remarks.

- (i) If  $f$  is not one-to-one, then  $f^{-1}$  isn't uniquely defined.
- (ii) (important!) The  $-1$  in  $f^{-1}$  is **NOT** an exponent. That is,  $f^{-1}(x)$  is **NOT** equal to  $\frac{1}{f(x)}$  in general. For example, if  $f(x) = x^3$  then what is  $f^{-1}(8)$ ? What is  $\frac{1}{f(8)}$ ? (However, we *can* write  $(f(x))^{-1} = \frac{1}{f(x)}$ .)
- (iii) Since  $f$  and  $f^{-1}$  “undo” each other, we have the rules  $f^{-1}(f(x)) = x$  for every  $x$  in  $A$ , and  $f(f^{-1}(y)) = y$  for every  $y$  in  $B$ .

# Inverse functions

## Remarks.

- (i) If  $f$  is not one-to-one, then  $f^{-1}$  isn't uniquely defined.
- (ii) (important!) The  $-1$  in  $f^{-1}$  is **NOT** an exponent. That is,  $f^{-1}(x)$  is **NOT** equal to  $\frac{1}{f(x)}$  in general. For example, if  $f(x) = x^3$  then what is  $f^{-1}(8)$ ? What is  $\frac{1}{f(8)}$ ? (However, we *can* write  $(f(x))^{-1} = \frac{1}{f(x)}$ .)
- (iii) Since  $f$  and  $f^{-1}$  “undo” each other, we have the rules  $f^{-1}(f(x)) = x$  for every  $x$  in  $A$ , and  $f(f^{-1}(y)) = y$  for every  $y$  in  $B$ .

So given  $f$ , how do we explicitly find the inverse function  $f^{-1}$ ? We first write  $y = f(x)$ , then try to solve for  $x$  in terms of  $y$ , then at the end switch the variables  $x$  and  $y$  to express  $f^{-1}$  as a function of  $x$ .

**Example.** Find a formula for the inverse of the function  $f$  given by the rule  $f(x) = 2x^3 + 3$ .

**Example.** Find a formula for the inverse of the function  $f$  given by the rule  $f(x) = 2x^3 + 3$ .

If  $f$  has an inverse, notice that  $f(a) = b$  if and only if  $f^{-1}(b) = a$ . That is,  $(a, b)$  is in the graph of  $f$  if and only if  $(b, a)$  is in the graph of  $f^{-1}$ . Since the point  $(b, a)$  is obtained from  $(a, b)$  by reflecting about the line  $y = x$ , we get the graph of  $f^{-1}$  by reflecting the graph of  $f$  about the line  $y = x$ .



**Example.** Find a formula for the inverse of the function  $f$  given by the rule  $f(x) = 2x^3 + 3$ .

If  $f$  has an inverse, notice that  $f(a) = b$  if and only if  $f^{-1}(b) = a$ . That is,  $(a, b)$  is in the graph of  $f$  if and only if  $(b, a)$  is in the graph of  $f^{-1}$ . Since the point  $(b, a)$  is obtained from  $(a, b)$  by reflecting about the line  $y = x$ , we get the graph of  $f^{-1}$  by reflecting the graph of  $f$  about the line  $y = x$ .

**Example.** Sketch the graph of  $f$  and  $f^{-1}$  if  $f(x) = \sqrt{x-2}$ .

# Inverse functions and calculus

If a function  $f$  is one-to-one then its inverse function  $g$  is also one-to-one. What other properties of  $f$  does its inverse function  $g$  inherit?

- If  $f$  is continuous, will  $g$  be continuous?

# Inverse functions and calculus

If a function  $f$  is one-to-one then its inverse function  $g$  is also one-to-one. What other properties of  $f$  does its inverse function  $g$  inherit?

- If  $f$  is continuous, will  $g$  be continuous?
- If  $f$  is differentiable, will  $g$  be differentiable?

# Inverse functions and calculus

If a function  $f$  is one-to-one then its inverse function  $g$  is also one-to-one. What other properties of  $f$  does its inverse function  $g$  inherit?

- If  $f$  is continuous, will  $g$  be continuous?
- If  $f$  is differentiable, will  $g$  be differentiable?

Intuitively, we expect that if the graph of  $f$  is a curve with no breaks, then the graph of  $f^{-1}$  should also be a curve with no breaks. We have the following nice theorem:

# Inverse functions and calculus

If a function  $f$  is one-to-one then its inverse function  $g$  is also one-to-one. What other properties of  $f$  does its inverse function  $g$  inherit?

- If  $f$  is continuous, will  $g$  be continuous?
- If  $f$  is differentiable, will  $g$  be differentiable?

Intuitively, we expect that if the graph of  $f$  is a curve with no breaks, then the graph of  $f^{-1}$  should also be a curve with no breaks. We have the following nice theorem:

## Theorem

*If  $f$  is a one-to-one continuous function defined on an interval, then its inverse function  $f^{-1}$  is also continuous.*

# Inverse function theorem

Intuitively, we know that if  $f$  is differentiable then the graph has no kinks or vertical tangent lines. If the graph of  $f^{-1}$  is obtained by reflecting about the line  $y = x$ , then if the graph of  $f$  has no kinks, then neither will the graph of  $f^{-1}$ . The only way to get vertical tangent lines to the graph of  $f^{-1}$  is if there are horizontal tangent lines to the graph of  $f$ . This motivates the following important theorem.

# Inverse function theorem

Intuitively, we know that if  $f$  is differentiable then the graph has no kinks or vertical tangent lines. If the graph of  $f^{-1}$  is obtained by reflecting about the line  $y = x$ , then if the graph of  $f$  has no kinks, then neither will the graph of  $f^{-1}$ . The only way to get vertical tangent lines to the graph of  $f^{-1}$  is if there are horizontal tangent lines to the graph of  $f$ . This motivates the following important theorem.

## Theorem (The inverse function theorem)

*Suppose  $f$  is a one-to-one differentiable function with inverse function  $f^{-1}$  and  $f'(f^{-1}(a)) \neq 0$ , then the inverse function is differentiable at  $a$  and*

$$(f^{-1})'(a) = \frac{1}{f'(f^{-1}(a))}.$$

# Inverse function theorem

## Remark.

- Note that this theorem tells you both that the inverse function **is** differentiable at the point  $a$ , and also gives you an easy formula to calculate its value.



# Inverse function theorem

## Remark.

- Note that this theorem tells you both that the inverse function **is** differentiable at the point  $a$ , and also gives you an easy formula to calculate its value.

**Example.** If  $f(x) = 2x + \cos x$ , find  $(f^{-1})'(1)$ .

- 1 Inverse functions
  - Inverse functions and calculus
  
- 2 The natural logarithm

# The natural logarithm

We now introduce a function of fundamental importance. We will define the natural logarithm using an integral!

# The natural logarithm

We now introduce a function of fundamental importance. We will define the natural logarithm using an integral!

We motivate our definition by considering the following problem:

# The natural logarithm

We now introduce a function of fundamental importance. We will define the natural logarithm using an integral!

We motivate our definition by considering the following problem:

The volume of a colony of bacteria on a plate is 1 cubic millimeter and doubles every day. Let  $V(t)$  denote the volume of bacteria after  $t$  days. It is clear that

$$V(0) = 1, \quad V(1) = 2, \quad V(2) = 4, \quad V(3) = 8$$

and so on. In general,  $V(t) = 2^t$  if  $t$  is a nonnegative integer.

# The natural logarithm

We now introduce a function of fundamental importance. We will define the natural logarithm using an integral!

We motivate our definition by considering the following problem:

The volume of a colony of bacteria on a plate is 1 cubic millimeter and doubles every day. Let  $V(t)$  denote the volume of bacteria after  $t$  days. It is clear that

$$V(0) = 1, \quad V(1) = 2, \quad V(2) = 4, \quad V(3) = 8$$

and so on. In general,  $V(t) = 2^t$  if  $t$  is a nonnegative integer.

**Question.** Can we graph  $V(t) = 2^t$  if  $t \in \mathbb{R}$ ?  
Only if we have defined  $2^t$  when  $t \in \mathbb{R}$ .

- What is  $2^{3/7}$ ?

# The natural logarithm

We now introduce a function of fundamental importance. We will define the natural logarithm using an integral!

We motivate our definition by considering the following problem:

The volume of a colony of bacteria on a plate is 1 cubic millimeter and doubles every day. Let  $V(t)$  denote the volume of bacteria after  $t$  days. It is clear that

$$V(0) = 1, \quad V(1) = 2, \quad V(2) = 4, \quad V(3) = 8$$

and so on. In general,  $V(t) = 2^t$  if  $t$  is a nonnegative integer.

**Question.** Can we graph  $V(t) = 2^t$  if  $t \in \mathbb{R}$ ?  
Only if we have defined  $2^t$  when  $t \in \mathbb{R}$ .

- What is  $2^{3/7}$ ?
- What is  $2^{\sqrt{3}}$ ?

# The natural logarithm

We now introduce a function of fundamental importance. We will define the natural logarithm using an integral!

We motivate our definition by considering the following problem:

The volume of a colony of bacteria on a plate is 1 cubic millimeter and doubles every day. Let  $V(t)$  denote the volume of bacteria after  $t$  days. It is clear that

$$V(0) = 1, \quad V(1) = 2, \quad V(2) = 4, \quad V(3) = 8$$

and so on. In general,  $V(t) = 2^t$  if  $t$  is a nonnegative integer.

**Question.** Can we graph  $V(t) = 2^t$  if  $t \in \mathbb{R}$ ?

Only if we have defined  $2^t$  when  $t \in \mathbb{R}$ .

- What is  $2^{3/7}$ ?
- What is  $2^{\sqrt{3}}$ ?
- What is  $2^\pi$ ?



# The natural logarithm

There is lots of work to be done here before we can draw the graph of  $V$ . At the moment we only have the following: If  $b$  is a positive real number and  $r$  is a rational number with  $r = p/q$  then  $b^r$  is defined to be the unique positive  $q$ th root of  $b^p$ . We do not yet have a definition for  $b^r$  when  $r \notin \mathbb{Q}$ .

# The natural logarithm

There is lots of work to be done here before we can draw the graph of  $V$ . At the moment we only have the following: If  $b$  is a positive real number and  $r$  is a rational number with  $r = p/q$  then  $b^r$  is defined to be the unique positive  $q$ th root of  $b^p$ . We do not yet have a definition for  $b^r$  when  $r \notin \mathbb{Q}$ .

Our strategy for improving this situation is the following.

- 1 Define the natural logarithm function  $\ln$  as a function from  $(0, \infty)$  to  $\mathbb{R}$  using an integral formula.

# The natural logarithm

There is lots of work to be done here before we can draw the graph of  $V$ . At the moment we only have the following: If  $b$  is a positive real number and  $r$  is a rational number with  $r = p/q$  then  $b^r$  is defined to be the unique positive  $q$ th root of  $b^p$ . We do not yet have a definition for  $b^r$  when  $r \notin \mathbb{Q}$ .

Our strategy for improving this situation is the following.

- 1 Define the natural logarithm function  $\ln$  as a function from  $(0, \infty)$  to  $\mathbb{R}$  using an integral formula.
- 2 Define the exponential function  $\exp = e^x$ , from  $\mathbb{R}$  to  $(0, \infty)$  as the inverse of  $\ln$ .

# The natural logarithm

There is lots of work to be done here before we can draw the graph of  $V$ . At the moment we only have the following: If  $b$  is a positive real number and  $r$  is a rational number with  $r = p/q$  then  $b^r$  is defined to be the unique positive  $q$ th root of  $b^p$ . We do not yet have a definition for  $b^r$  when  $r \notin \mathbb{Q}$ .

Our strategy for improving this situation is the following.

- 1 Define the natural logarithm function  $\ln$  as a function from  $(0, \infty)$  to  $\mathbb{R}$  using an integral formula.
- 2 Define the exponential function  $\exp = e^x$ , from  $\mathbb{R}$  to  $(0, \infty)$  as the inverse of  $\ln$ .
- 3 Define  $b^r$  when  $r \notin \mathbb{Q}$  using the  $\exp$  and  $\ln$  functions.

# The natural logarithm

## Definition

The function  $\ln$  with domain  $(0, \infty)$  is defined by the formula

$$\ln(x) = \int_1^x \frac{1}{t} dt.$$

# The natural logarithm

## Definition

The function  $\ln$  with domain  $(0, \infty)$  is defined by the formula

$$\ln(x) = \int_1^x \frac{1}{t} dt.$$

By interpreting  $\ln$  as the area under a curve, we can see that:

- $\ln x > 0$  if  $x > 1$  and  $\ln 1 = 0$ ,

# The natural logarithm

## Definition

The function  $\ln$  with domain  $(0, \infty)$  is defined by the formula

$$\ln(x) = \int_1^x \frac{1}{t} dt.$$

By interpreting  $\ln$  as the area under a curve, we can see that:

- $\ln x > 0$  if  $x > 1$  and  $\ln 1 = 0$ ,
- $\ln x < 0$  if  $0 < x < 1$ .

# The natural logarithm

By the Fundamental Theorem of Calculus, part 1, we see that

$$\frac{d}{dx}(\ln x) = \frac{d}{dx} \int_1^x \frac{1}{t} dt = \frac{1}{x}.$$



# The natural logarithm

By the Fundamental Theorem of Calculus, part 1, we see that

$$\frac{d}{dx}(\ln x) = \frac{d}{dx} \int_1^x \frac{1}{t} dt = \frac{1}{x}.$$

This implies the following (familiar) properties of the logarithm:

- (i)  $\ln(xy) = \ln(x) + \ln(y)$  for all positive real numbers  $x$  and  $y$ .

# The natural logarithm

By the Fundamental Theorem of Calculus, part 1, we see that

$$\frac{d}{dx}(\ln x) = \frac{d}{dx} \int_1^x \frac{1}{t} dt = \frac{1}{x}.$$

This implies the following (familiar) properties of the logarithm:

- (i)  $\ln(xy) = \ln(x) + \ln(y)$  for all positive real numbers  $x$  and  $y$ .
- (ii)  $\ln\left(\frac{x}{y}\right) = \ln(x) - \ln(y)$  for all positive real numbers  $x$  and  $y$ .

# The natural logarithm

By the Fundamental Theorem of Calculus, part 1, we see that

$$\frac{d}{dx}(\ln x) = \frac{d}{dx} \int_1^x \frac{1}{t} dt = \frac{1}{x}.$$

This implies the following (familiar) properties of the logarithm:

- (i)  $\ln(xy) = \ln(x) + \ln(y)$  for all positive real numbers  $x$  and  $y$ .
- (ii)  $\ln\left(\frac{x}{y}\right) = \ln(x) - \ln(y)$  for all positive real numbers  $x$  and  $y$ .
- (iii)  $\ln(x^r) = r \ln(x)$  whenever  $r$  is a rational number and  $x$  is a positive real number.

# The natural logarithm

**Example.** Expand the expression  $\ln \frac{(x^2+5)^4 \sin x}{x^3+1}$ .