

## Week 10: The logarithmic and exponential functions, exponential growth

October 9 – October 12, 2012

## 1 General exponential and logarithmic functions

- Exponential growth and decay

## 2 Inverse functions

- The inverse trigonometric functions
- The hyperbolic functions

▶ Lecture 18

▶ Lecture 19

## Recall from last class

- Since  $\ln$  and  $\exp$  are inverses and  $\ln(e^r) = r \ln(e) = r$ , this means that  $\exp(r) = e^r$ .

# Recall from last class

- Since  $\ln$  and  $\exp$  are inverses and  $\ln(e^r) = r \ln(e) = r$ , this means that  $\exp(r) = e^r$ .
- We *define*  $e^x = \exp(x)$  for all  $x \in \mathbb{R}$ . Therefore,

$$e^x = y \quad \text{and} \quad \ln y = x.$$

# Recall from last class

- Since  $\ln$  and  $\exp$  are inverses and  $\ln(e^r) = r \ln(e) = r$ , this means that  $\exp(r) = e^r$ .
- We *define*  $e^x = \exp(x)$  for all  $x \in \mathbb{R}$ . Therefore,

$$e^x = y \quad \text{and} \quad \ln y = x.$$

- $e^{x+y} = e^x e^y$

# Recall from last class

- Since  $\ln$  and  $\exp$  are inverses and  $\ln(e^r) = r \ln(e) = r$ , this means that  $\exp(r) = e^r$ .
- We *define*  $e^x = \exp(x)$  for all  $x \in \mathbb{R}$ . Therefore,

$$e^x = y \quad \text{and} \quad \ln y = x.$$

- $e^{x+y} = e^x e^y$
- $e^{x-y} = \frac{e^x}{e^y}$

# Recall from last class

- Since  $\ln$  and  $\exp$  are inverses and  $\ln(e^r) = r \ln(e) = r$ , this means that  $\exp(r) = e^r$ .
- We *define*  $e^x = \exp(x)$  for all  $x \in \mathbb{R}$ . Therefore,

$$e^x = y \quad \text{and} \quad \ln y = x.$$

- $e^{x+y} = e^x e^y$
- $e^{x-y} = \frac{e^x}{e^y}$
- $(e^x)^r = e^{rx}$ .

**Examples.** Find the limit:

(i)  $\lim_{x \rightarrow \infty} \frac{e^{3x} - e^{-3x}}{e^{3x} + e^{-3x}}.$



**Examples.** Find the limit:

(i)  $\lim_{x \rightarrow \infty} \frac{e^{3x} - e^{-3x}}{e^{3x} + e^{-3x}}.$

(ii)  $\lim_{x \rightarrow 2^+} e^{3/(2-x)}$

What is the derivative of  $e^x$ ? We know it is differentiable because it is the inverse function of  $\ln x$ , which is differentiable with nonzero derivative. Let's use implicit differentiation to answer this question:

What is the derivative of  $e^x$ ? We know it is differentiable because it is the inverse function of  $\ln x$ , which is differentiable with nonzero derivative. Let's use implicit differentiation to answer this question:

Let  $y = e^x$ . Then,  $\ln y = x$ .

What is the derivative of  $e^x$ ? We know it is differentiable because it is the inverse function of  $\ln x$ , which is differentiable with nonzero derivative. Let's use implicit differentiation to answer this question:

Let  $y = e^x$ . Then,  $\ln y = x$ . Differentiating both sides,

$$\frac{1}{y} \cdot \frac{dy}{dx} = 1.$$

What is the derivative of  $e^x$ ? We know it is differentiable because it is the inverse function of  $\ln x$ , which is differentiable with nonzero derivative. Let's use implicit differentiation to answer this question:

Let  $y = e^x$ . Then,  $\ln y = x$ . Differentiating both sides,

$$\frac{1}{y} \cdot \frac{dy}{dx} = 1.$$

Thus,  $\frac{dy}{dx} = y = e^x$ .

What is the derivative of  $e^x$ ? We know it is differentiable because it is the inverse function of  $\ln x$ , which is differentiable with nonzero derivative. Let's use implicit differentiation to answer this question:

Let  $y = e^x$ . Then,  $\ln y = x$ . Differentiating both sides,

$$\frac{1}{y} \cdot \frac{dy}{dx} = 1.$$

Thus,  $\frac{dy}{dx} = y = e^x$ .

**(Important) derivative.**  $\frac{d}{dx} e^x = e^x$ . Moreover, up to multiplication by a constant, this is the *only* function which is its own derivative!!

## Examples.

(i) Differentiate the function given by  $y = \sqrt{1 + xe^{-2x}}$ .

**Examples.**

- (i) Differentiate the function given by  $y = \sqrt{1 + xe^{-2x}}$ .
- (ii) Find an equation of the tangent line to the curve  $y = e^x/x$  at the point  $(1, e)$ .



**Examples.**

- (i) Differentiate the function given by  $y = \sqrt{1 + xe^{-2x}}$ .
- (ii) Find an equation of the tangent line to the curve  $y = e^x/x$  at the point  $(1, e)$ .
- (iii) On what interval is the curve  $y = xe^{3x}$  concave upward?

What about integration? Well, because  $e^x$  has a simple derivative, it also has a simple antiderivative! Namely

$$\int e^x dx = e^x + C.$$

What about integration? Well, because  $e^x$  has a simple derivative, it also has a simple antiderivative! Namely

$$\int e^x dx = e^x + C.$$

**Example.** Evaluate the integral

$$\int \frac{e^{1/x}}{x^2} dx.$$

## 1 General exponential and logarithmic functions

- Exponential growth and decay

## 2 Inverse functions

- The inverse trigonometric functions
- The hyperbolic functions

Now, as promised, we will finally be able to define  $a^x$ , where  $a$  is a positive real number and  $x$  is *any* real number! First, we observe that if  $r$  is rational, then

$$a^r = (e^{\ln a})^r = e^{r \ln a}.$$

Now, as promised, we will finally be able to define  $a^x$ , where  $a$  is a positive real number and  $x$  is *any* real number! First, we observe that if  $r$  is rational, then

$$a^r = (e^{\ln a})^r = e^{r \ln a}.$$

However, the right side of the equation makes sense even if  $r$  is not rational, because we have defined  $e^x$  for *all* real  $x$ . So if  $x$  is real, we make the following definition:

Now, as promised, we will finally be able to define  $a^x$ , where  $a$  is a positive real number and  $x$  is *any* real number! First, we observe that if  $r$  is rational, then

$$a^r = (e^{\ln a})^r = e^{r \ln a}.$$

However, the right side of the equation makes sense even if  $r$  is not rational, because we have defined  $e^x$  for *all* real  $x$ . So if  $x$  is real, we make the following definition:

### Definition

If  $a > 0$  and  $x \in \mathbb{R}$ , then

$$a^x = e^{x \ln a}.$$

Now, as promised, we will finally be able to define  $a^x$ , where  $a$  is a positive real number and  $x$  is *any* real number! First, we observe that if  $r$  is rational, then

$$a^r = (e^{\ln a})^r = e^{r \ln a}.$$

However, the right side of the equation makes sense even if  $r$  is not rational, because we have defined  $e^x$  for *all* real  $x$ . So if  $x$  is real, we make the following definition:

### Definition

If  $a > 0$  and  $x \in \mathbb{R}$ , then

$$a^x = e^{x \ln a}.$$

When we first introduced the problem of defining arbitrary exponential functions, we asked the question, what is  $2^{\sqrt{3}}$ ? Now we can answer it!

$$2^{\sqrt{3}} = e^{\sqrt{3} \ln 2}$$



Now, as promised, we will finally be able to define  $a^x$ , where  $a$  is a positive real number and  $x$  is *any* real number! First, we observe that if  $r$  is rational, then

$$a^r = (e^{\ln a})^r = e^{r \ln a}.$$

However, the right side of the equation makes sense even if  $r$  is not rational, because we have defined  $e^x$  for *all* real  $x$ . So if  $x$  is real, we make the following definition:

### Definition

If  $a > 0$  and  $x \in \mathbb{R}$ , then

$$a^x = e^{x \ln a}.$$

When we first introduced the problem of defining arbitrary exponential functions, we asked the question, what is  $2^{\sqrt{3}}$ ? Now we can answer it!

$$2^{\sqrt{3}} = e^{\sqrt{3} \ln 2}$$

We call a function defined by  $f(x) = a^x$  an **exponential function with base  $a$** . We can see that such a function is always positive because  $e^x$  is always positive.

We call a function defined by  $f(x) = a^x$  an **exponential function with base  $a$** . We can see that such a function is always positive because  $e^x$  is always positive.

**Example.** Solve for  $x$ :

①  $2^{x-5} = 3.$

Armed with this definition of an arbitrary exponential function, we can improve one of the properties of  $\ln x$ . We showed earlier that  $\ln(a^r) = r \ln a$  when  $r$  is *rational*. But now, if  $x$  is any *real* number, we notice that

$$\ln(a^x) = \ln(e^{x \ln a}) = x \ln a,$$

so the formula holds in fact for any *real* number!

Armed with this definition of an arbitrary exponential function, we can improve one of the properties of  $\ln x$ . We showed earlier that  $\ln(a^r) = r \ln a$  when  $r$  is *rational*. But now, if  $x$  is any *real* number, we notice that

$$\ln(a^x) = \ln(e^{x \ln a}) = x \ln a,$$

so the formula holds in fact for any *real* number!

We list this, and a few other properties. The book justifies a few of these properties more. In the following, we assume that  $x, y$  are real numbers and  $a, b$  are strictly positive.

- $\ln a^x = x \ln a$  for any real number  $x$ .

Armed with this definition of an arbitrary exponential function, we can improve one of the properties of  $\ln x$ . We showed earlier that  $\ln(a^r) = r \ln a$  when  $r$  is *rational*. But now, if  $x$  is any *real* number, we notice that

$$\ln(a^x) = \ln(e^{x \ln a}) = x \ln a,$$

so the formula holds in fact for any *real* number!

We list this, and a few other properties. The book justifies a few of these properties more. In the following, we assume that  $x, y$  are real numbers and  $a, b$  are strictly positive.

- $\ln a^x = x \ln a$  for any real number  $x$ .
- $a^{x+y} = a^x a^y$ .

Armed with this definition of an arbitrary exponential function, we can improve one of the properties of  $\ln x$ . We showed earlier that  $\ln(a^r) = r \ln a$  when  $r$  is *rational*. But now, if  $x$  is any *real* number, we notice that

$$\ln(a^x) = \ln(e^{x \ln a}) = x \ln a,$$

so the formula holds in fact for any *real* number!

We list this, and a few other properties. The book justifies a few of these properties more. In the following, we assume that  $x, y$  are real numbers and  $a, b$  are strictly positive.

- $\ln a^x = x \ln a$  for any real number  $x$ .
- $a^{x+y} = a^x a^y$ .
- $a^{x-y} = a^x / a^y$ .

Armed with this definition of an arbitrary exponential function, we can improve one of the properties of  $\ln x$ . We showed earlier that  $\ln(a^r) = r \ln a$  when  $r$  is *rational*. But now, if  $x$  is any *real* number, we notice that

$$\ln(a^x) = \ln(e^{x \ln a}) = x \ln a,$$

so the formula holds in fact for any *real* number!

We list this, and a few other properties. The book justifies a few of these properties more. In the following, we assume that  $x, y$  are real numbers and  $a, b$  are strictly positive.

- $\ln a^x = x \ln a$  for any real number  $x$ .
- $a^{x+y} = a^x a^y$ .
- $a^{x-y} = a^x / a^y$ .
- $(a^x)^y = a^{xy}$ .



Armed with this definition of an arbitrary exponential function, we can improve one of the properties of  $\ln x$ . We showed earlier that  $\ln(a^r) = r \ln a$  when  $r$  is *rational*. But now, if  $x$  is any *real* number, we notice that

$$\ln(a^x) = \ln(e^{x \ln a}) = x \ln a,$$

so the formula holds in fact for any *real* number!

We list this, and a few other properties. The book justifies a few of these properties more. In the following, we assume that  $x, y$  are real numbers and  $a, b$  are strictly positive.

- $\ln a^x = x \ln a$  for any real number  $x$ .
- $a^{x+y} = a^x a^y$ .
- $a^{x-y} = a^x / a^y$ .
- $(a^x)^y = a^{xy}$ .
- $(ab)^x = a^x b^x$ .

What do the graphs of arbitrary exponential functions look like?

What do the graphs of arbitrary exponential functions look like?

Finally, we see that  $\frac{d}{dx}(a^x) = a^x \ln a$ :

What do the graphs of arbitrary exponential functions look like?

Finally, we see that  $\frac{d}{dx}(a^x) = a^x \ln a$ :

### Examples.

(i) Write  $(\cos x)^x$  as a power of  $e$ .

What do the graphs of arbitrary exponential functions look like?

Finally, we see that  $\frac{d}{dx}(a^x) = a^x \ln a$ :

### Examples.

- (i) Write  $(\cos x)^x$  as a power of  $e$ .
- (ii) Differentiate the function  $h(t) = t^3 - 3^t$ .

How do we integrate a general exponential function?

How do we integrate a general exponential function? Since

$$\frac{d}{dx}(a^x) = a^x \ln a,$$

this implies that

$$\int a^x dx = \frac{a^x}{\ln a} + C, \quad a \neq 1.$$

How do we integrate a general exponential function? Since

$$\frac{d}{dx}(a^x) = a^x \ln a,$$

this implies that

$$\int a^x dx = \frac{a^x}{\ln a} + C, \quad a \neq 1.$$

**Example.** Calculate  $\int_0^5 3^x dx$ .



We can now state the **general power rule**, namely if  $n$  is *any* real number and  $f$  is given by  $f(x) = x^n$ , then

$$f'(x) = nx^{n-1}.$$

We can now state the **general power rule**, namely if  $n$  is *any* real number and  $f$  is given by  $f(x) = x^n$ , then

$$f'(x) = nx^{n-1}.$$

**Example.** Differentiate  $y = x^{e^x}$ .

**General Logarithmic Functions.** If  $a$  is positive and not equal to one, then  $a^x$  is one-to-one. Hence it has an inverse, denoted by  $\log_a$ , which is called the **logarithmic function with base  $a$** . That is,

$$\log_a x = y \iff a^y = x.$$

**General Logarithmic Functions.** If  $a$  is positive and not equal to one, then  $a^x$  is one-to-one. Hence it has an inverse, denoted by  $\log_a$ , which is called the **logarithmic function with base  $a$** . That is,

$$\log_a x = y \iff a^y = x.$$

We see that  $\log_e x = \ln x$ , and that

$$a^{\log_a x} = x \quad \text{and} \quad \log_a(a^x) = x.$$

**General Logarithmic Functions.** If  $a$  is positive and not equal to one, then  $a^x$  is one-to-one. Hence it has an inverse, denoted by  $\log_a$ , which is called the **logarithmic function with base  $a$** . That is,

$$\log_a x = y \iff a^y = x.$$

We see that  $\log_e x = \ln x$ , and that

$$a^{\log_a x} = x \quad \text{and} \quad \log_a(a^x) = x.$$

How does the graphs of the general logarithm look like?

It is easy to check the following important **change of base formula**, relating general logarithms to the natural logarithm: If  $a > 0$ ,  $a \neq 1$ , then

$$\log_a x = \frac{\ln x}{\ln a}.$$

It is easy to check the following important **change of base formula**, relating general logarithms to the natural logarithm: If  $a > 0$ ,  $a \neq 1$ , then

$$\log_a x = \frac{\ln x}{\ln a}.$$

This implies that

$$\frac{d}{dx}(\log_a x) = \frac{1}{x \ln a}.$$

It is easy to check the following important **change of base formula**, relating general logarithms to the natural logarithm: If  $a > 0$ ,  $a \neq 1$ , then

$$\log_a x = \frac{\ln x}{\ln a}.$$

This implies that

$$\frac{d}{dx}(\log_a x) = \frac{1}{x \ln a}.$$

**Example.** Calculate the derivative of the function given by  $f(x) = \log_{10}(2 + \sin x)$ .



We briefly derive a formula for the number  $e$  as a limit. You may be familiar with this definition of  $e$  from high school.

We briefly derive a formula for the number  $e$  as a limit. You may be familiar with this definition of  $e$  from high school.

We record this as

$$e = \lim_{x \rightarrow 0} (1 + x)^{1/x} \quad \text{or equivalently} \quad e = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n.$$

**True or False?.**

(i) If  $0 < a < b$ , then  $\ln a < \ln b$ .

**True or False?.**

(i) If  $0 < a < b$ , then  $\ln a < \ln b$ .

(ii)  $\pi^{\sqrt{5}} = e^{\sqrt{5} \ln \pi}$ .

**True or False?.**

- (i) If  $0 < a < b$ , then  $\ln a < \ln b$ .
- (ii)  $\pi^{\sqrt{5}} = e^{\sqrt{5} \ln \pi}$ .
- (iii) You can always divide by  $e^x$ .

**True or False?.**

- (i) If  $0 < a < b$ , then  $\ln a < \ln b$ .
- (ii)  $\pi^{\sqrt{5}} = e^{\sqrt{5} \ln \pi}$ .
- (iii) You can always divide by  $e^x$ .
- (iv) If  $a > 0$  and  $b > 0$  then  $\ln(a + b) = \ln a + \ln b$ .

**True or False?.**

- (i) If  $0 < a < b$ , then  $\ln a < \ln b$ .
- (ii)  $\pi^{\sqrt{5}} = e^{\sqrt{5} \ln \pi}$ .
- (iii) You can always divide by  $e^x$ .
- (iv) If  $a > 0$  and  $b > 0$  then  $\ln(a + b) = \ln a + \ln b$ .
- (v) If  $x > 0$ , then  $(\ln x)^6 = 6 \ln x$ .

**True or False?.**

- (i) If  $0 < a < b$ , then  $\ln a < \ln b$ .
- (ii)  $\pi^{\sqrt{5}} = e^{\sqrt{5} \ln \pi}$ .
- (iii) You can always divide by  $e^x$ .
- (iv) If  $a > 0$  and  $b > 0$  then  $\ln(a + b) = \ln a + \ln b$ .
- (v) If  $x > 0$ , then  $(\ln x)^6 = 6 \ln x$ .
- (vi)  $\frac{d}{dx}(10^x) = x10^{x-1}$ .



**True or False?.**

- (i) If  $0 < a < b$ , then  $\ln a < \ln b$ .
- (ii)  $\pi^{\sqrt{5}} = e^{\sqrt{5} \ln \pi}$ .
- (iii) You can always divide by  $e^x$ .
- (iv) If  $a > 0$  and  $b > 0$  then  $\ln(a + b) = \ln a + \ln b$ .
- (v) If  $x > 0$ , then  $(\ln x)^6 = 6 \ln x$ .
- (vi)  $\frac{d}{dx}(10^x) = x10^{x-1}$ .
- (vii)  $\frac{d}{dx}(\ln 10) = \frac{1}{10}$ .

**True or False?.**

- (i) If  $0 < a < b$ , then  $\ln a < \ln b$ .
- (ii)  $\pi^{\sqrt{5}} = e^{\sqrt{5} \ln \pi}$ .
- (iii) You can always divide by  $e^x$ .
- (iv) If  $a > 0$  and  $b > 0$  then  $\ln(a + b) = \ln a + \ln b$ .
- (v) If  $x > 0$ , then  $(\ln x)^6 = 6 \ln x$ .
- (vi)  $\frac{d}{dx}(10^x) = x10^{x-1}$ .
- (vii)  $\frac{d}{dx}(\ln 10) = \frac{1}{10}$ .
- (viii) The inverse function of  $y = e^{3x}$  is  $y = \frac{1}{3} \ln x$ .

## Exponential growth and decay.

We come to one of the most important applications of exponential functions. In many natural phenomena, quantities grow or decay at a rate proportional to their size.

## Exponential growth and decay.

We come to one of the most important applications of exponential functions. In many natural phenomena, quantities grow or decay at a rate proportional to their size. If  $y(t)$  is the value of such a quantity at time  $t$  then

$$\frac{dy}{dt} = ky,$$

where  $k$  is the constant of proportionality.

## Exponential growth and decay.

We come to one of the most important applications of exponential functions. In many natural phenomena, quantities grow or decay at a rate proportional to their size. If  $y(t)$  is the value of such a quantity at time  $t$  then

$$\frac{dy}{dt} = ky,$$

where  $k$  is the constant of proportionality.

### Theorem

*The only solutions of the differential equation  $\frac{dy}{dt} = ky$  are the exponential functions*

$$y(t) = y(0)e^{kt}.$$

**Example: Population growth.** The CIA World Fact Book estimated that the population of Australia in July 2009 was 21,262,641 and that its 2009 growth rate was 1.195%. Estimate the population of Australia in July 2015, assuming that its growth rate remains constant.

**Example: Population growth.** The CIA World Fact Book estimated that the population of Australia in July 2009 was 21,262,641 and that its 2009 growth rate was 1.195%. Estimate the population of Australia in July 2015, assuming that its growth rate remains constant.

**Remark.** In population studies, the constant of proportionality  $k$  is often called the *(relative) growth rate*.

**Radioactive decay.** Radioactive substances decay by spontaneously emitting radiation. If  $m(t)$  is the mass remaining from an initial mass  $m_0$  then the relative decay rate has been found experimentally to be constant.



**Radioactive decay.** Radioactive substances decay by spontaneously emitting radiation. If  $m(t)$  is the mass remaining from an initial mass  $m_0$  then the relative decay rate has been found experimentally to be constant. This means that

$$\frac{dm}{dt} = km$$

where  $k < 0$ . The positive number  $-k$  is called the decay rate.

**Radioactive decay.** Radioactive substances decay by spontaneously emitting radiation. If  $m(t)$  is the mass remaining from an initial mass  $m_0$  then the relative decay rate has been found experimentally to be constant. This means that

$$\frac{dm}{dt} = km$$

where  $k < 0$ . The positive number  $-k$  is called the decay rate. The *half-life* of a radioactive substance is the time required for half of any given quantity to decay.

## Some Half-Lives.

<b>Yield</b>	<b>Fission Product</b>	<b>Half-life</b>
6.8%	Cesium-134	2 years
6.3%	Iodine-135	7 hours
6.1%	Zirconium-93	1.5M years
6.1%	Cesium-137	30 years
5.8%	Technetium-99	200k years
2.8%	Iodine-131	8 days

**Some Half-Lives.**

<b>Yield</b>	<b>Fission Product</b>	<b>Half-life</b>
6.8%	Cesium-134	2 years
6.3%	Iodine-135	7 hours
6.1%	Zirconium-93	1.5M years
6.1%	Cesium-137	30 years
5.8%	Technetium-99	200k years
2.8%	Iodine-131	8 days

**Example: Carbon dating.** Carbon-14 has a half-life of about 5730 years.

- (a) Suppose that  $m(t)$  denotes the mass of Carbon-14 present in a fossil at time  $t$ . Find a formula for  $m(t)$ .

### Some Half-Lives.

Yield	Fission Product	Half-life
6.8%	Cesium-134	2 years
6.3%	Iodine-135	7 hours
6.1%	Zirconium-93	1.5M years
6.1%	Cesium-137	30 years
5.8%	Technetium-99	200k years
2.8%	Iodine-131	8 days

**Example: Carbon dating.** Carbon-14 has a half-life of about 5730 years.

- (a) Suppose that  $m(t)$  denotes the mass of Carbon-14 present in a fossil at time  $t$ . Find a formula for  $m(t)$ .
- (b) The proportion of Carbon-14 in an animal fossil is 0.6%. Assume that when the animal died the proportion was 1%. Use this information to estimate when the animal died.

**Compound interest.** Suppose that an amount  $A_0$  is invested at an interest rate of  $r$  per annum and is compounded  $n$  times a year. Then the investment  $A(t)$  in  $t$  years is given by

$$A(t) = A_0 \left(1 + \frac{r}{n}\right)^{nt}.$$

**Compound interest.** Suppose that an amount  $A_0$  is invested at an interest rate of  $r$  per annum and is compounded  $n$  times a year. Then the investment  $A(t)$  in  $t$  years is given by

$$A(t) = A_0 \left(1 + \frac{r}{n}\right)^{nt}.$$

*Example.* If  $A_0 = \$1000$  and  $r = 6\%$  per annum then the investment in 3 years is worth:

**Compound interest.** Suppose that an amount  $A_0$  is invested at an interest rate of  $r$  per annum and is compounded  $n$  times a year. Then the investment  $A(t)$  in  $t$  years is given by

$$A(t) = A_0 \left(1 + \frac{r}{n}\right)^{nt}.$$

*Example.* If  $A_0 = \$1000$  and  $r = 6\%$  per annum then the investment in 3 years is worth:

$$\$1000(1.06)^3 = \$1191.02 \text{ with annual compounding } (n = 1)$$

$$\$1000(1.03)^6 = \$1194.05 \text{ with half-yearly compounding } (n = 2)$$

$$\$1000(1.015)^{12} = \$1195.62 \text{ with quarterly compounding } (n = 4)$$

$$\$1000(1.005)^{36} = \$1196.68 \text{ with monthly compounding } (n = 12)$$

$$\$1000 \left(1 + \frac{0.06}{365}\right)^{1095} = \$1197.20 \text{ with daily compounding } (n = 365).$$



**Compound interest.** Suppose that an amount  $A_0$  is invested at an interest rate of  $r$  per annum and is compounded  $n$  times a year. Then the investment  $A(t)$  in  $t$  years is given by

$$A(t) = A_0 \left(1 + \frac{r}{n}\right)^{nt}.$$

*Example.* If  $A_0 = \$1000$  and  $r = 6\%$  per annum then the investment in 3 years is worth:

$$\$1000(1.06)^3 = \$1191.02 \text{ with annual compounding } (n = 1)$$

$$\$1000(1.03)^6 = \$1194.05 \text{ with half-yearly compounding } (n = 2)$$

$$\$1000(1.015)^{12} = \$1195.62 \text{ with quarterly compounding } (n = 4)$$

$$\$1000(1.005)^{36} = \$1196.68 \text{ with monthly compounding } (n = 12)$$

$$\$1000 \left(1 + \frac{0.06}{365}\right)^{1095} = \$1197.20 \text{ with daily compounding } (n = 365).$$

If we let  $n \rightarrow \infty$ , we will be compounding the interest *continuously*.

We derive a formula for continuous compounding:

We derive a formula for continuous compounding:

We return now to the example of \$1000 invested at an interest rate of 6% per annum. If the interest is compounded continuously then the amount after three years will be

$$A(3) = \$1000e^{(0.06)3} = \$1197.22.$$

We derive a formula for continuous compounding:

We return now to the example of \$1000 invested at an interest rate of 6% per annum. If the interest is compounded continuously then the amount after three years will be

$$A(3) = \$1000e^{(0.06)^3} = \$1197.22.$$

Notice how close this is to the amount for daily compounding. However, the formula using continuous compounding is much easier to manipulate mathematically.


- 1 General exponential and logarithmic functions
  - Exponential growth and decay
  
- 2 Inverse functions
  - The inverse trigonometric functions
  - The hyperbolic functions

# The inverse trig functions

We now leave behind  $\ln x$  and  $e^x$  and examine inverse functions to another familiar class of examples, the trigonometric functions. Consider the graph of  $\sin x$ :


# The inverse trig functions

We now leave behind  $\ln x$  and  $e^x$  and examine inverse functions to another familiar class of examples, the trigonometric functions. Consider the graph of  $\sin x$ :



# The inverse trig functions

We now leave behind  $\ln x$  and  $e^x$  and examine inverse functions to another familiar class of examples, the trigonometric functions. Consider the graph of  $\sin x$ :




Notice that this is not one-to-one, but if we restrict  $x$  to lie in  $[-\frac{\pi}{2}, \frac{\pi}{2}]$ , then  $\sin x$  is one-to-one, hence it has an inverse. We denote this inverse by  $\sin^{-1} x$ .




Note that  $\sin^{-1} x$  is **NOT** equal to  $\frac{1}{\sin x}$ . Here is the graph of  $\sin^{-1} x$ :

Note that  $\sin^{-1} x$  is **NOT** equal to  $\frac{1}{\sin x}$ . Here is the graph of  $\sin^{-1} x$ :




Note that  $\sin^{-1} x$  is **NOT** equal to  $\frac{1}{\sin x}$ . Here is the graph of  $\sin^{-1} x$ :



We have

$$\sin^{-1} x = y \iff \sin y = x \quad \text{and} \quad -\frac{\pi}{2} \leq y \leq \frac{\pi}{2}.$$

Note that  $\sin^{-1} x$  is **NOT** equal to  $\frac{1}{\sin x}$ . Here is the graph of  $\sin^{-1} x$ :



We have

$$\sin^{-1} x = y \iff \sin y = x \quad \text{and} \quad -\frac{\pi}{2} \leq y \leq \frac{\pi}{2}.$$

That is, if  $x \in [-1, 1]$ , then  $\sin^{-1} x$  is the number between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$  whose sine is  $x$ . The domain of  $\sin^{-1} x$  is  $[-1, 1]$  and the range is  $[-\frac{\pi}{2}, \frac{\pi}{2}]$ .

The cancellation formulas for  $\sin^{-1}$  are:

$$\begin{aligned}\sin^{-1}(\sin x) &= x && \text{for } -\frac{\pi}{2} \leq x \leq \frac{\pi}{2} \\ \sin(\sin^{-1} x) &= x && \text{for } -1 \leq x \leq 1.\end{aligned}$$

The cancellation formulas for  $\sin^{-1}$  are:

$$\begin{aligned}\sin^{-1}(\sin x) &= x && \text{for } -\frac{\pi}{2} \leq x \leq \frac{\pi}{2} \\ \sin(\sin^{-1} x) &= x && \text{for } -1 \leq x \leq 1.\end{aligned}$$

Let's use implicit differentiation to calculate the derivative of  $\sin^{-1}$ :

The cancellation formulas for  $\sin^{-1}$  are:

$$\begin{aligned}\sin^{-1}(\sin x) &= x && \text{for } -\frac{\pi}{2} \leq x \leq \frac{\pi}{2} \\ \sin(\sin^{-1} x) &= x && \text{for } -1 \leq x \leq 1.\end{aligned}$$

Let's use implicit differentiation to calculate the derivative of  $\sin^{-1}$ :  
Therefore we have

$$\frac{d}{dx}(\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}}, \quad -1 < x < 1.$$

The cancellation formulas for  $\sin^{-1}$  are:

$$\begin{aligned}\sin^{-1}(\sin x) &= x && \text{for } -\frac{\pi}{2} \leq x \leq \frac{\pi}{2} \\ \sin(\sin^{-1} x) &= x && \text{for } -1 \leq x \leq 1.\end{aligned}$$

Let's use implicit differentiation to calculate the derivative of  $\sin^{-1}$ :  
Therefore we have


$$\frac{d}{dx}(\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}}, \quad -1 < x < 1.$$

**Example.** If  $f(x) = \sin^{-1}(x^2 - 1)$ , find (a) the domain of  $f$ , (b) the derivative  $f'(x)$ , and (c) the domain of  $f'$ .




What about  $\cos^{-1}$ ? Here is the graph of  $y = \cos x$ :

What about  $\cos^{-1}$ ? Here is the graph of  $y = \cos x$ :




What about  $\cos^{-1}$ ? Here is the graph of  $y = \cos x$ :




We must again restrict the domain of  $\cos x$  so that it is one-to-one. We restrict  $x$  to lie in  $[0, \pi]$ , then define the inverse function, denoted by  $\cos^{-1} x$ .

Its graph looks like this:




Its graph looks like this:



We summarize properties of  $\cos^{-1}$ :

- $\cos^{-1} x$  has domain  $[-1, 1]$  and range  $[0, \pi]$ .


Its graph looks like this:



We summarize properties of  $\cos^{-1}$ :

- $\cos^{-1} x$  has domain  $[-1, 1]$  and range  $[0, \pi]$ .
- $\cos^{-1} x = y \iff \cos y = x \quad \text{and} \quad 0 \leq y \leq \pi.$


Its graph looks like this:



We summarize properties of  $\cos^{-1}$ :

- $\cos^{-1} x$  has domain  $[-1, 1]$  and range  $[0, \pi]$ .
- $\cos^{-1} x = y \iff \cos y = x \quad \text{and} \quad 0 \leq y \leq \pi.$
- $\cos^{-1}(\cos x) = x \quad \text{for } 0 \leq x \leq \pi.$

Its graph looks like this:




We summarize properties of  $\cos^{-1}$ :

- $\cos^{-1} x$  has domain  $[-1, 1]$  and range  $[0, \pi]$ .
- $\cos^{-1} x = y \iff \cos y = x \quad \text{and} \quad 0 \leq y \leq \pi.$
- $\cos^{-1}(\cos x) = x \quad \text{for } 0 \leq x \leq \pi.$
- $\cos(\cos^{-1} x) = x \quad \text{for } -1 \leq x \leq 1.$



Its graph looks like this:




We summarize properties of  $\cos^{-1}$ :

- $\cos^{-1} x$  has domain  $[-1, 1]$  and range  $[0, \pi]$ .
- $\cos^{-1} x = y \iff \cos y = x \quad \text{and} \quad 0 \leq y \leq \pi.$
- $\cos^{-1}(\cos x) = x \quad \text{for } 0 \leq x \leq \pi.$
- $\cos(\cos^{-1} x) = x \quad \text{for } -1 \leq x \leq 1.$
- $\frac{d}{dx}(\cos^{-1} x) = -\frac{1}{\sqrt{1-x^2}} \quad -1 < x < 1.$


What about  $\tan^{-1}$ ? Here is the graph of  $y = \tan x$ , where we restrict  $x$  to lie in  $(-\frac{\pi}{2}, \frac{\pi}{2})$ :

What about  $\tan^{-1}$ ? Here is the graph of  $y = \tan x$ , where we restrict  $x$  to lie in  $(-\frac{\pi}{2}, \frac{\pi}{2})$ :




With  $x$  restricted in this way,  $\tan x$  is one-to-one, hence it has an inverse function, denoted by  $\tan^{-1}$ .

With  $x$  restricted in this way,  $\tan x$  is one-to-one, hence it has an inverse function, denoted by  $\tan^{-1}$ . The graph of  $\tan^{-1}$  looks like this:




With  $x$  restricted in this way,  $\tan x$  is one-to-one, hence it has an inverse function, denoted by  $\tan^{-1}$ . The graph of  $\tan^{-1}$  looks like this:



We summarize some properties of  $\tan^{-1}$ :

- $\tan^{-1} x = y \iff \tan y = x \quad \text{and} \quad -\frac{\pi}{2} < y < \frac{\pi}{2}.$


With  $x$  restricted in this way,  $\tan x$  is one-to-one, hence it has an inverse function, denoted by  $\tan^{-1}$ . The graph of  $\tan^{-1}$  looks like this:



We summarize some properties of  $\tan^{-1}$ :

- $\tan^{-1} x = y \iff \tan y = x \quad \text{and} \quad -\frac{\pi}{2} < y < \frac{\pi}{2}.$
- $\lim_{x \rightarrow \infty} \tan^{-1} x = \frac{\pi}{2} \quad \lim_{x \rightarrow -\infty} \tan^{-1} x = -\frac{\pi}{2}.$

With  $x$  restricted in this way,  $\tan x$  is one-to-one, hence it has an inverse function, denoted by  $\tan^{-1}$ . The graph of  $\tan^{-1}$  looks like this:



We summarize some properties of  $\tan^{-1}$ :

- $\tan^{-1} x = y \iff \tan y = x \quad \text{and} \quad -\frac{\pi}{2} < y < \frac{\pi}{2}.$
- $\lim_{x \rightarrow \infty} \tan^{-1} x = \frac{\pi}{2} \quad \lim_{x \rightarrow -\infty} \tan^{-1} x = -\frac{\pi}{2}.$
- $\frac{d}{dx}(\tan^{-1} x) = \frac{1}{1+x^2}.$



**Remarks.** Note that  $\sin^{-1} x$ ,  $\cos^{-1} x$ ,  $\tan^{-1} x$ , etc are often also called arcsin, arccos, arctan, etc. Also, note that one can discuss the definition and basic properties (such as derivatives) of the inverse functions to sec, csc, and cot.

**Remarks.** Note that  $\sin^{-1} x$ ,  $\cos^{-1} x$ ,  $\tan^{-1} x$ , etc are often also called arcsin, arccos, arctan, etc. Also, note that one can discuss the definition and basic properties (such as derivatives) of the inverse functions to sec, csc, and cot.

**Example.** Simplify the expression  $\cos(\tan^{-1} x)$ .

**Remarks.** Note that  $\sin^{-1} x$ ,  $\cos^{-1} x$ ,  $\tan^{-1} x$ , etc are often also called arcsin, arccos, arctan, etc. Also, note that one can discuss the definition and basic properties (such as derivatives) of the inverse functions to sec, csc, and cot.

**Example.** Simplify the expression  $\cos(\tan^{-1} x)$ .

**Example.** Differentiate the function given by  $f(x) = x \tan^{-1} \sqrt{x}$ .

# The hyperbolic functions

There is another important and useful class of functions we will briefly study, known as the **hyperbolic functions**. They are similar to trigonometric functions, but are closely related to the hyperbola instead of to the circle. Here are the definitions:

# The hyperbolic functions

There is another important and useful class of functions we will briefly study, known as the **hyperbolic functions**. They are similar to trigonometric functions, but are closely related to the hyperbola instead of to the circle. Here are the definitions:

- $\sinh x = \frac{e^x - e^{-x}}{2}.$

# The hyperbolic functions

There is another important and useful class of functions we will briefly study, known as the **hyperbolic functions**. They are similar to trigonometric functions, but are closely related to the hyperbola instead of to the circle. Here are the definitions:

- $\sinh x = \frac{e^x - e^{-x}}{2}.$
- $\cosh x = \frac{e^x + e^{-x}}{2}.$

# The hyperbolic functions

There is another important and useful class of functions we will briefly study, known as the **hyperbolic functions**. They are similar to trigonometric functions, but are closely related to the hyperbola instead of to the circle. Here are the definitions:

- $\sinh x = \frac{e^x - e^{-x}}{2}.$
- $\cosh x = \frac{e^x + e^{-x}}{2}.$
- $\tanh x = \frac{\sinh x}{\cosh x}.$

# The hyperbolic functions

There is another important and useful class of functions we will briefly study, known as the **hyperbolic functions**. They are similar to trigonometric functions, but are closely related to the hyperbola instead of to the circle. Here are the definitions:

- $\sinh x = \frac{e^x - e^{-x}}{2}.$
- $\cosh x = \frac{e^x + e^{-x}}{2}.$
- $\tanh x = \frac{\sinh x}{\cosh x}.$
- $\operatorname{csch} x = \frac{1}{\sinh x}.$



# The hyperbolic functions

There is another important and useful class of functions we will briefly study, known as the **hyperbolic functions**. They are similar to trigonometric functions, but are closely related to the hyperbola instead of to the circle. Here are the definitions:

- $\sinh x = \frac{e^x - e^{-x}}{2}.$
- $\cosh x = \frac{e^x + e^{-x}}{2}.$
- $\tanh x = \frac{\sinh x}{\cosh x}.$
- $\operatorname{csch} x = \frac{1}{\sinh x}.$
- $\operatorname{sech} x = \frac{1}{\cosh x}.$

# The hyperbolic functions

There is another important and useful class of functions we will briefly study, known as the **hyperbolic functions**. They are similar to trigonometric functions, but are closely related to the hyperbola instead of to the circle. Here are the definitions:

- $\sinh x = \frac{e^x - e^{-x}}{2}.$
- $\cosh x = \frac{e^x + e^{-x}}{2}.$
- $\tanh x = \frac{\sinh x}{\cosh x}.$
- $\operatorname{csch} x = \frac{1}{\sinh x}.$
- $\operatorname{sech} x = \frac{1}{\cosh x}.$
- $\operatorname{coth} x = \frac{\cosh x}{\sinh x}.$

We draw the graphs of  $\sinh$ ,  $\cosh$ , and  $\tanh$ :




Figure 1:  $y = \sinh x$




Figure 2:  $y = \cosh x$




Figure 3:  $y = \tanh x$

**Properties of  $\cosh$  and  $\sinh$ .** Notice how similar the following properties are to those of the trigonometric functions:

**Properties of cosh and sinh.** Notice how similar the following properties are to those of the trigonometric functions:

- $\sinh(-x) = -\sinh x$ .

**Properties of cosh and sinh.** Notice how similar the following properties are to those of the trigonometric functions:

- $\sinh(-x) = -\sinh x$ .
- $\cosh(-x) = \cosh x$ .

**Properties of cosh and sinh.** Notice how similar the following properties are to those of the trigonometric functions:

- $\sinh(-x) = -\sinh x$ .
- $\cosh(-x) = \cosh x$ .
- $\cosh^2 x - \sinh^2 x = 1$ .

**Properties of cosh and sinh.** Notice how similar the following properties are to those of the trigonometric functions:

- $\sinh(-x) = -\sinh x$ .
- $\cosh(-x) = \cosh x$ .
- $\cosh^2 x - \sinh^2 x = 1$ .
- $1 - \tanh^2 x = \operatorname{sech}^2 x$ .



**Properties of cosh and sinh.** Notice how similar the following properties are to those of the trigonometric functions:

- $\sinh(-x) = -\sinh x$ .
- $\cosh(-x) = \cosh x$ .
- $\cosh^2 x - \sinh^2 x = 1$ .
- $1 - \tanh^2 x = \operatorname{sech}^2 x$ .

We prove the third property in the above list:

**Properties of cosh and sinh.** Notice how similar the following properties are to those of the trigonometric functions:

- $\sinh(-x) = -\sinh x$ .
- $\cosh(-x) = \cosh x$ .
- $\cosh^2 x - \sinh^2 x = 1$ .
- $1 - \tanh^2 x = \operatorname{sech}^2 x$ .

We prove the third property in the above list:

We now can get some insight into why these functions are called “hyperbolic”. Notice that if  $t$  is a real number, then the point  $(\cosh t, \sinh t)$  lies on the right branch of the hyperbola  $x^2 - y^2 = 1$ . It lies on the right branch because  $\cosh t \geq 1$ :

We now list formulas for the derivatives of the hyperbolic functions. Note the similarities to the derivatives of the trigonometric functions (and the differences!):

- $\frac{d}{dx}(\sinh x) = \cosh x.$

We now list formulas for the derivatives of the hyperbolic functions. Note the similarities to the derivatives of the trigonometric functions (and the differences!):

- $\frac{d}{dx}(\sinh x) = \cosh x.$
- $\frac{d}{dx}(\cosh x) = \sinh x.$

We now list formulas for the derivatives of the hyperbolic functions. Note the similarities to the derivatives of the trigonometric functions (and the differences!):

- $\frac{d}{dx}(\sinh x) = \cosh x.$
- $\frac{d}{dx}(\cosh x) = \sinh x.$
- $\frac{d}{dx}(\tanh x) = \operatorname{sech}^2 x.$

We now list formulas for the derivatives of the hyperbolic functions. Note the similarities to the derivatives of the trigonometric functions (and the differences!):

- $\frac{d}{dx}(\sinh x) = \cosh x.$
- $\frac{d}{dx}(\cosh x) = \sinh x.$
- $\frac{d}{dx}(\tanh x) = \operatorname{sech}^2 x.$
- $\frac{d}{dx}(\operatorname{csch} x) = -\operatorname{csch} x \coth x.$

We now list formulas for the derivatives of the hyperbolic functions. Note the similarities to the derivatives of the trigonometric functions (and the differences!):

- $\frac{d}{dx}(\sinh x) = \cosh x.$
- $\frac{d}{dx}(\cosh x) = \sinh x.$
- $\frac{d}{dx}(\tanh x) = \operatorname{sech}^2 x.$
- $\frac{d}{dx}(\operatorname{csch} x) = -\operatorname{csch} x \coth x.$
- $\frac{d}{dx}(\operatorname{sech} x) = -\operatorname{sech} x \tanh x.$

We now list formulas for the derivatives of the hyperbolic functions. Note the similarities to the derivatives of the trigonometric functions (and the differences!):

- $\frac{d}{dx}(\sinh x) = \cosh x.$
- $\frac{d}{dx}(\cosh x) = \sinh x.$
- $\frac{d}{dx}(\tanh x) = \operatorname{sech}^2 x.$
- $\frac{d}{dx}(\operatorname{csch} x) = -\operatorname{csch} x \coth x.$
- $\frac{d}{dx}(\operatorname{sech} x) = -\operatorname{sech} x \tanh x.$
- $\frac{d}{dx}(\coth x) = -\operatorname{csch}^2 x.$



We now list formulas for the derivatives of the hyperbolic functions. Note the similarities to the derivatives of the trigonometric functions (and the differences!):

- $\frac{d}{dx}(\sinh x) = \cosh x.$
- $\frac{d}{dx}(\cosh x) = \sinh x.$
- $\frac{d}{dx}(\tanh x) = \operatorname{sech}^2 x.$
- $\frac{d}{dx}(\operatorname{csch} x) = -\operatorname{csch} x \coth x.$
- $\frac{d}{dx}(\operatorname{sech} x) = -\operatorname{sech} x \tanh x.$
- $\frac{d}{dx}(\coth x) = -\operatorname{csch}^2 x.$

**Example.** Find the derivative:  $y = e^{\cosh 3x}.$

**Example.** If  $\tanh x = \frac{4}{5}$ , find the values of the other hyperbolic functions at  $x$ .

**Example.** If  $\tanh x = \frac{4}{5}$ , find the values of the other hyperbolic functions at  $x$ .

**Remark.** In the textbook there are similar formulas for the inverse hyperbolic functions and their derivatives. Although you will have webassign questions and a tutorial quiz on the hyperbolic functions and their inverses, I will not emphasize them on the final exam to the same extent as the trigonometric functions and their inverses. **That is, you are expected to be familiar with hyperbolic functions and their inverses. I may ask you questions involving these functions on the final exam, but if I do, they will be very basic questions.**

# Indeterminate Forms and L'Hospital's Rule

Earlier in the semester we discussed limits of the form  $\frac{\infty}{\infty}$ . Such limits could be

# Indeterminate Forms and L'Hospital's Rule

Earlier in the semester we discussed limits of the form  $\frac{\infty}{\infty}$ . Such limits could be

(a) zero, e.g.  $\lim_{x \rightarrow \infty} \frac{2x}{x^2 + 1} = 0$ ;

# Indeterminate Forms and L'Hospital's Rule

Earlier in the semester we discussed limits of the form  $\frac{\infty}{\infty}$ . Such limits could be

- (a) zero, e.g.  $\lim_{x \rightarrow \infty} \frac{2x}{x^2 + 1} = 0$ ;
- (b) infinite, e.g.  $\lim_{x \rightarrow \infty} \frac{x^2 + 6}{x} = \infty$ ;

# Indeterminate Forms and L'Hospital's Rule

Earlier in the semester we discussed limits of the form  $\frac{\infty}{\infty}$ . Such limits could be

(a) zero, e.g.  $\lim_{x \rightarrow \infty} \frac{2x}{x^2 + 1} = 0$ ;

(b) infinite, e.g.  $\lim_{x \rightarrow \infty} \frac{x^2 + 6}{x} = \infty$ ;

(c) or somewhere between, e.g.  $\lim_{x \rightarrow \infty} \frac{3x^2 + 6}{x^2 - x} = 3$ .

# Indeterminate Forms and L'Hospital's Rule

Earlier in the semester we discussed limits of the form  $\frac{\infty}{\infty}$ . Such limits could be

(a) zero, e.g.  $\lim_{x \rightarrow \infty} \frac{2x}{x^2 + 1} = 0$ ;

(b) infinite, e.g.  $\lim_{x \rightarrow \infty} \frac{x^2 + 6}{x} = \infty$ ;

(c) or somewhere between, e.g.  $\lim_{x \rightarrow \infty} \frac{3x^2 + 6}{x^2 - x} = 3$ .

Because one cannot say in advance exactly what a limit of the form  $\frac{\infty}{\infty}$  will be, we call it an *indeterminate form*.



# Indeterminate Forms and L'Hospital's Rule

Earlier in the semester we discussed limits of the form  $\frac{\infty}{\infty}$ . Such limits could be

- (a) zero, e.g.  $\lim_{x \rightarrow \infty} \frac{2x}{x^2 + 1} = 0$ ;
- (b) infinite, e.g.  $\lim_{x \rightarrow \infty} \frac{x^2 + 6}{x} = \infty$ ;
- (c) or somewhere between, e.g.  $\lim_{x \rightarrow \infty} \frac{3x^2 + 6}{x^2 - x} = 3$ .

Because one cannot say in advance exactly what a limit of the form  $\frac{\infty}{\infty}$  will be, we call it an *indeterminate form*. Other indeterminate forms include

$$0 \cdot \infty, \quad \frac{0}{0}, \quad \infty - \infty, \quad 0^0, \quad \infty^0 \quad \text{and} \quad 0^\infty.$$

Each of the above limits (a), (b) and (c) can be calculated using the standard trick of dividing the denominator through by the fastest growing term. It is easy to tell which terms grow fastest because we are working with polynomials. What if this is not the case?

Each of the above limits (a), (b) and (c) can be calculated using the standard trick of dividing the denominator through by the fastest growing term. It is easy to tell which terms grow fastest because we are working with polynomials. What if this is not the case? What about the following limits?

$$\lim_{x \rightarrow \infty} \frac{e^x}{x}$$

$$\lim_{x \rightarrow \infty} \frac{\ln x}{x}$$

Each of the above limits (a), (b) and (c) can be calculated using the standard trick of dividing the denominator through by the fastest growing term. It is easy to tell which terms grow fastest because we are working with polynomials. What if this is not the case? What about the following limits?

$$\lim_{x \rightarrow \infty} \frac{e^x}{x}$$

$$\lim_{x \rightarrow \infty} \frac{\ln x}{x}$$

In each case, does the denominator win out, or the numerator, or is there some kind of compromise?

One way to solve this problem is to look at the derivatives.

One way to solve this problem is to look at the derivatives.  
If, for example we could verify that

$$\lim_{x \rightarrow \infty} \frac{f'(x)}{g'(x)} = 0$$

then this would suggest that the *rate* of increase of  $g$  is greater than that of  $f$ .

One way to solve this problem is to look at the derivatives.  
If, for example we could verify that

$$\lim_{x \rightarrow \infty} \frac{f'(x)}{g'(x)} = 0$$

then this would suggest that the *rate* of increase of  $g$  is greater than that of  $f$ . This in turn would suggest that

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = 0.$$

In fact, this kind of intuitive reasoning is generally true!

## L'Hôpital's rule

Suppose that  $f$  and  $g$  are differentiable functions,  $a \in \mathbb{R}$ , and  $g'(x) \neq 0$ , except possibly at  $a$ . Suppose also that either one of the two following conditions hold:

- $f(x) \rightarrow 0$  and  $g(x) \rightarrow 0$  as  $x \rightarrow a$ ;



## L'Hôpital's rule

Suppose that  $f$  and  $g$  are differentiable functions,  $a \in \mathbb{R}$ , and  $g'(x) \neq 0$ , except possibly at  $a$ . Suppose also that either one of the two following conditions hold:

- $f(x) \rightarrow 0$  and  $g(x) \rightarrow 0$  as  $x \rightarrow a$ ;
- $f(x) \rightarrow \pm\infty$  and  $g(x) \rightarrow \pm\infty$  as  $x \rightarrow a$ .

## L'Hôpital's rule

Suppose that  $f$  and  $g$  are differentiable functions,  $a \in \mathbb{R}$ , and  $g'(x) \neq 0$ , except possibly at  $a$ . Suppose also that either one of the two following conditions hold:

- $f(x) \rightarrow 0$  and  $g(x) \rightarrow 0$  as  $x \rightarrow a$ ;
- $f(x) \rightarrow \pm\infty$  and  $g(x) \rightarrow \pm\infty$  as  $x \rightarrow a$ .

If

$$\lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

exists or is  $\pm\infty$ , then

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}.$$