

T H E

Diabetes ***Protocol***

GLUCOSE. BALANCE. CONTROL.

BY

Swetha Ashok

WWW.THEGUTGURU.CO.IN

GUTGURU PROTOCOL

Table of Contents

01 Understanding Blood Sugar

02 Glycaemic Index: Indian Foods

03 Smart Carb Swaps

04 The Diabetic Plate Formula

05 Common Diabetes Mistakes

06 Myth Busters

CHAPTER 01

Understanding Blood Sugar

Type 2 diabetes is largely a dietary condition — and dietary intervention is the most powerful tool available.



Insulin Resistance

When cells stop responding to insulin, glucose accumulates in the blood. This begins years before a diabetes diagnosis — during the pre-diabetic window.



Postprandial Spikes

It's not just fasting glucose that matters. Post-meal spikes (PP2BS > 140 mg/dL) cause vascular, nerve, and kidney damage even when fasting sugar is normal.



Gut Microbiome & Glucose

Gut bacteria directly influence how quickly glucose is absorbed. A diverse, fibre-fed microbiome naturally blunts blood sugar spikes.




Inflammation Link

Chronic blood sugar elevation triggers systemic inflammation — increasing risk of cardiovascular disease, kidney disease, and nerve damage.

CHAPTER 02

Glycaemic Index: *Indian Foods*



The glycaemic index measures how fast a food raises blood sugar. Pairing any carbohydrate with protein, fat, or fibre reduces its effective GI significantly.



KEY TAKEAWAY

The 15-minute walk rule: a brisk walk within 15 minutes of eating reduces post-meal blood glucose by 20–30% — as effective as some oral medications.

CHAPTER 03

High GI vs Low GI Swaps

Every high-GI meal is a blood sugar spike. Simple swaps — without giving up Indian food — make a measurable difference.

CHOOSE

Low-GI Indian Choices

These keep blood sugar stable, improve insulin sensitivity, and still feel satisfying.

Brown rice (small portion)

Ragi roti

Dalia upma

Rajma + rice (combo)

Oat idli

Moong dal cheela

Barley khichdi

Chana chaat

REDUCE

High-GI Triggers

These cause rapid glucose spikes — particularly dangerous when eaten alone in large portions.

White rice (large portion)

Maida roti / naan

Biscuits / toast breakfast

Instant noodles

White bread

Sweetened yoghurt

Fruit juice


Mango (large portions)

The key is not elimination but context: white rice with dal, ghee, and sabzi has a lower effective GI than white rice eaten alone.

CHAPTER 04

The Diabetic Plate Formula

THE GUTGURU BLOOD SUGAR METHOD

**Non-Starchy Vegetables**

Palak, methi, lauki, broccoli, karela (½ plate)

+

**Lean Protein**

Dal, paneer, eggs, fish, chana (¼ plate)

+

**Low-GI Carbohydrate**

Brown rice, ragi, rajma, dalia (¼ plate)

+

**Healthy Fat**

Ghee (small), mustard oil, walnuts, flaxseed

EXAMPLE MEAL

Palak + moong dal + small brown rice + 1 tsp ghee**FBS <100 · PP2BS <140 · HbA1c <6.5%**

CLINICAL BLOOD SUGAR TARGETS

CHAPTER 05

Common *Diabetes* Mistakes

**Switching to "sugar-free" products freely**

Many "sugar-free" products contain maltitol, sorbitol, or refined starch — all of which spike blood sugar in diabetics.

**Eating fruit freely because "it's natural sugar"**

Fruits contain fructose and glucose. Large portions of high-GI fruits (mango, grapes, banana) cause measurable blood sugar spikes.

**Skipping meals to lower blood sugar**

Meal skipping causes dangerous glucose fluctuations — particularly dangerous for those on insulin or sulphonylurea medications.

**Depending on medication without changing diet**

Diet is the most powerful HbA1c-lowering tool available — more effective than medication in early-stage Type 2 diabetes.

**Eliminating rice entirely from the diet**

Portion control and pairing rice with protein and fibre is the evidence-based approach — complete elimination is unsustainable and unnecessary.

CHAPTER 06

Quick Myth Busters

MYTH Diabetics can never eat rice again.

TRUTH **Portion size and food pairing are what matters.** A small serving of rice eaten with dal, sabzi, and ghee has a far lower glucose impact than rice alone.

MYTH Jaggery and honey are safe alternatives to sugar for diabetics.

TRUTH **Both spike blood glucose similarly to white sugar.** Jaggery is 65–80% sucrose. Neither is safe for uncontrolled blood sugar.

MYTH If you take diabetes medication, diet doesn't matter much.

TRUTH **Diet is the most effective HbA1c-reduction tool available** — more so than medication in early T2D. The medication dose required decreases significantly with dietary improvement.

TAKE THIS HOME

Your 3 Quick Wins

Three specific actions you can start this week — no equipment, no cost, measurable results.

1

Walk 15 Minutes After Every Meal

A **15-minute brisk walk within 30 minutes of eating** reduces post-meal blood glucose by 20–30% — as effective as some oral medications for PP2BS control. Do it after all three meals for maximum impact.

AFTER BREAKFAST, LUNCH, AND DINNER

2

Eat Your Protein and Vegetables First

At every meal, **eat your dal, sabzi, or protein first — then your rice or roti**. This simple sequence reduces the glucose spike of the entire meal because fibre and protein slow carbohydrate absorption when eaten first.

AT EVERY MEAL, STARTING TODAY

3

Test and Record Your Post-Meal Blood Sugar

Use a glucometer to **test your blood sugar 2 hours after your typical meals** and record the results for 7 days. You will discover exactly which of your meals are spiking you — and that knowledge is more valuable than any generic food list.

FOR 7 CONSECUTIVE DAYS

MORE AT www.thegutguru.co.in →



READY TO PERSONALIZE?

Book a 1-on-1 Consultation

Get a clinical nutrition plan tailored to your diabetes / blood sugar management, body, and lifestyle.

 www.thegutguru.co.in



+91 9769921409



gutguru.swetha@gmail.com