

# ASSIGNMENT COVER SHEET

ANU College of Engineering and  
Computer Science

Australian National University

Canberra ACT 0200 Australia

[www.anu.edu.au](http://www.anu.edu.au)

+61 2 6125 5254

This coversheet must be attached to the front of your assessment

The assessment is due on 28/08/2023, 9:05 AM unless otherwise specified in the course outline.

Student ID	U7540836
Student Name	Nanthawat Anancharoenpakorn
Course Code	COMP6261
Course Name	Information Theory
Assignment Item	Assignment 1
Due Date	28/08/2023,
Date Submitted	26/08/2023

I declare that this work:

- ☐ upholds the principles of academic integrity, as defined in the ANU Policy: [Code of Practice for Student Academic Integrity](#);
- ☐ is original, except where collaboration (for example group work) has been authorized in writing by the course convener in the course outline and/or Wattle site;
- ☐ is produced for the purposes of this assessment task and has not been submitted for assessment in any other context, except where authorized in writing by the course convener.
- ☐ gives appropriate acknowledgment of the ideas, scholarship, and intellectual property of others insofar as these have been used.
- ☐ in no part involves copying, cheating, collusion, fabrication, plagiarism, or recycling.

Signature	Nanthawat Anancharoenpakorn
-----------	-----------------------------

Question 1 : section II

• Random variable  $X = \begin{Bmatrix} XXXX \\ \dots XXXX \end{Bmatrix}$ ,  $A_X = \{A, M, S\}$ ,  $P_X = \frac{1}{3}$

• once win 3 rounds, game over, max round = 5

• random v.  $Y = \begin{Bmatrix} y \\ A_Y = \{3, 4, 5\} \end{Bmatrix}$ ,  $P_Y = \square$

→ Compute all relevant probability

3 teams play against each other until one of them win 3 rounds.

with 3 game | there are 3 teams. Hence 3 ways.

$$\text{with } (Y=3) = 3 \times \left(\frac{1}{3}\right)^3 = \frac{1}{3}$$

with 4 game | let fix A to be the winner then

$\square \rightarrow A$   ${}^3C_2$  since A must have won 2 out of 3 games

$$\text{then } {}^3C_2 = \frac{3!}{1!2!} = 3. \text{ since there are 3 teams,}$$

there are  $3 \times 3 = 9$  events. Since  $AA \neq AASA$

then we need to multiply by 2 which is  $9 \times 2 = 18$

ways. Hence, 18 ways with  $(Y=4) = 18 \times \left(\frac{1}{3}\right)^4 = \frac{2}{9}$

Summary

Y	n. events	each prob	game prob.
y=3	3	$(0.33)^3$	$\frac{1}{9}$
y=4	18	$(0.33)^4$	$\frac{2}{9}$
y=5	36	$(0.33)^5$	$\frac{4}{27}$

(a)

$$h(P=AMSSS) = \log \frac{1}{(1/3)^5} = 4.92 \text{ bH}$$

$$(b) H(P) = 3 \left(\frac{1}{27}\right) \log(27) + 18 \left(\frac{1}{81}\right) \log(81) +$$

$$36 \left(\frac{1}{243}\right) \log(243) = 3.11 \text{ bH}$$

$$(c) E(Y) = \frac{1}{9} \log(9) + \frac{2}{9} \log\left(\frac{9}{2}\right) + \frac{4}{27} \log\left(\frac{27}{4}\right)$$

$$= 1.24 \text{ bH}$$

$$(d) H(P|Q=4) = 18 \times \left(\frac{1}{3}\right)^4 \times \log\left(\frac{1}{(1/3)^4}\right) = 1.4 \text{ bH}$$

$$(e) H(P|Q) = H(P) + H(Q|P) - H(Q) \text{ since } H(Q|P) = 0$$

because Q is a deterministic f P Hence  $H(P|Q) = 1.84$

(f)  $H(Q|P) = 0$  because Q is deterministic of P, meaning that knowing P will allow to know Q with certainty example.  $P = A \square \square \square$ , then  $Q = 3$

(g) If (civilization win first round  $\neq$  civil win the last rnd)

then 33%  $\rightarrow$  N 66%  $\rightarrow$  normal

If (civil 101) 4 consecutive then 50%  $\rightarrow$  N

50%  $\rightarrow$  normal.

Summary

y=3	3	$(1/3)^3$	[same]
y=4	12	$(1/3)^4$	
	6	$(1/3)^4 \times \frac{2}{3} \cdot 66\%$	

y=5

$$H(Q) =$$

(h)



Question 2

- Value (v)  $V: \{A, 2, 3, \dots, K\} \quad |V| = 13$
- Suit (s)  $S: \{\text{heart, diamond, } \dots\} \quad |S| = 4$
- Color (c)  $C: \{\text{black, red}\} \quad |C| = 2$
- Face (f)  $F: \{0, 1\} \quad |F| = 2$

- (1) (a) since face card are  $\{J, Q, K\}$  which has 2 reds per value. Then  $\text{prob}(c=\text{red}, f=1) = \frac{6}{52}$   
 $h(c=\text{red}, f=1) = \log_2 \frac{1}{(6/52)} = 3.1134 \text{ bits}$

(b)  $h(V=K | f=1) = \log_2 \frac{1}{(4/12)} = 1.5849 \text{ bits}$   
 since there are 4 kings and (4/12) 12 face cards.

(c) since there are 4 suits  $H(S) = -\sum p(s) \log_2 p(s)$   
 $= 4 \times \frac{1}{4} \log_2 \left(\frac{1}{4}\right) = 2 \text{ bits}$  and  $H(V, S) = 5.7 \text{ bits}$

(d)  $I(V; S) = H(V) + H(S) - H(V, S) = 0$  there is independent.

(e) according to data processing inequality and as color is determined by  $S$   $I(V; C) \leq I(V; S)$  hence  $I(V; C) = 0$  since mutual information can't be negative

- (2) Remove 13 2s and 3 hearts of 2 to 5 and, A♦ and K♣  
 • Value =  $\{A, \dots, K\}$ , Face =  $\{0, 1\}$ , Suit =  $\{\text{heart, diamond, clubs}\}$   $|S| = 3$ , color =  $\{\text{black, red}\}$ .

(a)  $H(S) = \frac{8}{32} \log_2 \left(\frac{1}{8/32}\right) + \frac{11}{32} \log_2 \left(\frac{1}{11/32}\right) + \frac{13}{32} \log_2 \left(\frac{1}{13/32}\right)$

$H(S) = 1.5575 \text{ bits}$ ,  $H(V, S) = 9 \text{ bits}$

(b)  $I(V; S) = H(V) + H(S) - H(V, S) = 5 + 1.5575 - 9 = 1.5575$ . The reason it different because we have remove some of the card which lead to the result that suits are not uniform anymore. Then knowing  $S$  provide information about  $V$ .

(c)  $I(V; S|C) = 0.107$ , because we reduce the card a lot which lead to less possibility for black.

eg. if we know the card is black, then we know for sure that it must be club. Knowing color change the probability make it more certain when we know color of the card.

Question 3

(a)  $I(X; Y) = \frac{1}{2} \log(2) + \frac{1}{2} \log(2) = 2 \text{ bits}$  because it dependent to each other completely.

(b)  $I(X; Y) = H(X) - H(X|Y) = 0.58$

(c) since the first and last element are the same, we can compute by log sum inequality.  $H_2 - H_1 = -2 \frac{(p_1 + p_2)}{2} \log \frac{(p_1 + p_2)}{2} + p_1 \log p_1 + p_2 \log p_2 \geq 0$ .

$H_2 \geq H_1$  imply that entropy will be larger for most uniform distribution.

(d)  $X: \{X, X \in X = \{1, 2, \dots, m\}, P_X = \{q_1, \dots, q_m\} \mid q_i \text{ in decreasing order}\}$  • new r.v.  $Y$  with 2 more outcome that each prob is divided by 3 and the last term is  $\frac{2}{3}$ .

$H(X) = -\sum_{i=1}^m q_i \log q_i$ ,  $H(Y) = -\sum_{i=1}^m \frac{q_i}{3} \log \frac{q_i}{3} + \frac{2}{3} \log \frac{2}{3}$   
 write  $H(Y)$  in term of  $H(X)$ :

$H(Y) = -\sum_{i=1}^m \frac{q_i}{3} \log \frac{q_i}{3} + \frac{2}{3} \log \frac{2}{3}$

$= -\frac{1}{3} \sum_{i=1}^m q_i \log q_i + \frac{2}{3} \log \frac{2}{3}$

$= -\frac{1}{3} H(X) + \frac{2}{3} \log \frac{2}{3}$

Question 4

$$Y : (y, \{0, 2\}, \{0.5, 0.5\})$$

$$X : (x, \{a, b, c\}, p_X)$$

$y \backslash x$	$a$	$b$	$c$
0	$\frac{1}{5}$	$\frac{2}{5}$	$\frac{1}{5}$
2	$\frac{2}{5}$	$\frac{1}{5}$	$\frac{1}{5}$

$$(a) I(X; Y) = H(X) + H(Y) - H(X, Y)$$

$$H(X) = 1.58 \text{ bit}$$

$$H(Y) = 1 \text{ bit}$$

$$H(X|Y) = 0.5(1.37) + 0.5(1.37) = 1.37 \text{ bit}$$

$$H(X, Y) = 1 + 1.37 = 2.37$$

$$\text{hence, } I(X; Y) = 1.58 + 1 - 2.37 = 0.21$$

$$(b) D_{KL}(p||q) = \sum_x p(x) \log \frac{p(x)}{q(x)}$$

$$I(X; Y) = \sum_{x,y} p(x,y) \log_2 \frac{p(x,y)}{p(x)p(y)}$$

let  $y$  be a binary random v. ( $y, y \in \{0, 2\}$   $p_y = \{\frac{1}{2}, \frac{1}{2}\}$ )

&

$$I(X; y) = \frac{1}{2} \sum_x p(x|Y=1) \log \frac{p(x|Y=1)}{m(x)} + \frac{1}{2} \sum_x p(x|Y=0) \log \frac{p(x|Y=0)}{m(x)}$$

$$m(x) = \frac{p(x|Y=1) + p(x|Y=0)}{2}$$

$$I(X; y) = \frac{1}{2} D_{KL}(p||m) + \frac{1}{2} D_{KL}(q||m)$$

$$(c) \text{ As } p(Z=X|Y=y) = p(X=x|Y=1-y), \text{ it}$$

means that conditional of  $Z|Y$  is equal to  $X|1-y$

then  $I(Z; Y) = \frac{1}{2} D_{KL}(q||m) + \frac{1}{2} D_{KL}(p||m)$ . Therefore

$I(Z; Y) = I(X; Y)$  according to the formula above.

Since  $Z$  and  $X$  have same conditional distribution, if we know  $Y$ , it will also reduce uncertainty about  $Z$  in the same as reduce uncertainty in  $X$ .

(d)

Question 5