

Computing Residues:

Working Rules:

- (a) Obtain an entire Laurent expansion of $f(z)$ about $z = z_0$ to identify a_{-1} , the coefficient of $(z - z_0)^{-1}$ in the expansion.
- (b) If $f(z)$ has a simple pole at $z = z_0$, then with a_n the coefficients in the expansion of $f(z)$,

$$(z - z_0)f(z) = a_{-1} + a_0(z - z_0) + a_1(z - z_0)^2 + \dots,$$
 and, recognizing that $(z - z_0)f(z)$ may not have a form permitting an obvious cancellation of the factor $z - z_0$, we take the limit as $z \rightarrow z_0$:

$$a_{-1} = \lim_{z \rightarrow z_0} ((z - z_0)f(z)).$$

- (c) If there is a pole of order $n > 1$ at $z = z_0$, then $(z - z_0)^n f(z)$ must have the expansion

$$(z - z_0)^n f(z) = a_{-n} + \dots + a_{-1}(z - z_0)^{n-1} + a_0(z - z_0)^n + \dots.$$

We see that a_{-1} is the coefficient of $(z - z_0)^{n-1}$ in the Taylor expansion of $(z - z_0)^n f(z)$ and there we can identify it as satisfying

$$a_{-1} = \frac{1}{(n-1)!} \lim_{z \rightarrow z_0} \left[\frac{d^{n-1}}{dz^{n-1}} ((z - z_0)^n f(z)) \right],$$

where a limit is indicated to take account of the fact that the expression involved may be indeterminate.

- (d) Suppose the function $f(z)$ can be represented by

$$f(z) = \frac{g(z)}{h(z)}$$

where $g(z)$ and $h(z)$ are analytic functions. If $g(z_0) \neq 0$ and $h(z_0) = 0$ but $h'(z_0) \neq 0$, the function $1/h(z)$ has a simple pole at $z = z_0$ and $f(z)$ has a simple pole at $z = z_0$. The residue of $f(z)$ at $z = z_0$ is

$$\begin{aligned} a_{-1} &= \lim_{z \rightarrow z_0} \left[\frac{g(z)}{h(z)} (z - z_0) \right] = \lim_{z \rightarrow z_0} g(z) \lim_{z \rightarrow z_0} \left[\frac{z - z_0}{h(z)} \right] = g(z_0) \left[\frac{1}{h'(z_0)} \right] \\ \Rightarrow a_{-1} &= \frac{g(z_0)}{h'(z_0)}. \end{aligned}$$

- (e) Sometimes, the general formula is found to be more complicated than the judicious use of power series expansion. See items 4 and 5 in the example below.
- (f) Essential singularities will also have well-defined residues but finding them may be more difficult. In principle, one can use

$$a_n = \frac{1}{2\pi i} \oint_C \frac{f(z') dz'}{(z' - z_0)^{n+1}}$$

with $n = -1$, but the integral involved may seem intractable. Sometimes the easiest route to the residue is by first finding the Laurent expansion.

Examples: Here are some examples:

1. The residue of $\frac{1}{4z+1}$ at $z = -\frac{1}{4}$ is $\lim_{z \rightarrow -\frac{1}{4}} \left(\frac{z + \frac{1}{4}}{4z+1} \right) = \frac{1}{4}$.

2. Residue of $\frac{1}{\sin z}$ at $z = 0$ is $\lim_{z \rightarrow 0} \left(\frac{z}{\sin z} \right) = 1$.

3. Residue of $\frac{\ln z}{z^2 + 4}$ at $z = 2i$ is

$$\lim_{z \rightarrow 2i} \left(\frac{(z - 2i) \ln z}{z^2 + 4} \right) = \frac{\ln 2 + \pi i/2}{4i} = \frac{\pi}{8} - \frac{i \ln 2}{4}.$$

4. Residue of $\frac{z}{\sin^2 z}$ at $z = \pi$; the pole is second order, and the residue is given by

$$\frac{1}{1!} \lim_{z \rightarrow \pi} \left(\frac{d}{dz} \frac{z(z - \pi)^2}{\sin^2 z} \right).$$

However, it may be easier to make the substitution $w = z - \pi$, note that $\sin^2 z = \sin^2 w$, and to identify the residue as the coefficient of $1/w$ in the expansion of $(w + \pi)/\sin^2 w$ about $w = 0$. This expansion can be written

$$\frac{w + \pi}{\left(w - \frac{w^3}{3!} + \dots \right)^2} = \frac{w + \pi}{w^2 - \frac{w^4}{3} + \dots}.$$

The denominator expands entirely into even powers of w , so the π in the numerator cannot contribute to the residue. Then from the w in the numerator and the leading term of denominator, we find the residue to be 1.

5. Residue of $f(z) = \frac{\cot \pi z}{z(z + 2)}$ at $z = 0$.

The pole at $z = 0$ is second-order, and direct application of

$$a_n = \frac{1}{2\pi i} \oint_C \frac{f(z') dz'}{(z' - z_0)^{n+1}}$$

leads to a complicated indeterminate expression requiring multiple applications of L'Hôpital's rule. Perhaps easier is to introduce the initial terms of the expansion about $z = 0$:

$$\cot \pi z = (\pi z)^{-1} + \mathcal{O}(z), \quad \frac{1}{z + 2} = \frac{1}{2} \left[1 - \frac{z}{2} + \mathcal{O}(z^2) \right], \text{ reaching}$$

$$f(z) = \frac{1}{z} \left[\frac{1}{\pi z} + \mathcal{O}(z) \right] \left(\frac{1}{2} \right) \left[1 - \frac{z}{2} + \mathcal{O}(z^2) \right],$$

from which we can read out the residue as the coefficient of z^{-1} , namely, $-1/4\pi$.

6. Residue of $e^{-1/z}$ at $z = 0$.

This is an essential singularity; from the Taylor series of e^w with $w = -1/z$, we have

$$e^{-1/z} = 1 - \frac{1}{z} + \frac{1}{2!} \left(-\frac{1}{z} \right)^2 + \dots,$$

from which we read out the value of the residue, -1 .

Examples: Calculate the residue(s) of the functions:

(a) $f(z) = \frac{1}{z^2 - 1}$

(b) $f(z) = \frac{1}{(z^2 + a^2)^2}$, where $a > 0$

(c) $f(z) = \frac{\sin z}{z^4}$

(d) $f(z) = \frac{\sin z}{z^6}$

(e) $f(z) = ze^{1/z}$

(f) $f(z) = z^2 \sin \frac{1}{z}$

$$(g) f(z) = \frac{1 + e^z}{\sin z + z \cos z}$$

$$(h) f(z) = \frac{1}{z(e^z - 1)}$$

$$(i) f(z) = \frac{A(z)}{\sin z}, \text{ where } A(z) \text{ is analytic and contains no zeros.}$$