

## QFT ASSIGNMENT II

DUE DATE: 16.09.2023

---

- (1) Consider two complex Klein-Gordon fields of the same mass. Label each of the fields as  $\phi_a(x)$ , where  $a = 1, 2$ .

- (a) Show that you have four conserved charges given by

$$Q^\mu = \int d^3x \frac{i}{2} (\phi_a^\star(A^\mu)_{ab} \pi_b^\star - \pi_a(A^\mu)_{ab} \phi_b)$$

where  $\mu = 0, \dots, 3$  and  $A^0 = 1, A^i = \sigma^i$  with  $\sigma^i$  being the Pauli sigma matrices.

- (b) Show that the three charges  $Q^i$  with  $i = 1, 2, 3$  have the commutation relations of angular momentum (SU(2)).  
(c) Generalise the result for  $n$  identical scalar fields.

- (2) Evaluate the function

$$\langle 0 | \phi(x) \phi(y) | 0 \rangle = D(x - y) = \int \frac{d^3p}{(2\pi)^3} \frac{1}{2E_p} e^{-ip \cdot (x - y)},$$

for spacelike  $(x - y)$  so that  $(x - y)^2 = -r^2$ , explicitly in terms of Bessel functions.

- (3) Consider the spinless Schrodinger field given by the Lagrangian

$$\mathcal{L} = i\psi^\dagger \frac{\partial \psi}{\partial t} - \frac{1}{2m} \nabla \psi^\dagger \nabla \psi - V(\mathbf{r}) \psi^\dagger \psi$$

- (a) Express the free fields  $\psi, \psi^\dagger$  in terms of creation and annihilation operators and find the commutation relations between them.

- (b) Calculate the Green's function

$$G(x_0, \mathbf{x}, y_0, \mathbf{y}) = -i \langle 0 | \psi(x_0, \mathbf{x}) \psi^\dagger(y_0, \mathbf{y}) | 0 \rangle \theta(x_0 - y_0).$$

- (c) Show that this satisfies the equation

$$\left( i \frac{\partial}{\partial t} + \frac{1}{2m} \nabla^2 \right) G(t, \mathbf{x}, 0, 0) = \delta(t) \delta^{(3)}(\mathbf{x}).$$

- (4) Calculate the vacuum expectation value

$$\langle 0 | \{ \phi(x), \phi(y) \} | 0 \rangle$$

for a massless free scalar field, where  $\{, \}$  is the anticommutator. Show that this satisfies the KG equation.

- (5) Calculate

$$\langle 0 | \phi(x_1) \phi(x_2) \phi(x_3) \phi(x_4) | 0 \rangle$$

for a free scalar field by plugging in the expansion of the fields in terms of creation and annihilation operators.