Activity: DAGs to Joint Probabilities

Pooja Panwar[1]
Anant Deshwal[2]


[1]University of Jyväskylä
[2]Bradley University

Credit to:


- Tom Hobbs (Colorado State University)
- Mevin Hooten (University of Texas)
- Becky Tang (Middlebury College)
- Alison Ketz (Funga)
- Slides adopted or based on the 2024 Short course on Bayesian Modeling for Ecologists, funded by NSF (*Award DEB 2042028 to Colorado State University*)

Convert following DAGs to Joint Distributions


1)



2)




3)




Convert following DAGs to Joint Distributions


1)



2)




3)



[A,B,C] = [A|B,C] [B] [C]


Convert following DAGs to Joint Distributions

1)




[A,B,C] = [A|B,C] [B][C]

2)




3)



[A,B,C,D,E] = [A|B,C] [C|D,E] [B] [D] [E]


Convert following DAGs to Joint Distributions

1)




[A,B,C] = [A|B,C] [B][C]

2)



[A,B,C,D,E] = [A|B,C][C|D,E] [B][D][E]

3)



[A,B,C,D,E] = [A|B,C] [C|D,E] [B|D [D] [E]

Convert following Joint Probabilities to DAG

1) $P(A,B) = P(A|B)P(B)$

2) $P(A,B, C) = P(A|B,C) \times P(B|C) \times P(C)$

3) $P(A,B,C,D) = P(A|C) \times P(B|C) \times P(C|D) \times P(D)$

Convert following Joint Probabilities to DAG

1) P(A,B) = P(A|B)P(B)




2) P(A,B, C) = P(A|B,C) x P(B|C) x P(C)

3) P(A,B,C,D) = P(A|C) x P(B|C) x P(C|D) x P(D)

Convert following Joint Probabilities to DAG

2) $P(A,B,C) = P(A|B,C) \times P(B|C) \times P(C)$




1) $P(A,B) = P(A|B)P(B)$

3) $P(A,B,C,D) = P(A|C) \times P(B|C) \times P(C|D) \times P(D)$

Convert following Joint Probabilities to DAG

3) $P(A,B,C,D) = P(A|C) \times P(B|C) \times P(C|D) \times P(D)$



1) $P(A,B) = P(A|B)P(B)$

2) $P(A,B,C) = P(A|B,C) \times P(B|C) \times P(C)$

Ecological Example

$$\frac{1}{N}\frac{dN}{dt} = r - \frac{r}{K}N,$$

- Per-capita rate of population Growth
  - r is intrinsic growth rate
  - K is the Carrying capacity. At N=K dN/dt =0
  - r/K is the strength of feedback from population size to population growth rate

Ecological Example


- Per-capita rate of population Growth
  - $y = 1/N_i \, dN_i/dt$
  - $x_i = N_i$
  - $\mu_i = r - rx_i/K$; where μ_i is the mean for y
  - What distribution would y take?

$$\frac{1}{N}\frac{dN}{dt} = r - \frac{r}{K}N,$$


Ecological Example

- Per-capita rate of population Growth
  - $y = 1/N_i \, dN_i/dt$
  - $x_i = N_i$
  - $\mu_i = r - rx_i/K$; where μ_i is the mean for y
  - What distribution would y take - Normal
  - Draw DAG for it

$$\frac{1}{N}\frac{dN}{dt} = r - \frac{r}{K}N,$$



OR



$$[y,r,K] = [y|r,K]\,[r]\,[K]$$

- $[r, K, \tau|y] \propto \Pi_{i=1}^{n}$ normal $(y_i \mid \mu_i, \tau)$ gamma $(K| 0.001, 0.001)$ gamma $(\tau| 0.001, 0.001)$ gamma $(r|0.001, 0.001)$ … eqn 1
- $\mu_i = f(r, K, x_i) = r - rx_i/K$ …eqn2
- Eqn 1 can also be written as:
- $[r, K, \tau|y] \propto \Pi_{i=1}^{n}$ normal $(y_i \mid f(r, K, x_i), \tau)$ gamma $(K| 0.001, 0.001)$ gamma $(\tau| 0.001, 0.001)$ gamma $(r|0.001, 0.001)$
- Eqn 1 over here is the basis for your JAGs model
  - We are using τ (precision) instead of σ because JAGs uses τ
  - $\tau = 1/\sigma^2$

- $[r, K, \tau|y] \propto \Pi_{i=1}^{n}$ normal $(y_i \mid \mu_i, \tau)$ gamma $(K| 0.001, 0.001)$ gamma $(\tau|$ $0.001, 0.001)$ gamma $(r|0.001, 0.001)$ … eqn 1

- Structure: probability distribution (RV | parameters)

Per-Capita Growth Rate equation is Linear Regression

- $\mu_i = r - rx_i/K$
  - Is a Linear Regression equation $y = mx+b$,
    - $b = r$ and $m = -r/K$

JAGs Model

```
 ## Growth Model
Model {
        #Priors
        K ~ dgamma (0.001, 0.001)
        r ~ dgamma (0.001, 0.001)
        sigma ~ dunif (0, 2)
        tau <- 1/sigma^2 # Calculating precision from SD as JAGs understands Tau

        #Likelihood
        for(i in 1:n){
        mu[i] <- r- r/K* x[i]
        y[i] ~ dnorm(mu[i], tau)

        }
    }
```

$[r, K, \tau|y] \propto \Pi_{i=1}^{n}$ normal $(y_i \mid \mu_i, \tau)$ gamma $(K| 0.001, 0.001)$ gamma $(\tau| 0.001, 0.001)$ gamma $(r|0.001, 0.001)$ … eqn 1

$\mu_i = f(r, K, x_i) = r - rx_i/K$ …eqn2

- Write your JAGs model in a separate R script and save it eg: LogisticsJAGS. R

- You will use “LogisticsJAGS. R” in a separate R code to call to this JAGs model

Structure for JAGs Model

- Code for priors
- Code for deterministic model
- Code for likelihoods
- Priors and liklihoods are specified using ~
- We chose vague gamma priors for r, K, τ becuase they must be non-negative
- Loop replaces $\Pi_{i=1}^{n}$
-