

Introduction to AI and ML

Matrix Project

Ananthapadmanabhan M, EE17BTECH11047
S.Yogesh, EE17BTECH11039

February 13, 2019

Question

The area(in sq.unit) of the quadrilateral formed by the tangents at the ends of the latera recta of the ellipse $\frac{x^2}{9} + \frac{y^2}{5} = 1$, is:

Given ellipse in matrix form:

$$x^T \begin{bmatrix} 1/9 & 0 \\ 0 & 1/5 \end{bmatrix} x = 1$$

Conic equation is matrix form:

$$x^T V x + 2u^T x + F = 0$$

Here,

$$V = \begin{bmatrix} 1/9 & 0 \\ 0 & 1/5 \end{bmatrix}$$

$$u = 0$$

$$F = -1$$

The length of semi major axis $a = 3$,the length of semi minor axis $b = \sqrt{5}$

$$F_1 = \begin{bmatrix} \sqrt{a^2 - b^2} \\ 0 \end{bmatrix}$$

$$F_2 = \begin{bmatrix} -\sqrt{a^2 - b^2} \\ 0 \end{bmatrix}$$

$$F_1 = \begin{bmatrix} 2 \\ 0 \end{bmatrix}$$

$$F_2 = \begin{bmatrix} -2 \\ 0 \end{bmatrix}$$

The end points of the latera recta can be found from

$$\begin{bmatrix} 2 & y \end{bmatrix} \begin{bmatrix} 1/9 & 0 \\ 0 & 1/5 \end{bmatrix} \begin{bmatrix} 2 \\ y \end{bmatrix} = 1$$

$$\begin{bmatrix} -2 & y \end{bmatrix} \begin{bmatrix} 1/9 & 0 \\ 0 & 1/5 \end{bmatrix} \begin{bmatrix} -2 \\ y \end{bmatrix} = 1$$

From the above equations we get,

$$P_1 = \begin{bmatrix} 2 \\ \frac{5}{3} \end{bmatrix}$$

$$P_3 = \begin{bmatrix} -2 \\ -\frac{5}{3} \end{bmatrix}$$

$$P_2 = \begin{bmatrix} 2 \\ -\frac{5}{3} \end{bmatrix}$$

$$P_4 = \begin{bmatrix} -2 \\ \frac{5}{3} \end{bmatrix}$$

Tangent of a conic is given by

$$(p^T V + u^T)x + p^T u + F = 0$$

Tangent at P_1

$$\begin{bmatrix} \frac{2}{9} & \frac{1}{3} \end{bmatrix} x = 1$$

Tangent at P_2

$$\begin{bmatrix} \frac{2}{9} & -\frac{1}{3} \end{bmatrix} x = 1$$

Tangent at P_3

$$\begin{bmatrix} -\frac{2}{9} & -\frac{1}{3} \end{bmatrix} x = 1$$

Tangent at P_4

$$\begin{bmatrix} -\frac{2}{9} & \frac{1}{3} \end{bmatrix} x = 1$$

The point of intersection of two lines is given by:

$$x = N^{-T} p$$

Where $N = (n_1 \ n_2)$ Intersection points of tangents at P_1 and P_2 is

$$A = \begin{bmatrix} \frac{9}{4} & \frac{9}{4} \\ \frac{3}{2} & \frac{3}{2} \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} \frac{9}{2} \\ 0 \end{bmatrix}$$

$$B = \begin{bmatrix} \frac{9}{4} & -\frac{9}{4} \\ -\frac{3}{2} & -\frac{3}{2} \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ -3 \end{bmatrix}$$

$$C = \begin{bmatrix} -\frac{9}{4} & -\frac{9}{4} \\ \frac{3}{2} & \frac{3}{2} \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} -\frac{9}{2} \\ 0 \end{bmatrix}$$

$$D = \begin{bmatrix} -\frac{9}{4} & \frac{9}{4} \\ \frac{3}{2} & \frac{3}{2} \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 3 \end{bmatrix}$$

Area of quadrilateral formed by tangents,

$$\text{Area} = 4 * \text{Area}(\triangle AOB)$$

$$\text{Area} = 4 * \frac{27}{4}$$

$$\text{Area} = 27 \text{sq. units}$$

