

# *Ch 1. The Foundations: Logic and Proofs*

## Logic and Mathematics

Sungwon Kang

### Acknowledgement

- [Rosen 19] Kenneth H. Rosen, for Discrete Mathematics & Its Applications (8th Edition), Lecture slides
- [Hunter 11] David J. Hunter, Essentials of Discrete Mathematics, 2nd Edition, Jones & Bartlett Publishers, 2011, Lecture Slides

Historically,

Logic

is the basis of

Mathematics

We want to make  
logic mathematical.

Historically,


Logic

is the basis of

Mathematics



We want to make  
logic mathematical.




Formal System

Historically,


Logic

is the basis of

Mathematics



We want to make  
logic mathematical.



Formal System

**Gödel proved in 1929**  
**Ph.D. Dissertation**


Successful  
≤ Completeness  
of the Formal System  
for Predicate Logic

Historically,

Logic

is the basis of

Mathematics



We want to make  
logic mathematical.

**Gödel proved in 1929**  
**Ph.D. Dissertation**


Successful

$\leq$  Completeness  
of the Formal System  
for Predicate Logic

**Gödel proved in 1931**

Unsuccessful

$\leq$  Incompleteness  
of Formalizing  
Arithmetic



Formal System

## Incompleteness Theorem [Gödel 1931]

Any number-theoretic formal system (i.e. a consistent formal system to contain arithmetic) is incomplete (i.e. there is a true statement or a theorem of arithmetic that cannot be proved within the system).

**Kurt Gödel**  
(1906 – 1978)




Historically,

Logic

is the basis of

Mathematics

We want to make  
logic mathematical.



Formal System  $\Rightarrow$  Automated Computation

Enabled by Modern Computer

# Logic and Mathematics

## => Linking Logic and Mathematics

- The Role of Definitions in Mathematics
- Other Types of Mathematical Statements
- Counterexamples
- Axiomatic Systems



# Definitions in mathematics

Dictionary definitions are usually *descriptive*.

**Example:** “mortadella” - Any of several types of Italian sausage.

- Definitions in mathematics are *stipulative*.  
(= demand or require)

Example:

## Definition

Two lines are parallel if they have no points in common.

Does this agree with your concept image?

- No. It specifies what exactly we “shall” mean when we use the word “parallel”.

# Using definitions: even and odd

## Definition

An integer  $n$  is *even* if  $n = 2k$  for some integer  $k$ .

## Definition

An integer  $n$  is *odd* if  $n = 2k + 1$  for some integer  $k$ .

To show that “17 is odd” we note that  $17 = 2 \cdot 8 + 1$ .

(Let  $k = 8$ . Then  $17 = 2 \cdot 8 + 1$ . Therefore 17 is odd.)

How can you show that 10 is even?

# Definitions are reversible

A definition of the form

*[Object]  $x$  is [defined term] if [defining property about  $x$ ].*

means

$$(\forall x) (D(x) \leftrightarrow P(x))$$

where

$D(x)$  =  $x$  is [defined term] - D: definiendum

$P(x)$  = [defining property about  $x$ ] - P: definiens

**Example**

$$\forall n (Even(n) \leftrightarrow \exists k (n = 2 \times k))$$

# Mathematical statements

- Definitions
- Axioms (a.k.a. postulates)
- Theorems
- Corollaries
- Lemmas
- Propositions : statements
- Claims
- Conjectures

} **Theorems**

Theory: the set of all statements that can be proved from a set of axioms.

# Axiomatic systems

- Axioms

- Undefined terms



An ***axiomatic system*** consists of axioms and undefined terms.

# Axioms of the Formal Number Theory

$$(S1) \ x = y \rightarrow (x = z \rightarrow y = z)$$

$$(S2) \ x = y \rightarrow x' = y'$$

$$(S3) \ 0 \neq x'$$

$$(S4) \ x' = y' \rightarrow x = y$$

$$(S5) \ x + 0 = x$$

$$(S6) \ x + y' = (x + y)'$$

$$(S7) \ x \cdot 0 = 0$$

$$(S8) \ x \cdot (y') = (x \cdot y) + x$$

$$(S9) \text{ For any well-formed formula } \mathcal{A}(x) \text{ of } S,$$

$$\mathcal{A}(0) \Rightarrow (((\forall x) \mathcal{A}(x) \Rightarrow \mathcal{A}(x')) \Rightarrow (\forall x) \mathcal{A}(x))$$

# Axioms of the Formal Set Theory

**Extensionality Axiom** If two sets have exactly the same members, then they are equal.

$$\forall A \forall B [\forall x (x \in A \Leftrightarrow x \in B) \Rightarrow A = B]$$

**Empty Set Axiom** There is a set having no members.

$$\exists B \forall x \ x \notin B.$$

**Pairing Axiom** For any sets  $u$  and  $v$ , there is a set having as members just  $u$  and  $v$ .

$$\forall u \forall v \exists B \forall x [x \in B \Leftrightarrow x = u \vee x = v]$$

**Union Axiom** For any sets  $a$  and  $b$ , there is a set whose members are those sets belonging either to  $a$  or to  $b$  (or both)

$$\forall a \forall b \exists B \forall x [x \in B \Leftrightarrow x \in a \vee x \in b]$$

**Power Set Axiom** For any set  $a$ , there is a set whose members are exactly the subsets of  $a$

$$\forall a \exists B \forall x [x \in B \Leftrightarrow x \subseteq a]$$

**Subset Axioms** For each formula  $P(x)$  not containing  $B$ ,

$$\forall t_1 \dots \forall t_k \forall c \exists B \forall x [x \in B \Leftrightarrow x \in c \ \& \ P(x) ]$$

# The Axioms of Euclidean Plane Geometry

1. A straight line may be drawn between any two points.
2. Any terminated straight line may be extended indefinitely.
3. A circle may be drawn with any given point as center and any given radius.
4. All right angles are equal.
5. For any given point not on a given line, there is exactly one line through the point that does not meet the given line.



# Axiomatic system for four-point geometry

## Examples.

*Undefined terms:* point, line, is on

*Axioms:*

- 1 For every pair of distinct points  $x$  and  $y$ , there is a unique line  $l$  such that  $x$  is on  $l$  and  $y$  is on  $l$ .
- 2 Given a line  $l$  and a point  $x$  that is not on  $l$ , there is a unique line  $m$  such that  $x$  is on  $m$  and no point on  $l$  is also on  $m$ .
- 3 There are exactly four points.
- 4 It is impossible for three points to be on the same line.

# Simple four-point theorem and proof

## Examples.

### Theorem

*In the axiomatic system for four-point geometry, there are at least two distinct lines.*

### Proof.

# Simple four-point theorem and proof

## Examples.

### Theorem


*In the axiomatic system for four-point geometry, there are at least two distinct lines.*

### Proof.

By Axiom 3, there are distinct points  $x$ ,  $y$ , and  $z$ . By Axiom 1, there is a line  $l_1$  through  $x$  and  $y$ , and a line  $l_2$  through  $y$  and  $z$ . By Axiom 4,  $x$ ,  $y$ , and  $z$  are not on the same line, so  $l_1$  and  $l_2$  must be distinct lines. □

# Model for four-point geometry


Point P1 lies on  
L1 and L2.



*Axioms:*

- 1 For every pair of distinct points  $x$  and  $y$ , there is a unique line  $l$  such that  $x$  is on  $l$  and  $y$  is on  $l$ .
- 2 Given a line  $l$  and a point  $x$  that is not on  $l$ , there is a unique line  $m$  such that  $x$  is on  $m$  and no point on  $l$  is also on  $m$ .
- 3 There are exactly four points.
- 4 It is impossible for three points to be on the same line.

# Model for four-point geometry



A standard interpretation

*Axioms:*

- 1 For every pair of distinct points  $x$  and  $y$ , there is a unique line  $l$  such that  $x$  is on  $l$  and  $y$  is on  $l$ .
- 2 Given a line  $l$  and a point  $x$  that is not on  $l$ , there is a unique line  $m$  such that  $x$  is on  $m$  and no point on  $l$  is also on  $m$ .
- 3 There are exactly four points.
- 4 It is impossible for three points to be on the same line.

# Badda-Bing axiomatic system

*Undefined terms:* badda, bing, hit

*Axioms:*

- 1 Every badda hits exactly four bings.
- 2 Every bing is hit by exactly two baddas.
- 3 If  $x$  and  $y$  are distinct baddas, each hitting bing  $q$ , then there are no other bings hit by both  $x$  and  $y$ .
- 4 There is at least one bing.

# Badda-Bing axiomatic system

*Undefined terms:* badda, bing, hit

*Axioms:*

- 1 Every badda hits exactly four bings.
- 2 Every bing is hit by exactly two baddas.
- 3 If  $x$  and  $y$  are distinct baddas, each hitting bing  $q$ , then there are no other bings hit by both  $x$  and  $y$ .
- 4 There is at least one bing.


**One possible interpretation:**

**badda:** square

**bing:** corner of a square

**hit:** a square hits corner if the corner belongs to the square

# A model for badda-bing system




*Axioms:*

- 1 Every badda hits exactly four bings.
- 2 Every bing is hit by exactly two baddas.
- 3 If  $x$  and  $y$  are distinct baddas, each hitting bing  $q$ , then there are no other bings hit by both  $x$  and  $y$ .
- 4 There is at least one bing.



# Model for badda-bing system using fractals

hit



## Axiomatic system

- All its theorems can be generated from the axioms with logical or mathematical inference rules.
- Can we capture all mathematical theorems in a finite set of axioms? (☛ Hilbert's Program in early 1920s.)
  - No !
  - This idea, although failed for the whole mathematics, provides an ideal way of organizing and developing computer programs

# Quiz 08-1

---

Which of the following is FALSE ?

- [1] A formal system can be inconsistent (or unsound).
- [2] An inconsistent system has no model.
- [3] A consistent system has one or more models.
- [4] A consistent and complete system has a model.
- [5] A consistent and incomplete system has no model.