

Ch 9. Discrete Probability (3)

Sungwon Kang

Acknowledgement

- Michael P. Frank, University of Florida, COT 3100, Applications of Discrete Structures, Lecture Slides
- [Rosen 19] Kenneth H. Rosen, for Discrete Mathematics & Its Applications (8th Edition), Lecture slides
- [Hunter 11] David J. Hunter, Essentials of Discrete Mathematics, 2nd Edition, Jones & Bartlett Publishers, 2011, Lecture Slides
- Milos Hauskrecht, Pittsburgh University, Lecture Slides

Ch 7. Discrete Probability

7.1 An Introduction to Discrete Probability

7.2 Probability Theory

7.3 Bayes' Theorem

7.4 Expected Value and Variance 

Table of Contents

1. Finite probability space, events
2. Basic concepts of probability theory
3. Conditional probability, Bayes' theorem
4. Independence
- 5. Integer random variables, Bernoulli trial**
- 6. Expected value, Linearity of expectation, Variance**

5. Integer random variables, Bernoulli trial

Random Variable

- **Random process**: an experiment in which we know what outcomes could happen (sample space), but we don't know which particular outcome will happen

Definition A **random variable** is a function $f: S \rightarrow R$ from the **sample space of an experiment** to the set of real numbers.

- A **random variable** is a numerical result of a random experiment or process.
=> A random variable assigns a number to each possible outcome.

The **probability distribution** of a random variable X on the sample space S is a set of pairs $(r, P(X=r))$ for all r in S
where r is a number **representing an outcome**
and $P(X=r)$ is the probability that X takes the value r .

Example 4.42 Rolling two standard six-sided dice is a random experiment. The sum of the values on the two dice is a random variable X . Recall the following example.

Example: If you roll two standard six-sided dice, what is the probability you roll 10 or less?

Solution

Compute the probability of rolling more than 10: There are two ways to roll 11 and one way to roll 12, so the probability of rolling more than 10 is $(2 + 1)/36 = 1/12$. Thus the probability of rolling 10 or less is $1 - 1/12 = 11/12$.

$$P(X = 11) = 2/36$$

$$P(X = 12) = 1/36$$

$$P(X \leq 10) = P(2 \leq X \leq 12) - \{P(X = 11) + P(X = 12)\}$$

Random Variable

Example

Let S be the outcomes of a two-dice roll.

Let random variable X denote the sum of outcomes.

$$(1,1) \rightarrow X = 2$$

$$(1,2) \text{ and } (2,1) \rightarrow X = 3$$

$$(1,3), (3,1) \text{ and } (2,2) \rightarrow X = 4$$

...

Probability Distribution of X :

$$2 \rightarrow P(X=2) = 1/36,$$

$$3 \rightarrow P(X=3) = 2/36,$$

$$4 \rightarrow P(X=4) = 3/36,$$

...

$$12 \rightarrow P(X=12) = 1/36$$

Bernoulli trial

- Suppose that an experiment can have only two possible outcomes.

Example. When a coin is flipped, the possible outcomes are heads and tails.

- Each performance of an experiment with two possible outcomes is called a *Bernoulli trial*.
 - Possible outcome of a Bernoulli trial is called a **success** or a **failure**.
 - If p is the probability of a success and q is the probability of a failure,
$$p + q = 1$$
- Many problems can be solved by determining *the probability of k successes* when an experiment consists of n mutually independent Bernoulli trials.

Suppose we flip a coin flip repeatedly.

$P(\text{heads}) = 0.6$ and $P(\text{tails}) = 0.4$.

Each coin flip is independent of the previous flip.

- What is the probability of seeing HHHHH?

$$P(\text{HHHHH}) = 0.6^5$$

- What is the probability of seeing TTHHT?

$$P(\text{TTHHT}) = 0.4^2 \times 0.6^2 \times 0.4 = 0.6^2 \times 0.4^3$$

- What is the probability of seeing two heads and three tails?

The number of two heads and three tails combinations = $C(5,2)$

$$P(\text{two-heads-three-tails}) = C(5,2) \times 0.6^2 \times 0.4^3$$

A variant of a repeated coin flip problem.

Sample space: The number of occurrences of heads in 5 coin flips.

For example,

TTTTT yields outcome 0

HTTTT or TTHTT yields 1

HTHHT yields 3 ...

What is the probability of an outcome i when i ranges from 0 to 5?

$$P(\text{outcome} = 0) = C(5,0) \times 0.6^0 \times 0.4^5$$

$$P(\text{outcome} = 1) = C(5,1) \times 0.6^1 \times 0.4^4$$

$$P(\text{outcome} = 2) = C(5,2) \times 0.6^2 \times 0.4^3$$

$$P(\text{outcome} = 3) = C(5,3) \times 0.6^3 \times 0.4^2$$

$$P(\text{outcome} = 4) = C(5,4) \times 0.6^4 \times 0.4^1$$

$$P(\text{outcome} = 5) = C(5,5) \times 0.6^5 \times 0.4^0$$

What is

$$\sum_{0 \leq i \leq 5} P(\text{outcome} = i) ?$$

$$(0.6 + 0.4)^5 = ?$$

Recall

The Binomial Theorem

Theorem

Let j and k be nonnegative integers such that $j + k = n$. The coefficient of the $a^j b^k$ term in the expansion of $(a + b)^n$ is $C(n, j)$.

Corollary

The Binomial Theorem.

$$\begin{aligned}(a + b)^n = & \binom{n}{0} a^n + \binom{n}{1} a^{n-1} b + \binom{n}{2} a^{n-2} b^2 + \dots \\ & + \binom{n}{j} a^{n-j} b^j + \dots + \binom{n}{n} b^n\end{aligned}$$

Theorem

In n independent Bernoulli trials,
with probability of success p
and probability of failure $q = 1 - p$,

the probability of exactly k successes is

$$C(n, k) \times p^k \times q^{n-k}$$

Exercise. A coin is biased so that the probability of heads is $2/3$.
What is the probability that exactly **four heads** come up
when the coin is **flipped seven times**, assuming that the flips are independent?

Exercise. A coin is biased so that the probability of heads is $2/3$.
What is the probability that exactly **four heads** come up
when the coin is **flipped seven times**, assuming that the flips are independent?

$$C(7,4) (2/3)^4 (1/3)^3 = \frac{35 \cdot 16}{3^7} = \frac{560}{2187}$$

Exercise. A coin is biased so that the probability of heads is $2/3$.
What is the probability that exactly **four heads** come up
when the coin is **flipped seven times**, assuming that the flips are independent?

Solution

When a coin is flipped 7 times, #possible outcomes = $2^7 = 128$

#ways 4 out of 7 flips are heads = $C(7, 4)$.

Because the 7 flips are independent,
the probability of each such outcome (four heads and three tails) = $(2/3)^4(1/3)^3$.

Consequently, the probability that exactly four heads appear is

$$C(7,4) (2/3)^4(1/3)^3 = \frac{35 \cdot 16}{3^7} = \frac{560}{2187}$$

6. Expected Value, Linearity of Expectation, Variance

Expected value

Definition

Let x_1, x_2, \dots, x_n be all of the possible values of a random variable X . Then X 's expected value $E(X)$ is the sum

$$\sum_{i=1}^n x_i \cdot P(X = x_i).$$

That is,

$$E(X) = x_1 \cdot P(X = x_1) + x_2 \cdot P(X = x_2) + \dots + x_n \cdot P(X = x_n).$$

What is $\sum_{i=1}^n P(X = x_i)$?

Example 1. Outcomes of rolling a die: 1, 2, 3, 4, 5, 6

$E(X) = ?$

$$\begin{aligned} E(X) &= 1 \cdot P(X=1) + 2 \cdot P(X=2) + 3 \cdot P(X=3) + 4 \cdot P(X=4) + 5 \cdot P(X=5) + 6 \cdot P(X=6) \\ &= 1 \cdot 1/6 + 2 \cdot 1/6 + 3 \cdot 1/6 + 4 \cdot 1/6 + 5 \cdot 1/6 + 6 \cdot 1/6 \\ &= 7/2 \end{aligned}$$

Example 1. Outcomes of rolling a die: 1, 2, 3, 4, 5, 6

$E(X) = ?$

$$\begin{aligned} E(X) &= 1 \cdot P(X=1) + 2 \cdot P(X=2) + 3 \cdot P(X=3) + 4 \cdot P(X=4) + 5 \cdot P(X=5) + 6 \cdot P(X=6) \\ &= 1 \cdot 1/6 + 2 \cdot 1/6 + 3 \cdot 1/6 + 4 \cdot 1/6 + 5 \cdot 1/6 + 6 \cdot 1/6 \\ &= 7/2 \end{aligned}$$

Example 2.

Flip a fair coin 3 times.

The outcome X of the trial is the number of heads.

What is the expected value of the trial?

Possible results = {HHH, HHT, HTH, THH, HTT, THT, TTH, TTT}

$X = 3, 2, 2, 2, 1, 1, 1, 0$

$E(X) = ?$


$$E(X) = 1/8 \times (3 \times 1 + 2 \times 3 + 1 \times 3 + 0 \times 1) = 12/8 = 3/2$$

Average

- The term ‘average’ expresses that something is statistically the norm.
- Various measures of ‘average’:
 - The *(arithmetic) mean* of a finite set of real numbers is the sum of the real numbers in the set divided by the number of real numbers in the set.
 - Given two positive real numbers x and y ,
 - their *arithmetic mean* is $(x+y)/2$.
 - their *geometric mean* is \sqrt{xy} .
 - The *median* of a finite set of real numbers is the middle element in the list when these integers are listed in the order of increasing size.

Average(= Arithmetic Mean) vs. Expected Value

"As the number of trials increases,
the **average** of the results obtained from them
gets close to the **expected value**." - The Law of Large Numbers



Investment problem

You have 100 dollars and can invest into a stock.
The returns are volatile and
you may get either \$120 with probability of 0.4, or
\$90 with probability of 0.6.

What is the expected value of your investment?

$$\begin{aligned} E(X) &= 0.4 \times 120 + 0.6 \times 90 \\ &= 48 + 54 \\ &= 102 \end{aligned}$$

Is it OK to invest?

Application: playing the lottery (More complicated problem)

In the “Both Ways” version of Australia’s “Cash 3” lottery, you pick a three digit number from 000 to 999, and a randomly chosen winning three-digit number is announced every evening at 5:55pm. If you pick a number with three distinct digits, you win \$580 if your number matches the winning number exactly, and you win \$80 if the digits of your number match the digits of the winning number, but in a different order. What is the expected value of the amount of money you win?

Solution

Let X be the amount of money you win. The possible values of X are 0, 580, and 80. Out of 1000 possible winning numbers, only one matches your number exactly, and five ($3! - 1$) have the same digits, but in a different order. Therefore your expected winnings are


$$\begin{aligned} E(X) &= 0 \cdot P(X = 0) + 580 \cdot P(X = 580) + 80 \cdot P(X = 80) \\ &= 0 \cdot \frac{994}{\underline{1000}} + 580 \cdot \frac{1}{1000} + 80 \cdot \frac{5}{\underline{1000}} \\ &= 0.98 \end{aligned}$$

How much would you pay to play?

Decision making

We will invest \$100 for 6 months.


We need to make a choice whether to invest in Stock 1 or Stock 2, put money into bank or keep them at home.



Deterministic outcome (1/2)

Assume the following simplified problem with the Bank and Home choices only.


The result is guaranteed – the outcome is deterministic.



What is the rational choice assuming our goal is to make money?

Deterministic outcome (2/2)


Assume the simplified problem with the Bank and Home choices only.
The result is guaranteed – the outcome is deterministic




What is the rational choice assuming our goal is to make money?

Answer: Put money into the bank.
The choice is strictly better in terms of the outcome.


Expected value for the outcome of the Stock 1 option (1/2)



Expected value for the outcome of the Stock 1 option (2/2)




Expected value for the outcome of the Stock 2 option



Selection based on expected values

The optimal action is the option that maximizes the expected outcome:



Linearity of Expectation

Theorem If X_i , $i=1,2,3, \dots, n$, with a positive integer n , are random variables on S , and a and b are real numbers, then

$$(1) E(X_1 + X_2 + \dots + X_n) = E(X_1) + E(X_2) + \dots + E(X_n)$$

$$(2) E(aX + b) = aE(X) + b$$

Example

Roll a pair of dice. What is the expected value of the sum of outcomes?

Approach 1:

Outcomes: (1,1),(1,2), ..., (1,6), (2,1),(2,2), ..., (2,6), ..., (6,1),(6,2), ..., (6,6)

X= 2, 3, ..., 7, 3, 4, ..., 8, ..., 7, 8, ..., 12

Expected value: $\sum_{i=1}^n x_i \times P(X = x_i) =$

$$1/36 \times ((2+3+\dots+7)+(3+4+\dots+8)+\dots+(7+8+\dots+12)) = 7$$

Approach 2 (Using the Linearity Theorem):

$$E(x_1+x_2) = E(x_1) + E(x_2)$$

$$E(x_1) = E(x_2) = 7/2$$

$$E(x_1+x_2) = 7$$

Variance

- Variance measures how far a data set is spread out.

- The *variance* of a random variable X is :

$$\mathbf{V}[X] = \sum_{s \in S} (X(s) - \mathbf{E}[X])^2 P(s)$$

What is the following?


$$\sum_{s \in S} (X(s) - \mathbf{E}[X]) \cdot P(s)$$

- The *standard deviation* of X is:

$$\sigma(X) := \mathbf{V}[X]^{1/2}.$$


- A *normal distribution*, sometimes called the *bell curve*, is a distribution that occurs naturally in many situations.

Examples SAT scores, GRE scores, CS204 exam scores.




Normal distribution

Normal distributions with various variances



68-95-99.7 rule



Percentages of the values that lie within a band around the mean in a normal distribution

Six Sigma in the manufacturing world

6 σ : Defect ratio of less than 0.002 out of 100Million

Theorem If X is a random variable on a sample space S , then

$$V(X) = E(X^2) - E(X)^2$$

Proof

$$\begin{aligned} V(X) &= \sum_{s \in S} (X(s) - E(X))^2 p(s) && \text{- by definition of } V(X) \\ &= \sum_{s \in S} X(s)^2 p(s) - 2E(X) \sum_{s \in S} X(s) p(s) + E(X)^2 \sum_{s \in S} p(s) && \text{- by definition of } E(X) \\ &= E(X^2) - 2E(X)E(X) + E(X)^2 \\ &= E(X^2) - E(X)^2. \end{aligned}$$

We have used the fact that $\sum_{s \in S} p(s) = 1$ in the next-to-last step.



Example

Two tetrahedral dice are rolled.

(A tetrahedral die has four faces, which are numbered 1, 2, 3, 4.)

Let $X(i,j) = i + j$, where the first die shows i and the second die shows j .

Find $E(X)$ and $V(X)$.

Solution

The sample space S consists of 16 outcomes: $S = \{(i,j) \mid i,j = 1,2,3,4\}$. We have the following probabilities:

$$p(2) = 1/16, \quad p(3) = 2/16, \quad p(4) = 3/16, \quad p(5) = 4/16, \quad p(6) = 3/16, \quad p(7) = 2/16, \quad p(8) = 1/16.$$

Therefore,

$$E(X) = 2 \cdot \frac{1}{16} + 3 \cdot \frac{2}{16} + 4 \cdot \frac{3}{16} + 5 \cdot \frac{4}{16} + 6 \cdot \frac{3}{16} + 7 \cdot \frac{2}{16} + 8 \cdot \frac{1}{16} = \frac{80}{16} = 5.$$

To find $V(X)$, we use the equality $V(X) = E(X^2) - E(X)^2$:



Example

Two tetrahedral dice are rolled.

(A tetrahedral die has four faces, which are numbered 1, 2, 3, 4.)

Let $X(i,j) = i + j$, where the first die shows i and the second die shows j .

Find $E(X)$ and $V(X)$.

Solution

The sample space S consists of 16 outcomes: $S = \{(i, j) \mid i, j = 1, 2, 3, 4\}$. We have the following probabilities:

$$p(2) = 1/16, \quad p(3) = 2/16, \quad p(4) = 3/16, \quad p(5) = 4/16, \quad p(6) = 3/16, \quad p(7) = 2/16, \quad p(8) = 1/16.$$

Therefore,

$$E(X) = 2 \cdot \frac{1}{16} + 3 \cdot \frac{2}{16} + 4 \cdot \frac{3}{16} + 5 \cdot \frac{4}{16} + 6 \cdot \frac{3}{16} + 7 \cdot \frac{2}{16} + 8 \cdot \frac{1}{16} = \frac{80}{16} = 5.$$

To find $V(X)$, we use the equality $V(X) = E(X^2) - E(X)^2$:

$$\sum_{s \in S} (X - E(X))^2 \times P(s)$$

$$= (2-5)^2 \times \frac{1}{16} + (3-5)^2 \times \frac{2}{16} + (4-5)^2 \times \frac{3}{16} + (5-5)^2 \times \frac{4}{16} + (6-5)^2 \times \frac{3}{16} + (7-5)^2 \times \frac{2}{16} + (8-5)^2 \times \frac{1}{16}$$

$$= \frac{1 \times 9}{16} + \frac{2 \times 4}{16} + \frac{3 \times 1}{16} + \frac{4 \times 0}{16} + \frac{3 \times 1}{16} + \frac{2 \times 4}{16} + \frac{1 \times 9}{16} = \frac{40}{16} = 2.5$$

Example

Two tetrahedral dice are rolled.

(A tetrahedral die is a die with four faces, which are numbered 1, 2, 3, 4.)

Let $X(i,j) = i + j$, where the first die shows i and the second die shows j .

Find $E(X)$ and $V(X)$.

Solution

$$E(X) = 5$$

To find $V(X)$, we use the equality $V(X) = E(X^2) - E(X)^2$:

$$\begin{aligned} V(X) &= E(X^2) - E(X)^2 \\ &= \left(4 \cdot \frac{1}{16} + 9 \cdot \frac{2}{16} + 16 \cdot \frac{3}{16} + 25 \cdot \frac{4}{16} + 36 \cdot \frac{3}{16} + 49 \cdot \frac{2}{16} + 64 \cdot \frac{1}{16} \right) - 5^2 \\ &= \frac{1}{16}(440) - 5^2 = 27.5 - 25 = 2.5. \end{aligned}$$

Quiz 22-1

What is the smallest sigma value we can use when we want a less than 0.3% defect ratio?

- 1) 1 σ
- 2) 2 σ
- 3) 3 σ
- 4) 4 σ
- 5) 6 σ