

## 2020 Fall CS300 Homework # 4

TA: Yoonsung Choi  
E-mail: giantsol2@kaist.ac.kr  
Due: 11/11(Wed), 18:00:00 KST

· Total 100 points

### Problem 1. [ $3 \times 10 = 30$ Points]

We use Huffman's algorithm to obtain an encoding of alphabet  $\{a, b, c, d\}$  with frequencies  $f_a, f_b, f_c, f_d$ . In each of the following cases, either give an example of frequencies  $(f_a, f_b, f_c, f_d)$  that would yield the specified code, or explain why the code cannot possibly be obtained (no matter what the frequencies are).

- (a) Code:  $\{0, 10, 110, 111\}$
- (b) Code:  $\{1, 00, 01, 110\}$
- (c) Code:  $\{00, 01, 10, 11\}$

### Problem 2. [40 Points]

Let  $G$  be a graph with  $n$  vertices. Suppose there are two vertices  $x, y \in V(G)$  such that the every simple path between  $x$  and  $y$  has length strictly greater than  $n/2$  (at least one path exists between  $x$  and  $y$ ). Prove that there exists some vertex  $z (\neq x, y)$  such that deleting  $z$  from  $G$  destroys all  $x$ - $y$  paths (in other words, no path from  $x$  to  $y$ ).

**Problem 3. [ $2 \times 15 = 30$  Points]**

For each of the following statements, decide whether it is true or false. If it is true, give a short explanation. If it is false, give a counterexample.

- (a) Suppose we are given an instance of the Minimum Spanning Tree Problem on a graph  $G$ , with edge costs that are all positive. Let  $T$  be a minimum spanning tree for this instance. Suppose we replace each edge cost  $c_e$  by adding 1,  $c_e + 1$ , thereby creating a new instance of the problem with the same graph but different costs.

True or false? “ $T$  must still be a minimum spanning tree for this new instance.”

- (b) Suppose we are given an instance of the Shortest  $s$ - $t$  Path Problem on a directed graph  $G$ . We assume that all edge costs are positive. Let  $P$  be a minimum-cost  $s$ - $t$  path for this instance. Suppose we replace each edge cost  $c_e$  by adding 1,  $c_e + 1$ , thereby creating a new instance of the problem with the same graph but different costs.

True or false? “ $P$  must still be a minimum-cost  $s$ - $t$  path for this new instance.”