

1 Introduction

Recall, that 'eigenvalues' distribution of Gaussian β -ensemble of size N has density

$$\frac{1}{Z_N^{(\beta)}} \exp \left(-\frac{1}{2} \sum_{i=1}^N x_i^2 \right) |V_N(x_1, \dots, x_N)|^\beta,$$

where V_N is a Vandermonde determinant:

$$V_N(x_1, \dots, x_N) = \prod_{i=1}^N \prod_{j=i+1}^N |x_i - x_j|$$

and $Z_N^{(\beta)}$ is just a normalization coefficient:

$$Z_N^{(\beta)} = \int_{\mathbb{R}^N} \exp \left(-\frac{1}{2} \sum_{i=1}^N x_i^2 \right) |V_N(x_1, \dots, x_N)|^\beta.$$

We denote this density by $g_N^{(\beta)}(x_1, \dots, x_N)$.

The goal of this paper is, in certain sense, describe the asymptotic behaviour of a random vector

$$(\lambda_1^{(\beta)}, \dots, \lambda_N^{(\beta)}) \propto g_N^{(\beta)},$$

when $N \rightarrow +\infty$ while $N \cdot \beta \rightarrow 2c$, where c is some constant.

To be more precise, we introduce the random measure

$$L_N^{(\beta)} = \frac{1}{N} \sum_{i=1}^N \delta_{\lambda_i^{(\beta)}/\sqrt{N}},$$

where δ_a is a Dirac measure.

Then fix any sequence of positive integers $\{N_m\}_{m=1}^\infty$ and positive reals $\{\beta_m\}_{m=1}^\infty$ such that

$$\lim_{m \rightarrow \infty} N_m = +\infty \text{ and } \lim_{m \rightarrow \infty} N_m \beta_m = 2c,$$

for some constant c and set $L_m := L_{N_m}^{(\beta_m)}$. Denote by \bar{L}_m the mean of L_m : $\forall f \in C_b(\mathbb{R}), \langle \bar{L}_m, f \rangle = \mathbb{E}[\langle L_m, f \rangle]$.

We prove that moments of \bar{L}_m converge to the following sequence:

$$\begin{cases} M_{2k+1} = 0 & k \geq 0 \\ M_{2k} = \sum_{r+s=k-1} M_{2r} M_{2s} + c(2k-1)M_{2k-2} & k > 0 \\ M_{2k} = 1 & k = 0. \end{cases}$$

Proposition 1. *For every non-negative integer k :*

$$\lim_{m \rightarrow \infty} \langle \bar{L}_m, x^k \rangle = M_k$$

We also prove, that the variance of moments of L_m tends to 0:

Proposition 2. *For every positive integer k :*

$$\lim_{m \rightarrow \infty} \text{Var} [\langle L_m, x^k \rangle] = 0$$

These two propositions together imply the main result of this paper:

Theorem 1. *There is a unique measure L , whose k -th moments are M_k and L_m converges weakly, in probability to L . Precisely:*

For all $\epsilon > 0$ and $f \in C_b(\mathbb{R})$:

$$\lim_{m \rightarrow \infty} \text{P} [|\langle L_m, f \rangle - \langle L, f \rangle| > \epsilon] = 0$$

Coefficients $h_{\mu, \nu}^\tau$, in turn, give a connection of Jack polynomials to the maps enumeration.

2 Connections to h -coefficients

First we state the main result of [Oko97], which connects β -distribution to Jack polynomials.

Denote by $[|y|^{|\lambda|}] J_\lambda^{(\alpha)}(y)$ the coefficient of the polynomial $(\sum x_i^2)^{|\lambda|/2}$ in the power-sum expansion of $J_\lambda^{(\alpha)}(y)$.

Then the following holds true:

$$\int_{\mathbb{R}^n} J_\lambda^{(\alpha)}(t) g_N^{(2/\alpha)}(t) dt = J_\lambda^{(\alpha)}(1, \dots, 1) \left([|y|^{|\lambda|}] J_\lambda^{(\alpha)}(y) \right). \quad (1)$$

We rewrite it as follows:

$$\int_{\mathbb{R}^n} J_\lambda^{(\alpha)}(t) g_N^{(2/\alpha)}(t) dt = J_\lambda^{(\alpha)}(x) J_\lambda^{(\alpha)}(y) \Big|_{\substack{p_r(x)=N \\ p_r(y)=\delta_{r,2}}}, \quad (2)$$

meaning, that we expand $J_\lambda^{(\alpha)}(x)$ and $J_\lambda^{(\alpha)}(y)$ in the power-sum basis and for every integer $r \geq 0$ we substitute N instead of $p_r(x)$ and $\delta_{r,2}$ instead of $p_r(y)$, where δ is a Kronecker delta.

Lemma 1. *For any partition $\theta \vdash m$:*

$$\int_{\mathbb{R}^N} p_\theta(t) g_N^{(2/\alpha)}(t) dt = z_\theta \alpha^{l(\theta)} [p_\theta(z)] \left(\sum_\lambda \frac{J_\lambda^{(\alpha)}(x) J_\lambda^{(\alpha)}(y) J_\lambda^{(\alpha)}(z)}{\langle J_\lambda^{(\alpha)} J_\lambda^{(\alpha)} \rangle_\alpha} \right) \Big|_{\substack{p_r(x)=N \\ p_r(y)=\delta_{r,2}}} \quad (3)$$

Proof. Note, that:

$$\frac{1}{z_\theta \alpha^{l(\theta)}} \int_{\mathbb{R}^N} p_\theta(t) g_N^{(2/\alpha)}(t) dt = [p_\theta(z)] \int_{\mathbb{R}^N} \left(\sum_{\lambda \vdash m} \frac{p_\lambda(z) p_\lambda(t)}{z_\lambda \alpha^{l(\lambda)}} \right) g_N^{(2/\alpha)}(t) dt. \quad (4)$$

Now we use the Cauchy identity:

$$\prod_{i,j} (1 - x_i y_j)^{-\frac{1}{\alpha}} = \sum_{\lambda} \frac{p_\lambda(x) p_\lambda(y)}{z_\lambda \alpha^{l(\lambda)}} = \sum_{\lambda} \frac{J_\lambda^{(\alpha)}(x) J_\lambda^{(\alpha)}(y)}{\langle J_\lambda^{(\alpha)}, J_\lambda^{(\alpha)} \rangle_\alpha},$$

to switch from power sum symmetric functions to Jack polynomials and apply (2):

$$\begin{aligned} \int_{\mathbb{R}^N} \left(\sum_{\lambda \vdash m} \frac{p_\lambda(t) p_\lambda(z)}{z_\lambda \alpha^{l(\lambda)}} \right) g_N^{(2/\alpha)}(t) dt &= \int_{\mathbb{R}^N} \left(\sum_{\lambda \vdash m} \frac{J_\lambda^{(\alpha)}(t) J_\lambda^{(\alpha)}(z)}{\langle J_\lambda^{(\alpha)}, J_\lambda^{(\alpha)} \rangle_\alpha} \right) g_N^{(2/\alpha)}(t) dt = \\ &= \sum_{\lambda \vdash m} \frac{J_\lambda^{(\alpha)}(z)}{\langle J_\lambda^{(\alpha)}, J_\lambda^{(\alpha)} \rangle_\alpha} \int_{\mathbb{R}^N} J_\lambda^{(\alpha)}(t) g_N^{(2/\alpha)}(t) dt = \\ &= \sum_{\lambda \vdash m} \frac{J_\lambda^{(\alpha)}(x) J_\lambda^{(\alpha)}(y) J_\lambda^{(\alpha)}(z)}{\langle J_\lambda^{(\alpha)}, J_\lambda^{(\alpha)} \rangle_\alpha} \Big|_{\substack{p_r(x)=N \\ p_r(y)=\delta_{r,2}}} \end{aligned} \quad (5)$$

Combining (4) and (5) gives the desired result. \square

Denote

$$\phi(x, y, z; t, \alpha) = \sum_{n \geq 0} t^n \sum_{\lambda \vdash n} \frac{J_\lambda^{(\alpha)}(x) J_\lambda^{(\alpha)}(y) J_\lambda^{(\alpha)}(z)}{\langle J_\lambda^{(\alpha)}, J_\lambda^{(\alpha)} \rangle_\alpha}$$

and

$$\psi(x, y, z; t, \alpha) = \alpha t \frac{\partial}{\partial t} \log(\phi) = \sum_{n \geq 1} t^n \left(\sum_{\mu, \nu, \tau \vdash n} h_{\mu, \nu}^\tau (\alpha - 1) p_\mu(x) p_\nu(y) p_\tau(z) \right). \quad (6)$$

Now we are ready to prove the crucial lemma:

Lemma 2. *For any partition $\theta \vdash m$:*

$$\begin{aligned} \int_{\mathbb{R}^N} p_\theta(t) g_N^{(2/\alpha)}(t) dt &= \\ z_\theta \alpha^{l(\theta)} [t^m p_\theta(z)] \exp \left(\frac{1}{\alpha} \sum_{n \geq 1} \frac{t^{2n}}{2n} \sum_{\mu, \tau \vdash 2n} h_{\mu, (2)^n}^\tau (\alpha - 1) N^{l(\mu)} p_\tau(z) \right). \end{aligned} \quad (7)$$

In particular

$$\int_{\mathbb{R}^N} p_\theta(t) g_N^{(2/\alpha)}(t) dt = 0,$$

unless m is even.

Proof. First, we note that:

$$\begin{aligned} [p_\theta(z)] \left(\sum_{\lambda} \frac{J_{\lambda}^{(\alpha)}(x) J_{\lambda}^{(\alpha)}(y) J_{\lambda}^{(\alpha)}(z)}{\langle J_{\lambda}^{(\alpha)} J_{\lambda}^{(\alpha)} \rangle_{\alpha}} \right) \Big|_{\substack{p_r(x)=N \\ p_r(y)=\delta_{r,2}}} &= \\ &= [t^m p_\theta(z)] \phi(x, y, z; t, \alpha) \Big|_{\substack{p_r(x)=N \\ p_r(y)=\delta_{r,2}}} \end{aligned} \quad (8)$$

From (6) we deduce:

$$\begin{aligned} \phi(x, y, z; \alpha) &= \exp \left(\int \frac{\psi(x, y, z; \alpha)}{\alpha t} dt \right) = \\ &= \exp \left(\frac{1}{\alpha} \sum_{n \geq 1} \frac{t^n}{n} \left(\sum_{\mu, \nu, \tau \vdash n} h_{\mu, \nu}^{\tau} (\alpha - 1) p_{\mu}(x) p_{\nu}(y) p_{\tau}(z) \right) \right) \end{aligned} \quad (9)$$

Note, that:

$$p_{\mu}(x) \Big|_{p_r(x)=N} = N^{l(\mu)} \text{ and } p_{\nu}(y) \Big|_{p_r(y)=\delta_{r,2}} = \begin{cases} 1 & \nu = (2^{l(\nu)/2}) \\ 1 & \text{otherwise} \end{cases}.$$

In particular, $p_{\nu}(y) \Big|_{p_r(y)=\delta_{r,2}} = 0$ unless the size of ν is even. Now we are ready to do the corresponding substitutions in (9) to obtain:

$$\phi(x, y, z; t, \alpha) \Big|_{\substack{p_r(x)=N \\ p_r(y)=\delta_{r,2}}} = \exp \left(\frac{1}{\alpha} \sum_{n \geq 1} \frac{t^{2n}}{2n} \sum_{\mu, \tau \vdash 2n} h_{\mu, (2)^n}^{\tau} (\alpha - 1) N^{l(\mu)} p_{\tau}(z) \right). \quad (10)$$

The rest is just combining (8), (10) and **Lemma 1**. \square

Proposition 3. For all positive integers k the following holds:

$$\begin{cases} \langle \bar{L}_N^{(2/\alpha)}, x^{2k+1} \rangle = 0 \\ \langle \bar{L}_N^{(2/\alpha)}, x^{2k} \rangle = \frac{1}{N^{k+1}} \sum_{\mu \vdash 2k} h_{\mu, (2)^k}^{(2k)} (\alpha - 1) N^{l(\mu)} \end{cases} \quad (11)$$

Proof. Note, that:

$$\langle \bar{L}_N^{(2/\alpha)}, x^r \rangle = \mathbb{E} \left[\frac{1}{N} \sum_{i=1}^N \left(\frac{\lambda_i}{\sqrt{N}} \right)^r \right] = \frac{1}{N^{1+r/2}} \int_{\mathbb{R}^N} p_{(r)}(t) g_N^{(2/\alpha)}(t) dt.$$

It follows from **Lemma 2**, that:

$$\langle \bar{L}_N^{(2/\alpha)}, x^{2k+1} \rangle = 0$$

Now for brevity, denote:

$$A(t) = \frac{1}{\alpha} \sum_{n \geq 1} \frac{t^{2n}}{2n} \sum_{\mu, \tau \vdash 2n} h_{\mu, (2)^n}^\tau (\alpha - 1) N^{l(\mu)} p_\tau(z).$$

Lemma 2 says, that:

$$\int_{\mathbb{R}^N} p_{(r)}(t) g_N^{(2/\alpha)} dt = 2k\alpha [t^{2k} p_{(2k)}(z)] \left(1 + \frac{A(t)}{1!} + \frac{A(t)^2}{2!} + \dots \right)$$

Note, that the term $p_{(2k)}(t)$ can not appear in the part:

$$\frac{A(t)^2}{2!} + \frac{A(t)^3}{3!} + \dots,$$

since all power-sum polynomials appearing there are indexed by partitions having length at least 2.

So we have:

$$\int_{\mathbb{R}^N} p_{(r)}(t) g_N^{(2/\alpha)} dt = 2k\alpha [t^{2k} p_{(2k)}(z)] A(t) = \sum_{\mu \vdash 2k} h_{\mu, (2^k)}^{(2k)} (\alpha - 1) N^{l(\mu)}. \quad (12)$$

□

Proposition 4. *For all positive integers k the following holds:*

$$\begin{cases} \text{Var}[\langle L_N^{(2/\alpha)}, x^{2k+1} \rangle] = 0 \\ \text{Var}[\langle L_N^{(2/\alpha)}, x^{2k} \rangle] = \frac{2k\alpha}{N^{2+2k}} \sum_{\mu \vdash 4k} h_{\mu, (2^{2k})}^{(2k, 2k)} (\alpha - 1) N^{l(\mu)}. \end{cases} \quad (13)$$

Proof. Similarly to the previous proposition, we note

$$\begin{aligned} \text{Var}[\langle L_N^{(2/\alpha)}, x^r \rangle] &= \mathbb{E}[L_N^{(2/\alpha)}, x^r] = \mathbb{E}[(\langle L_N^{(2/\alpha)}, x^r \rangle)^2] - \left(\mathbb{E}[\langle L_N^{(2/\alpha)}, x^r \rangle] \right)^2 = \\ &= \frac{1}{N^{2+2k}} \left(\int_{\mathbb{R}^N} p_{(r,r)} g_N^{(2/\alpha)}(t) dt - \left(\int_{\mathbb{R}^N} p_{(r)} g_N^{(2/\alpha)}(t) dt \right)^2 \right). \end{aligned} \quad (14)$$

It follows from **Lemma 2**, that:

$$\text{Var}[\langle L_N^{(2/\alpha)}, x^{2k+1} \rangle] = 0.$$

Similarly to the proof of the **Prop 3** we denote:

$$A(t) = \frac{1}{\alpha} \sum_{n \geq 1} \frac{t^{2n}}{2n} \sum_{\mu, \tau \vdash 2n} h_{\mu, (2)^n}^\tau (\alpha - 1) N^{l(\mu)} p_\tau(z),$$

and, using **Lemma 2**, note, that

$$\begin{aligned}
\int_{\mathbb{R}^N} p_{(2k,2k)} g_N^{(2/\alpha)}(t) dt &= z_{(2k,2k)} \alpha^2 [t^{4k} p_{(2k,2k)}(z)] \left(\frac{A(t)}{1!} + \frac{A(t)}{2!} \right) = \\
&= 8k^2 \alpha^2 \left(\frac{1}{4k\alpha} \sum_{\mu \vdash 4k} h_{\mu, (2^{2k})}^{(2k,2k)} (\alpha - 1) N^{l(\mu)} + \frac{1}{8k^2 \alpha^2} \left(\sum_{\mu \vdash 2k} h_{\mu, (2^k)}^{(2k)} (\alpha - 1) N^{l(\mu)} \right)^2 \right) = \\
&= 2k\alpha \sum_{\mu \vdash 4k} h_{\mu, (2^{2k})}^{(2k,2k)} (\alpha - 1) N^{l(\mu)} + \left(\int_{\mathbb{R}^N} p_{(r)} g_N^{(2/\alpha)}(t) dt \right)^2. \quad (15)
\end{aligned}$$

The last equality follows from (12). The rest is just combining (14) and (15). \square

3 Proofs of the main results

We borrow the notions of *bridge*, *border*, *twisted edge*, *handle* from [Dol16] and denote a number of corresponding type root edges appearing during the root deletion process of the map by $br(M)$, $bo(M)$, $t(M)$, $h(M)$. Also we'll use $\eta(M)$ – one of the measures of non-orientability, also introduced in [Dol16]. By $F(M)$ we denote the face degree partition of M .

Lemma 3. *The following holds true:*

- $v(M) = br(M) + 1$
- $e(M) = bo(M) + t(M) + h(M) + br(M)$
- $f(M) = bo(M) - h(M) + 1$

Proof. During the root deletion process the number of connected components increases only if the border deleted. Since the initial number of connected components of M is 1 and the final number is $v(M)$. We get $v(M) = br(M) + 1$.

The second identity is obvious.

Suppose at some point of the root deletion process there are k connected maps M_1, \dots, M_k . We are interested in the characteristic $\left(\sum_{i=1}^k f(M_i) \right) - k$. We note, that the deletion of borders and twisted edges doesn't change the characteristic, the deletion of handles – increases by 1, the deletion of borders – decreases by 1. At the beginning it's value is $f(M) - 1$ and in the end – 0, so we get $f(M) - 1 = bo(M) - h(M)$. \square

Lemma 4.

$$\eta(M) \leq 2g(M),$$

and the equality holds if and only if M is unhandled ($h(M) = 0$).

Proof. We rewrite Euler's identity, using **Lemma 3**:

$$\begin{aligned} 2g(M) &= 2 - v(M) + e(M) - f(M) = \\ &= 2 - (br(M) + 1) + (bo(M) + t(M) + h(M) + br(M)) - (bo(M) - h(M) + 1) = \\ &= t(M) + 2h(M) \end{aligned}$$

By the definition of η :

$$t(M) \leq \eta(M) \leq t(M) + h(M) \leq t(M) + 2h(M) = 2g(M),$$

which proves the statement. \square

*Proof of the **Proposition 2**.* We begin with the reformulation of **Proposition 4** in terms of maps. The key ingredient for that is [La 09] Corollary 4.17:

$$\sum_{l(\mu)=v} h_{\mu, (2^n)}^\tau(b) = \sum_{F(M)=\tau, v(M)=v} b^{\eta(M)},$$

It's enough to prove the proposition for even moments. We see, that:

$$\begin{aligned} \sum_{\mu \vdash 4k} h_{\mu, (2^{2k})}^{(2k, 2k)}(\alpha - 1) N^{l(\mu)} &= \sum_{v=1}^{4k} N^v \sum_{\mu \vdash 4k, l(\mu)=v} h_{\mu, (2^{2k})}^{(2k, 2k)}(\alpha - 1) = \\ &= \sum_{v=1}^{4k} N^v \sum_{\substack{F(M)=(2k, 2k), \\ v(M)=v}} (\alpha - 1)^{\eta(M)} = \sum_{F(M)=(2k, 2k)} (\alpha - 1)^{\eta(M)} N^{v(M)}. \quad (16) \end{aligned}$$

Now note, that if $F(M) = (2k, 2k)$, then $e(M) = 2k$ and $f(M) = 2$ and hence $v(M) - 2k = -2g(M)$. We get:

$$\begin{aligned} \text{Var}[\langle L_N^{(2/\alpha)}, x^{2k} \rangle] &= \frac{2k\alpha}{N^{2+2k}} \sum_{F(M)=(2k, 2k)} (\alpha - 1)^{\eta(M)} N^{v(M)} = \\ &= \frac{2k\alpha}{N^2} \sum_{F(M)=(2k, 2k)} (\alpha - 1)^{\eta(M)} N^{-2g(M)}. \quad (17) \end{aligned}$$

Note that $L_m = L_{N_m}^{2/(2/\beta_m)}$, hence:

$$\text{Var}[\langle L_m, x^{2k} \rangle] = \frac{4k}{\beta_m N_m^2} \sum_{F(M)=(2k, 2k)} \left(\frac{2}{\beta_m} - 1 \right)^{\eta(M)} N_m^{-2g(M)}.$$

Since $\eta(M) \leq 2g(M)$:

$$\lim_{m \rightarrow \infty} \left(\frac{2}{\beta_m} - 1 \right)^{\eta(M)} N_m^{-2g(M)} \leq (2c)^{\eta(M)},$$

and:

$$\lim_{m \rightarrow \infty} \frac{4k}{\beta_m N_m^2} = 0,$$

we get:

$$\lim_{m \rightarrow \infty} \text{Var}[\langle L_m, x^{2k} \rangle] = 0.$$

□

Lemma 5. *Routed unhandled, unicellular maps with e edges and v vertices are in bijection with rooted oriented maps with the same number of edges and vertices.*

Proof. The root deletion process defines an ordering of edges. Reversing this ordering and adding ribbons one by one, starting from v disconnected vertices, defines the *root adding* process. During the process, added ribbons connect the same corners they connected initially, but we're allowed to twist ribbons, as shown on the picture:

INSERT PIC

We describe algorithm of choosing the twists, which will constitute the desired bijection.

We start by exploring how twist affects the type of the root in different settings.

Case 1 *The root connects corners from two disconnected components.*

The root twist results in the isomorphic map. In any case the added root is a **bridge**. Also note (we'll need this fact later), that connecting two unicellular components gives a unicellular component too.

INSERT PIC.

Case 2 *The root connects corners from the same face.* First, we depict the traversed face before adding the root:

INSERT PIC

Then depending on the twist, we either add one more face (in that case the added root is a **border**) or we keep the number of faces unchanged (the added root is a **twist**):

INSERT PIC

Note, that the border doesn't change the orientability of the map, while adding a twist always gives a non-orientable map.

Case 3 *The root connects corners from different faces but from the same connected component.*

We diagrammatically depict the traversed faces, containing the added root:

INSERT PIC

In both cases we decrease the number of faces by one and the added root is a **handle**. But the twist changes the traversing direction of one of the faces. So if the map without the added root is oriented, only one of the resulting maps is oriented.

First, we show how to make an oriented rooted map from unicellular unhandled one. We start the root adding process from v trivial components and by induction we assume, that on each step all the connected components are orientable.

- If we encounter **Case 1** during the root addition process, we choose any of the twists.
- If we encounter **Case 2**, we twist the ribbon in a way to get a border.
- If we encounter **Case 3**, we twist the ribbon in unique available way to keep the corresponding component orientable.

Now we give an algorithm of getting an unicellular unhandled map from an oriented rooted map. Again, we do the root addition process. Our strategy is to avoid the **Case 3**, since it always results in adding a handle. Inductively we maintain the following property: every connected component is unicellular.

- If we encounter **Case 1**, we choose any of the twists. Note, that connecting to unicellular components results in an unicellular component too.
- If we encounter **Case 2**, we twist the ribbon in a way that gives a twisted root, thus keeping the number of faces of the component equal to 1.
- We can not encounter **Case 3**, since it implies, that there is a component with at least 2 faces, contradicting our induction hypothesis.

Careful analysis of both algorithms shows, that these transformations are inverse to each other. \square

Lemma 6.

$$\lim_{m \rightarrow \infty} \langle \bar{L}_m, x^{2k} \rangle = c^{k+1} \sum_{\substack{M - \text{rooted,} \\ \text{oriented, } e(M)=k}} c^{-v(M)} \quad (18)$$

Proof. Similarly to the proof of **Proposition 2**, we use [La 09] **Corollary 4.17** to reformulate **Proposition 3** in terms of maps:

$$\begin{aligned} \langle \bar{L}_N^{(2/\alpha)}, x^{2k} \rangle &= \frac{1}{N^{k+1}} \sum_{\mu \vdash 2k} h_{\mu, (2k)}^{(2k)} (\alpha - 1) N^{l(\mu)} = \\ &= \frac{1}{N^{k+1}} \sum_{F(M)=(2k)} (\alpha - 1)^{\eta(M)} N^{v(M)} = \sum_{F(M)=(2k)} (\alpha - 1)^{\eta(M)} N^{-2g(M)} \quad (19) \end{aligned}$$

And hence:

$$\langle \bar{L}_m, x^{2k} \rangle = \sum_{F(M)=(2k)} \left(\frac{2}{\beta_m} - 1 \right)^{\eta(M)} N_m^{-2g(M)}.$$

From **Lemma 4** it follows, that:

$$\begin{cases} \lim_{m \rightarrow \infty} \left(\frac{2}{\beta_m} - 1 \right)^{\eta(M)} N_m^{-2g(M)} = c^{-2g(M)} & \text{if } M \text{ is unhandled} \\ \lim_{m \rightarrow \infty} \left(\frac{2}{\beta_m} - 1 \right)^{\eta(M)} N_m^{-2g(M)} = 0 & \text{otherwise} \end{cases}$$

Using **Lemma 5**:

$$\begin{aligned} \lim_{m \rightarrow \infty} \langle \bar{L}_m, x^{2k} \rangle &= \sum_{\substack{M - \text{unhandled,} \\ \text{unicellular, } e(M)=k}} c^{-2g(M)} = \\ &= \sum_{\substack{M - \text{rooted,} \\ \text{unicellular,} \\ \text{unhandled, } e(M)=k}} c^{-k+v(M)-1} = c^{-k-1} \sum_{\substack{M - \text{rooted,} \\ \text{oriented, } e(M)=k}} c^{v(M)}. \end{aligned}$$

□

*Proof of the **Proposition 1**.* Let $M(k)$ denote the set of all rooted, oriented, connected maps with e edges. By deleting the root edge, we see, for $k \geq 1$:

$$M(k) \equiv \bigsqcup_{r+s=k-1} M(r) \times M(s) \sqcup (2k-1)M(k-1). \quad (20)$$

Now denote:

$$a_k = c^{k+1} \sum_{\substack{M - \text{rooted,} \\ \text{oriented, } e(M)=k}} c^{-v(M)}.$$

and combining it with (20), we get the recurrence relation for $k > 1$:

$$\begin{cases} a_k = \sum_{r+s=k-1} a_r a_s + c(2k-1)a_{k-1} & k \geq 1 \\ a_k = 1 & k = 0 \end{cases}, \quad (21)$$

which coincides with the recurrence for M_{2k} . □

References

- [Dol16] Maciej Dolega. Top degree part in b -conjecture for unicellular bipartite maps. *arXiv preprint arXiv:1604.03288*, 2016.
- [La 09] La Croix, Michael Andrew. *The combinatorics of the Jack parameter and the genus series for topological maps*. PhD thesis, 2009.
- [Oko97] Andrei Okounkov. Proof of a conjecture of goulden and jackson. *Canadian Journal of Mathematics*, 49(5):883–886, 1997.