

## Appendix

### Supplemental Materials for Chapter 1: Assessing the Resistance and Resilience of Recreationally Important Fish Species to Extreme Events in Coastal Texas

#### Time series analysis plots




Figure S1: Red Drum monthly averaged CPUE over the last ten years parsed out by bay system. Dotted lines signifying two extreme events, Hurricane Harvey and the 2021 Texas Freeze, left to right.

### Spotted Seatrout CPUE Monthly Average since 2013




Figure S2: Spotted Seatrout monthly averaged CPUE over the last ten years parsed out by bay system. Dotted lines signifying two extreme events, Hurricane Harvey and the 2021 Texas Freeze, left to right.




Figure S3: Black Drum monthly averaged CPUE over the last ten years parsed out by bay system. Dotted lines signifying two extreme events, Hurricane Harvey and the 2021 Texas Freeze, left to right.




Figure S4: Southern Flounder montly averaged CPUE over the last ten years parsed out by bay system. Dotted lines signifying two extreme events, Hurricane Harvey and the 2021 Texas Freeze, left to right.

## Recruitment Windows




Figure S5: Red Drum CPUE monthly average raw data points plots over julian day establishing the recruitment window for Red Drum.




Figure S6: Spotted Seatrout CPUE monthly average raw data points plots over julian day establishing the recruitment window for Spotted Seatrout.




Figure S7: Black Drum CPUE monthly average raw data points plots over julian day establishing the recruitment window for Black Drum.




Figure S8: Southern Flounder CPUE monthly average raw data points plots over julian day establishing the recruitment window for Southern Flounder.



# Hurricane Harvey Percent Change Tables

Table S1: Red Drum percent changes after Hurricane Harvey

MAJOR_AREA_CODE	SPECIES	10 yrs	1st year	2nd year	first_perChange	second_perChange
SabineLake	Red Drum	18.46	25.79	7.595	39.7	-58.9
GalvestonBay	Red Drum	27.26	22.58	24.109	-17.2	-11.5
MatagordaBay	Red Drum	12.57	8.75	12.998	-30.4	3.44
SanAntonioBay	Red Drum	35.42	93.75	93.333	165	163
AransasBay	Red Drum	34.80	53.96	43.542	55	25.1
CorpusChristiBay	Red Drum	28.69	43.31	37.317	51	30.1
UpperLagunaMadre	Red Drum	21.65	12.58	6.458	-41.9	-70.2
LowerLagunaMadre	Red Drum	35.50	37.08	50.208	4.45	41.4

Table S2: Spotted Seatrout percent changes after Hurricane Harvey

MAJOR_AREA_CODE	SPECIES	10 yrs	1st year	2nd year	first_perChange	second_perChange
SabineLake	Spotted Seatrout	5.926	4.048	1.667	-31.7	-71.9
GalvestonBay	Spotted Seatrout	22.470	13.333	35.523	-40.7	58.1
MatagordaBay	Spotted Seatrout	10.218	9.592	8.571	-6.12	-16.1
SanAntonioBay	Spotted Seatrout	29.361	44.197	48.333	50.5	64.6
AransasBay	Spotted Seatrout	60.999	40.476	25.899	-33.6	-57.5
CorpusChristiBay	Spotted Seatrout	21.365	13.909	24.088	-34.9	12.7
UpperLagunaMadre	Spotted Seatrout	26.590	20.144	18.095	-24.2	-31.9
LowerLagunaMadre	Spotted Seatrout	11.350	4.048	12.143	-64.3	6.99

Table S3: Black Drum percent changes after Hurricane Harvey

MAJOR_AREA_CODE	SPECIES	10 yrs	1st year	2nd year	first_perChange	second_perChange
SabineLake	Black Drum	4.402	1.1905	2.3810	-73	-45.9
GalvestonBay	Black Drum	8.214	1.4388	3.3573	-82.5	-59.1
MatagordaBay	Black Drum	4.608	5.2381	3.5714	13.7	-22.5
SanAntonioBay	Black Drum	9.971	43.4568	23.5714	336	136
AransasBay	Black Drum	5.906	1.9048	1.4286	-67.8	-75.8
CorpusChristiBay	Black Drum	2.269	0.2398	0.0000	-89.4	-100
UpperLagunaMadre	Black Drum	96.679	76.9784	41.1905	-20.4	-57.4
LowerLagunaMadre	Black Drum	1.073	0.7143	0.4762	-33.4	-55.6

Table S4: Southern Flounder percent changes after Hurricane Harvey

MAJOR_AREA_CODE	SPECIES	10 yrs	1st year	2nd year	first_perChange	second_perChange
SabineLake	Southern Flounder	7.673	5.0420	5.6497	-34.3	-26.4
GalvestonBay	Southern Flounder	8.842	7.5362	5.0847	-14.8	-42.5
MatagordaBay	Southern Flounder	4.762	3.0556	1.3889	-35.8	-70.8
SanAntonioBay	Southern Flounder	3.818	0.8696	2.7778	-77.2	-27.3
AransasBay	Southern Flounder	3.122	1.9444	0.0000	-37.7	-100
CorpusChristiBay	Southern Flounder	2.846	0.5556	4.3103	-80.5	51.5
UpperLagunaMadre	Southern Flounder	3.005	3.3898	2.7778	12.8	-7.56
LowerLagunaMadre	Southern Flounder	2.613	3.0556	0.5556	16.9	-78.7

# Texas 2021 Freeze Percent Change Tables

Table S5: Red Drum percent changes after freeze event

MAJOR_AREA_CODE	SPECIES	10 yrs	1st year	2nd year	first_perChange	second_perChange
SabineLake	Red Drum	16.40	10.417	5.208	-36.46	-68.23
GalvestonBay	Red Drum	26.37	28.721	11.458	8.911	-56.55
MatagordaBay	Red Drum	11.68	6.289	5.063	-46.13	-56.63
SanAntonioBay	Red Drum	49.00	49.476	21.250	0.9738	-56.63
AransasBay	Red Drum	39.48	54.060	22.917	36.92	-41.96
CorpusChristiBay	Red Drum	29.86	33.333	25.417	11.64	-14.88
UpperLagunaMadre	Red Drum	17.07	18.868	16.667	10.51	-2.384
LowerLagunaMadre	Red Drum	34.67	20.833	18.125	-39.91	-47.72

Table S6: Spotted Seatrout percent changes after freeze event

MAJOR_AREA_CODE	SPECIES	10 yrs	1st year	2nd year	first_perChange	second_perChange
SabineLake	Spotted Seatrout	5.675	4.524	5.952	-20.28	4.892
GalvestonBay	Spotted Seatrout	23.084	21.905	30.476	-5.11	32.02
MatagordaBay	Spotted Seatrout	10.279	11.511	36.869	11.99	258.7
SanAntonioBay	Spotted Seatrout	37.269	25.476	24.762	-31.64	-33.56
AransasBay	Spotted Seatrout	56.129	27.857	33.809	-50.37	-39.76
CorpusChristiBay	Spotted Seatrout	19.396	27.857	17.986	43.63	-7.27
UpperLagunaMadre	Spotted Seatrout	23.861	10.476	22.619	-56.09	-5.204
LowerLagunaMadre	Spotted Seatrout	10.595	5.238	10.476	-50.56	-1.124

Table S7: Black Drum percent changes after freeze event

MAJOR_AREA_CODE	SPECIES	10 yrs	1st year	2nd year	first_perChange	second_perChange
SabineLake	Black Drum	2.718	1.429	6.429	-47.44	136.5
GalvestonBay	Black Drum	5.631	16.905	5.952	200.2	5.699
MatagordaBay	Black Drum	5.580	31.667	38.095	467.6	582.8
SanAntonioBay	Black Drum	17.831	76.191	68.571	327.3	284.6
AransasBay	Black Drum	5.499	58.571	7.381	965.1	34.22
CorpusChristiBay	Black Drum	1.886	9.286	3.357	392.4	78.02
UpperLagunaMadre	Black Drum	88.841	3.333	5.714	-96.25	-93.57
LowerLagunaMadre	Black Drum	1.524	22.381	5.000	1369	228.1

Table S8: Southern Flounder percent changes after freeze event

MAJOR_AREA_CODE	SPECIES	10 yrs	1st year	2nd year	first_perChange	second_perChange
SabineLake	Southern Flounder	7.386	4.2424	0.3236	-42.56	-95.62
GalvestonBay	Southern Flounder	7.911	10.5769	5.3872	33.7	-31.9
MatagordaBay	Southern Flounder	2.339	7.1197	7.9208	204.4	238.6
SanAntonioBay	Southern Flounder	3.440	6.0897	6.6007	77.05	91.9
AransasBay	Southern Flounder	2.673	0.3333	0.9709	-87.53	-63.68
CorpusChristiBay	Southern Flounder	2.597	0.3175	0.0000	-87.77	-100
UpperLagunaMadre	Southern Flounder	3.175	1.6181	0.0000	-49.03	-100
LowerLagunaMadre	Southern Flounder	2.446	6.4815	2.7273	164.9	11.48