

| Investigato | r Sign: |  |
|-------------|---------|--|
|-------------|---------|--|

# Montgomery Asberg Depression Rating Scale (MADRS)

| Investigator: | Dr.Mohsin Qayyum                |
|---------------|---------------------------------|
| eCRF ID:      | A103-013                        |
| Filled By:    | Investigator                    |
| Scale:        | Double-blind Treatment (Day 28) |
| Date:         | 7/30/2025                       |

### 1. Apparent Sadness

Representing despondency, gloom and despair, (more than just ordinary transient low spirits) reflected in speech, facial expression, and posture. Rate by depth and inability to brighten up.

Answer: 4 Appears sad and unhappy most of the time.

## 2. Reported sadness

Representing reports of depressed mood, regardless of whether it is reflected in appearance or not. Includes low spirits, despondency or the feeling of being beyond help and without hope. Rate according to intensity, duration and the extent to which the mood is reported to be influenced by events.

Answer: 0 Occasional sadness in keeping with the circumstances.

#### 3. Inner tension

Representing feeling of ill-defined discomfort, edginess, inner turmoil, mental tension mounting to either panic, dread or anguish. Rate according to intensity, frequency, duration and the extent of reassurance called for.

Answer: 1

### 4. Reduced sleep

Representing the experience of reduced duration or depth of sleep compared with the subject's own normal pattern when well.

Answer: 2 Slight difficulty dropping off to sleep or slightly reduced, light or fitful sleep.

### 5. Reduced appetite

Representing the feeling of a loss of appetite compared with when well. Rate by loss of desire for food or the need to force oneself to eat.

Answer: 0 Normal or increased appetite.

#### 6. Concentration difficulties

Representing difficulties in collecting one's thoughts mounting to incapacitating lack of concentration. Rate according to intensity, frequency, and degree of incapacity produced.

Answer: 2 Occasional difficulties in collecting one's thoughts.

#### 7. Lassitude

Representing a difficulty getting started or slowness initiating and performing everyday activities.

Answer: 2 Difficulties in starting activities.

# 8. Inability to feel

Representing the subjective experience of reduced interest in the surroundings, or activities that normally give pleasure. The ability to react with adequate emotion to circumstances or people is reduced.

Answer: 1

# 9. Pessimistic thoughts

Representing thoughts of guilt, inferiority, self-reproach, sinfulness, remorse and ruin.

Answer: 1

# 10. Suicidal thoughts

Representing the feeling that life is not worth living, that a natural death would be welcome, suicidal thoughts and preparations for suicide. Suicidal attempts should not in themselves influence the rating.

Answer: 0 Enjoys life or takes it as it comes.

**Total Score: 13** 

Disclaimer: By signing this document electronically, I acknowledge that I have reviewed its contents, understand its implications, and confirm its accuracy. I understand that my electronic signature is legally binding, the content of this document is confidential, and will not be shared with third parties without authorization.