

Scatter plot showing VL sES (Y-axis) versus VL no stim (X-axis) for the MOp group (n=282, p=2.1e-29). The plot includes a dashed diagonal line representing the identity line (y=x). Data points are colored by group: red (top-left), purple (middle), and green (bottom-left). The red group shows higher VL sES values compared to VL no stim, while the green and purple groups show lower VL sES values.

CA1
n=229, p=1.5e-24

Scatter plot showing the relationship between distance from sES (mm) and $-\log_{10}(\text{p-value})$ for various brain regions. The x-axis represents distance from sES / mm (0 to 4), and the y-axis represents $-\log_{10}(\text{p-value})$ (0 to 25). A red dashed line indicates a significance threshold at approximately $-\log_{10}(\text{p-value}) \approx 2.5$. Most regions show a positive correlation, with MOp and MOs having the highest p-values.

Region	Distance from sES / mm (approx.)	$-\log_{10}(\text{p-value})$ (approx.)
MOp	1.8	28
MOs	2.4	28
CA1	1.3	23
VISp	0.8	18
RSPd	2.5	15
CL	2.7	13
DG	1.7	14
CP	2.7	10
ACAv	3.2	9
AV	3.4	6
AMd	3.6	3
MD	3.0	3
SS-1	2.2	3
ccb	2.6	3
APN	2.1	3
cing	2.3	3
RSPagl	2.6	3
RSPAd	2.8	3
CA3	2.9	3
LP	2.4	3
VISpm	0.7	4
VISa	0.9	3
GP	0.8	2
HMP	1.9	3
ECT	2.0	3