

## Violence against women II

# Health consequences of intimate partner violence

Jacquelyn C Campbell

**Intimate partner violence, which describes physical or sexual assault, or both, of a spouse or sexual intimate, is a common health-care issue. In this article, I have reviewed research on the mental and physical health sequelae of such violence. Increased health problems such as injury, chronic pain, gastrointestinal, and gynaecological signs including sexually-transmitted diseases, depression, and post-traumatic stress disorder are well documented by controlled research in abused women in various settings. Intimate partner violence has been noted in 3–13% of pregnancies in many studies from around the world, and is associated with detrimental outcomes to mothers and infants. I recommend increased assessment and interventions for intimate partner violence in health-care settings.**

In reviews of US and Canadian population-based surveys during 1985–98, between 8 and 14% of women of all ages reported physical assault in the previous year by a husband, boyfriend, or ex-partner; the lifetime prevalence was between 25 and 30%.<sup>1,2</sup> Such assault is most often termed intimate partner violence in North America, with the definition usually including acts of forced sex as well as other forms of physical violence. It is also recognised that psychological coercion and degradation almost always accompany such violence and are included with physical and sexual assault under the broader terms of spouse abuse, domestic violence, or violence against women.

Most investigations into health-care setting and health consequences of violence against women have been from the USA, and have analysed intimate partner violence; some studies have measured emotional abuse separately. In this article, I have reviewed research published in English during the past decade, and have paid special attention to findings that have been replicated outside the USA. The review includes findings from population-based investigations or studies with sufficient sample size, minimal selection or response bias, controlled comparisons, or rigorous qualitative methods that have been replicated in more than one sample.

### Health-care settings

In investigations in health-care settings (mainly from the USA), yearly prevalence of intimate partner violence has varied between 4 and 23%, with middle-level socioeconomic and well educated groups having the lowest prevalence, and poorer women the highest (figure 1).<sup>1,3</sup> Intimate partner violence might be higher in health-care settings than in most US population-based samples because samples would be expected to include a substantial proportion of battered women. Furthermore, the same studies showed a narrower range of lifetime prevalence (33–39%) that was less affected by socioeconomic status and more similar to population-based studies, suggesting at least two hypotheses. Although health consequences of intimate partner violence can linger long after the violence has ended, seeking help for those symptoms, especially from accident and emergency departments, might subside

over time. Another possibility is that economic and educational resources might not protect women from being abused, but make it easier for them to relatively quickly escape or end the violence, thus rendering them less likely to be currently abused.

### Physical health effects

Women who are abused are frequently treated within health-care systems, however, they generally do not present with obvious trauma, even in accident and emergency departments.<sup>3</sup> Intimate partner violence has long-term negative health consequences for survivors, even after the abuse has ended.<sup>4,5</sup> These effects can manifest as poor health status, poor quality of life, and high use of health services.<sup>6–8</sup> Battering is a significant direct and indirect risk factor for various physical health problems frequently seen in health-care settings. Intimate partner violence is one of the most common causes of injury in women.<sup>9</sup> In several large studies in US accident and emergency departments, 11–30% of injured women whose mechanisms of injury and relationship to the perpetrator had been recorded had been battered.<sup>9</sup> Battered women were more likely to have been injured in the head, face, neck, thorax, breasts, and abdomen than women injured in other ways (figure 2).<sup>10</sup> In two large case-control studies of women in US accident and emergency departments, risk factors for injury by an intimate partner were male rather than female characteristics including histories of arrest, substance abuse, poor education, unemployment, and ex-partner status.<sup>10,11</sup> These results, and findings that a higher proportion of women who have been battered in the past year (12–17%) present to accident and emergency departments than women with acute trauma from abuse (2–3%)<sup>3</sup> suggest the need for universal rather than incident-based screening in emergency-care settings. Although most battered women in the USA state that they have been injured as a result of the abuse, less than half say that they sought health care specifically for those injuries.<sup>12</sup>

40–60% of murders of women in North America are done by intimate partners.<sup>13,14</sup> In less-industrialised countries, percentages might be even higher, although global data on murder of women is sparse.<sup>12,15,16</sup> More research and surveillance is needed to ascertain the global proportion of female deaths, including maternal deaths, that should be attributed to intimate partners.<sup>17</sup> As well as murder, mortality associated with domestic violence includes suicide of women in non-industrialised as well as industrialised societies.<sup>18–20</sup>

*Lancet* 2002; **359**: 1331–36

Johns Hopkins University School of Nursing, 525 North Wolfe Street, Baltimore, MD 21205–2110, USA (Prof J C Campbell PhD) (e-mail: jcampbel@son.jhmi.edu)



Nancy Durrell-McKenna/The Hutchinson Library

## Forced sex

Gynaecological problems are the most consistent, longest lasting, and largest physical health difference between battered and non-battered women. Differential symptoms and conditions include sexually-transmitted diseases, vaginal bleeding or infection, fibroids, decreased sexual desire, genital irritation, pain on intercourse, chronic pelvic pain, and urinary-tract infections.<sup>5,6,8,22–25,27,30–32</sup> In one of the best-sampled (good sampling techniques, high representation, good randomisation, &c) US population-based studies of self-reported data, the odds of having a gynaecological problem were three times greater than average for victims of spouse abuse;<sup>8</sup> evidence of a dose-response effect with severity of physical assault has also been reported.<sup>5</sup> The combination of physical and sexual abuse that

characterises the experience of at least 40–45% of battered women puts these women at an even higher risk for health problems than women only physically assaulted.<sup>24,32</sup> Forced sex has consequences that could explain the higher prevalence of gynaecological problems, although few studies have measured forced sex separately and those that did were not population-based.<sup>23,30</sup> Possible mechanisms of increased risk include the shame and stress reported with forced sex manifesting as especially high levels of stress and depression known to depress the immune system; vaginal, anal, and urethral trauma from forced sex (direct force or lack of lubrication) leading to increased transmission of micro-organisms through direct transmission into the bloodstream or back flow of bacteria in the urethra; and men forcing sex on partners and having unprotected sex with other partners. Although descriptive studies and work in animals implicate all these mechanisms, no largescale controlled studies indicate which factors are most salient.

As well as forced sex, in several well designed, indepth, interview-based, and longitudinal studies battered women have described sexually abusive and controlling acts such as verbal sexual degradation, refusal to use condoms, or refusal to use contraception.<sup>30,33</sup> These issues might partly explain links between intimate partner violence and sexually-transmitted diseases, HIV, and unintended pregnancy in population-based studies in the USA and developing countries.<sup>8,33–35,37–39</sup> Qualitative data from indepth interviews show how abuse interacts with complex social, psychological, and cultural factors involved in decisions and actions to prevent pregnancy or sexually-transmitted diseases, including HIV and AIDS and the difficulty of negotiation of use of condoms or contraception in violent relationships—especially in countries and cultures in which societal norms do not accord women equal say in such decisions.<sup>16,33–35,39–41</sup> However, studies in the USA, South Africa, and India have revealed much secret use of contraception, although there are both health and abuse risks involved with this strategy.<sup>37,42,43</sup> Additionally, from the USA and developing countries come narratives of requests for condom use, HIV testing, or notification of positive-HIV status resulting in abusive incidents, often with accusations of infidelity.<sup>16,42,43</sup>

Figure 1: Domestic violence is more prevalent than most health-care providers realise

The injuries, fear, and stress associated with intimate partner violence can result in chronic health problems such as chronic pain (eg, headaches, back pain) or recurring central nervous system symptoms including fainting and seizures.<sup>2,6,8,21–25</sup> The exact mechanism of such effects has not been established but could include recurrent injury or stress, alterations in neurophysiology, or both. For instance, abused women frequently (10–44%) report choking—incomplete strangulation—and blows to the head resulting in loss of consciousness,<sup>8,26</sup> both of which can lead to serious medical problems including neurological sequelae.

Battered women also have significantly more than average self-reported gastrointestinal symptoms (eg, loss of appetite, eating disorders) and diagnosed functional gastrointestinal disorders (eg, chronic irritable bowel syndrome) associated with chronic stress.<sup>21–24</sup> These disorders may begin during an acutely violent and thus stressful relationship, be related to child sexual abuse, or both.<sup>22</sup> The consequent functional damage to the bowel can last far longer than the violent relationship. Similarly, self-reported cardiac symptoms such as hypertension and chest pain have also been associated with intimate partner violence.<sup>6,25</sup> It is plausible to postulate interactions between genetic tendencies for hypertension, lifestyle risk behaviours (such as smoking),<sup>27</sup> and stress from violent relationships but mechanisms have not been thoroughly investigated. Most studies have been cross-sectional and included analysis of bivariate and multivariate associations rather than mediated or moderated models, such as those reporting an increased incidence of self-reported colds and influenza.<sup>27</sup> Suppression of the immune system as a result of stress, mental-health disorders such as depression, or both is again another reasonable but untested causal hypothesis.

The table presents findings for physical health symptoms and intimate partner violence from five roughly comparable samples of women from the general population, health-care insurance databases, and different primary-care settings in the USA.<sup>8,22–25</sup> There is not absolute congruence across these studies, perhaps because of the different somatic responses people can have to trauma and the different types of injuries abused women can sustain, as well as subtle differences in study methods. Despite this variability, there is agreement that battering has significant short-term and long-term physical health effects.<sup>5,25,28,29</sup>

### Abuse during pregnancy

In a large review of US studies of prevalence of abuse during pregnancy, Gazmararian and colleagues<sup>44</sup> noted a range of 0·9–20·1%; in most studies a range of 3·9–8·3% was recorded. Prevalence of such abuse in other industrialised and non-industrialised nations are similar: 6·4% during the past year and 2·5% overall in the UK (7·8% including threats),<sup>45</sup> 5·5–6·6% in Canada,<sup>46,47</sup> at least 6·8% in South Africa,<sup>48</sup> 11% in Sweden (21% including symbolic violence and threats),<sup>49</sup> and 13% in Nicaragua.<sup>50</sup> Although higher prevalence is associated with more inclusive definitions of abuse, more than one question being asked about abuse, adolescence,<sup>51</sup> and low income of respondents, prevalence across countries is remarkably similar.

The main health effect specific to abuse during pregnancy is the threat to health and risk of death of the mother, fetus, or both from trauma.<sup>17,52,53</sup> Another cause of fetal death, elective pregnancy termination, has also been related to intimate partner violence in large but uncontrolled studies in the USA.<sup>54</sup> Physical abuse in pregnancy is associated with health problems during pregnancy such as sexually-transmitted diseases including HIV-1,<sup>39</sup> urinary-tract infections, substance abuse, depression, and other mental-health symptoms.<sup>55,56</sup> In-depth interviews of women<sup>57</sup> suggest that abuse during pregnancy could be a factor in the overlap of intimate partner violence and child abuse seen in US research.<sup>58</sup>

Although many US studies have noted associations of abuse during pregnancy with infant outcomes such as preterm delivery, fetal distress, antepartum haemorrhage, and pre-eclampsia, evidence is inconsistent across studies.<sup>59</sup> The focus of the most numerous and best-controlled (using all or most appropriate control variables, or well matched case-control studies) investigations has been on intimate partner violence as a risk factor for low birthweight. Although evidence from individual studies has been contradictory, a meta-analysis of 14 published studies from North America and Europe showed a weak but significant association between abuse during pregnancy and low birthweight (odds ratio 1·4, 95% CI 1·1–1·8).<sup>60</sup> Variations between samples and subsamples and potential causal pathways could account for some of the differences in individual studies. Results from one study suggested a stronger relation between abuse and birthweight in women of middle socioeconomic status than in poor women,<sup>61</sup> perhaps because the proportion of low birthweight attributable to intimate partner violence is greater in populations with fewer competing causes. Low birthweight might be the outcome of premature delivery caused by trauma resulting from intimate partner violence in rare situations or might occur in fullterm infants as a result of more complex causal pathways.<sup>60</sup> Maternal low weight gain, smoking, or both were mediators of the connection between abuse and low birthweight in several studies.<sup>62,63</sup> Abusive partners might pressure their wives or girlfriends not to gain weight, or abuse could cause stress, which has in turn been associated with smoking, low weight gain, and consequent low birthweight.<sup>62–64</sup> Only one controlled study of abuse during pregnancy has been done in a developing country; results showed a significant relation with low




Figure 2: **Poster showing physical effects of abuse**  
From: San Francisco District Attorney, Family Violence Project.

birthweight.<sup>16</sup> Thus, there is much to be explained about interactions between risk factors for low birthweight, ethnic origin, and intimate partner violence.<sup>64</sup>

### Mental-health effects

Depression and post-traumatic stress disorder, which have substantial comorbidity, are the most prevalent mental-health sequelae of intimate partner violence (figure 3).<sup>2,8,20,65–69</sup> In a comprehensive meta-analysis of mainly US studies, Golding<sup>20</sup> showed that the risk for depression and post-traumatic stress disorder associated with intimate partner violence was even higher than that resulting from childhood sexual assault. Depression in battered women has also been associated with other life stressors that often accompany domestic violence, such as childhood abuse, daily stressors, many children, changes in residence, forced sex with an intimate partner, marital separations, negative life events, and child behaviour problems.<sup>65,66,68</sup> Some battered women might have chronic depression that is exacerbated by the stress of a violent relationship, but there is also evidence that first episodes of depression can be triggered by such violence, and longitudinal evidence of depression lessening with decreasing intimate partner violence.<sup>68,69</sup> Even so, the relations between violent experiences and mental-health problems needs further study. Much of the sex differences in global incidence of depression could be attributable to the sex difference in intimate partner violence, although this premise has never been specifically tested.



	Reference number				
	McCauley et al <sup>18</sup>	Campbell et al <sup>23</sup>	Coker et al <sup>24</sup>	Leserman et al <sup>22</sup>	Plichta <sup>25</sup>
<b>Symptoms</b>					
General health lower or number of symptoms higher than average	S	S	S	S	S
Digestive problems: diarrhoea, spastic colon, constipation, nausea	S	S	S	NM	NM
Loss of appetite, eating binges, making self vomit	S	S	NM	NS	NM
Abdominal pain, stomach pain	S	S	NM	NM	NM
UTI: bladder/kidney infection, pain, problems with urination	S	S	S	NS	S
Vaginal infection: discharge, itching	S	S	NM	S	NM
Sexually transmitted disease	NM	S	S	NM	NM
AIDS or HIV-1	NM	NS	NM	NM	NM
Vaginal bleeding, severe menstrual problems, dysmenorrhoea	NM	S	NM	NM	NM
Pelvic pain, genital area pain	S	S	S	S	NM
Fibroids or hysterectomy	NM	NS	S	NM	NM
Painful intercourse, sexual dysfunction	NM	S	S	S	NM
Headaches, migraines	S	S	S	S	S
Fainting, passing out	S	NS	NM	S	S
Seizures, convulsions	NM	NS	S	NM	NM
Back pain, chronic neck pain	NS	S	S	S	S
Influenza or cold, stuffy or runny nose	NM	NS	NM	S	NM
Hypertension	NM	NS	S	NM	NM

Table design by J Dienemann (reference 23). NM=not measured in study. S=significant ( $p<0.05$ ) or relative risk=1.0 or higher. NS=not significant. UTI=urinary tract infection.

### Physical symptoms in battered women from US health-care settings or population-based health surveys

Post-traumatic stress disorder has also been extensively studied in North America as a sequelae of intimate partner violence; prevalence in battered women is much higher (weighted mean odds ratio 3.74) than in non-abused women.<sup>20,69</sup> Severity of abuse, previous trauma, and partner dominance have been identified as important precursors of post-traumatic stress disorder developing from intimate partner violence.<sup>67,69</sup> Suicidal tendencies, although less often studied than post-traumatic stress disorder, have also been associated with intimate partner violence in the USA, Scandinavia, and Papua New Guinea.<sup>18–20</sup>

In a Canadian population-based study, Ratner found that in addition to depression, abused women had significantly more anxiety, insomnia, and social dysfunction than those not abused, with physical violence having a stronger effect than psychological abuse.<sup>2</sup> These sleep disturbances seem related to a complex interaction of physical, psychological, and environmental mechanisms, although there is only one known sleep laboratory study of intimate partner violence.<sup>70</sup> Women in developing countries also report mental-health problems from abuse, with 70% of cases of emotional distress in Nicaragua attributed to intimate partner violence, and depression and anxiety reported in battered women in Pakistan.<sup>71,72</sup>

Finally, alcohol and drug abuse is the other mental-health problem most frequently seen in battered women in industrialised countries.<sup>2,8,20,55,56</sup> In at least two comparisons between abused and non-abused women,

controlling for demographic differences, partner substance use, or both, led to the risk of alcohol abuse in battered women disappearing,<sup>11,25</sup> but in another population-based controlled study, the association persisted.<sup>73</sup> Although causal pathways between intimate partner violence and substance abuse are difficult to establish, results from a study set in accident and emergency departments suggested that intimate partner violence preceded both alcohol and drug abuse in most cases.<sup>74</sup> A postulated explanation of substance use as an outcome of intimate partner violence is through post-traumatic stress disorder.<sup>75</sup> Women with post-traumatic stress disorder might use drugs or alcohol to calm or cope with the specific groups of symptoms associated with post-traumatic stress disorder: intrusion, avoidance, and hyperarousal. In a population-based study, substance use was both a risk factor for, and effect of, post-traumatic stress disorder and all forms of violence, especially repeated violence and childhood trauma.<sup>76</sup> Women can also begin to abuse substances through their relationships with men or from wanting to escape the reality of intimate partner violence. It is important to address and understand these complex relations between intimate partner violence, mental health, and behaviour to diagnose accurately and intervene in substance-abuse problems.

### Use of medical care

Analysis of the relations between partner abuse, health status, and use of medical care in women in population-based and clinical studies has shown poorer overall mental and physical health, more injuries, and more consumption of medical care including prescriptions and admissions to hospital in abused than non-abused women.<sup>5–7,22</sup> In a Canadian population-based study,<sup>2</sup> battered women sought care from accident and emergency departments and saw a medical professional about 3 times more often than non-battered women. Furthermore, strong evidence suggests that use of medical services increases with the severity of physical assault.<sup>5</sup> Several estimates have been made of the health-care costs of intimate partner violence, but only a few rigorous cost analyses have been done. In a well designed comparison of health plans, battered women generated around 92% more costs per year than non-battered women, with mental-health services accounting for most of the increased costs.<sup>7</sup>



Figure 3: WHO poster highlighting mental effects of abuse

## Conclusions

Battered women present in all health-care settings with many different physical and mental-health problems and demographic characteristics. Women may present to health-care settings before they present to criminal justice or social service agencies, and if abuse is identified they can receive interventions that increase their safety and improve their health. Thus, assessment for intimate partner violence of all women should be done in all health-care settings;<sup>77</sup> several studies have validated brief clinical screens and chart prompts in inpatient and outpatient settings.<sup>78,79</sup>

The intertwined relations of abuse and physical and mental-health outcomes should be of interest to health-care practitioners as well as researchers. Abuse is a risk factor for many health-care problems, but the causes and extent of such risk are only beginning to be understood. If abuse contributes to factors such as smoking, poor nutrition, substance abuse, and stress, interventions aimed at these problems will not succeed without addressing intimate partner violence. An important first step in establishing evidence-based practice<sup>80</sup> in the area of intimate partner violence is to test interventions such as that described by the March of Dimes Birth Defects Foundation.<sup>81,82</sup> This protocol is not specific to abuse during pregnancy and could be used in any health-care setting in any country by any health-care provider. Research in countries other than the USA, especially developing countries, must be done after establishing interventions that result in improved safety and better mental and physical health for abused women, while recognising that improvement may take time.

It is important that we also recognise the resourcefulness and many strengths that battered women use in addressing their health problems and domestic violence.<sup>41,83</sup> Any assessment and intervention with abused women needs to respect their autonomy and be offered in a spirit of advocacy and collaboration as well as concern for their health and safety. Worldwide research, prevention, and interventions in injury, maternal child-health, mental health, and HIV and AIDS must recognise the role of violence against women.

## Acknowledgment

I thank the Johns Hopkins University Population Information Program's Media/Materials Clearinghouse for their help in finding some of the images used in this publication. The images are from their End Violence Against Women website, [www.endvaw.org](http://www.endvaw.org).

## References

- Jones AS, Campbell JC, Schollenberger J, et al. Annual and lifetime prevalence of partner abuse in a sample of female HMO enrollees. *Womens Health Issues* 1999; **9**: 295–305.
- Ratner PA. The incidence of wife abuse and mental health status in abused wives in Edmonton, Alberta. *Can J Public Health* 1993; **84**: 246–49.
- Dearwater SR, Coben JH, Nah G, et al. Prevalence of domestic violence in women treated at community hospital emergency department. *JAMA* 1998; **280**: 433–38.
- Campbell JC, Lewandowski L. Mental and physical health effects of intimate partner violence on women and children. *Psychiatr Clin North Am* 1997; **20**: 353–74.
- Koss MP, Koss PG, Woodruff WJ. Deleterious effects of criminal victimization on women's health and medical utilization. *Arch Intern Med* 1991; **151**: 342–47.
- Tollestrup K, Sklar D, Frost FJ, et al. Health indicators and intimate partner violence among women who are members of a managed care organization. *Prev Med* 1999; **29**: 431–40.
- Wisner CL, Gilmer TP, Saltzman LE, Zink TM. Intimate partner violence against women: do victims cost health plans more? *J Fam Pract* 1999; **48**: 439–43.
- McCauley J, Kern DE, Kolodner K, et al. The "battering syndrome": prevalence and clinical characteristics of domestic violence in primary care internal medicine practices. *Ann Intern Med* 1995; **123**: 737–46.
- Rand MR. Violence-related injuries treated in hospital emergency departments. Bureau of Justice Statistics special report. Washington, DC: US Department of Justice, 1997.
- Grisso JA, Schwarz DF, Hirschinger N, et al. Violent injuries among women in an urban area. *N Engl J Med* 1999; **341**: 1899–1905.
- Kyriacou DN, Anglin D, Taliaferro E, et al. Risk factors for injury to women from domestic violence against women. *N Engl J Med* 1999; **341**: 1892–98.
- Bachman R, Saltzman LE. Violence against women: estimates from the redesigned survey. Washington, DC: Bureau of Justice Statistics, National Institute of Justice, 1995.
- Crawford M, Gartner R, Dawson M. Intimate femicide in Ontario, 1991–1994. Toronto: Women We Honour Action Committee, 1997.
- Brock K, Stenzel A. When men murder women: an analysis of 1997 homicide data—females murdered by males in single victim/single offender incidents. Washington, DC: Violence Policy Center, 1999.
- Gartner R. The victims of homicide: a temporal and cross national comparison. *Am Soc Rev* 1990; **55**: 92–106.
- Heise LL, Ellsberg MC, Gottemoeller M. Ending violence against women. *Popul Rep L* 1999; **27**: 1–43.
- Parsons LH, Harper MA. Violent maternal deaths in North Carolina. *Obstet Gynecol* 1999; **94**: 990–93.
- Counts DA. Female suicide and wife abuse in cross cultural perspective. *Suicide Life Threat Behav* 1987; **17**: 194–204.
- Bergman B, Brismar B. Suicide attempts by battered wives. *Acta Psychiatr Scand* 1991; **83**: 380–84.
- Golding JM. Intimate partner violence as a risk factor for mental disorders: a meta-analysis. *J Fam Viol* 1999; **14**: 99–132.
- Diaz-Olavarrieta C, Campbell JC, Garcia de la Cadena C, Paz F, Villa A. Domestic violence against patients with chronic neurologic disorders. *Arch Neurol* 1999; **56**: 681–85.
- Leserman J, Li D, Drossman DA, Hu YJB. Selected symptoms associated with sexual and physical abuse among female patients with gastrointestinal disorders: the impact on subsequent health care visits. *Psychol Med* 1998; **28**: 417–25.
- Campbell JC, Dienemann J, Jones AS, et al. Intimate partner violence and physical health consequences in a sample of female HMO enrollees. *Arch Intern Med* (in press).
- Coker AL, Smith PH, Bethea L, King MR, McKeown RE. Physical health consequences of physical and psychological intimate partner violence. *Arch Fam Med* 2000; **9**: 451–57.
- Plichta SB. Violence and abuse: implications for women's health. In Falik MK, Collins KS, eds. *Women's Health, the Commonwealth Fund Survey*. Baltimore, MD: The Johns Hopkins University Press, 1996.
- Sharps PW, Campbell JC, Campbell DW, Gary FA, Webster DW. The role of alcohol use in intimate partner femicide. *Am J Addictions* 2001; **10**: 122–35.
- Letourneau EJ, Holmes M, Chasendunn-Roark J. Gynecologic health consequences to victims of interpersonal violence. *Womens Health Issues* 1999; **9**: 115–20.
- Drossman DA, Talley NJ, Leserman J, Olden KW, Barreiro MA. Sexual and physical abuse and gastrointestinal illness: review and recommendations. *Ann Intern Med* 1995; **123**: 782–94.
- Sutherland C, Bybee D, Sullivan C. The long-term effects of battering on women's health. *Womens Health Res Gen Behav Pol* 1998; **4**: 41–70.
- Campbell JC, Soeken K. Forced sex and intimate partner violence: effects on women's health. *Violence Against Women* 1999; **5**: 1017–35.
- Schei B, Bakketeig LS. Gynaecological impact of sexual and physical abuse by spouse: a study of a random sample of Norwegian women. *Br J Obstet Gynaecol* 1989; **96**: 1379–83.
- Collett BJ, Cordle CJ, Stuart CR, Jagger C. A comparative study of women with chronic pelvic pain, chronic nonpelvic pain and those with no history of pain attending general practitioners. *Br J Obstet Gynaecol* 1998; **105**: 87–92.
- Champion JD, Shain RN. The context of sexually transmitted disease: life histories of woman abuse. *Issues MHN*; **19**: 463–80.
- Coker AL, Richter DL. Violence against women in Sierra Leone: frequency and correlates of intimate partner violence and forced sexual intercourse. *Afr J Reprod Health* 1998; **2**: 61–72.
- Wood K, Jewkes R. Violence, rape, and sexual coercion: everyday love in a South African township. *Gender Dev* 1997; **5**: 41–46.
- Gazmararian JA, Adams M, Saltzman L, et al. The relationship 38 between intendedness and physical violence in mothers of newborns. *Obstet Gynecol* 1995; **85**: 131–38.
- Wingood G, DiClemente R. The effects of an abusive primary partner on the condom use and sexual negotiation practices of African-American women. *Am J Public Health* 1997; **87**: 1016–18.
- Watts C, Ndlovu M, Njovana E, Keogh E. Women, violence and HIV/AIDS in Zimbabwe. *SAfAIDS* 1997; **5**: 2–6.
- Maman S, Campbell JC, Sweat M, Gielen AC. The intersection of HIV and violence: directions for future research and interventions. *Soc Sci Med* 2000; **50**: 459–78.

- 40 Martin SL, Kilgallen B, Tsui AO, et al. Sexual behaviors and reproductive health outcomes: associations with wife abuse in India. *JAMA* 1999; **282**: 1967–72.
- 41 Lempert LB. Women's strategies for survival: developing agency in abusive relationships. *J Fam Violence* 1996; **11**: 269–90.
- 42 Zierler S, Cunningham W, Andersen R, et al. Violence victimization after HIV infection in a US probability sample of adult patients in primary care. *Am J Public Health* 2000; **90**: 208–15.
- 43 Gielen AC, McDonnell KA, Burke JG, O'Campo P. Women's lives after an HIV-positive diagnosis: disclosure and violence. *Matern Child Health J* 2000; **4**: 111–20.
- 44 Gazmararian JA, Lazorki S, Spitz A, et al. Prevalence of violence against pregnant women: a review of the literature. *JAMA* 1996; **275**: 1915–20.
- 45 Mezey GC, Bacchus L, Bewley S, Haworth A. An exploration of the prevalence, nature and effects of domestic violence in pregnancy. Surrey: Violence Research Program, University of London, 2001.
- 46 Stewart DE, Cecutti A. Physical abuse during pregnancy. *Can Med Assoc J* 1993; **149**: 1257–63.
- 47 Muhajarine N, D'Arcy C. Physical abuse during pregnancy: prevalence and risk factors. *Can Med Assoc J* 1999; **160**: 1007–11.
- 48 Jewkes R, Penn-Kekana L, Levin J, Ratsaka M, Schriber M. "He must give me money, he musn't beat me": violence against women in three South African provinces. CERSA (Women's Health) technical report. Pretoria: Medical Research Council, 1999.
- 49 Widding L, Grimstad H, Moller A, Schei B, Janson PO. Prevalence of physical and sexual abuse before and during pregnancy among Swedish couples. *ACTA Obstet Gynecol Scand* 1999; **78**: 310–15.
- 50 Ellsberg MC, Pena R, Herrera A, Liljestrand J, Winkvist A. Candies in hell: women's experience of violence in Nicaragua. *Soc Sci Med* (in press).
- 51 Parker B, McFarlane J, Soeken K, Torres S, Campbell D. Physical and emotional abuse in pregnancy: a comparison of adult and teenage women. *Nurs Res* 1993; **42**: 173–78.
- 52 Jejeebhoy SJ. Associations between wife-beating and fetal and infant death: impressions from a survey in rural India. *Stud Fam Plann* 1998; **29**: 300–08.
- 53 Pearlman MD, Tintinalli JE, Lorenz RP. Blunt trauma during pregnancy. *N Engl J Med* 1990; **323**: 1609–13.
- 54 Glander SS, Moore ML, Michielutte R, Parsons LH. The prevalence of domestic violence among women seeking abortion. *Obstet Gynecol* 1998; **91**: 1002–06.
- 55 Amaro H, Fried L, Cabral H, et al. Violence during pregnancy and substance use. *Am J Public Health* 1990; **80**: 575–79.
- 56 Martin SL, Kilgallen B, Dee DL, Dawson S, Campbell JC. Women in a prenatal care/substance abuse treatment program: links between domestic violence and mental health. *Matern Child Health J* 1998; **2**: 85–94.
- 57 Campbell JC, Oliver C, Bullock L. Why battering during pregnancy? *AWHONNS Clin Issues Perinat Womens Health Nurs* 1993; **4**: 343–49.
- 58 Straus MA, Gelles RJ. Physical violence in American families: risk factors and adaptations to family violence in 8,145 families. New Brunswick, NJ: Transaction Publishers, 1990.
- 59 Gazmararian JA, Petersen R, Spitz AM, Goodwin MM, Saltzman LE, Marks JS. Violence and reproductive health: current knowledge and future research directions. *Matern Child Health J* 2000; **4**: 79–84.
- 60 Murphy CC, Schei B, Myhr TL, Du Mont J. Abuse: a risk factor for low birth weight? A systematic review and meta-analysis. *Can Med Assoc J* 2001; **164**: 1567–72.
- 61 Bullock LF, McFarlane J. Higher prevalence of low birthweight infants born to battered women. *Am J Nurs* 1989; **89**: 1153–55.
- 62 Campbell JC, Ryan J, Campbell DW, et al. Physical and nonphysical abuse and other risk factors for low birthweight among term and preterm babies: a multiethnic case control study. *Am J Epidemiol* 1999; **150**: 714–26.
- 63 McFarlane J, Parker B, Soeken K. Abuse during pregnancy: associations with maternal health and infant birth weight. *Nurs Res* 1996; **45**: 37–42.
- 64 Curry MA, Harvey SM. Stress related to domestic violence during pregnancy and infant birthweight. In: J Campbell, ed. Empowering survivors of abuse: health care for battered women and their children. Thousand Oaks, CA: Sage, 1998.
- 65 Cascardi M, O'Leary KD, Schlee KA. Co-occurrence and correlates of posttraumatic stress disorder and major depression in physically abused women. *J Fam Violence* 1999; **14**: 227–50.
- 66 Campbell JC, Kub J, Belknap RA, Templin T. Predictors of depression in battered women. *Violence Against Women* 1997; **3**: 271–93.
- 67 Campbell R, Sullivan CM, Davidson WS. Women who use domestic violence shelters: changes in depression over time. *Psych Women Q* 1995; **19**: 237–55.
- 68 Campbell JC, Soeken K. Women's responses to battering over time: an analysis of change. *J Interpers Viol* 1999; **14**: 21–40.
- 69 Silva C, McFarlane J, Soeken K, Parker B, Reel S. Symptoms of post-traumatic stress disorder in abused women in a primary care setting. *J Womens Health* 1997; **6**: 543–52.
- 70 Humphreys JC, Lee K, Neylan T, Marmar CR. Sleep patterns of sheltered battered women. *Image J Nurs Sch* 1999; **31**: 139–43.
- 71 Ellsberg M, Caldera T, Herrera A, Winkvist A, Kullgren G. Domestic violence and emotional distress among Nicaraguan women: results from a population-based study. *Am Psychol* 1999; **54**: 30–36.
- 72 Fikree FF, Bhatt LI. Domestic violence and health of Pakistani women. *Internat J Gynaecol Obstet* 1999; **65**: 195–201.
- 73 Brookoff D, O'Brien KK, Cook CS, Thompson TD, Williams C. Characteristics of participants in domestic violence. *JAMA* 1997; **277**: 1369–73.
- 74 Stark E, Flitcraft AH. Women at risk: domestic violence and women's health. Thousand Oaks, CA: Sage, 1996.
- 75 Kulka RA, Schlenger WE, Fairbank JA, et al. Trauma and the Vietnam war generation: report of findings from the national Vietnam veterans readjustment study. New York: Bruner-Mazel, 1990.
- 76 Kilpatrick DG, Acierno R, Resnick HS, Saunders BE, Best CL. A two-year longitudinal analysis of the relationships between violent assault and substance use in women. *J Consult Clin Psychol* 1997; **65**: 834–47.
- 77 Family Violence Prevention Fund. Preventing domestic violence: clinical guidelines on routine screening. San Francisco, CA: Family Violence Prevention Fund, 1999.
- 78 Soeken K, Parker B, McFarlane J, et al. The abuse assessment screen: a clinical instrument to measure frequency, severity, and perpetrator of abuse against women. In: Campbell J, ed. Empowering survivors of abuse: health care for battered women and their children. Thousand Oaks, CA: Sage, 1998.
- 79 Wiist WH, McFarlane J. The effectiveness of an abuse assessment protocol in public health prenatal clinics. *Am J Public Health* 1999; **89**: 1217–21.
- 80 Chalk R, King P. Health care interventions in family violence: Washington, DC: National Academy Press, 1998.
- 81 Parker B, McFarlane J, Soeken K, Silva C, Reel S. Testing an intervention to prevent further abuse to pregnant women. *Res Nurs Health* 1999; **22**: 59–66.
- 82 March of Dimes Birth Defects Foundation. Protocol of care for the battered woman. New York: March of Dimes, 2001.
- 83 Campbell JC, Rose L, Kub J, Nedd, D. Voices of strength and resistance: a contextual and longitudinal analysis of women's responses to battering. 1998. *J Interpers Violence* 1998; **13**: 743–61.