


PGSChat Test Case-1: Information Retrieval




PGSChat Test Case-2: Information Retrieval & Visualization



PGSChat Test Case-3: Information Retrieval & Visualization







..Contd..



PGSChat Test Case-4: Information Retrieval & Visualization



PGSChat Test Case-5: Query Chat GPt Directly



Functional annotation is the process of assigning biological functions to genes based on experimental evidence or computational predictions. Here is a brief functional annotation of the genes you mentioned:

- 1. TREM2 (Triggering Receptor Expressed on Myeloid Cells 2): It is involved in the regulation of immune responses and phagocytosis. Mutations in TREM2 have been associated with increased risk of Alzheimer's disease.
- SORL1 (Sortilin-Related Receptor 1): It plays a role in the trafficking and processing of amyloid precursor protein (APP), which is involved in the production of amyloid-beta plaques
 in Alzheimer's disease.
- 3. APOC1 (Apolipoprotein C1): It is a component of lipoprotein particles and plays a role in lipid metabolism. Variations in APOC1 have been associated with increased risk of Alzheimer's disease.
- 4. APOE (Apolipoprotein E): It is involved in lipid transport and metabolism. APOE has been extensively studied in relation to Alzheimer's disease, as certain variants of this gene are associated with increased risk.
- 5. BIN1 (Bridging Integrator 1): It is involved in membrane remodeling and endocytosis. BIN1 has been implicated in Alzheimer's disease, as it interacts with tau protein and affects its aggregation.
- 6. CYP27C1 (Cytochrome P450 Family 27 Subfamily C Member 1): It encodes an enzyme involved in the metabolism of vitamin D. Its specific role in relation to Alzheimer's disease is not well understood.
- 7. ABI3 (AbI Interactor 3): It is involved in signal transduction and regulation of cell adhesion. ABI3 has been associated with immune-related disorders, but its role in Alzheimer's disease is not well characterized.
- 8. DSG2 (Desmoglein 2): It is a component of desmosomes, which are cell adhesion structures. DSG2 mutations have been associated with arrhythmogenic right ventricular cardiomyopathy, but its role in Alzheimer's disease is unclear.

Please note that the functional annotations provided here are a general overview and may not encompass all known functions or associations of these genes. Further research and experimental evidence are required to fully understand the functions and roles of these genes in specific biological processes or diseases.

(iii) Chat GPT/GPT-4 provides the answer to the query using its vast pretrained knowledge