

Explore

Notes

Output Created		17-MAY-2026 00:53:33
Comments		
Input	Data	C:\Users\user\Desktop\paper.sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	270
Missing Value Handling	Definition of Missing	User-defined missing values for dependent variables are treated as missing.
	Cases Used	Statistics are based on cases with no missing values for any dependent variable or factor used.
Syntax		EXAMINE VARIABLES=pcq_total_m ean uwes_total_mean msq_total_mean /PLOT BOXPLOT HISTOGRAM NPLOT /STATISTICS DESCRIPTIVES EXTREME /CINTERVAL 95 /MISSING LISTWISE /NOTOTAL.
Resources	Processor Time	00:00:01.83
	Elapsed Time	00:00:01.02

[DataSet1] C:\Users\user\Desktop\paper.sav

Case Processing Summary

	Valid		Cases Missing		Total	
	N	Percent	N	Percent	N	Percent
PCQ total mean	270	100.0%	0	0.0%	270	100.0%
UWES total mean	270	100.0%	0	0.0%	270	100.0%
MSQ total mean	270	100.0%	0	0.0%	270	100.0%

Descriptives

		Statistic	Std. Error
PCQ total mean	Mean	4.1307	.03889
	95% Confidence Interval for Mean	Lower Bound	4.0541
		Upper Bound	4.2073
	5% Trimmed Mean	4.1491	
	Median	4.2083	
	Variance	.408	
	Std. Deviation	.63909	
	Minimum	2.04	
	Maximum	5.33	
	Range	3.29	
	Interquartile Range	.93	
	Skewness	-.359	.148
	Kurtosis	-.283	.295
UWES total mean	Mean	4.2263	.05530
	95% Confidence Interval for Mean	Lower Bound	4.1175
		Upper Bound	4.3352
	5% Trimmed Mean	4.2487	
	Median	4.3333	
	Variance	.826	
	Std. Deviation	.90869	
	Minimum	1.44	
	Maximum	6.00	
	Range	4.56	
	Interquartile Range	1.33	
	Skewness	-.371	.148
	Kurtosis	-.252	.295
MSQ total mean	Mean	3.6941	.04207

Descriptives

			Statistic	Std. Error
	95% Confidence Interval for Mean	Lower Bound	3.6112	
		Upper Bound	3.7769	
	5% Trimmed Mean		3.7130	
	Median		3.7500	
	Variance		.478	
	Std. Deviation		.69133	
	Minimum		1.65	
	Maximum		4.95	
	Range		3.30	
	Interquartile Range		.91	
	Skewness		-.396	.148
	Kurtosis		-.312	.295

Extreme Values

			Case Number	Value
PCQ total mean	Highest	1	182	5.33
		2	198	5.33
		3	66	5.29
		4	250	5.29
		5	222	5.25
	Lowest	1	139	2.04
		2	210	2.38
		3	157	2.54
		4	241	2.67
		5	81	2.67
UWES total mean	Highest	1	140	6.00
		2	163	6.00
		3	74	5.89
		4	178	5.89
		5	198	5.89 ^a
	Lowest	1	241	1.44
		2	153	1.56

Extreme Values

			Case Number	Value
MSQ total mean	Highest	3	139	2.00
		4	53	2.00
		5	210	2.22 ^b
	Lowest	1	17	4.95
		2	163	4.95
		3	251	4.95
		4	178	4.90
		5	250	4.90 ^c
		1	139	1.65
		2	210	1.80
3		172	2.10	
4	113	2.10		
5	2	2.15		

a. Only a partial list of cases with the value 5.89 are shown in the table of upper extremes.

b. Only a partial list of cases with the value 2.22 are shown in the table of lower extremes.



c. Only a partial list of cases with the value 4.90 are shown in the table of upper extremes.



Tests of Normality

	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
PCQ total mean	.055	270	.046	.984	270	.003
UWES total mean	.080	270	.000	.984	270	.004
MSQ total mean	.067	270	.005	.980	270	.001



a. Lilliefors Significance Correction



PCQ total mean









UWES total mean





MSQ total mean





Oneway

Notes

Output Created		17-MAY-2026 00:53:34
Comments		
Input	Data	C: \Users\user\Desktop\paper .sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	270
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each analysis are based on cases with no missing data for any variable in the analysis.
Syntax		ONEWAY pcq_total_mean uwes_total_mean msq_total_mean BY gender /STATISTICS HOMOGENEITY.
Resources	Processor Time	00:00:00.03
	Elapsed Time	00:00:00.01

Tests of Homogeneity of Variances

		Levene Statistic	df1	df2	Sig.
PCQ total mean	Based on Mean	1.216	1	268	.271
	Based on Median	1.034	1	268	.310
	Based on Median and with adjusted df	1.034	1	267.676	.310
	Based on trimmed mean	1.168	1	268	.281
UWES total mean	Based on Mean	.022	1	268	.881
	Based on Median	.003	1	268	.954
	Based on Median and with adjusted df	.003	1	265.218	.954
	Based on trimmed mean	.011	1	268	.915
MSQ total mean	Based on Mean	1.103	1	268	.294
	Based on Median	.681	1	268	.410
	Based on Median and with adjusted df	.681	1	262.286	.410
	Based on trimmed mean	.976	1	268	.324

ANOVA

		Sum of Squares	df	Mean Square	F	Sig.
PCQ total mean	Between Groups	.048	1	.048	.116	.733
	Within Groups	109.820	268	.410		
	Total	109.868	269			
UWES total mean	Between Groups	.001	1	.001	.002	.968
	Within Groups	222.118	268	.829		
	Total	222.119	269			
MSQ total mean	Between Groups	.728	1	.728	1.525	.218
	Within Groups	127.838	268	.477		
	Total	128.566	269			

Correlations

Notes

Output Created		17-MAY-2026 00:53:35
Comments		
Input	Data	C: \Users\user\Desktop\paper .sav
	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	270
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each pair of variables are based on all the cases with valid data for that pair.
Syntax		CORRELATIONS VARIABLES=pcq_total_m ean uwes_total_mean msq_total_mean /PRINT=TWOTAIL NOSIG /MISSING=PAIRWISE.
Resources	Processor Time	00:00:00.03
	Elapsed Time	00:00:00.01

Correlations

		PCQ total mean	UWES total mean	MSQ total mean
PCQ total mean	Pearson Correlation	1	.529**	.482**
	Sig. (2-tailed)		.000	.000
	N	270	270	270
UWES total mean	Pearson Correlation	.529**	1	.651**
	Sig. (2-tailed)	.000		.000
	N	270	270	270
MSQ total mean	Pearson Correlation	.482**	.651**	1
	Sig. (2-tailed)	.000	.000	
	N	270	270	270

** . Correlation is significant at the 0.01 level (2-tailed).