


CS4102 Algorithms

Nate Brunelle

Spring 2018

Warm up


Given any 5 points on the unit square, show
there's always a pair distance $\leq \frac{\sqrt{2}}{2}$ apart



If points p_1, p_2 in same quadrant, then $\delta(p_1, p_2) \leq \frac{\sqrt{2}}{2}$

Given 5 points, two must share the same quadrant

Pigeonhole Principle!



Today's Keywords

- Solving recurrences
- Cookbook Method
- Master Theorem
- Substitution Method

CLRS Readings

- Chapter 4

Homeworks

- Hw1 due 11pm Friday, February 9
 - Written (use Latex!)
 - Asymptotic notation
 - Recurrences
 - Divide and conquer
- HW2 released Wednesday, February 7
 - Programming assignment (use Python)
 - Divide and Conquer

Recurrence Solving Techniques



Tree



Guess/Check



“Cookbook”



Substitution


Observation

- **Divide:** $D(n)$ time,
- **Conquer:** recurse on small problems, size s
- **Combine:** $C(n)$ time
- **Recurrence:**
 - $T(n) = D(n) + \sum T(s) + C(n)$
- Many D&C recurrences are of form:
 - $T(n) = aT\left(\frac{n}{b}\right) + f(n)$

General

$$T(n) = aT\left(\frac{n}{b}\right) + f(n)$$


$$T(n) = \sum_{i=0}^{\log_b n} a^i f\left(\frac{n}{b^i}\right)$$



3 Cases

$$T(n) = f(n) + af\left(\frac{n}{b}\right) + a^2f\left(\frac{n}{b^2}\right) + a^3f\left(\frac{n}{b^3}\right) + \dots + a^L f\left(\frac{n}{b^L}\right)$$


Case 1:
Most work
happens at
the leaves



Case 2:
Work happens
consistently
throughout



Case 3:
Most work
happens at
top of tree



Master Theorem

$$T(n) = aT\left(\frac{n}{b}\right) + f(n)$$

- **Case 1:** if $f(n) = O(n^{\log_b a - \varepsilon})$ for some constant $\varepsilon > 0$, then $T(n) = \Theta(n^{\log_b a})$
- **Case 2:** if $f(n) = \Theta(n^{\log_b a})$, then $T(n) = \Theta(n^{\log_b a} \log n)$
- **Case 3:** if $f(n) = \Omega(n^{\log_b a + \varepsilon})$ for some constant $\varepsilon > 0$, and if $af\left(\frac{n}{b}\right) \leq cf(n)$ for some constant $c < 1$ and all sufficiently large n , then $T(n) = \Theta(f(n))$

Proof of Case 1

$$T(n) = \sum_{i=0}^{\log_b n} a^i f\left(\frac{n}{b^i}\right),$$
$$f(n) = O(n^{\log_b a - \varepsilon}) \Rightarrow f(n) \leq c \cdot n^{\log_b n - \varepsilon}$$

Insert math here...

Conclusion: $T(n) = O(n^{\log_b a})$

[see lecture supplemental](#)

Master Theorem Example 1

$$T(n) = aT\left(\frac{n}{b}\right) + f(n)$$

- **Case 1:** if $f(n) = O(n^{\log_b a - \varepsilon})$ for some constant $\varepsilon > 0$, then $T(n) = \Theta(n^{\log_b a})$
- **Case 2:** if $f(n) = \Theta(n^{\log_b a})$, then $T(n) = \Theta(n^{\log_b a} \log n)$
- **Case 3:** if $f(n) = \Omega(n^{\log_b a + \varepsilon})$ for some constant $\varepsilon > 0$, and if $af\left(\frac{n}{b}\right) \leq cf(n)$ for some constant $c < 1$ and all sufficiently large n , then $T(n) = \Theta(f(n))$


$$T(n) = 2T\left(\frac{n}{2}\right) + n$$

Case 2

$$\Theta(n^{\log_2 2} \log n) = \Theta(n \log n)$$

Tree method

$$T(n) = 2T\left(\frac{n}{2}\right) + n$$



Master Theorem Example 2

$$T(n) = aT\left(\frac{n}{b}\right) + f(n)$$

- **Case 1:** if $f(n) = O(n^{\log_b a - \varepsilon})$ for some constant $\varepsilon > 0$, then $T(n) = \Theta(n^{\log_b a})$
- **Case 2:** if $f(n) = \Theta(n^{\log_b a})$, then $T(n) = \Theta(n^{\log_b a} \log n)$
- **Case 3:** if $f(n) = \Omega(n^{\log_b a + \varepsilon})$ for some constant $\varepsilon > 0$, and if $af\left(\frac{n}{b}\right) \leq cf(n)$ for some constant $c < 1$ and all sufficiently large n , then $T(n) = \Theta(f(n))$


$$T(n) = 4T\left(\frac{n}{2}\right) + 5n$$

Case 1

$$\Theta(n^{\log_2 4}) = \Theta(n^2)$$

Tree method

$$T(n) = 4T\left(\frac{n}{2}\right) + 5n$$




$5n$



$\frac{4}{2} \cdot 5n$



$\frac{16}{4}$



$2^{\log_2 n}$

Master Theorem Example 3

$$T(n) = aT\left(\frac{n}{b}\right) + f(n)$$

- **Case 1:** if $f(n) = O(n^{\log_b a - \varepsilon})$ for some constant $\varepsilon > 0$, then $T(n) = \Theta(n^{\log_b a})$
- **Case 2:** if $f(n) = \Theta(n^{\log_b a})$, then $T(n) = \Theta(n^{\log_b a} \log n)$
- **Case 3:** if $f(n) = \Omega(n^{\log_b a + \varepsilon})$ for some constant $\varepsilon > 0$, and if $af\left(\frac{n}{b}\right) \leq cf(n)$ for some constant $c < 1$ and all sufficiently large n , then $T(n) = \Theta(f(n))$


$$T(n) = 3T\left(\frac{n}{2}\right) + 8n$$

Case 1

$$\Theta(n^{\log_2 3}) \approx \Theta(n^{1.5})$$

Karatsuba

$$T(n) = 3T\left(\frac{n}{2}\right) + 8n$$



$$8 \cdot 1n$$

$$\frac{8}{2} \cdot 3n$$

$$\frac{8}{4} \cdot 9n$$

$$\frac{8}{2^{\log_2 n}} \cdot 3^{\log_2 n}$$

Master Theorem Example 4

$$T(n) = aT\left(\frac{n}{b}\right) + f(n)$$

- **Case 1:** if $f(n) = O(n^{\log_b a - \varepsilon})$ for some constant $\varepsilon > 0$, then $T(n) = \Theta(n^{\log_b a})$
- **Case 2:** if $f(n) = \Theta(n^{\log_b a})$, then $T(n) = \Theta(n^{\log_b a} \log n)$
- **Case 3:** if $f(n) = \Omega(n^{\log_b a + \varepsilon})$ for some constant $\varepsilon > 0$, and if $af\left(\frac{n}{b}\right) \leq cf(n)$ for some constant $c < 1$ and all sufficiently large n , then $T(n) = \Theta(f(n))$


$$T(n) = 2T\left(\frac{n}{2}\right) + 15n^3$$

Case 3

$$\Theta(n^3)$$

Tree method

$$T(n) = 2T\left(\frac{n}{2}\right) + 15n^3$$



Recurrence Solving Techniques



Tree



Guess/Check



“Cookbook”




Substitution

Substitution Method

- Idea: take a “difficult” recurrence, re-express it such that one of our other methods applies.
- Example: $T(n) = 2T(\sqrt{n}) + \log_2 n$

Tree method

$$T(n) = 2T(\sqrt{n}) + \log_2 n$$



$$T(n) = O(\log_2 n \cdot \log_2 \log_2 n)$$

Substitution Method

- Idea: take a “difficult” recurrence, re-express it such that one of our other methods applies.
- Example: $T(n) = 2T(\sqrt{n}) + \log_2 n$

Let $n = 2^m$, i.e. $m = \log_2 n$

$$T(2^m) = 2T\left(2^{\frac{m}{2}}\right) + m \quad \text{Rewrite in terms of exponent!}$$

$$\text{Let } S(m) = 2S\left(\frac{m}{2}\right) + m \quad \text{Case 2!}$$


$$\text{Let } S(m) = \Theta(m \log m) \quad \text{Substitute Back}$$

$$\text{Let } T(n) = \Theta(\log n \log \log n)$$

Tree method

$$n = 2^m$$


$$T(2^m) = 2T(2^{\frac{m}{2}}) + m$$



Tree method

$$n = 2^m$$

$$T(2^m) = 2T(2^{m-1}) + \log_2 n$$




Tree method

$$n = 2^m$$

$$T(2^m) = S(m)$$

$$S(m) = 2S\left(\frac{m}{2}\right) + m$$



$$T(n) = O(m \cdot \log_2 m) = O(\log_2 n \cdot \log_2 \log_2 n)$$