From Fermat's Equation to Modularity: A Structural Understanding of Fermat's Last Theorem

Sungjun Lee

Feburary 20, 2026

Abstract

Fermat's Last Theorem states that the Diophantine equation

$$x^n + y^n = z^n$$

has no nontrivial solutions in positive integers for any exponent $n > 2$. Its statement is elementary enough to be understood in a first course in algebra, yet its proof belongs to some of the deepest parts of modern number theory. The theorem was ultimately proved by Andrew Wiles through a chain of ideas linking Diophantine equations, elliptic curves, modular forms, Galois representations, and deformation theory.

The purpose of this paper is not to reproduce the full technical proof, which lies far beyond the scope of an introductory exposition, but to understand why the theorem is so deep and why a simple-looking equation forces us into such advanced mathematics. The main thesis of this paper is that Fermat's Last Theorem became solvable only after the original arithmetic question was transformed into a structural problem about symmetry, geometry, and automorphic behavior.

1 About the Author

The first time I encountered Fermat's Last Theorem, I was surprised by how different its appearance was from its reputation. The statement itself looked almost too simple. If the Pythagorean equation

$$x^2 + y^2 = z^2$$

has many integer solutions, then it seemed natural to ask whether

$$x^3 + y^3 = z^3, \quad x^4 + y^4 = z^4,$$

and so on, might also have solutions. At first, the theorem looked like a natural extension of something already familiar. However, the more I learned about it, the more I realized that the actual difficulty of the problem was not computational. The problem was structural.

What fascinated me most was the enormous distance between the statement and the proof. The equation is accessible to anyone who understands exponents, but the proof leads through elliptic curves, modular forms, Hecke algebras, and Galois representations. That gap made me want to understand the theorem not merely as a historical fact, but as an example of how modern mathematics changes the language of a problem until the problem becomes solvable.

2 The Statement and Its First Mystery

Fermat's Last Theorem asserts that for every integer $n > 2$, the equation

$$x^n + y^n = z^n$$

has no nonzero integer solutions. In modern language, there are no nontrivial solutions in positive integers.

The first mystery comes from the contrast with the case $n = 2$. When $n = 2$, we have the classical Pythagorean triples:

$$3^2 + 4^2 = 5^2, \qquad 5^2 + 12^2 = 13^2,$$

and in fact infinitely many more. One can parametrize all primitive Pythagorean triples by setting

$$x = m^2 - n^2, \qquad y = 2mn, \qquad z = m^2 + n^2,$$

for integers $m > n > 0$ with suitable parity and coprimality conditions. So the exponent 2 is not only solvable, but richly solvable.

This immediately raises a natural question: what changes when the exponent becomes 3, or 4, or any higher integer? Why should the equation behave so differently merely because the exponent increases by 1? If one approaches the equation naïvely, it is tempting to imagine that higher powers should simply make the numbers larger, while the basic nature of the problem stays the same. But this is false. The arithmetic structure changes drastically.

3 Why the Problem Is Not Merely Computational

One might think that Fermat's Last Theorem is difficult because the numbers involved can be large. That is not the real reason. A computer can check many examples and find no solutions, but such a search says very little about all integers. The true difficulty is that the theorem concerns exact integer solutions, and integers are rigid objects. In analysis, one often studies continuity and approximation. In Diophantine equations, approximation is not enough. Either an exact integer solution exists, or it does not.

This rigidity explains why elementary numerical evidence is weak. The failure of small examples such as

$$2^3 + 3^3 \neq 4^3, \qquad 3^4 + 4^4 \neq 5^4,$$

does not come close to explaining why no examples can ever occur. The theorem is not asking for a pattern in decimal expansions or a rough asymptotic inequality. It is asking for an impossibility statement in the discrete arithmetic universe.

Historically, mathematicians proved special exponents one at a time. Fermat himself proved the case $n = 4$ using infinite descent. Euler proved the case $n = 3$. Over time, many individual exponents were settled. But the general theorem resisted proof because there was no uniform method strong enough to handle all exponents at once. The eventual solution did not come from extending elementary methods. It came from changing the problem into a different kind of problem.

4 Reduction to Prime Exponents

A standard first step is to observe that it is enough to prove the theorem for prime exponents. The reason is simple but important. Suppose there were a solution for some composite exponent n. If $n = mp$ where p is prime, then

$$x^n + y^n = z^n$$

can be rewritten as

$$(x^m)^p + (y^m)^p = (z^m)^p.$$

Thus a solution for a composite exponent would induce a solution for a prime exponent. Therefore, to prove Fermat's Last Theorem, it suffices to rule out solutions for prime exponents $p > 2$.

This reduction already shows a common theme in mathematics: even before solving a problem, one tries to identify its essential core. The theorem is not really about all integers $n > 2$ separately. It is about prime exponents.

5 A New Philosophy: Arithmetic Through Geometry

The decisive conceptual shift in modern number theory is the realization that arithmetic questions can sometimes be studied more effectively through geometric objects. Instead of directly attacking the equation

$$a^p + b^p = c^p,$$

one asks whether such a hypothetical solution would force the existence of another object with impossible properties.

This is where elliptic curves enter. An elliptic curve over $\mathbb{Q}$ can be given, after a suitable change of variables, by an equation of the form

$$E : y^2 = x^3 + Ax + B,$$

where $A, B \in \mathbb{Q}$ and the discriminant

$$\Delta(E) = -16(4A^3 + 27B^2)$$

is nonzero. The condition $\Delta(E) \neq 0$ ensures that the cubic has no repeated root, so the curve is nonsingular.

At first sight, there is no obvious reason why the equation $a^p + b^p = c^p$ should have anything to do with a cubic curve in two variables. Yet this is exactly the kind of unexpected bridge that modern mathematics exploits. The theorem was solved not because mathematicians learned to manipulate $a^p + b^p = c^p$ more cleverly in isolation, but because they discovered that a hypothetical solution would create an elliptic curve with very special arithmetic properties.

6 The Frey Curve

Suppose, for contradiction, that there exists a nontrivial solution

$$a^p + b^p = c^p$$

with $p > 2$ prime and a, b, c pairwise coprime integers. Gerhard Frey suggested associating to such a solution the elliptic curve

$$E_{a,b,p} : y^2 = x(x - a^p)(x + b^p).$$

This is now called the *Frey curve*.

Why is this curve important? The answer is not that it directly solves Fermat's equation. Rather, it packages the arithmetic information of the alleged solution into a geometric object. The coefficients of the curve are built out of a^p and b^p, so any unusual arithmetic feature of the equation has a chance of reappearing as an unusual geometric or Galois-theoretic feature of the curve.

A remarkable aspect of the Frey curve is that it appears to be semistable and to have a very special conductor. Roughly speaking, the conductor measures the primes at which the curve has bad reduction and how badly it degenerates there. Frey's insight was that if a Fermat solution existed, then the associated elliptic curve would have such peculiar properties that it ought to violate the modularity expectations known at the time. This suggestion was not yet a proof, but it transformed the problem. Fermat's equation was no longer being attacked directly. It was being used to generate a test object in the theory of elliptic curves.

7 Elliptic Curves as Arithmetic Objects

To understand why the Frey curve matters, one must understand elliptic curves not merely as cubic equations but as arithmetic objects carrying multiple layers of structure.

First, elliptic curves have rational points. In fact, the set $E(\mathbb{Q})$ of rational points forms an abelian group under a geometrically defined addition law. This already makes elliptic curves far richer than a generic algebraic curve. The group law means that an elliptic curve simultaneously belongs to algebraic geometry, group theory, and arithmetic.

Second, for each prime ℓ, one can consider the ℓ-power torsion points of the curve. These torsion points are not just combinatorial decorations. They are acted on by the absolute Galois group

$$G_{\mathbb{Q}} = \mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}),$$

which encodes the arithmetic symmetries of all algebraic extensions of $\mathbb{Q}$. This action produces Galois representations, and those representations allow us to study the arithmetic of the curve via linear algebra over finite fields or p-adic rings.

Third, elliptic curves have L-functions and local reduction data at every prime. Thus they carry local-global information in a very precise way. One can ask how the curve reduces modulo different primes, how many points it has over finite fields, and how those point counts vary. These quantities eventually connect the curve to modular forms.

8 What a Modular Form Is

At this stage modular forms enter, and the term can sound mysterious if left unexplained. A modular form is, at one level, a holomorphic function on the upper half-plane

$$\mathfrak{H} = \{z \in \mathbb{C} : \mathrm{Im}(z) > 0\}$$

that transforms in a specific way under the action of a subgroup of $SL_2(\mathbb{Z})$. For a modular form of weight k, one requires a transformation law of the form

$$f\left(\frac{az+b}{cz+d}\right) = (cz+d)^k f(z)$$

for matrices

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

in the appropriate subgroup, together with suitable growth conditions at cusps.

This definition can seem analytic and remote from number theory, but the arithmetic content appears through the Fourier expansion

$$f(z) = \sum_{n=0}^{\infty} a_n q^n, \qquad q = e^{2\pi i z}.$$

The coefficients a_n often encode deep arithmetic information. In particular, certain normalized cusp forms of weight 2 produce Hecke eigenvalues that behave like point-count data of elliptic curves over finite fields.

That parallel is not superficial. It is the central bridge in the story. The modularity philosophy says that certain geometric objects, namely elliptic curves over $\mathbb{Q}$, should correspond to certain analytic objects, namely modular forms. Arithmetic and analysis are being linked through shared local data.

9 Modularity of Elliptic Curves

The Taniyama–Shimura–Weil conjecture, now part of the modularity theorem, asserts that elliptic curves over $\mathbb{Q}$ are modular. In a simplified form, this means that to an elliptic curve $E/\mathbb{Q}$ one can associate a modular form f such that the arithmetic data of E match the Hecke eigenvalues of f.

More concretely, for almost all primes q, if

$$a_q(E) = q + 1 - \#E(\mathbb{F}_q),$$

then there should exist a modular form

$$f(z) = \sum_{n=1}^{\infty} a_n q^n$$

such that

$$a_q(E) = a_q(f)$$

for those primes. Thus the number of points on the curve modulo q is controlled by the Fourier coefficients of a modular form. This is astonishing because it says that a geometric object defined by an algebraic equation and an analytic object defined on the upper half-plane share the same arithmetic fingerprints.

The reason this matters for Fermat is now visible. If one can show that every semistable elliptic curve is modular, and if the Frey curve is semistable, then the Frey curve should be modular. But if one can also prove that the Frey curve would have to be non-modular, then one gets a contradiction.

10 Ribet's Theorem and the Contradiction Strategy

Frey's original suggestion was turned into a rigorous theorem through work of Serre and Ribet. Serre formulated a precise modularity conjecture for mod p Galois representations, and Ribet proved a crucial level-lowering theorem that applied to the Frey curve.

The conceptual logic is as follows. Suppose the Fermat equation had a nontrivial solution. Then the Frey curve would exist. From this curve, one obtains a mod p Galois representation coming from its p-torsion. Ribet's theorem shows that if this representation were modular, then it would arise from a modular form of a level so small that no such form can exist. Therefore, the Frey curve would have to be non-modular.

This was an extraordinary reduction. Ribet did not prove Fermat's Last Theorem directly. Instead, he proved something structurally stronger:

a Fermat solution would imply the existence of a non-modular semistable elliptic curve.

From that point onward, Fermat's Last Theorem would follow if one could prove that all semistable elliptic curves over $\mathbb{Q}$ are modular.

So the equation

$$a^p + b^p = c^p$$

had effectively disappeared from center stage. The problem had been rephrased as a theorem about elliptic curves. This is one of the most beautiful examples in mathematics of solving a problem by embedding it into a richer theory.

11 Why Galois Representations Appear

The next major ingredient is Galois representation theory. This is the point where the proof usually feels most abstract, because the original equation seems to have been completely left behind. But there is a clear reason these objects appear.

The absolute Galois group $G_{\mathbb{Q}}$ is too complicated to study directly. One way to understand it is to examine how it acts on arithmetic objects. Elliptic curves provide such objects. If E is an elliptic curve and p is a prime, then the p-power torsion points of E form modules on which $G_{\mathbb{Q}}$ acts. Passing to inverse limits, one obtains the p-adic Tate module $T_p(E)$, and hence a representation

$$\rho_{E,p} : G_{\mathbb{Q}} \to GL_2(\mathbb{Z}_p).$$

Reducing modulo p, one obtains

$$\overline{\rho}_{E,p} : G_{\mathbb{Q}} \to GL_2(\mathbb{F}_p).$$

These representations encode deep information about the elliptic curve. For example, the trace of Frobenius at a prime $q \neq p$ is related to the quantity $a_q(E)$, which is also the number that appears in modular forms. In this way, Galois representations form the language in which elliptic curves and modular forms can be compared.

This is the crucial point: modularity is not only a statement about functions and curves. It is also a statement that certain Galois representations coming from elliptic curves coincide with those arising from modular forms.

12 The Strategy of Wiles

Andrew Wiles did not prove the full modularity theorem for all elliptic curves. What he proved, with the crucial additional work of Richard Taylor, was enough for Fermat's Last Theorem: the modularity of semistable elliptic curves over $\mathbb{Q}$.

The broad strategy was to begin with a residual representation

$$\overline{\rho} : G_{\mathbb{Q}} \to GL_2(\mathbb{F}_p),$$

which was already known or assumed to be modular, and then study its lifts to p-adic representations. The question became: if we deform the representation in controlled ways, can every suitable lift be shown to come from a modular form?

This is not a minor technical question. It is the heart of the proof. Wiles had to compare two worlds:

1. the world of all deformations of a fixed residual Galois representation, and
2. the world of Galois representations actually arising from modular forms.

If those two worlds could be shown to coincide, then any Galois representation satisfying the right deformation conditions would automatically be modular.

13 What Deformation Theory Means Here

The word "deformation" can sound geometric or physical, but here it has a specific algebraic meaning. Suppose one has a mod p representation $\overline{\rho}$. A deformation of $\overline{\rho}$ is, roughly, a lift of it to a representation over a larger coefficient ring, typically a complete local Noetherian p-adic ring, such that reducing the lift modulo the maximal ideal recovers the original representation.

The idea is to understand all such lifts that satisfy prescribed local conditions, such as ramification restrictions and determinant conditions. One can then ask whether there exists a universal ring R that parameterizes all these deformations. Mazur's deformation theory gives a framework in which such universal deformation rings often exist.

Thus R represents, in one object, the entire family of admissible Galois representations lifting $\overline{\rho}$. It is a ring of possibilities.

On the other hand, modular forms give rise to Hecke algebras. The Hecke operators act on spaces of modular forms, and their eigenvalues encode arithmetic information. One can complete the relevant Hecke algebra at a maximal ideal corresponding to $\overline{\rho}$, obtaining a local Hecke algebra T. This ring captures the modular side.

So the proof is set up as a comparison between two rings:

- R, the universal deformation ring, which parameterizes all admissible lifts, and
- T, the Hecke algebra, which parameterizes lifts known to be modular.

14 The Meaning of $R = T$

The famous equation

$$R = T$$

is not a slogan but the central structural statement of the proof. It means that the ring parameterizing all allowable deformations is isomorphic to the ring generated by Hecke operators coming from modular forms. In less compressed language, it says that the entire deformation space one wants to study is already exhausted by modular objects.

Why is this powerful? Because a priori, R could be larger than T. There could be admissible Galois representations that satisfy all the expected local conditions but do not come from modular forms. If that happened, one would not be able to conclude modularity. Proving $R = T$ rules out exactly that possibility.

So when people say "Wiles proved modularity by showing $R = T$," what they mean is this:

every deformation that ought to exist from the Galois-theoretic side already arises from the modular side.

This is a breathtaking statement because it identifies two objects that originate in very different mathematical worlds.

15 Why $R = T$ Is Hard

The difficulty of proving $R = T$ lies in the fact that neither ring is easy to control. The deformation ring R depends on subtle local-global arithmetic conditions. The Hecke algebra T depends on the structure of modular forms and congruences between them. To prove that they are isomorphic, one must show that they have matching size and structure.

Wiles' method involved studying tangent spaces, Selmer groups, and certain numerical criteria. A key part of the argument is to establish enough control over both sides so that a surjective map

$$R \twoheadrightarrow T$$

can be upgraded to an isomorphism. This requires a very delicate balance between commutative algebra and arithmetic geometry.

The Taylor–Wiles innovation introduced auxiliary primes and patching methods that allow the deformation problem to be modified in controlled ways so that the comparison becomes tractable.

These methods are technical, but their conceptual role is clear: they create enough flexibility to compare the Galois side and the modular side systematically.

16 How the Contradiction Finally Works

Now the pieces come together cleanly.

Assume there exists a nontrivial solution to Fermat's equation for some prime exponent $p > 2$. From this solution, construct the Frey curve. By the work of Frey, Serre, and Ribet, this curve would give rise to a non-modular semistable elliptic curve. But Wiles and Taylor–Wiles prove that every semistable elliptic curve over $\mathbb{Q}$ is modular. Therefore, the Frey curve must both be modular and not modular. This contradiction shows that the original Fermat solution cannot exist.

What is remarkable here is how indirect the proof is. The theorem is not proved by factoring $x^p + y^p$, nor by a clever manipulation of exponents, nor by an inequality. It is proved by showing that a hypothetical solution would force an impossible mismatch between geometry and analysis.

17 What This Reveals About Mathematics

Fermat's Last Theorem teaches several lessons about modern mathematics.

First, simplicity of statement does not imply simplicity of proof. A problem stated in elementary language may secretly belong to a much deeper structural framework.

Second, arithmetic is no longer isolated from geometry and analysis. The theorem was solved only because mathematicians learned how to move between these domains. Elliptic curves turned an equation into geometry; modular forms brought in complex analysis and automorphic symmetry; Galois representations converted arithmetic information into linear algebra; deformation theory and commutative algebra supplied the machinery needed to compare families of such objects.

Third, the proof exemplifies a central habit of advanced mathematics: one often solves a problem not by attacking it in its original form, but by embedding it into a richer category of objects where hidden structure becomes visible.

18 Conclusion

Fermat's Last Theorem is often remembered as a historical puzzle finally solved after more than three centuries. But from a mathematical point of view, its greatest significance may be the route by which it was solved. The theorem became approachable only after the original Diophantine equation was translated into the arithmetic of elliptic curves, then into the language of modular forms and Galois representations, and finally into a comparison between deformation rings and Hecke algebras.

The deepest point, in my view, is that the proof does not merely settle one equation. It reveals that arithmetic problems are governed by hidden symmetries and that those symmetries can often be detected only through structures that seem, at first, completely unrelated to the original problem. In that sense, Fermat's Last Theorem is not just a theorem about integers. It is a theorem about the unity of mathematics.

References

[1] T. M. Apostol, *Modular Functions and Dirichlet Series in Number Theory*, 2nd ed., Springer, 1990.

[2] G. Cornell, J. H. Silverman, and G. Stevens (eds.), *Modular Forms and Fermat's Last Theorem*, Springer-Verlag, 1997.

[3] F. Diamond and J. Shurman, *A First Course in Modular Forms*, Springer, 2005.

[4] G. Frey, Links between stable elliptic curves and certain Diophantine equations, *Annales Universitatis Saraviensis. Series Mathematicae*, 1(1), 1–40, 1986.

[5] Y. Hellegouarch, *Invitation to the Mathematics of Fermat-Wiles*, Academic Press, 2001.

[6] K. Ireland and M. Rosen, *A Classical Introduction to Modern Number Theory*, 2nd ed., Springer, 1990.

[7] N. Koblitz, *Introduction to Elliptic Curves and Modular Forms*, 2nd ed., Springer, 1993.

[8] S. Lang, *Introduction to Modular Forms*, Springer, 1995.

[9] T. Miyake, *Modular Forms*, Springer-Verlag, 1989.

[10] J. Neukirch, *Algebraic Number Theory*, Springer-Verlag, 1999.

[11] K. A. Ribet, On modular representations of $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ arising from modular forms, *Inventiones mathematicae*, 100(2), 431–476, 1990.

[12] J.-P. Serre, *A Course in Arithmetic*, Springer-Verlag, 1973.

[13] J.-P. Serre, Sur les représentations modulaires de degré 2 de $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$, *Duke Mathematical Journal*, 54(1), 179–230, 1987.

[14] J. H. Silverman, *The Arithmetic of Elliptic Curves*, 2nd ed., Springer, 2009.

[15] J. H. Silverman and J. Tate, *Rational Points on Elliptic Curves*, Springer, 1992.

[16] R. Taylor and A. Wiles, Ring-theoretic properties of certain Hecke algebras, *Annals of Mathematics*, 141(3), 553–572, 1995.

[17] A. Wiles, Modular elliptic curves and Fermat's Last Theorem, *Annals of Mathematics*, 141(3), 443–551, 1995.