

Bias Correction and Uncertainty Quantification for Small-Sample Ecosystem Condition Assessments

Anders Lorentzen Kolstad^{a,*}, Matthew Grainger^a, Marianne Evju^b

^aNorwegian Institute for Nature Research, Department of Terrestrial Ecology, Pb 5685 Torgarden, Trondheim, 7485

^bNorwegian Institute for Nature Research, NINA Oslo, Sognsveien 68, Oslo, NO-0855

Abstract

Ecosystem Condition Accounts (ECAs) require spatially representative indicators to accurately reflect ecosystem integrity, yet data constraints—particularly small sample sizes and spatial bias—often hinder robust assessments. While national-scale ECAs can mitigate these issues through modeling, local ECAs face unique challenges, as spatial extrapolation beyond the accounting area risks compromising local relevance. In this study, we address two critical gaps in local-scale ecosystem condition accounting: (1) correcting for spatial bias without external data and (2) uncertainty quantification for small samples. Using three Norwegian municipalities as test cases, we demonstrate a workflow that combines post-stratification (control bias via Homogenous Impact Areas) and empirical Bayes variance smoothing (to stabilize uncertainty estimates for 1-inflated beta distributions). Point estimates are derived through direct, area-weighted averaging within strata, ensuring local fidelity. Our approach enables local ECAs to leverage imperfect datasets while maintaining transparency and local relevance. We provide a scalable solution for small-area ecosystem assessments where traditional modeling is infeasible. This work advances the practical implementation of ecosystem accounting under real-world data constraints.

Keywords: alien species, disturbance, ecosystem accounting, ecosystem condition, horizontal aggregation, indicator, mire, peatlands, SEEA EA, small area estimation, wetlands

1. Introduction

Ecosystem condition accounting (ECA) is the structured compiling and communication of data on the status, trends and qualities of ecosystems (Comte et al., 2022; Edens et al., 2022). Its purpose is to make it easier to account for nature in policy by making the environmental costs of certain policies and practices visible to decision-makers.

The System of Environmental-Economic Accounting - Ecosystem Accounting (SEEA EA) is a statistical standard for ecosystem accounting, including ecosystem condition accounting, that was developed by the UN and adopted by the UN Statistical Commission in 2021 (Nations, 2024). This framework provides a set of rules, principles and best practices for compiling ecosystem accounts, primarily at a national level (United Nations, 2022). At its core is the idea that data on ecosystem condition can be summarised in the form of metrics (indicators and variables) presented in generalised tables that describe the condition of a larger ecosystem accounting area (EAA).

*Corresponding author

Email addresses: anders.kolstad@nina.no (Anders Lorentzen Kolstad), matthew.grainger@nina.no (Matthew Grainger), marianne.evju@nina.no (Marianne Evju)

With intentional similarities to national accounts, which deal with the economic activity at the scale of nations, SEEA EA accounts have mainly been developed for and used at national scales. However, there is a surge of examples of more local accounts appearing, motivated at least partly by the need to match the scale of the information to the scale of decision-making processes (REF!). Compared to accounts with large EAAs, local accounts face increasing challenges with data resolution and availability.

An important requirement of SEEA EA is that condition metrics must give an unbiased representation of the condition for a given area (table 1 in Czucz et al., 2021; see also Nations, 2024, §2.87). These metrics should ideally be recorded (or modeled) uniquely for all contiguous spaces (patches) of a specific ecosystem type (a common definition of an *ecosystem asset*), enabling the calculation of area-weighted means for the EAA (Nations, 2024, §5.54). While unbiased estimation is essential, this requirement severely limits the types of usable data. Local ECAs are particularly constrained by data availability, leading to pragmatic (and potentially unrepresentative) selections of variables and indicators. Being able to make use of spatially biased data would greatly alleviate data shortage problems in local ECAs.

The issue of representativeness extends beyond spatial biases to include thematic biases—such as the under-representation of key ecosystem components like insects or soil biota. Also, having data from only one or a subset of nature types inside what is defined as the ecosystem in the assessment, is a typical thematic bias in ecosystem accounting. However, this paper focuses on spatial (geographic and environmental) bias.

Wall-to-wall raw data, such as satellite imagery, is spatially representative through the property of being spatially exhaustive, although grain size also plays a role [cf. the ecological fallacy; Robinson (2009)]. However, many critical ecosystem characteristics - such as species composition, soil properties, or fine-scale habitat features - cannot be derived remotely and still require field surveys. These surveys, however, often suffer from sparse or biased sampling, leaving gaps in ecosystem asset coverage. Field data are especially problematic if sampling intensity correlates with anthropogenic pressures (and thus ecosystem condition) (Aubry et al., 2024). A poor solution, which doesn't attempt to control for sampling biases, is to ignore unsampled areas by directly estimating values for the entire EAA via central tendencies (e.g. means). Other options include imputing values to unsampled areas using some type of model or spatial generalisation. United Nations (2022) list six relevant modeling techniques for ecosystem condition metrics that can also be used for projecting metric values to ecosystem assets without field data, and simultaneously control for sampling biases:

1. Look-up table
2. Spatial interpolation (e.g. inverse distance weighting)
3. Geostatistical models (e.g. kriging)
4. Statistical models (e.g. regression)
5. Dynamic system models (many types of process-based models)
6. Machine learning (e.g. random forest)

Depending on the data that goes into them, models can make very accurate estimates, especially when the areas are large (e.g. regions or nations), but their reliability diminishes for smaller areas (e.g., municipalities) due to data scarcity and model uncertainty. Small area estimation is a branch in statistics aimed at finding ways to overcome this issue by “borrowing strength” from related areas (Ghosh and Rao, 1994; Guldin, 2021). Ecosystem accounts should ideally reflect conditions only inside the accounting area and are arguable more useful when reflecting local management and land use policies, unaffected by the actions of neighboring or ecologically similar administrative units (i.e. direct estimation). There might therefore be a limit to how much data one should “borrow” across ecosystem assets or EAAs.

Look-up tables, the simplest approach, assign the same value across categorical strata (e.g. ecosystem type within a region), often with no uncertainty estimate (although this is quite possible to include). Values may derive from expert opinion or aggregated data (e.g. stratum means). If strata align with relevant gradients in pressures or environmental conditions, this method can mitigate sampling bias via post-stratification. The level of spatial or environmental extrapolation is subject to the size of the strata, with fine-grained

typologies leading to less extrapolation. Extrapolating non-continuous spatial data (e.g. points) over a surface is a type of spatial generalisation. The challenge of spatial generalisation is not unique to ecosystem condition accounts, and it is much discussed in the ecosystem service accounting community of practice via the related term *value transfer* (United Nations, 2022; NCAVES and MAIA; Grammatikopoulou et al., 2023; Barton, 2023). Its key advantages are simplicity and the possibility to rely solely on local data. A major drawback is the high data requirements, especially for fine-grained strata.

After extrapolating data, the spatial grain of the data (and thus also the information) is essentially still the same. However, a critical step in ecosystem accounting is upscaling—synthesising fine-grained data to characterise larger areas, effectively reducing grain size or increasing support (Bierkens et al., 2002) (Kolstad et al. in prep). This is typically achieved by aggregation (e.g. arithmetic means), sampling (Brus and de Gruijter, 1997) or resampling techniques (Jakobsson et al., 2021). For simplicity we also refer to this as aggregation, as the end result is the same - a synthesis of information. Upscaling is required to make the information available for decision makers or monitoring frameworks at the appropriate spatial scale (King et al., 2024).

Spatial aggregation allows for the quantification of spatial variance in the estimate. In addition, there can be measurement or modelling errors that could potentially also be propagated and reported alongside the upscaled estimate (e.g. Heris et al., 2021). Currently, SEEA EA lacks guidelines for estimating or disclosing any type of numerical uncertainty. This could potentially undermine the credibility and finally the uptake of knowledge from ecosystem accounting (Bagstad et al., 2021, Venter et al. (2024)). Local ECAs face a unique side of this problem because sample sizes are frequently so small that traditional arithmetic estimates of variance are unreliable. In addition, calculating variances is even more data demanding when metrics are truncated, such as for indicators bound between zero and one. In such cases, a small sample size can by chance contain only or predominantly ones (or zeros), resulting in an artificially small variance which will then paradoxically increase as sample sizes increase due to the inclusion of more data points from the tail of the distribution.

In this study we present two techniques for direct estimation of local ecosystem condition indicators using scarce and spatially biased data. Firstly, we demonstrate a stratified aggregation technique, using homogeneous impact areas, to reduce the negative consequences of spatial and environmental bias in the data. With this we get direct indicator point estimates using area weighting and local data only. Second, we use an empirical Bayes variance smoothing approach to predict indicator uncertainties where we borrow strength from a larger national dataset. This allows us to get a realistic picture of indicator uncertainty even sample sizes are very low. We demonstrate these techniques with three indicators from the same spatially biased nature type mapping data set from Norwegian mires, comparing results across three municipalities in Norway to illustrate its potential for local applications.

2. Material and Methods

Mires cover about 10-15% of mainland Norway and play a disproportional role in climate mitigation due to their large carbon stores. To delineate open mire ecosystems we used a remote-sensing derived map which estimates 12.7% of the area in southern Norway is mire (Bakkestuen et al., 2023). There has not been a national assessment of the ecosystem condition of mires in Norway, but the authors recently contributed to a report which presented several new indicators that can be used in future national assessments for this ecosystem (Nybø et al., 2023; see also Kolstad et al., 2023). The current study builds on the work in that report.

2.1. Data source and survey design

We used a spatial dataset of nature types in Norway (Norwegian Environment Agency, 2024), surveyed in the field following a standardised protocol (Norwegian Environment Agency, 2022). The dataset comprises over 140,000 polygons (as of 2023), each scored on several variables describing ecological condition and qualities. The data collection is spatially biased, typically towards areas with high human impact or expected

impact, reflecting their primary use in land-use decisions. In addition, the sampling protocol introduces thematic and sizes biases. For mires, this implies an elevated focus on calcareous and medium to large mires. The survey maps the following mire types which we use in our analyses:

- Southern bogs (ombrotrophic mires) $> 2500 \text{ m}^2$
- Northern bogs $> 10.000 \text{ m}^2$
- All semi-natural mires $> 1000 \text{ m}^2$
- Calcareous southern fens (minerotrophic mires) $> 500 \text{ m}^2$
- Calcareous northern fens $> 1000 \text{ m}^2$

Here, *southern* refers to boreonemoral and southern boreal zones, and *northern* refers to mid-boreal, north boreal, and alpine zones (Moen, 1998). Northern fens require larger size and higher calcareousness to be surveyed. Data from 2018 to 2023 were included.

We assume the surveyed mires are representative for the broader mire ecosystem inside the surveyed areas, despite undersampling of smaller and less calcareous mires. We posit that our selected variables (see below) are relatively unaffected by this bias, with a possible exception for alien plant occurrences, which are probably less common on bogs, but the variable is recorded only in fens. The other variables are recorded in all delineated mires.

2.2. Indicator development

Six raw variables (Table 1, variables 1–6) were normalised and turned into three condition indicators (sensu Nations, 2024):

- *Alien species* (7FA)
- Anthropogenic disturbance to soil and vegetation (*ADSV*; combined from variables 7SE, PRSL, 7TK and PRTK)
- *Trenching* (7GR-GI).

Normalisation followed a piece-wise linear transformation with three reference levels [Czúcz et al.; @jakobsson2020]

- X_{100} : Best possible condition (minimal human influence)
- X_0 : Worst possible condition
- X_{60} : defining the breaking point between good and poor condition.

The reference levels make up a numerical representation of the reference condition. We define the reference condition as one where ecosystems are subject to little or no human influence, with a climate as in the period 1961-1990 and a native species pools similar as today (Jakobsson et al., 2021). For details about the normalisation process, see Appendix A.




Figure 1: Map of the three focal municipalities. For each municipality the maps in the top row show ocean in blue and non-ocean in green. The survey coverage maps are in grey, and the mapped mires are in red, with polygon borders made extra thick to make them visible, but then also exaggerating their size. The bottom row shows the delineation for homogeneous impact areas which is an ordinal gradient from 0–3 with increasing presence of human infrastructure.

2.3. Homogenous Impact Areas

To alleviate issues of spatial bias we used spatial information about human pressures, presumably a major structuring variable, to post-stratify the data. The Norwegian infrastructure index for the year 2022 is a continuous variable representing the frequency of different infrastructure types inside 500 m radius circles around each 100×100 m pixel (Erikstad et al., 2023a). We categorised this data into four non-overlapping Homogeneous Impact Areas (HIAs; Table 1), with boundaries chosen to balance gradient resolution, data stratum area, and computational limitations (Table S1). Data was aggregated to 1×1 km pixels to ease computations. HIAs align conceptually with the concept of Homogeneous Ecosystem Areas, *sensu* Vallecillo et al. (2022). The nature type polygons with the indicator values were intersected with a vectorised HIA map and the relationship between the HIA classes and indicator values was examined visually across the entire data set.

2.3.1. Municipal-level analyses and spatial aggregation

Three municipalities were selected test the indicators, varying in mire area, survey coverage, and infrastructure prevalence (Table 2, Figure 1, S3, S4). For each HIA — municipality (strata) combination we calculated the area weighted arithmetic mean of the indicator and used a look-up table approach to transfer these values to all ecosystem assets in the same strata. Ignoring strata with no data, we then took another area-weighted mean across all ecosystem assets inside the municipality to get the point estimate.

The three indicators were all highly left skewed with strong 1-inflation. It was therefore essential to describe the indicator uncertainty using non-symmetrical errors. It also makes sense for a bound indicator not to have errors exceed the bounds. The sample size for each stratum, excluding those with no data, ranged between 1 and 145, yet we wanted a generic approach that allowed us to include reliable uncertainty estimates for all cases. Our solution was to use the entire national dataset, divided into the four HIAs, to describe the indicator distributions as zero and one inflated beta distributions. We recorded the frequency of zeros and ones in the data as a measure of zero- and one inflation, respectively. For values between, but not including, zero and one, we used maximum likelihood estimation to get the two shape parameters for a beta distribution. We did the same for the local strata, although the estimation of the shape parameters requires a certain sample size and dispersion of data values to converge, making it impossible to describe

the samples for some strata as a distribution. Using an empirical Bayes variance smoothing technique for small area estimation we combined prior information about the shape of the national-level data with the local strata data using a prior weight of 20, meaning the national data informed about the shape of the local indicator distribution with similar weight as 20 local data points. The prior weight was chosen after some trial and error to find a suitable weight that balanced local autonomy with realistic distributions. With less 1-inflation, a smaller prior weight could have been used. Still, the prior weight was low enough that all point estimates (calculated independently from the predicted distributions) fell inside the 95% credible intervals. The predicted distributions were transferred over to the ecosystem assets of the corresponding strata, and from there, values were sampled proportional to asset area to obtain a probability distribution for the EAA (Figure 2 - C).

All the data preparation and analyses were done in RStudio (Posit team, 2024) and R version 4.3.0 (R Core Team, 2023), relying heavily on the packages tidyverse (Wickham et al., 2019), sf (Pebesma, 2018), terra (Hijmans, 2023) and stars (Pebesma and Bivand, 2023).

Table 1: Variables used in this study

id	Variable code	Variable name	Measurement unit	Description	Reference
1	7FA	Prevalence of alien species	Unit-less, ordinal, non-linear scale from 1 (no alien species) to 7 (only alien species)	The fraction of the species composition made up from alien species	Halvorsen and Bratli (2019)
2	7SE	Human caused abrasion or abrasion-caused erosion	Unit-less, ordinal, non-linear scale from 1 to 4.	Measures the frequency of imagined 4 m ² quadrats laid over the area that has some sign of abrasion	Halvorsen and Bratli (2019)
3	PRSL	<i>as above</i>	Unit-less, ordinal, non-linear scale from 0 to 7.	Same as 7SE, but recorded at a higher resolution	Norwegian Environment Agency (2022)
4	7TK	Tracks from large vehicles	Unit-less, ordinal, non-linear scale from 1 to 4.	Measures the frequency of imagined 100 m ² quadrats laid over the area that has some signs of vehicle tracks	Halvorsen and Bratli (2019)
5	PRTK	<i>as above</i>	Unit-less, ordinal, non-linear scale from 0 to 7.	Same as 7TK, but recorded at a higher resolution	Norwegian Environment Agency (2022)
6	7GR-GI	Trenching intensity	Unit-less, ordinal scale from 1 to 5	Describes the effect that drainage ditches is estimate to have on the species composition and environmental variables ones the system reached its new equilibrium	Norwegian Environment Agency (2022)
7	Infra-structure Index	Infrastructure Index	Unit-less linear scale from 0 to 13.2	Unit-less index ranging from from 0 to 13.2	Erikstad et al. (2023b)
8	HIA	Homogeneous Impact Area	Ordinal, non-linear scale from 1 to 4	A categorical representation of the infrastructure index	this paper

Table 2: Information for the three target municipalities in Norway

Municipality	Total terrestrial area (km ²)	% of terrestrial area surveyed	% open mires in relation to total terrestrial area	Total mire area (km ²)	%of mire area inside survey area	Number of mire polygons in survey	Mean Infrastructure Index value
Nordre Follo	203	40	0.3	0.6	18.6	6	1.8
Gran	756	21	2.7	20.2	0.5	56	1.2
Nord-Aurdal	906	26	11.1	100.7	18.1	236	1

3. Results

3.1. Indicator validity and trends in Norway

The three indicators generally showed a worsening of condition with increasing presence of human infrastructure when we looked at data from all of Norway with 9026 individual mires (Figure 3). This relationship was much stronger for the indicator *Trenching*, and weakest for the indicator *Alien species*. *Alien species* also had the highest indicator values overall, with most mires having no alien species recorded. For *Trenching*, 57% of mires in HIA-3 had some trenches, whereas in HIA-0 this number was 10%.

3.2. The three focal municipalities

The indicator values were comparable across HISs in the three municipalities, although there were some differences, as for the *Trenching* indicator where in Gran, HIA-2 had worse condition than HIA-1 (Figure 4). Conversely, the *ADSV* indicator in Gran showed worse condition in HIA-1 compared to HIA-2, but the predicted indicator values had a wide credible interval (horizontal bars in Figure 4). Nordre-Follo had considerably fewer data points compared to the other two municipalities and the 95% credible intervals are therefore mainly reflecting the indicator distribution nationally.

None of the municipalities had survey data from all HIAs. Therefore, when we transferred the point estimates and the predicted indicator values for each HIA (Figure 4) over to the ecosystem delineation map (Figure 5), some mire polygons in the ecosystem delineation map did not get assigned any data, and were therefore not included (i.e. did not have any influence on) the following aggregation to ecosystem accounting areas (EAA) (Figure 2, E).

At the municipality level (i.e. the EAA level) the indicators generally showed good ecological condition (i.e. values >0.6) and point estimates ranged between 0.79 and 0.98, with the exception of *Trenching* in Nordre Follo which had 0.24 (Figure 6; Table S2). Nordre Follo also had the highest (the best) indicator values for one of the indicators (*ADSV*); probably reflecting stochasticity due to low sample size.

4. Discussion

This paper presents a generalizable workflow to address two critical challenges in local ecosystem condition assessments (ECAs): (1) mitigating spatial and environmental bias in scarce field data, and (2) quantifying realistic uncertainties for small-sample indicators (Figure 2). Our approach combines direct estimation (to preserve local representativeness) with empirical Bayes variance smoothing (to stabilize uncertainty estimates), offering a pragmatic solution for municipalities where data limitations hinder robust assessments. Below, we discuss the strengths and limitations of our two core innovations: post-stratification using Homogeneous Impact Areas (HIAs) and variance smoothing for small-area statistics. The two approaches can also be used independently.

4.1. Post-stratification using HIAs

Local ECAs often rely on opportunistic datasets that lack spatial representativity. Our post-stratification method addresses this by partitioning data into HIAs—homogeneous impact areas (e.g., infrastructure prevalence)—ensuring indicators are estimated only within comparable contexts. This minimizes extrapolation bias while retaining interpretability. A similar concept called homogeneous ecosystem areas was introduced by Vallecillo et al. (2022) and refer to parts of an ecosystem type that can be allowed to share common reference levels. HIAs on the other hand, refer to areas that can be allowed to share common indicator values.

The effectiveness of HIAs depends on their alignment with indicator dynamics. For instance, infrastructure prevalence strongly predicted trenching impacts in mires but weakly explained alien species occurrence (Figure 3). For the latter, other stratification variables could have been explored, e.g. latitude, as there are generally more alien species in the south compared to the north in Norway. Even though suboptimal HIAs reduce sample size per stratum, they still mitigate geographic bias by limiting inference to relatively




Figure 2: Simplified schematic showing the workflow (Panels A-E) for spatial generalisation of a spatially biased ecosystem condition indicator. Panel A shows in red the data coverage of a patchy and spatially biased indicator. Surveyed areas are grey and unsurveyed areas are green. Together they make up the ecosystem accounting area (EEA). Panel B shows the location of four homogeneous impact areas (HIAs) inside the EEA, where the prevalence of human infrastructure increases from 0 to 3. The indicator values in Panel A are used, in combination with the HIA map in Panel B, to calculate the area-weighted point estimates shown as grey circles in Panel C. The errors surrounding the point estimate are calculated independently from the point estimate via an empirical Bayes approach as predicted distributions (shown in Panel C as 95% credible intervals (CI)) informed with prior knowledge about the shape of the distribution of indicator values (a zero- and one-inflated beta distribution in this case) for a larger area, i.e. outside the EEA. Note that because in this example there were no indicator data for HIA-3, indicator values are only aggregated for HIAs 0-2. In Panel D the point estimates and sampled distributions are transferred to individual polygons in an ecosystem delineation map, for each HIA separately. The colours indicate different point estimates (highlighted in the inset). Panel E shows area-weighted indicator values aggregated to the EAA level. Blue vertical bars show the distribution of real samples values (samples here along an ordinal 5-step scale). The orange density curve shows the predicted indicator values informed both by local samples from inside the EAA and from a larger national dataset (indicator weight was 20, meaning the national data had the same influence on the predicted distribution as 20 local samples). The area-weighted mean and 95% CI are shown at the bottom as a thick vertical bar and thin horizontal line, respectively. The mean is informed by the local data only (blue bars) and the 95% CI is a summary of orange distribution.




Figure 3: Proportion of localities with different indicator values for three ecosystem condition indicators along an ordinal gradient of increasing infrastructure densities (Homogeneous Impact Areas 0–3). The data is from a national nature type survey in Norway.




Figure 4: Indicator values (circles = mean; bars = 95% credible intervals) for three mire ecosystem condition indicators (rows) in three Norwegian municipalities (columns). The indicator values are calculated uniquely for each Homogeneous Impact Area (HIA) in each municipality. The numbers to the left of each bar is the sample size, i.e. the number of surveyed mires.




Figure 5: An indicator for mire trenching shown for Nord-Aurdal municipality. Individual mire polygons are coloured by the mean indicator values for the homogenous impact area where it lies. Colours are chosen to best reflect categorical differences and exaggerates the absolute difference between areas. The inset it just a visual aid. Coordinate reference system is EPSG 25832. Axis are in meters.




Figure 6: Distributions for three ecosystem condition indicators in three Norwegian municipalities. Blue vertical bars show the distribution of real samples values. Orange density curves show the predicted indicator values informed both by local samples from inside the EAA and from a larger national dataset. The area-weighted mean and 95% credible intervals are shown at the bottom as a thick vertical bar and thin horizontal line, respectively. The mean is informed by the local data only (blue bars) and the 95% CI is a summary of orange distribution.

spatially cohesive areas. We see little risk with using HIA use even when relationships are weak, but not accounting for other, potentially more relevant predictors of indicator values, will leave the indicator as spatially biased as the original variable.

HIAs are particularly valuable for local ECAs, where small samples preclude complex geostatistical methods (e.g., kriging). Such methods are also more complicated and therefore at a compromise with the ideal for simplicity (Czúcz et al. (2021)). HIAs also provide actionable insights by revealing spatial patterns in degradation (Figure 4). However, trade-offs exist between HIA granularity and data availability; our pilot municipalities had insufficient data for some strata, suggesting broader definitions or imputation could be explored. Alternatively one can chose to impute values *post hoc* instead of simply dropping strata with no data.

4.2. Uncertainty quantification for small samples

An indicator estimate is of little value unless it is accompanied by a measure of uncertainty. However, there is a general lack of guidelines for estimation and disclosure of uncertainties in the SEEA EA statistical standard, and indicator values are often reported simply as opening and closing values. Traditional variance estimates fail for truncated or skewed indicators, where limited sampling can artificially inflate precision. Our empirical Bayes solution circumvents this by borrowing strength from a national dataset to model uncertainty distributions without affecting the point estimates which are estimated directly using only data from inside the EAA to ensure indicators reflect local condition.

Our approach enables uncertainty reporting even for tiny samples (e.g., 1–3 data points), which would otherwise be discarded. While it cannot eliminate the inherent stochasticity of sparse data, it contextualizes estimates within plausible value ranges, supporting decision-making. As sample sizes grow, local data dominate the uncertainty estimates, making the method adaptable across scales.

However, our results point to where the data shortages are most pressing (HIAs with little or no data), and this information could be used to guide future data collection. For example, Gran and Nord-Aurdal have not recorded any field data from their most urban areas (HIA 3; Figure 4) and could make these areas a priority in upcoming field campaigns in order to get more complete spatial indicator coverage.

4.3. Indicator values in pilot municipalities

The values for our three indicators across the three municipalities showed in general a good condition in the mire ecosystems (Figure 6). An exception was *Trenching* in Nordre Follo (indicator: 0.24, $n = 4$) which was well below the threshold for good ecosystem condition (0.6). The predicted indicator distribution is higher, suggesting the value might rise with additional sampling, but the direct estimate—uninfluenced by national data—reflects local realities. Nordre Follo has surveyed more of its total land area compared to the other two municipalities, but has less mire area to begin with (Table 2), which is the reason for this low sample size. Nordre Follo’s limited mire area underscores the value of our small-area methods.

Mire condition in Norway has not been assessed at a national scale. Still, we know that they are subject to multiple threats, especially land conversion into agriculture, forestry, or development, and we see this reflected in the Norwegian Red List for Nature types where many ombrotrophic bog types, and rich southern minerotrophic fens are listed (Centre, 2018). Land conversion of mires is often initiated with trenching to lower the water table but note that our *Trenching* indicator does not document the effect of trenching on old mires that are today transitioned into another ecosystem. This is in line with the UN standard for ecosystem accounting, where these effects should be captured by the extent accounts (United Nations, 2021). Therefore, the negative consequences of trenching on all Norwegian mires, old and current, must be higher than what our indicator is made to show, most likely considerably higher.

Alien species risk is not considered a threat to any of the Norwegian mire types, and our results confirm this, at least for the present (Figure 6). Motorised vehicle traffic and other forms of human transportation are also threats listed for some mire types in the red list assessment, and we see some reduction in the ADSV indicator, most so in Gran and in areas closer to infrastructure (Figure 4). Still, the indicator values are consistently above the threshold value for good ecological condition (Figure 6).

5. Conclusion

By integrating post-stratification and empirical Bayes variance smoothing, we enable local ECAs to leverage imperfect datasets while maintaining transparency and local relevance via direct estimation. Our methods address fundamental challenges in ecosystem accounting: spatial bias correction and uncertainty quantification for small samples. As SEEA EA matures, we urge greater emphasis on uncertainty reporting and robust aggregation techniques. We have demonstrated approaches that are new or relatively new to the ECA field of practice but well established in other fields of statistics. This work demonstrates that even sparse data can yield actionable insights when paired with statistically principled approaches.

CRediT authorship contribution statement

Anders L. Kolstad: Conceptualization, Methodology, Software, Validation, Formal analysis, Data Curation, Writing - Original Draft, Visualization, Project administration, Funding acquisition. **Matthew Grainger:** Validation, Writing - Review & Editing. **Marianne Evju:** Conceptualization, Writing - Review & Editing.

Declaration of Competing Interest

We have no competing interests to declare.

Acknowledgements

We wish to thank Kwaku Peprah Adjei for statistical help, and Hanno Sandvik for code review. We acknowledge partial funding from the Research Council of Norway project EcoGaps (contract #320042).

Data availability

This manuscript is written in Quarto, and the source file (Appendix B) also contains code for all the analyses underlying this study, including data exploration and cleaning. For a rendered version of the source file, with all code and calculation visible, see Appendix C. The source files and the data are also located on GitHub (<https://github.com/anders-kolstad/HIAs>) with an archived stable version on Zenodo (doi:10.5281/zenodo.11504703). The exception is the large nature type survey data, which was downloaded locally, but which is freely available (see [Norwegian Environment Agency \(2024\)](#)).

References

- Aubry, P., Francesiaz, C., Guillemain, M., 2024. On the impact of preferential sampling on ecological status and trend assessment. *Ecological Modelling* 492, 110707. URL: <https://www.sciencedirect.com/science/article/pii/S0304380024000954>, doi:10.1016/j.ecolmodel.2024.110707.
- Bagstad, K.J., Ingram, J.C., Shapiro, C.D., La Notte, A., Maes, J., Vallecillo, S., Casey, C.F., Glynn, P.D., Heris, M.P., Johnson, J.A., Lauer, C., Matuszak, J., Oleson, K.L.L., Posner, S.M., Rhodes, C., Voigt, B., 2021. Lessons learned from development of natural capital accounts in the united states and european union. *Ecosystem Services* 52, 101359. URL: <https://www.sciencedirect.com/science/article/pii/S2212041621001170>, doi:10.1016/j.ecoser.2021.101359.
- Bakkestuen, V., Venter, Z., Ganerød, A.J., Framstad, E., 2023. Delineation of Wetland Areas in South Norway from Sentinel-2 Imagery and LiDAR Using TensorFlow, U-Net, and Google Earth Engine. *Remote Sensing* 15, 1203. URL: <https://www.mdpi.com/2072-4292/15/5/1203>, doi:10.3390/rs15051203. number: 5 Publisher: Multidisciplinary Digital Publishing Institute.
- Barton, D.N., 2023. Value ‘generalisation’ in ecosystem accounting - using bayesian networks to infer the asset value of regulating services for urban trees in oslo. *One Ecosystem* 8, e85021. URL: <https://oneecosystem.pensoft.net/article/85021/>, doi:10.3897/oneeco.8.e85021. publisher: Pensoft Publishers.
- Bierkens, M.F., Finke, P.A., de Willigen, P., 2002. Choosing appropriate upscaling and downscaling methods for environmental research. *International Association of Hydrological Sciences, Publication* , 405–409.
- Brus, D.J., de Gruijter, J.J., 1997. Random sampling or geostatistical modelling? choosing between design-based and model-based sampling strategies for soil (with discussion). *Geoderma* 80, 1–44. URL: <https://www.sciencedirect.com/science/article/pii/S0016706197000724>, doi:10.1016/S0016-7061(97)00072-4.
- Centre, N.B.I., 2018. Norsk rødliste for naturtyper 2018. URL: <https://www.artsdatabanken.no/rodlisteformaturtyper>.
- Comte, A., Sylvie Campagne, C., Lange, S., Bruzón, A., Hein, L., Santos-Martín, F., Levrel, H., 2022. Ecosystem accounting: Past scientific developments and future challenges. *Ecosystem Services* 58, 101486. URL: <https://www.sciencedirect.com/science/article/pii/S2212041622000821>, doi:10.1016/j.ecoser.2022.101486.
- Czúcz, B., Keith, H., Maes, J., Driver, A., Jackson, B., Nicholson, E., Kiss, M., Obst, C., 2021. Selection criteria for ecosystem condition indicators. *Ecological Indicators* 133, 108376. URL: <https://linkinghub.elsevier.com/retrieve/pii/S1470160X21010414>, doi:10.1016/j.ecolind.2021.108376. publisher: Elsevier B.V.
- Czúcz, B., Rusch, G., Sandvik, H., Kolstad, A., Lappalainen, M., Jernberg, S., Korpinen, S., Santos-Martin, F., Rendón, P., Vallecillo, S., Lange, S., Tanács, E., . A general framework for defining reference levels for ecosystem condition. a working paper for selina task 3.3 (v1.3). .
- Edens, B., Maes, J., Hein, L., Obst, C., Siikamaki, J., Schenau, S., Javorsek, M., Chow, J., Chan, J.Y., Steurer, A., Alfieri, A., 2022. Establishing the sea ecosystem accounting as a global standard. *Ecosystem Services* 54, 101413. URL: <https://www.sciencedirect.com/science/article/pii/S2212041622000092>, doi:10.1016/j.ecoser.2022.101413.
- Erikstad, L., Simensen, T., Bakkestuen, V., Halvorsen, R., 2023a. Index measuring land use intensity—a gradient-based approach. *Geomatics* 3, 188–204. URL: <https://www.mdpi.com/2673-7418/3/1/10>, doi:10.3390/geomatics3010010. number: 1 Publisher: Multidisciplinary Digital Publishing Institute.
- Erikstad, L., Simensen, T., Bakkestuen, V., Halvorsen, R., 2023b. Index Measuring Land Use Intensity—A Gradient-Based Approach. *Geomatics* 3, 188–204. URL: <https://www.mdpi.com/2673-7418/3/1/10>, doi:10.3390/geomatics3010010. number: 1 Publisher: Multidisciplinary Digital Publishing Institute.
- Ghosh, M., Rao, J.N.K., 1994. Small area estimation: An appraisal. *Statistical Science* 9, 55–76. URL: <https://projecteuclid.org/journals/statistical-science/volume-9/issue-1/Small-Area-Estimation-An-Appraisal/10.1214/ss/1177010647.full>, doi:10.1214/ss/1177010647. publisher: Institute of Mathematical Statistics.
- Grammatikopoulou, I., Badura, T., Johnston, R.J., Barton, D.N., Ferrini, S., Schaafsma, M., La Notte, A., 2023. Value transfer in ecosystem accounting applications. *Journal of Environmental Management* 326, 116784. URL: <https://www.sciencedirect.com/science/article/pii/S030147972202357X>, doi:10.1016/j.jenvman.2022.116784.
- Guldin, R.W., 2021. A systematic review of small domain estimation research in forestry during the twenty-first century from outside the united states. *Frontiers in Forests and Global Change* 4. URL: <https://www.frontiersin.org/journals/forests-and-global-change/articles/10.3389/ffgc.2021.695929/full>, doi:10.3389/ffgc.2021.695929. publisher: Frontiers.
- Halvorsen, R., Bratli, H., 2019. Dokumentasjon av NiN versjon 2.2 tilrettelagt for praktisk naturkartlegging: utvalgte variabler fra beskrivelsessystemet. – Natur i Norge, Artikkel 11 (versjon 2.2.0). Technical Report. Trondheim.
- Heris, M., Bagstad, K.J., Rhodes, C., Troy, A., Middel, A., Hopkins, K.G., Matuszak, J., 2021. Piloting urban ecosystem accounting for the united states. *Ecosystem Services* 48. doi:10.1016/j.ecoser.2020.101226. publisher: Elsevier B.V.
- Hijmans, R.J., 2023. terra: Spatial Data Analysis. URL: <https://CRAN.R-project.org/package=terra>. r package version 1.7-46.
- Jakobsson, S., Evju, M., Framstad, E., Imbert, A., Lyngstad, A., Sickel, H., Sverdrup-Thygeson, A., Töpper, J.P., Vandvik, V., Velle, L.G., Aarrestad, P.A., Nybø, S., 2021. Introducing the index-based ecological condition assessment framework (ibeca). *Ecological Indicators* 124, 107252. URL: <https://linkinghub.elsevier.com/retrieve/pii/S1470160X20311912>, doi:10.1016/j.ecolind.2020.107252. publisher: Elsevier B.V.
- King, S., Agra, R., Zolyomi, A., Keith, H., Nicholson, E., de Lamo, X., Portela, R., Obst, C., Alam, M., Honzák, M., Valbuena, R., Nunes, P.A.L.D., Santos-Martin, F., Equihua, M., Pérez-Maqueo, O., Javorsek, M., Alfieri, A., Brown, C., 2024. Using the system of environmental-economic accounting ecosystem accounting for policy: A case study on forest ecosystems. *Environmental Science & Policy* 152, 103653. URL: <https://www.sciencedirect.com/science/article/pii/S1462901123003027>, doi:10.1016/j.envsci.2023.103653.
- Kolstad, A.L., Grainger, M., Sandvik, H., Venter, Z., Speed, J.D., Nater, C., Evju, M., Åstrøm, J., 2023. Indicators for ecosystem condition in norway. URL: <https://ninanor.github.io/ecosystemCondition/index.html>.

- Moen, A., 1998. Vegetation. National Atlas for Norway, Statens Kartverk, Hønefoss.
- Nations, U., 2024. System of Environmental-Economic Accounting-Ecosystem Accounting. Technical Report. URL: <https://seea.un.org/ecosystem-accounting>.
- NCAVES, MAIA, . Monetary valuation of ecosystem services and ecosystem assets for ecosystem accounting: Interim Version 1st edition. Technical Report. New York.
- Norwegian Environment Agency, 2022. Mapping manual – Mapping of terrestrial Ecosystem Types following NiN2. Technical Report M-2209. Norwegian Environment Agency.
- Norwegian Environment Agency, 2024. Naturtyper etter Miljødirektoratets Instruks. URL: <https://kartkatalog.miljodirektoratet.no/Dataset/Details/2031>.
- Nybø, S., Kolstad, A.L., Sandvik, H., Bakkestuen, V., Evju, M., Framstad, E., Grainger, M., Jepsen, J.U., Nater, C.R., Speed, J.D., Töpper, J.P., Venter, Z., Åström, J., Åström, S.C., Bargmann, T.B., Silvennoinen, H.M., Sydenham, M.A., 2023. Indikatorer for økologisk tilstand i våtmark, semi-naturlig mark og naturlig åpne områder under skoggrensa. Technical Report. Norwegian Institute for Nature Research. URL: <https://brage.nina.no/nina-xmlui/handle/11250/3096724>.
- Pebesma, E., 2018. Simple features for r: Standardized support for spatial vector data. The R Journal 10, 439–446. URL: <https://doi.org/10.32614/RJ-2018-009>, doi:10.32614/RJ-2018-009.
- Pebesma, E., Bivand, R., 2023. Spatial Data Science: With applications in R. Chapman and Hall/CRC, London. URL: <https://r-spatial.org/book/>, doi:10.1201/9780429459016. DOI: 10.1201/9780429459016.
- Posit team, 2024. RStudio: Integrated Development Environment for R. Posit Software, PBC, Boston, MA. URL: <http://www.posit.co/>.
- R Core Team, 2023. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. URL: <https://www.R-project.org/>.
- Robinson, W., 2009. Ecological correlations and the behavior of individuals*. International Journal of Epidemiology 38, 337–341. URL: <https://doi.org/10.1093/ije/dyn357>, doi:10.1093/ije/dyn357.
- United Nations, 2021. System of Environmental-Economic Accounting-Ecosystem Accounting White cover (pre-edited) version. Technical Report. URL: <https://seea.un.org/ecosystem-accounting>.
- United Nations, 2022. Guidelines on biophysical modelling for ecosystem accounting .
- Vallecillo, S., Maes, J., Teller, A., Babí Almenar, J., Barredo, J.I., Trombetti, M., Abdul Malak, D., Paracchini, M.L., Carré, A., Addamo, A.M., Czúcz, B., Zulian, G., Marando, F., Erhard, M., Liqueste, C., Romao, C., Polce, C., Pardo Valle, A., Jones, A., Zurbaran-Nucci, M., Nocita, M., Vysna, V., Cardoso, A.C., Gervasini, E., Magliozzi, C., Baritz, R., Barbero, M., Andre, V., Kokkoris, I.P., Dimopoulos, P., Kovacevic, V., Gumbert, A., 2022. EU-wide methodology to map and assess ecosystem condition: towards a common approach consistent with a global statistical standard. Publications Office of the European Union, LU. URL: <https://data.europa.eu/doi/10.2760/13048>.
- Venter, Z.S., Czúcz, B., Stange, E., Nowell, M.S., Simensen, T., Immerzeel, B., Barton, D.N., 2024. ‘uncertainty audit’ for ecosystem accounting: Satellite-based ecosystem extent is biased without design-based area estimation and accuracy assessment. Ecosystem Services 66, 101599. URL: <https://www.sciencedirect.com/science/article/pii/S2212041624000056>, doi:10.1016/j.ecoser.2024.101599.
- Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemond, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T.L., Miller, E., Bache, S.M., Müller, K., Ooms, J., Robinson, D., Seidel, D.P., Spinu, V., Takahashi, K., Vaughan, D., Wilke, C., Woo, K., Yutani, H., 2019. Welcome to the tidyverse. Journal of Open Source Software 4, 1686. doi:10.21105/joss.01686.