

Status of Big Skate (*Beringraja binoculata*) Off the U.S. Pacific Coast in 2019



Ian G. Taylor¹
Vladlena Gertseva¹
Joseph Bizzarro²
Andi Stephens³

¹Northwest Fisheries Science Center, U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, 2725 Montlake Boulevard East, Seattle, Washington 98112

²Southwest Fisheries Science Center, U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, 110 Shaffer Road, Santa Cruz, California 95060

³Northwest Fisheries Science Center, U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, 2032 S.E. OSU Drive Newport, Oregon 97365

DRAFT SAFE

Disclaimer: This information is distributed solely for the purpose of pre-dissemination peer review under applicable information quality guidelines. It has not been formally disseminated by NOAA Fisheries. It does not represent and should not be construed to represent any agency determination or policy.

²² This report may be cited as:

²³ Taylor, I.G., Gertseva, V., Bizzarro, J., and Stephens, A. Status of Big Skate (*Beringraja*
²⁴ *binoculata*) Off the U.S. West Coast, 2019. Pacific Fishery Management Council, Portland, OR.
²⁵ Available from <http://www.pcouncil.org/groundfish/stock-assessments/>

Status of Big Skate (*Beringraja binoculata*) Off the U.S. Pacific Coast in 2019

Contents

Executive Summary	1
Stock	1
Catches	1
Data and Assessment	3
Stock Biomass	5
Recruitment	8
Exploitation status	10
Ecosystem Considerations	12
Reference Points	12
Management Performance	13
Unresolved Problems and Major Uncertainties	13
Decision Table	14
Research and Data Needs	18
1 Introduction	19
1.1 Distribution and Life History	19
1.2 Biology	19
1.3 Map	20
1.4 Ecosystem Considerations	21
1.5 Fishery Information	21
1.6 Stock Status and Management History	22
1.7 Management Performance	22
1.8 Fisheries Off Alaska, Canada and Mexico	22

51	2 Fishery Data	24
52	2.1 Data	24
53	2.2 Commercial Fishery Landings	24
54	2.2.1 Catch reconstructions for WA, OR, and CA	24
55	2.2.2 Tribal Catch in Washington	26
56	2.2.3 Commercial Discards	26
57	2.2.4 Commercial Fishery Length and Age Data	27
58	References	

59 **Executive Summary**

executive-summary

60 **Stock**

stock

61 This assessment reports the status of the Big Skate (*Beringraja binoculata*) resource in U.S.
62 waters off the West Coast using data through 2018.

63 **Catches**

catches

64 Landings and estimated discards of Big Skate were reconstructed for this assessment from
65 historical records of other species and from species composition data collected in the recent
66 fishery. These reflect the fishery from 1916-1994. The current fishery started in 1995. For
67 records from 1995-2017, Big Skate landings were estimated from species-composition samples
68 and the landings of “Unspecified Skates”. Beginning in 2017, Big Skate have been recorded
69 in species-specific landings.

70 (Table [a](#)).

71 (Figures ??-??)

72 (Figure [a](#))

73 In the current fishery (since 1995), annual total landings of Big Skate have ranged between
74 135-528 mt, with landings in 2018 totaling 173 mt.

Table a: Recent Big Skate landings (mt)

tab:Exec_catch

Year	Landings
2008	366.00
2009	205.70
2010	196.20
2011	268.40
2012	269.60
2013	135.00
2014	372.40
2015	331.50
2016	411.50
2017	277.60
2018	172.60




Figure a: Catch history of Big Skate in the model. ^{fig:r4ss_catches}

75 **Data and Assessment**

data-and-assessment

76 This the first full assessment for Big Skate, which was last assessed as part of the “Other
77 Species” complex. This assessment uses the newest version of Stock Synthesis (3.30.13). The
78 model begins in 1916, and assumes the stock was at an unfished equilibrium that year.

79 (Figure [b](#)).




Figure b: This is NOT the map depicting the distribution of Big Skate out to 600 ft. The stock assessment is bounded at Pt. Conception in the South to the U.S./Canada border in the north. fig:assess_region_map

80 **Stock Biomass**

stock-biomass

81 (Figure c and Table b).

82 The 2018 estimated spawning biomass relative to unfished equilibrium spawning biomass is
83 above the target of 40% of unfished spawning biomass at 99.8% (95% asymptotic interval:
84 \pm 99.8%-99.8%) (Figure d). Approximate confidence intervals based on the asymptotic
85 variance estimates show that the uncertainty in the estimated spawning biomass is high.

Table b: Recent trend in beginning of the year spawning output and depletion for the model for Big Skate.

tab:SpawningDeplete_mod1				
Year	Spawning Output (million eggs)	~ 95% confidence interval	Estimated depletion	~ 95% confidence interval
2010	70693.200	(70693.2- 70693.2)	0.998	(0.998-0.998)
2011	70697.500	(70697.5- 70697.5)	0.998	(0.998-0.998)
2012	70699.900	(70699.9- 70699.9)	0.998	(0.998-0.998)
2013	70702.400	(70702.4- 70702.4)	0.998	(0.998-0.998)
2014	70709.200	(70709.2- 70709.2)	0.998	(0.998-0.998)
2015	70708.700	(70708.7- 70708.7)	0.998	(0.998-0.998)
2016	70708.900	(70708.9- 70708.9)	0.998	(0.998-0.998)
2017	70706.000	(70706-70706)	0.998	(0.998-0.998)
2018	70706.500	(70706.5- 70706.5)	0.998	(0.998-0.998)
2019	70709.900	(70709.9- 70709.9)	0.998	(0.998-0.998)




Figure c: Time series of spawning biomass trajectory (circles and line: median; light broken lines: 95% credibility intervals) for the base case assessment model. fig:Spawnbi8_all




Figure d: Estimated relative depletion with approximate 95% asymptotic confidence intervals (dashed lines) for the base case assessment model. fig:RelDeplete_all

87 Recruitment deviations were estimated from 1916-2018 (Figure e and Table c).

Table c: Recent recruitment for the model.

Year	Estimated Recruitment (millions)	~ 95% confidence interval
2010	749.57	(749.57 - 749.57)
2011	749.59	(749.59 - 749.59)
2012	749.60	(749.6 - 749.6)
2013	749.61	(749.61 - 749.61)
2014	749.64	(749.64 - 749.64)
2015	749.63	(749.63 - 749.63)
2016	749.63	(749.63 - 749.63)
2017	749.62	(749.62 - 749.62)
2018	749.62	(749.63 - 749.63)
2019	749.64	(749.64 - 749.64)




Figure e: Time series of estimated Big Skate recruitments for the base-case model with 95% confidence or credibility intervals. `fig:Recruits_all`

88 Exploitation status

exploitation-status

89 Harvest rates estimated by the base model. management target levels (Table d and Figure
90 f).

Table d: Recent trend in spawning potential ratio and exploitation for Big Skate in the model. Fishing intensity is $(1-SPR)$ divided by 50% (the SPR target) and exploitation is F divided by F_{SPR} .

tab:SPR_Exploit_mod1				
Year	Fishing intensity	~ 95% confidence interval	Exploitation rate	~ 95% confidence interval
2009	0.00	(0-0)	0.00	(0-0)
2010	0.00	(0-0)	0.00	(0-0)
2011	0.00	(0-0)	0.00	(0-0)
2012	0.00	(0-0)	0.00	(0-0)
2013	0.00	(0-0)	0.00	(0-0)
2014	0.00	(0-0)	0.00	(0-0)
2015	0.00	(0-0)	0.00	(0-0)
2016	0.00	(0-0)	0.00	(0-0)
2017	0.00	(0-0)	0.00	(0-0)
2018	0.00	(0-0)	0.00	(0-0)




Figure f: Estimated spawning potential ratio (SPR) for the base-case model. One minus SPR is plotted so that higher exploitation rates occur on the upper portion of the y-axis. The management target is plotted as a red horizontal line and values above this reflect harvests in excess of the overfishing proxy based on the $\text{SPR}_{50\%}$ harvest rate. The last year in the time series is 2018. fig:SPR_all

Ecosystem Considerations

ecosystem-considerations

In this assessment, ecosystem considerations were not explicitly included in the analysis. This is primarily due to a lack of relevant data and results of analyses (conducted elsewhere) that could contribute ecosystem-related quantitative information for the assessment.

Reference Points

reference-points

This stock assessment estimates that Big Skate in the model is above the biomass target ($SB_{40\%}$), and well above the minimum stock size threshold ($SB_{25\%}$). The estimated relative depletion level for the base model in 2019 is 99.8% (95% asymptotic interval: $\pm 99.8\%$ -99.8%, corresponding to an unfished spawning biomass of 70709.9 million eggs (95% asymptotic interval: 70709.9-70709.9 million eggs) of spawning biomass in the base model (Table e). Unfished age 1+ biomass was estimated to be 2,814 mt in the base case model. The target spawning biomass ($SB_{40\%}$) is 2,834 million eggs, which corresponds with an equilibrium yield of 5,906 mt. Equilibrium yield at the proxy F_{MSY} harvest rate corresponding to $SPR_{50\%}$ is 5,070 mt (Figure g).

Table e: Summary of reference points and management quantities for the base case model.

Quantity	Estimate	tab:Ref_pts_mod1	
		Low 2.5% limit	High 2.5% limit
Unfished spawning output (million eggs)	7,086	7,086	7,086
Unfished age 1+ biomass (mt)	2,814	2,814	2,814
Unfished recruitment (R_0)	7,502	7,502	7,502
Spawning output(2018 million eggs)	7,071	7,071	7,071
Depletion (2018)	0.998	0.998	0.998
Reference points based on $SB_{40\%}$			
Proxy spawning output ($B_{40\%}$)	2,834	2,834	2,834
SPR resulting in $B_{40\%}$ ($SPR_{B_{40\%}}$)	0.625	0.625	0.625
Exploitation rate resulting in $B_{40\%}$	0.04	0.04	0.04
Yield with $SPR_{B_{40\%}}$ at $B_{40\%}$ (mt)	5,906	5,906	5,906
Reference points based on SPR proxy for MSY			
Spawning output	1,417	1,417	1,417
SPR_{proxy}	0.5		
Exploitation rate corresponding to SPR_{proxy}	0.058	0.058	0.058
Yield with SPR_{proxy} at SB_{SPR} (mt)	5,070	5,070	5,070
Reference points based on estimated MSY values			
Spawning output at MSY (SB_{MSY})	2,578	2,578	2,578
SPR_{MSY}	0.602	0.602	0.602
Exploitation rate at MSY	0.043	0.043	0.043
Dead Catch MSY (mt)	5,939	5,939	5,939
Retained Catch MSY (mt)	5,939	5,939	5,939

Management Performance

management-performance

Table f

Unresolved Problems and Major Uncertainties

unresolved-problems-and-major-uncertainties

Table f: Recent trend in total catch and commercial landings (mt) relative to the management guidelines. Big skate was managed in the Other Species complex in 2013 and 2014, designated an Ecosystem Component species in 2015 and 2016, and managed with stock-specific harvest specifications since 2017.

tab:mnmgt_perform				
Year	OFL (mt; ABC prior to 2011)	ABC (mt)	ACL (mt; OY prior to 2011)	Estimated total catch (mt)
2009				205.70
2010				196.20
2011				268.40
2012				269.60
2013	458.00	317.90	317.90	135.00
2014	458.00	317.90	317.90	372.40
2015				331.50
2016				411.50
2017	541.00	494.00	494.00	277.60
2018	541.00	494.00	494.00	172.60
2019	541.00	494.00	494.00	
2020	541.00	494.00	494.00	

108 Decision Table

decision-table

Table g: Projections of potential OFL (mt) for each model, using the base model forecast.

tab:OFL_projection	
Year	OFL
2019	158932.00
2020	149035.00
2021	141655.00
2022	136395.00
2023	132529.00
2024	129293.00
2025	126187.00
2026	122991.00
2027	119650.00
2028	116197.00
2029	112719.00
2030	109333.00




Figure g: Equilibrium yield curve for the base case model. Values are based on the 2018 fishery selectivity and with steepness fixed at 0.718. fig:Yield_all

Table h: Summary of 10-year projections beginning in 2020 for alternate states of nature based on an axis of uncertainty for the model. Columns range over low, mid, and high states of nature, and rows range over different assumptions of catch levels. An entry of "–" indicates that the stock is driven to very low abundance under the particular scenario.

tab:Decision_table_mod1

		States of nature					
		Low M 0.05		Base M 0.07		High M 0.09	
	Year	Catch	Spawning Output	Depletion	Spawning Output	Depletion	Spawning Output
40-10 Rule, Low M	2019	-	-	-	-	-	-
	2020	-	-	-	-	-	-
	2021	-	-	-	-	-	-
	2022	-	-	-	-	-	-
	2023	-	-	-	-	-	-
	2024	-	-	-	-	-	-
	2025	-	-	-	-	-	-
	2026	-	-	-	-	-	-
	2027	-	-	-	-	-	-
	2028	-	-	-	-	-	-
40-10 Rule	2019	-	-	-	-	-	-
	2020	-	-	-	-	-	-
	2021	-	-	-	-	-	-
	2022	-	-	-	-	-	-
	2023	-	-	-	-	-	-
	2024	-	-	-	-	-	-
	2025	-	-	-	-	-	-
	2026	-	-	-	-	-	-
	2027	-	-	-	-	-	-
	2028	-	-	-	-	-	-
40-10 Rule, High M	2019	-	-	-	-	-	-
	2020	-	-	-	-	-	-
	2021	-	-	-	-	-	-
	2022	-	-	-	-	-	-
	2023	-	-	-	-	-	-
	2024	-	-	-	-	-	-
	2025	-	-	-	-	-	-
	2026	-	-	-	-	-	-
	2027	-	-	-	-	-	-
	2028	-	-	-	-	-	-
Average Catch	2019	-	-	-	-	-	-
	2020	-	-	-	-	-	-
	2021	-	-	-	-	-	-
	2022	-	-	-	-	-	-
	2023	-	-	-	-	-	-
	2024	-	-	-	-	-	-
	2025	-	-	-	-	-	-
	2026	-	-	-	-	-	-
	2027	-	-	-	-	-	-
	2028	-	-	-	-	-	-

Table i: Base case results summary.

Quantity	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
	tab:base summary									
Landings (mt)										
Total Est. Catch (mt)										
OFL (mt)										
ACL (mt)										
(1-SPR)(1-SPR _{50%})	0	0	0	0	0	0	0	0	0	0
Exploitation rate	0	0	0	0	0	0	0	0	0	0
Age 1+ biomass (mt)	2654110	2654240	2654360	2654400	2654430	2654570	2654490	2654470	2654390	2654450
Spawning Output	70693.2	70697.5	70699.9	70702.4	70709.2	70708.7	70708.9	70706.0	70706.5	70709.9
95% CI	(70693.2-70693.2)	(70697.5-70697.5)	(70699.9-70699.9)	(70702.4-70702.4)	(70709.2-70709.2)	(70708.7-70708.7)	(70708.9-70708.9)	(70706-70706)	(70706.5-70706.5)	(70709.9-70709.9)
Depletion	1	1	1	1	1	1	1	1	1	1
95% CI	(0.998-0.998)	(0.998-0.998)	(0.998-0.998)	(0.998-0.998)	(0.998-0.998)	(0.998-0.998)	(0.998-0.998)	(0.998-0.998)	(0.998-0.998)	(0.998-0.998)
Recruits	749.57	749.59	749.60	749.61	749.64	749.63	749.63	749.62	749.62	749.64
95% CI	(749.57-749.57)	(749.59-749.59)	(749.6-749.6)	(749.61-749.61)	(749.64-749.64)	(749.63-749.63)	(749.63-749.63)	(749.62-749.62)	(749.63-749.63)	(749.64-749.64)

109 **Research and Data Needs**

research-and-data-needs

110 We recommend the following research be conducted before the next assessment:

111 1. **xxxx:**

112 2. **xxxx:**

113 3. **xxxx:**

114 4. **xxxx:**

115 5. **xxxx:**

1 Introduction

introduction

1.1 Distribution and Life History

distribution-and-life-history

Big Skate (*Raja binoculata*) is the largest of the skate species in North America with a documented maximum length of 244 cm total length and a maximum weight of 91 kg (Eschmeyer and Herald 1983). The species name “binoculata” (two-eyed) refers to the prominent ocellus at the base of each pectoral fin. Big skate range from the Bering Sea to Cedros Island in Baja California, but are uncommon south of Pt. Conception. Big skate have a shallow depth distribution; they occur in coastal bays, estuaries, and over the continental shelf, usually on sandy or muddy bottoms, but occasionally on low strands of kelp.

Skates are the largest and most widely distributed group of batoid fish with approximately 245 species ascribed to two families (Ebert and Compagno 2007)(McEachran and Miyake 1990). Skates are benthic fish that are found in all coastal waters but are most common in cold temperatures and polar waters (Ebert and Compagno 2007).

There are eleven species of skates in three genera (Amblyraja, Bathyrja, and Raja) present in the Northeast Pacific Ocean off California, Oregon and Washington (Ebert 2003). Of that number, just three species (Longnose Skate, *Raja rhina*; Big Skate, *Raja binoculata*; and Sandpaper Skate, *Bathyrja interrupta*) make up over 95 percent of West Coast Groundfish Bottom Trawl Survey (WCG BTS) catches in terms of biomass and numbers, with the Longnose Skate leading in both categories (with 62 percent of biomass and 56 percent of numbers).

Big Skates are usually seen buried in sediment with only their eyes showing. They feed on polychaete worms, mollusks, crustaceans, and small benthic fishes. Polychaetes and mollusks comprise a slightly greater percentage of the diet of younger individuals. The eyespots on the skates’ wings are believed to serve as decoys to confuse predators. A known predator of big skates is the Broadnose Sevengill Shark (*Notorhynchus cepedianus*). Juvenile Northern Elephant Seals (*Mirounga angustirostris*) are known to consume the egg cases of the Big Skate. Known parasites include the copepod *Lepeophtheirus cuneifer*.

In 2012, the Big Skate was moved from genus *Raja* to the new genus *Beringraja* together with the Mottled Skate (*B. pulchra*) (Ishihara et al. 2012). These are the only two skates with multiple embryos per egg case, and they are very similar morphologically and genetically (Bizzarro 2019).

1.2 Biology

biology

The Big Skate is broadly distributed, occurring from the southeastern Bering Sea (Mecklenburg, CW and Mecklenburg, TA and Thorsteinson, LK 2002) to southern Baja California

(22.90° N, 110.03° W; (Castro-Aguirre et al. 1993)) and the Gulf of California (Castro-Aguirre and Pérez 1996). It has been reported at depths of 2–501 m (min: Miller et al. (1980); max: Farrugia et al. (2016)) but is most common on the inner continental shelf (< 100 m; (Love, Milton S 2011); (Bizzarro 2015)). Big Skates are highly mobile and capable of long range (> 2000 km) movements ((King and McFarlane 2009); (Farrugia et al. 2016)).

Big Skate is oviparous, and is one of two skate species that have multiple embryos per egg case (Ebert et al. 2008). From 1–8 embryos can be contained in a single, large egg capsule, but most have 3–4 (DeLacy and Chapman 1935, Hitz 1964, Ford 1971). Eggs are deposited year-round on sand or mud substrates at depths of ~50–150 m (Hitz 1964, Ebert and Compagno 2007). Embryos hatch from eggs after 6–20 months, with shorter developmental periods associated with warmer temperatures (Hoff 2007). In captivity, Big Skate females may produce > 350 eggs/year (average of 2 embryos/egg case; Chiquillo, Kelcie L and Ebert, David A and Slager, Christina J and Crow, Karen D (2014)) from long-term sperm storage (???). Size at birth is 18–23 cm TL (Ebert 2003). Maximum size is 244 cm TL [Eschmeyer and Herald (1983), with females growing to larger sizes.

Size at maturity has been variably estimated for Big Skate populations off California, British Columbia, and Alaska. Off central California, Zeiner and Wolf (1993) reported sizes at first maturity of ~129 cm TL (females) and ~100 cm TL (males). A similar size at maturity was estimated for females from the Gulf of Alaska (first = 126 cm TL, 50% = 149 cm TL), but male estimates were considerably greater (first = 124 cm TL, 50% = 119 cm TL; Ebert et al. (2008)). Much smaller sizes at first (female = 60 cm TL, male = 50 cm TL) and 50% (female = 90 cm TL, male = 72 cm TL) maturity were generated for the Longnose Skate populations off British Columbia (McFarlane GA and King JR 2006); however, maturity evaluation criteria were flawed (subadults were considered to be mature), and these results are therefore not considered valid.

Age and growth parameters have been established from California, British Columbia, and the Gulf of Alaska. Maximum ages off central California (females = 12, males = 11; Zeiner, S.J. and P. Wolf. (1993)) and in the Gulf of Alaska (females = 14, males = 15; Gburski et al. 2007) were similar, but estimates off British Columbia were much greater (females = 26, males = 25; McFarlane and King 2006). It is important to note that age estimates are based on an unvalidated method and geographic differences in size or age may reflect differences in sampling or ageing criteria. In the Gulf of Alaska, Big Skates reach 50% maturity at 10 years and 7 years for females and males, respectively (Gburski, C.M. and Gaichas, S.K. and Kimura, D.K. (2007), Ebert et al. (2008)). Generation length estimates range from 11.5 (Zeiner, S.J. and P. Wolf. 1993) to 17 years (McFarlane GA and King JR 2006).

1.3 Map

map

A map showing the scope of the assessment and depicting boundaries for fisheries or data collection strata is provided in Figure 1.

1.4 Ecosystem Considerations

ecosystem-considerations-1

In this assessment, ecosystem considerations were not explicitly included in the analysis. This is primarily due to a lack of relevant data and results of analyses (conducted elsewhere) that could contribute ecosystem-related quantitative information for the assessment.

1.5 Fishery Information

fishery-information

Big Skate are caught in commercial and recreational fisheries on the West Coast using line and trawl gears. There is a limited market for pectoral fins (skate wings).

The history of Big Skate is not well documented. They were used as a food source by the native Coastal and Salish Tribes (Batdorf, C 1990) long before Europeans settled in the Pacific Northwest and then as fertilizer by the settlers (Bowers, G. M. 1909). No directed fishery for Big Skate has been documented; rather, they were taken along with other skates and rays as “scrap fish” and used for fertilizer, fish meal and oil (Lippert 2019).

Skates have been regarded as a predator on desirable market species such as Dungeness crab, and were thought of as nuisance fish with no appeal as a food item save for small local markets. They had been discarded or harvested at a minimal level until their livers became valued along with those of other cartilaginous fishes for the extraction of vitamin A in the 1940s. Chapman (Chapman, WM 1944) recorded that “At present they are being fished heavily, in common with the other elasmobranchs of the coast, for the vitamins in their livers. The carcasses are either thrown away at sea or made into fish meal. Little use is made of the excellent meat of the wings”.

Little information is available about the historic Washington fishery for Big Skate. In records before 2000, they are lumped together with other skates or in market categories (Lippert 2019); this necessitates considerable attention to reconstructing the fishery by observing the composition of skate catches in the modern fishery and applying those to the recently reconstructed historical records.

Very little information is known about the Big Skate historical fishery in Oregon. The information we do have is mainly from historical landing data and species composition samples starting in the mid-nineties. The bulk of the catch is from the bottom trawl and longline fisheries, with smaller amounts as by-catch in mid-water trawl and the shrimp trawl fishery. Big Skate was lumped into the nominal “Skate” category until 2015 when it was separated into its own market category. Species composition data have been vitally important in reconstructing the pre-2015 historical catch (Calavan 2019).

1.6 Stock Status and Management History

stock-status-and-management-history

Big Skate were managed in the “Other Fish” complex until 2015 when they were designated an Ecosystem Component (EC) species. Catches of Big Skate are estimated to have averaged 95 mt from 2007–2011, along with large landings of “Unspecified Skate”. Analysis of Oregon port-sampling data indicates that about 98 percent of the recent Unspecified Skate landings in Oregon were comprised of Big Skate. Such large landings indicates targeting of Big Skate has occurred and an EC designation was not warranted. Based on this evidence, Big Skate was redesignated as an actively-managed species in the fishery. Big skate have been managed with stock-specific harvest specifications since 2017.

The recent OFL of 541 mt was calculated by applying approximate MSY harvest rates to estimates of stock biomass from the Northwest Fisheries Science Center (NWFSC) West Coast Groundfish Bottom Trawl Survey. This survey-based biomass estimate is likely underestimated since Big Skate are distributed all the way to the shoreline and no West Coast trawl surveys have been conducted in water shallower than 55 meters. This introduces an extra source of uncertainty to management and suggests that increased precaution is needed to reduce the risk of overfishing the stock.

There has been consideration for managing Big Skate in a complex with Longnose Skate, the other actively-managed West Coast skate species, but the two species have disparate distributions and fishery interactions (Longnose Skate is much more deeply distributed than Big Skate) and that option was not endorsed. The Pacific Fishery Management Council has chosen to set the Annual Catch Limit (ACL) equal to the Allowable Biological Catch (ABC) with a buffer for management uncertainty (P^*) of 0.45.

1.7 Management Performance

management-performance-1

Table [f](#)

1.8 Fisheries Off Alaska, Canada and Mexico

fisheries-off-alaska-canada-and-mexico

** Alaska **

In Alaska, skates were primarily taken as bycatch in both longline and trawl fisheries until 2003, when a directed skate fishery developed in the Gulf of Alaska, where Longnose and Big skates comprise the majority of the skate biomass.

The Gulf of Alaska (GOA) skate complex is managed as three units. Big skates and Longnose Skates each have separate harvest specifications, with acceptable biological catches (ABCs)

specified for each GOA regulatory area (western, central, and eastern). A single gulfwide overfishing level (OFL) is specified for each stock. All remaining skate species are managed as an “Other Skates” group with gulfwide harvest specifications. All GOA skates are managed as Tier 5 stocks, where OFL and ABC are based on survey biomass estimates and natural mortality rate.

In the Bering Sea and Aleutian Islands, skates are assessed as a group rather than as separate species.

**** Canada ****

In Canada historic information regarding skate catches goes back to the 1950’s. Prior to 1990’s skates were taken mostly as bycatch and landings were reported as part of a skate complex (not by species). As with the West Coast, the trawl fishery is responsible for the largest amount of bycatch. Skate catches off British Columbia accelerated in the early 1990’s, partly due to emerging Asian markets. Since 1996, longnose skate has been targeted by the B.C. trawl fishery and, as a result, catches have been more accurately reported.

Assessments of Longnose Skate and Big Skate were conducted by Canada’s Division of Fisheries and Oceans in 2015(King, J.R., Surry, A.M., Garcia, S., and P.J. Starr 2015). For Big Skate, a Bayesian surplus production model failed to provide plausible results, and two data-limited approaches were investigated: Depletion-Corrected Average Catch Analysis (DCAC), and a Catch-MSY (maximum sustainable yield) Approach.

DCAC produced a range of potential yield estimates that were above the long-term average catch, with an upper bound that was three orders of magnitude larger than the long-term average catch. The Catch-MSY approach was found to be quite sensitive to assumptions and was not recommended as the sole basis of advice to managers.

The recommendation for management for both skate species was that they should be managed with harvest yields based on mean historic catch, with consideration given to survey trends and to the ranges of maximum sustainable yield estimates identified by the Catch-MSY Approach. However, the analysis found no significant trends in abundance indices for Big Skate, and mean historical catches were below the maximum MSY estimate from the catch-MSY results.

2 Fishery Data

fishery-data

2.1 Data

data

Data used in the Big Skate assessment are summarized in Figure 2. Descriptions of the data sources are in the following sections.

2.2 Commercial Fishery Landings

commercial-fishery-landings

2.2.1 Catch reconstructions for WA, OR, and CA

catch-reconstructions-for-wa-or-and-ca

Washington Commercial Skate Landings Reconstruction

Information for Big Skate is very limited, in part because the requirement to sort landings of Big Skate in the shore-based Individual Fishing Quota fishery from landings in the “Unidentified Skate” category was not implemented until June 2015. The historical catch of Big Skate therefore relies on the historical reconstruction of Longnose Skate.

For the 2019 assessment, a new approach has been developed for estimating the catch history for Longnose Skate based on a linear regression model that predicts the catch of Longnose Skate from the catch of Dover sole, for which historical catch estimates are available (Gertseva, V. 2019). The dependent variable for the linear regression model was the West Coast Groundfish Observer Program (WCGOP) annual estimates of the coastwide total catch (landings plus discards) of Longnose Skate for the period 2009 to 2017 and the independent variable was the corresponding WCGOP annual estimates of coastwide total catch (landings plus discards) of Dover sole. The regression model has good predictive power ($R^2 = 95.7\%$) over the range of the Dover sole catches (6,500 to 12,500 mt).

The discard component of the catch reconstruction for Big Skate may be based either on the catch reconstruction for Longnose Skate and the assumption that the two species experience similar discard rates (discard / total catch) or on a similar analysis with links to species that co-occur with big skate. Data from the Pikitch discard study (1985-1987) and from WCGOP (2015-2017) support the idea that discard rates for the two species are very similar. Also, market demand for skates does not seem to distinguish between the two species. There are insufficient years of data from the WCGOP to develop a regression model for Big Skate as was done for Longnose Skate.

Oregon Commercial Skate Landings Reconstruction

Oregon Department of Fish and Wildlife (ODFW) provided newly reconstructed commercial landings for all observed skate species for the 2019 assessment cycle (1978 – 2018). In

addition, the methods were reviewed at a pre-assessment workshop. Historically, skates were landed as a single skate complex in Oregon. In 2009, longnose skates were separated into their own single-species landing category, and in 2014, big skates were also separated. The reconstruction methodology differed by these three time blocks in which species composition collections diverged (1978 – 2008; 2009 – 2014; 2015 – 2018).

Species compositions of skate complexes from commercial port sampling are available throughout this time period but are generally limited, which precluded the use of all strata for reconstructing landings. Quarter and port were excluded, retaining gear type, PMFC area, and market category for stratifying reconstructed landings within the three time blocks. Bottom trawl gear types include multiple bottom trawl gears, and account for greater than 98% of skate landings. Minor gear types include primarily bottom longline gear, but also include mid-water trawl, hook and line, shrimp trawl, pot gear and scallop dredge.

For bottom trawl gears, trawl logbook areas and adjusted skate catches were matched with strata-specific species compositions. In Time Block 1 (1978 – 2008), all bottom trawl gear types were aggregated due to a lack of specificity in the gear recorded on the fish tickets. However, in Time Blocks 2 and 3, individual bottom trawl gear types were retained. Some borrowing of species compositions was required (31% of strata) and when necessary, borrowed from the closest area or from the most similar gear type. Longline gear landings were reconstructed in a similar fashion as to bottom trawl and required some borrowing among strata as well (25%).

Due to insufficient species compositions, mid-water trawl landings were reconstructed using a novel depth-based approach. Available compositions indicate that the proportion by weight of big skates within a composition drops to zero at approximately 100 fathoms, and an inverse relationship is observed for longnose skate, where the proportion by weight is consistently one beyond 100 – 150 fathoms. Complex-level landings were assigned a depth from logbook entries and these species specific depth associations were used to parse out landings by species. The approach differed somewhat by time block. Landings from shrimp trawls were handled using a similar methodology. Finally, very minor landings from hook and line, pot gear and scallop dredges were assigned a single aggregated species composition, as they lack any gear-specific composition samples. Landings from within a time block were apportioned by year using the proportion of the annual ticket landings.

Results indicate that the species-specific landings from this reconstruction are very similar to those from Oregon’s commercial catch reconstruction (Karnowski et al. 2014) during the overlapping years but cover a greater time period with methodology more applicable to skates in particular. ODFW intends to incorporate reconstructed skate landings into PacFIN in the future (A. Whitman, ODFW; pers. comm.).

California Catch Reconstruction

A reconstruction of historical skate landings from California waters was developed for the 1916–2017 time period using a combination of commercial catch data (spatially explicit block

summary catches and port sample data from 2009-2017) and fishery-independent survey data (Bizzarro 2019). Virtually all landings in California were of “unspecified skate” until species-composition sampling of skate market categories began in 2009.

From 2009 through 2017, catch estimates were based on these market category species-composition samples, and the average of those species-compositions was hindcast to 2002, based on the assumption that those data were representative of the era of large area closures in the post-2000 period.

For the period from 1936-1980, spatially explicit landings data (the California Department of Fisheries and Wildlife (CDFW) block summary data) were merged with survey data to provide species-specific estimates.

For years 1981-2001, a “blended” product of these two approaches was taken, in which a linear weighting scheme blended the two sets of catch estimates through that period. Landings estimates were also scaled upwards by an expansion factor for skates landed as “dressed” based on fish ticket data. Prior to 1981 these data had not been reported and skate landings were scaled by the “average” percentage landed as dressed in the 1981-1985 time period, but by the late 1980s nearly all skates were landed round.

As no spatial information on catch is available from 1916-1930, and the block summary data were very sparse in the first few years of the CDFW fish ticket program (1931–1934), spatial information from the late 1930’s was used to hindcast to the 1916–1935 time period.

2.2.2 Tribal Catch in Washington

tribal-catch-in-washington

2.2.3 Commercial Discards

commercial-discards

Commercial discards of Big Skate are highly uncertain. The method used to estimate discards for Longnose Skate was based on a strong correlation between total mortality of that species, and total mortality of Dover Sole for the years 2009–2017 during which Longnose were landed separately from other skates. In contrast, the sorting requirement for Big Skate occurred too recently to provide an adequate range of years for this type of correlation. Furthermore, there is greater uncertainty in the total mortality for the shallow-water species with which Big Skate most often co-occurs, such as Sand Sole and Starry Flounder, than there is for Dover Sole, which has been the subject of recurring stock assessments.

However, those involved in the fishery for both skate species report that discarding for Big Skate and Longnose Skate in the years prior to 1995 were driven by the same market forces and the discard rates were similar. primarily lack of markets or fish processors accepting only skate wings that had been separated at-sea, as well as the quantitative have more uncertainty in their own catch estimates have no stock assessment and more uncertain mortality estimated total mortality and Dover Sole for which a correlation between relationship (Gertseva, V. 2019),

2.2.4 Commercial Fishery Length and Age Data

commercial-fishery-length-and-age-data

The input sample sizes were calculated via the Stewart Method (Ian Stewart, personal communication, IPHC):

$$\text{Input effN} = N_{\text{trips}} + 0.138 * N_{\text{fish}} \text{ if } N_{\text{fish}}/N_{\text{trips}} \text{ is } < 44$$

$$\text{Input effN} = 7.06 * N_{\text{trips}} \text{ if } N_{\text{fish}}/N_{\text{trips}} \text{ is } \geq 44$$

Sport Fishery Removals and Discards

Biological samples from the recreational fleets are described in the sections below.

Fishery-Dependent Indices of Abundance

Data Source 1

Data Source 1 Index Standardization

Data Source 1 Length Composition

Data Source 2

Data Source 3

Fishery-Independent Data Sources

Alaska Fisheries Science Center (AFSC) Triennial Shelf Survey

Research surveys have been used since the 1970s to provide fishery-independent information about the abundance, distribution, and biological characteristics of Big Skate. A coast-wide survey was conducted in 1977 (Gunderson, Donald Raymond and Sample, Terrance M. 1980) by the Alaska Fisheries Science Center, and repeated every three years through 2001. The final year of this survey, 2004, was conducted by the NWFSC according to the AFSC protocol. We refer to this as the **Triennial Survey**.

The survey design used equally-spaced transects from which searches for tows in a specific depth range were initiated. The depth range and latitudinal range was not consistent across years, but all years in the period 1980-2004 included the area from 40° 10'N north to the Canadian border and a depth range that included 55-366 meters, which spans the range where the vast majority of Big Skate encountered in all trawl surveys. Therefore the index was based on this depth range. The survey as conducted in 1977 had incomplete coverage and is not believe to be comparable to the later years, and is not used in the index.

An index of abundance was estimated based on the VAST delta-GLMM model as described for the NWFSC Combo Index above. In this case as well, Q-Q plots indicated slightly better performance of the gamma over lognormal models for positive tows (Figure ??).

Northwest Fisheries Science Center West Coast Groundfish Bottom Trawl Survey

In 2003, the NWFSC took over an ongoing slope survey the AFSC had been conducting, and expanded it spatially to include the continental shelf. This survey, referred to in this document as the **NWFSC Combo Survey**, has been conducted annually since. It uses a random-grid design covering the coastal waters from a depth of 55 m to 1,280 m from late-May to early-October (Bradburn, M.J. and Keller, A.A. and Horness, B.H. 2011, Keller, A.A. and Wallace, J.R. and Methot, R.D. 2017). Four chartered industry vessels are used each year (with the exception of 2013 when the U.S. federal-government shutdown curtailed the survey). Yellowtail catches in the NWFSC Combo Survey are shown in ??.

The data from the NWFSC Combo survey was analyzed using a spatio-temporal delta-model (Thorson, J. T. and Shelton, A. O. and Ward, E. J. and Skaug, H. J. 2015), implemented as an R package VAST (Thorson, James T. and Barnett, Lewis A. K. 2017) and publicly available online (<https://github.com/James-Thorson/VAST>). Spatial and spatio-temporal variation is specifically included in both encounter probability and positive catch rates, a logit-link for encounter probability, and a log-link for positive catch rates. Vessel-year effects were included for each unique combination of vessel and year in the database.

Data Source 1 Index Standardization VAST

Data Source 1 Length Composition

Triennial Survey *Data Source 2 Index Standardization VAST*

International Pacific Halibut Commission Longline Survey

The IPHC has conducted an annual longline survey for Pacific Halibut off the coast of Oregon and Washington since 1997 (no surveys were performed in 1998 or 2000). Beginning in 1999, this has been a fixed station design, with 84 locations in this area (station locations differed in 1997, and are therefore not comparable with subsequent surveys). 400 to 800 hooks have been deployed at each station in 100-hook groups (typically called “skates” although that term will be avoided here to avoid confusion). The gear used to conduct the survey was designed to efficiently sample Pacific Halibut and used 16/0 (#3) circle hooks baited with Chum Salmon.

In some years from 2011 onward, additional stations were added to the survey to sample Yelloweye Rockfish. These stations were excluded from the analysis, as were additional stations added in 2013, 2014, and 2017, off the coast of California (south of 42 degrees latitude). Some variability in exact sampling location is practically unavoidable, and leeway is given in the IPHC methods to center the set on the target coordinates while allowing wind and currents to dictate the actual direction in which the gear is deployed. This can result in different habitats being accessed at each fixed deployment location across years. One station that was very close to the U.S. Canada border had the mid-point of the set in Canada in 2

out of the 19 years of the survey. For consistency among years, all samples from this station were included in the analysis, including those in Canada.

In most years, bycatch of non-halibut species has been recorded during this survey on the first 20 hooks of each 100-hook group, although in 2003 only 10% of the hooks were observed for bycatch, and starting in 2012, some stations had 100% of the hooks observed for bycatch. Combining these observation pattern with the number of hooks deployed each year, resulted in most stations having 80, 100, 120, 140, or 160 hooks observed, with a mean of 144 hooks and a maximum of 800 hooks observed. The depth range of the 84 stations considered was 42–530 m, thus extending beyond the range of Big Skate, but 74% of the stations were shallower than 200 m. Big Skate have been observed at 51 of the 84 the standard stations that were retained for this analysis, but no station had Big Skates observed in more than 12 out of the 19 years of survey data, and only 10% of the station/year combinations had at least one observed Big Skate (Figure X). Of those station/year combinations with at least one Big Skate observed, the Big Skates were observed on an average of 1.3% of the hooks observed. The highest proportion was 10 Big Skates out of 81 hooks observed at one station.

The IPHC longline survey catch data were standardized using a Generalized Linear Model (GLM) with binomial error structure. Catch-per-hook was modeled, rather than catch per station due to the variability in the number of hooks deployed and observed each year. The binomial error structure was considered logical, given the binary nature of capturing (or not) a Longnose Skate on each longline hook. The modeling approach is identical to that which has been applied in the past for Yelloweye Rockfish (*Stewart et al. 2009*), and Spiny Dogfish (*Gertseva et al., 2011*). MCMC sampling of the GLM parameters was used to estimate the variability around each index estimate. The median index estimates themselves were approximately equal to the observed mean catch rate in each year (Figure Y). In recent years, the IPHC standardization of the index of halibut abundance has included an adjustment to account for missing baits on hooks returned empty in an effort to account for reduced catchability of the gear that may result from the lost bait. This adjustment was not included in the analysis for Big Skate although it could be considered in future years.

483 ###Biological Parameters and Data

484 Measurement Details and Conversion Factors

485 Disc width to total length (estimated by Ian on Apr 15, similar to Ebert 2008 estimates
486 for Alaska) $L = 1.3399 * W$ estimated from 95 samples from WCGBTS where both mea-
487 surements collected (R-squared = 0.9983). Little sex difference observed, so using single
488 relationship for both sexes. Inter-spiracle width to total length from Downs & Cheng (2013):
489 $L = 12.111 + 9.761ISW$ (*females*) $L = 3.824 + 10.927ISW$ (*males*)

490 Love et al. ([1987](#))

491 Length and Age Compositions

492 Length comps (some based on widths)

493 WCGBTS Lengths from all years except 2006 and 2007 Widths in 2006 and 2007

494 Triennial Survey Sample sizes: 3 in 1998 (all widths), 84 in 2001 (3 widths, 81 lengths), 100
495 in 2004 (all lengths) Triennial survey About 90+ samples in each of 2001 and 2004 Only 3
496 unsexed fish from 1998

497 Commercial fisheries In process Discard comps from 2010-2015

498 Length compositions were provided from the following sources:

- 499 • Source 1 (*type, e.g., commercial dead fish, research, recreational, yyyy-yyyy*)
- 500 • Source 2 (*type, yyyy-yyyy*)
- 501 • Source 3 (*research, yyyy, yyyy, yyyy, yyyy*)

502 The length composition of all fisheries aggregated across time by fleet is in Figure [3](#). De-
503 scriptions and details of the length composition data are in the above section for each fleet
504 or survey.

505 Age Structures

506 von Bertalanffy growth curve (von Bertalanffy, L [1938](#)), $L_i = L_{\infty}e^{(-k[t-t_0])}$, where L_i is the
507 length (cm) at age i , t is age in years, k is rate of increase in growth, t_0 is the intercept, and
508 L_{∞} is the asymptotic length.

509 Ages WCGBTS Currently only 333 ages from 2010 present in data warehouse as of Apr
510 15 Patrick submitting an 300 additional ages from 2016 and 2017 to Beth on Apr 2 and
511 promised further additions during the week of Apr 15.

512 Triennial Survey No ages

513 Commercial fisheries 2009 samples from WA were stratified by length, so should be treated
514 as conditionals

515 **Aging Precision and Bias**

516 **Weight-Length**

517 Estimated by Ian based on WCGBT samples (n = 1159) $Weight = 0.0000074924 * Length^2.9925$ (Figure 4).

519 **Sex Ratio, Maturity, and Fecundity**

520 The female maturity relationship was based on visual maturity estimates from port sam-
521 plers (n = 278, of which 241 were from Oregon and 37 from Washington, with 24 mature
522 specimens) as well as 55 samples from the WCGBTs (of which 4 were mature). The result-
523 ing relationship was $L_{50\%} = 148.2453$ with a slope parameter of $Beta = -0.13155$ in the
524 relationship $M = (1 + Beta(L - L_{50\%}))^{-1}$ (Figure 5).

525 **Natural Mortality**

526 The Hamel prior for M is $\text{lognormal}(\ln(5.4/\text{max age}), .438)$, which based on 1 age-15 fish out
527 of 1034 observed in the WCGBTs results in $\text{lognormal}(-1.021651, 0.438)$

528 If it needs to be fixed, it should be set to $M = 5.4/\text{max age} = 5.4/15 = 0.36$

529 ###Environmental or Ecosystem Data Included in the Assessment In this assessment,
530 neither environmental nor ecosystem considerations were explicitly included in the analysis.
531 This is primarily due to a lack of relevant data and results of analyses (conducted elsewhere)
532 that could contribute ecosystem-related quantitative information for the assessment.

533 ##Previous Assessments

534 ###History of Modeling Approaches Used for this Stock

535 Deriving estimates of OFL for species in the “Other Fish” complex or potential alternative
536 complexes

537 The current “Other Fish” complex and proposed alternatives include a number of species for
538 which estimates of OFL contributions are not available from stock assessments or data-poor
539 methods. Four of the species had OFL contributions for the 2013–2014 management cycle
540 calculated by applying approximate MSY harvest rates to estimates of stock biomass from
541 the NWFSC West Coast Bottom Trawl Survey (Bradburn et al., 2012). This approach is
542 described in detail in Cope et al. (2012).

543 ###yyyy Assessment Recommendations

544 **Recommendation 1:**

545

546 STAT response: xxxxx

547 **Recommendation 2:**

548

549 STAT response: xxxxx

550 **Recommendation 3:**

551

552 STAT response: xxxx

553 ##Model Description

554 ###Transition to the Current Stock Assessment

555 ###Summary of Data for Fleets and Areas There are xxx fleets in the base model. They
556 include:

557 *Commercial:* The commercial fleets include ...

558 *Recreational:* The recreational fleets include ...

559 *Research:* There are xx sources of fishery-independent data available ...

560 ###Other Specifications

561 **###Modeling Software** The STAT team used Stock Synthesis 3 version 3.30.05.03 by
562 Dr. Richard Methot at the NWFSC. This most recent version was used, since it included
563 improvements and corrections to older versions. The r4SS package (GitHub release number
564 v1.27.0) was used to post-processing output data from Stock Synthesis.

565 **###Data Weighting**

566 **###Priors** The log-normal prior for female natural mortality were based on a meta-analysis
567 completed by Hamel (2015), as described under “Natural Mortality.” Female natural mor-
568 tality was fixed at the median of the prior, 0.xxx for an assumed maximum age of xx. An
569 uninformative prior was used for the male offset natural mortality, which was estimated.

570 The prior for steepness (h) assumes a beta distribution with parameters based on an update
571 for the Thorson-Dorn rockfish prior (Dorn, M. and Thorson, J., pers. comm.), which was
572 endorsed by the Science and Statistical Committee in 2018. The prior is a beta distribution
573 with $\mu=0.xxx$ and $\sigma=0.xxx$. Steepness is fixed in the base model at the mean of the
574 prior. The priors were applied in sensitivity analyses where these parameters were estimated.

575 **###Estimated and Fixed Parameters** A full list of all estimated and fixed parameters is
576 provided in Tables ??.

577 The base model has a total of xxx estimated parameters in the following categories:

- 578 • xxx,
- 579 • xxx
- 580 • xxx, and
- 581 • xxx selectivity parameters

582 The estimated parameters are described in greater detail below and a full list of all estimated
583 and parameters is provided in Table ??.

584 *Growth.*

585 *Natural Mortality.*

586 *Selectivity.*

587 *Other Estimated Parameters.*

588 *Other Fixed Parameters.*

589 **##Model Selection and Evaluation** **###Key Assumptions and Structural Choices**

590 ###Alternate Models Considered

591 ###Convergence

592 ##Response to the Current STAR Panel Requests

593 **Request No. 1:**

594

595 **Rationale:** xxx

596 **STAT Response:** xxx

597 **Request No. 2:**

598

599 **Rationale:** xxx

600 **STAT Response:** xxx

601 **Request No. 3:**

602

603 **Rationale:** x.

604 **STAT Response:** xxx

605 **Request No. 4:**

606

607 **Rationale:** xxx

608 **STAT Response:** xxx

609 **Request No. 5:**

610

611 **Rationale:** xxx

612 **STAT Response:** xxx

613 ##Base Case Model Results The following description of the model results reflects a base
614 model that incorporates all of the changes made during the STAR panel (see previous sec-
615 tion). The base model parameter estimates and their approximate asymptotic standard
616 errors are shown in Table ?? and the likelihood components are in Table ?. Estimates of
617 derived reference points and approximate 95% asymptotic confidence intervals are shown in
618 Table e. Time-series of estimated stock size over time are shown in Table ??.

619 ###Parameter Estimates

620 The additional survey variability (process error added directly to each year's input variabil-
621 ity) for all surveys was estimated within the model.

622 (Figure 7).

623 The stock-recruit curve ... Figure 8 with estimated recruitments also shown.

624 ###Fits to the Data Model fits to the indices of abundance, fishery length composition,
625 survey length composition, and conditional age-at-length observations are all discussed be-
626 low.

627 ###Uncertainty and Sensitivity Analyses A number of sensitivity analyses were conducted,
628 including:

629 1. Sensitivity 1

630 2. Sensitivity 2

631 3. Sensitivity 3

632 4. Sensitivity 4

633 5. Sensitivity 5, etc/

634 ###Retrospective Analysis

635 ###Likelihood Profiles

636 ###Reference Points Reference points were calculated using the estimated selectivities and
637 catch distribution among fleets in the most recent year of the model, (2017). Sustainable
638 total yield (landings plus discards) were 5,070 mt when using an $SPR_{50\%}$ reference harvest
639 rate and with a 95% confidence interval of 5,070 mt based on estimates of uncertainty. The
640 spawning biomass equivalent to 40% of the unfished level ($SB_{40\%}$) was 2,834 mt.

641 (Figure 12

642 The 2018 spawning biomass relative to unfished equilibrium spawning biomass is
643 above/below the target of 40% of unfished levels (Figure 13). The relative fishing intensity,
644 $(1 - SPR)/(1 - SPR_{50\%})$, has been xxx the management target for the entire time series
645 of the model.

646 Table e shows the full suite of estimated reference points for the base model and Figure 14
647 shows the equilibrium curve based on a steepness value xxx.

648 #Harvest Projections and Decision Tables The forecasts of stock abundance and yield were
649 developed using the final base model, with the forecasted projections of the OFL presented
650 in Table [g](#).

651 The forecasted projections of the OFL for each model are presented in Table [h](#).

653 #Research Needs There are a number of areas of research that could improve the stock
654 assessment for Big Skate. Below are issues identified by the STAT team and the STAR
655 panel:

- 656 1. xxxx:
- 657 2. xxxx:
- 658 3. xxxx:
- 659 4. xxxx:
- 660 5. xxxx:

661 #Acknowledgments

662 #Figures




Figure 1: Map showing the state boundary lines for management of the recreational fishing fleets | `fig:boundary_map`




Figure 2: Summary of data sources used in the model. fig:data_plot




Figure 3: Length comp data, aggregated across time by fleet. Labels ‘retained’ and ‘discard’ indicate discarded or retained sampled for each fleet. Panels without this designation represent the whole catch.
 fig:comp_lendat_aggregated_across_time




Figure 4: Estimated relationship between length and weight (left) and disc-width and length (right) for Big Skate. Colored points show observed values and the black line indicates the estimated relationship $W = 0.0000074924L^{2.9925}$. fig:weight-length




Figure 5: Estimated maturity relationship for female Big Skate. Gray points indicate average observed functional maturity within each length bin with point size proportional to the number of samples (indicated by text within each point).




Figure 6: Selectivity at length for all of the fleets in the base model. fig:sel01_multiple_fleets




Figure 7: Estimated time-series of recruitment for Big Skate. `fig:ts11_Age-0_recruits_(1`





Figure 8: Estimated recruitment (red circles) and the assumed stock-recruit relationship (black line) for Big Skate. The green line shows the effect of the bias correction for the lognormal distribution. fig:SR_curve2



Length (cm)

Figure continued from previous page




Figure 9: Conditional AAL plot, retained, Fishery_current (plot 1 of 2) These plots show mean age and std. dev. in conditional AAL. Left plots are mean AAL by size_class (obs. and pred.) with 90% CIs based on adding 1.64 SE of mean to the data. Right plots in each pair are SE of mean AAL (obs. and pred.) with 90% CIs based on the chi-square distribution. fig:mod1_4_co




Figure 10: Mean age from conditional data (aggregated across length bins) for WCGBTS with 95% confidence intervals based on current samples sizes. Francis data weighting method TA1.8: thinner intervals (with capped ends) show result of further adjusting sample sizes based on suggested multiplier (with 95% interval) for conditional age_at_length data from WCGBTS: 1.3806 (0.8289_39.92) For more info, see Francis, R.I.C.C. (2011). Data weighting in statistical fisheries stock assessment models. Can. J. Fish. Aquat. Sci. 68: 1124_1138. [fig:mod1_6_com




Figure 11: Conditional AAL plot, whole catch, WCGBTS (plot 1 of 2) These plots show mean age and std. dev. in conditional AAL. Left plots are mean AAL by size_class (obs. and pred.) with 90% CIs based on adding 1.64 SE of mean to the data. Right plots in each pair are SE of mean AAL (obs. and pred.) with 90% CIs based on the chi-square distribution. fig:mod1_7_co




Figure continued from previous page




Figure 12: Estimated spawning biomass (mt) with approximate 95% asymptotic intervals. fig:ts7_Spawn




Figure 13: Estimated spawning depletion with approximate 95% asymptotic intervals. fig:ts9_Spawni




Figure 14: Equilibrium yield curve for the base case model. Values are based on the 2018 fishery selectivity and with steepness fixed at 0.718. `fig:yield1_yield_curve`

667 #Appendix A. Detailed fits to length composition data {-}




Figure A15: Length comps, retained, Fishery_current. 'N adj.' is the input sample size after data-weighting adjustment. N eff. is the calculated effective sample size used in the McAllister-Lannelli tuning method.
 fig:mod1_1_comp_lenfit_fit1mkt2




Figure A16: Length comps, discard, Fishery_current. 'N adj.' is the input sample size after data-weighting adjustment. N eff. is the calculated effective sample size used in the McAllister-Lannelli tuning method.
 fig:mod1_2_comp_lenfit_fit1mkt1




Figure A17: Length comps, whole catch, WCG BTS. 'N adj.' is the input sample size after data-weighting adjustment. N eff. is the calculated effective sample size used in the McAllister-Lannelli tuning method.
 fig:mod1_3_comp_lenfit_fit5mkt0





Figure A18: Length comps, whole catch, Triennial. 'N adj.' is the input sample size after data-weighting adjustment. N eff. is the calculated effective sample size used in the McAllister-Lannelli tuning method. `fig:mod1_4_comp_lenfit_fit6mkt0`



Proportion

Length (cm)

Figure A19: Length comps, retained, IPHC. ‘N adj.’ is the input sample size after data_weighting adjustment. N eff. is the calculated effective sample size used in the McAlister-Iannelli tuning method.
 fig:mod1_5_comp_lenfit_fit7mkt2

References

references

- Batdorf, C. 1990. Northwest Native Harvest. Hancock House Publishers Ltd.; Surrey, B.C., Canada.
- Bizzarro, J. 2015. Comparative resource utilization of eastern north pacific skates (rajiformes: Rajidae) with applications for ecosystem-based fisheries management. WA: University of Washington.
- Bizzarro, J. 2019. National Marine Fisheries Service; UC Santa Cruz; Personal Communication, Santa Cruz, CA, USA.
- Bowers, G. M. 1909. Report of The Commissioner For the Year Ending June 30, 1909. Part XXVIII. Washington Printing Office.
- Bradburn, M.J. and Keller, A.A and Horness, B.H. 2011. The 2003 to 2008 US West Coast bottom trawl surveys of groundfish resources off Washington, Oregon, and California: estimates of distribution, abundance, length, and age composition. NOAA Technical Memorandum NMFS NOAA-TM-NMFS-NWFSC-114: 323 pp.
- Calavan, T. 2019. Oregon Department of Fisheries; Wildlife; Personal Communication, Newport, OR, USA.
- Castro-Aguirre, J., and Pérez, H. 1996. Catálogo sistemático de las rayas y especies afines de México: Chondrichthyes: Elasmobranchii: Rajiformes: Batoideomorpha. Unam.
- Castro-Aguirre, J., Schmitter, J., Balart, E., and Torres-Orozco, R. 1993. On the geographical distribution of some bentonics from the west coast of Baja California Sur, México, with ecological considerations y evolutivas. *In* Anales de la escuela nacional de ciencias biológicas, México. pp. 75–102.
- Chapman, W.M. 1944. The Latent Fisheries of Washington and Alaska. Washington State Department of Fisheries.
- Chiquillo, Kelcie L and Ebert, David A and Slager, Christina J and Crow, Karen D. 2014. The secret of the mermaid's purse: Phylogenetic affinities within the Rajidae and the evolution of a novel reproductive strategy in skates. *Molecular Phylogenetics and Evolution* **75**: 245–251. Elsevier.
- DeLacy, A.C., and Chapman, W.M. 1935. Notes on some elasmobranchs of Puget Sound, with descriptions of their egg cases. *Copeia* **1935**(2): 63–67. JSTOR.
- Ebert, D. 2003. Sharks, rays, and chimaeras of California. Univ of California Press.

- 699 Ebert, D.A., and Compagno, L.J. 2007. Biodiversity and systematics of skates (chondrichthyes: Rajiformes: Rajoidei). *In* Biology of skates. Springer. pp. 5–18.
- 701 Ebert, D.A., Smith, W.D., and Cailliet, G.M. 2008. Reproductive biology of two commercially exploited skates, (*Raja binoculata*) and (*Raja rhina*), in the western gulf of alaska. Fisheries Research **94**(1): 48–57. Elsevier.
- 704 Eschmeyer, W.N., and Herald, E.S. 1983. A field guide to pacific coast fishes: North america. Houghton Mifflin Harcourt.
- 706 Farrugia, T.J., Goldman, K.J., Tribuzio, C., and Seitz, A.C. 2016. First use of satellite tags to examine movement and habitat use of big skates (*Beringraja binoculata*) in the gulf of alaska. Marine Ecology Progress Series **556**: 209–221.
- 709 Ford, P. 1971. Differential growth rate in the tail of the pacific big skate, (*Raja binoculata*). Journal of the Fisheries Board of Canada **28**(1): 95–98. NRC Research Press.
- 711 Gburski, C.M. and Gaichas, S.K. and Kimura, D.K. 2007. Age and growth of big skate (*Raja binoculata*) and longnose skate (*Raja rhina*) in the Gulf of Alaska. *In* Biology of Skates. Springer, Dordrecht.
- 714 Gertseva, V. 2019. Manuscript in preparation.
- 715 Gunderson, Donald Raymond and Sample, Terrance M. 1980. Distribution and abundance of rockfish off Washington, Oregon and California during 1977. Northwest and Alaska Fisheries Center, National Marine Fisheries Service. Available from {<http://spo.nmfs.noaa.gov/mfr423-4/mfr423-42.pdf>}.
- 719 Hamel, Owen S. 2015. A method for calculating a meta-analytical prior for the natural mortality rate using multiple life history correlates. ICES Journal of Marine Science: Journal du Conseil **72**(1): 62–69. doi: {[10.1093/icesjms/fsu131](https://doi.org/10.1093/icesjms/fsu131)}.
- 722 Hitz, C. 1964. Observations on egg cases of the big skate (*Raja binoculata girard*) found in oregon coastal waters. Journal of the Fisheries Board of Canada **21**(4): 851–854. NRC Research Press.
- 725 Hoff, G.R. 2007. Reproductive biology of the alaska skate, bathyraja parmifera, with regard to nursery sites, embryo development, and predation. PhD thesis, University of Washington.
- 727 Ishihara, H., Treloar, M., Bor, P., Senou, H., and Jeong, C. 2012. The comparative morphology of skate egg capsules (Chondrichthyes: Elasmobranchii: Rajiformes). Bulletin of the Kanagawa Prefectural Museum (Natural Science) **41**: 9–25.
- 730 Keller, A.A. and Wallace, J.R. and Methot, R.D. 2017. The Northwest Fisheries Science Center's West Coast Groundfish Bottom Trawl Survey: History, Design, and Description. NOAA Technical Memorandum NMFS NOAA-TM-NMFS-NWFSC-136: 38 pp.

- King, J., and McFarlane, G. 2009. Biological results of the strait of georgia spiny dogfish (squalus acanthias) longline survey, october 10-22, 2008. Fisheries; Oceans Canada, Science Branch, Pacific Region.
- King, J.R., Surry, A.M., Garcia, S., and P.J. Starr. 2015. Big skate (*Raja binoculata*) and longnose skate (*Raja rhina*) stock assessments for British Columbia. Ottawa : Canadian Science Advisory Secretariat.
- Lippert, G. 2019. Washington Department of Fisheries; Wildlife; Personal Communication, Olympia, Washington, USA.
- Love, Milton S. 2011. Certainly more than you want to know about the fishes of the Pacific Coast: a postmodern experience. Really Big Press.
- Love, Milton S and Axell, Brita and Morris, Pamela and Collins, Robson and Brooks, Andrew. 1987. Life history and fishery of the California scorpionfish, *Scorpaena guttata*, within the Southern California Bight. Fishery Bulletin **85**: 99–116.
- McEachran, J., and Miyake, T. 1990. Zoogeography and bathymetry of skates (chondrichthyes, rajidae). Elasmobranchs as living resources. Advances in biology, Ecology, Systematics and the status of the fisheries: 305–326.
- McFarlane GA and King JR. 2006. Age and growth of big skate (*Raja binoculata*) and longnose skate (*Raja rhina*) in British Columbia waters. Fisheries Research **May 1 (2-3)**: 169–78.
- Mecklenburg, CW and Mecklenburg, TA and Thorsteinson, LK. 2002. Fishes of Alaska. American Fisheries Society, Bethesda, Maryland.
- Miller, B.S., Cross, J.N., Steinfert, S.N., Fresh, K.L., and Simenstad, C.A. 1980. Nearshore fish and macroinvertebrate assemblages along the strait of juan de fuca including food habits of the common nearshore fish.
- Thorson, James T. and Barnett, Lewis A. K. 2017. Comparing estimates of abundance trends and distribution shifts using single- and multispecies models of fishes and biogenic habitat. ICES Journal of Marine Science: Journal du Conseil: fsw193. doi: {10.1093/icesjms/fsw193}.
- Thorson, J. T. and Shelton, A. O. and Ward, E. J. and Skaug, H. J. 2015. Geostatistical delta-generalized linear mixed models improve precision for estimated abundance indices for West Coast groundfishes. ICES Journal of Marine Science **72(5)**: 1297–1310. doi: {10.1093/icesjms/fsu243}.
- von Bertalanffy, L. 1938. A quantitative theory of organic growth. Human Biology **10**: 181–213.

767 Zeiner, S.J. and P. Wolf. 1993. Growth characteristics and estimates of age at maturity of
768 two species of skates (*Raja binoculata*) and (*Raja rhina*) from Monterey Bay, California.