

Integrated Course Ib

Theoretical quantum mechanics

lecture by

DR. JUAN-DIEGO URBINA

during the summer semester 2012

revision and layout in LyX by

ANDREAS VÖLKLEIN



Last changed: May 25, 2012

ATTENTION

This script does *not* replace the lecture.

Therefore it is recommended *strongly* to attend the lecture.

These notes are under constant revision, and the level of rigor is not uniform.

Copyright Notice

Copyright © 2012 ANDREAS VÖLKLEIN

Permission is granted to copy, distribute and/or modify this document under the terms of the GNU Free Documentation License, Version 1.3 or any later version published by the Free Software Foundation;

with no Invariant Sections, no Front-Cover Texts, and no Back-Cover Texts.

A copy of the license is included in the section entitled “GNU Free Documentation License”.

Disclaimer of Warranty

UNLESS OTHERWISE MUTUALLY AGREED TO BY THE PARTIES IN WRITING AND TO THE EXTENT NOT PROHIBITED BY APPLICABLE LAW, **THE COPYRIGHT HOLDERS AND ANY OTHER PARTY, WHO MAY DISTRIBUTE THE DOCUMENT AS PERMITTED ABOVE, PROVIDE THE DOCUMENT “AS IS”, WITHOUT WARRANTY OF ANY KIND**, EXPRESSED, IMPLIED, STATUTORY OR OTHERWISE, INCLUDING, BUT NOT LIMITED TO, THE IMPLIED WARRANTIES OF MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, NON-INFRINGEMENT, THE ABSENCE OF LATENT OR OTHER DEFECTS, ACCURACY, OR THE ABSENCE OF ERRORS, WHETHER OR NOT DISCOVERABLE.

Limitation of Liability

IN NO EVENT UNLESS REQUIRED BY APPLICABLE LAW OR AGREED TO IN WRITING WILL THE COPYRIGHT HOLDERS, OR ANY OTHER PARTY, WHO MAY DISTRIBUTE THE DOCUMENT AS PERMITTED ABOVE, BE LIABLE TO YOU FOR ANY DAMAGES, INCLUDING, BUT NOT LIMITED TO, ANY GENERAL, SPECIAL, INCIDENTAL, CONSEQUENTIAL, PUNITIVE OR EXEMPLARY DAMAGES, HOWEVER CAUSED, REGARDLESS OF THE THEORY OF LIABILITY, ARISING OUT OF OR RELATED TO THIS LICENSE OR ANY USE OF OR INABILITY TO USE THE DOCUMENT, EVEN IF THEY HAVE BEEN ADVISED OF THE POSSIBILITY OF SUCH DAMAGES.

IN NO EVENT WILL THE COPYRIGHT HOLDERS’/DISTRIBUTOR’S LIABILITY TO YOU, WHETHER IN CONTRACT, TORT (INCLUDING NEGLIGENCE), OR OTHERWISE, EXCEED THE AMOUNT YOU PAID THE COPYRIGHT HOLDERS/DISTRIBUTOR FOR THE DOCUMENT UNDER THIS AGREEMENT.

Links

The text of the “GNU Free Documentation License” can also be read on the following site:

<https://www.gnu.org/licenses/fdl-1.3.en.html>

A transparent copy of the recent version of this document can be downloaded from:

<https://github.com/andiv/IK1b>

Literature

- JUN J. SAKURAI, JIM NAPOLITANO: *Modern quantum mechanics*, Addison-Wesley, 2011
ISBN: 0-321-50336-8
- LEONARD I. SCHIFF: *Quantum mechanics*, McGraw-Hill, 1968
ISBN: 0-07-085643-5
- Any introductory book on quantum mechanics.
- RICHARD P. FEYNMAN, ROBERT B. LEIGHTON, MATTHEW L. SANDS: *Lectures on physics*

Contents

1	The Formalism and its interpretation	1
1.1	The kinematical aspects of quantum mechanics	1
1.2	Observables and measurements in quantum mechanics	3
1.3	What does quantum actually predict?	8
1.4	The probabilistic interpretation of quantum mechanics (Born's rule)	9
1.5	“Quantization”:	
	Introducing the “fundamental” or basic observables	11
1.5.1	General approach	11
1.5.2	Algebraic properties of \hat{q} and \hat{p}	12
1.5.3	The Schrödinger equation in position and momentum representation	13
1.5.4	The Heisenberg and the Schrödinger “pictures”	14
2	Basic applications	16
2.1	Boxes	16
2.2	Square well	16
2.3	The harmonic oscillator (in one dimension)	17
2.3.1	The number representation	20
2.3.2	The dynamics of the harmonic oscillator	22
2.4	The tunneling effect	23
2.5	Two formal consequences of the position representation of the Schrödinger equation	25
2.5.1	Oscillation theorem (1D, bounded)	25
2.5.2	The continuity equation	25
3	Symmetries, Rotations and Angular momentum	27
3.1	The tensor product in quantum mechanics	27
3.2	Rotations and angular momentum	28
3.3	Eigenstates and eigenvalues of \hat{j}	30
3.4	Irreducible representations of the rotation operators	32
3.5	Orbital angular momentum	32
3.5.1	General considerations	32
3.5.2	Eigenstates in the position representation	33
3.5.3	Relation with the kinetic energy	35
3.5.4	Central forces	36
3.6	Addition of angular momentum	36
3.6.1	Formal theory (2 “particles”)	37
3.6.2	The formal theory for many particles	38
3.6.3	Example: Two spin $\frac{1}{2}$ particles	38

4	Perturbation theory (time-independent)	40
5	Perturbation theory (time-dependent)	41
6	Many-particle systems	42
7	Scattering theory	43
Appendix		45
	Danksagungen	45
	GNU Free Documentation License	46

1 The Formalism and its interpretation

As any theory about physical phenomenons, quantum mechanics requires

- i) kinematical aspects: The (mathematical) space where physical states are represented.
Example from classical mechanics: points in phase space
- ii) the definition of observables: Which quantities can be measured and how to represent them in the space of states?
Example from classical mechanics: any function $f(\vec{r}, \vec{p})$
(This is more complicated in quantum mechanics.)
- iii) an dynamical law: How do states evolve in “time”?
Example from classical mechanics: Hamilton’s equations determine $(\vec{q}(t), \vec{p}(t))$ depending on $(\vec{q}(t_0), \vec{p}(t_0))$.

1.1 The kinematical aspects of quantum mechanics

The FIRST POSTULATE says:

“The state of a quantum mechanical system is represented as a normalized vector in a (complex) Hilbert space.

Vectors, which differ only by a phase, represent the same state.”

To understand what this means, we introduce some concepts:

- a) *Hilbert space*: A (finite or infinite dimensional) complete vector space with a positive definite scalar product. Following Dirac, elements in the Hilbert space \mathcal{H} are called *kets* and we represent them by $|\psi\rangle, |\phi\rangle \in \mathcal{H}$.

For complex numbers $a, b \in \mathbb{C}$ also $a|\psi\rangle + b|\phi\rangle \in \mathcal{H}$ is a element of the Hilbert space.

The sum is associative and the product by scalars is distributive.

- b) *Scalar product*: The operation associating a complex number to each pair of states

$$\langle \cdot, \cdot \rangle : \mathcal{H} \times \mathcal{H} \rightarrow \mathbb{C}$$

$$(|\psi\rangle, |\phi\rangle) \mapsto \langle \psi, \phi \rangle$$

with the properties

$$\begin{aligned} \langle \eta, a\phi + b\psi \rangle &= a \langle \eta, \phi \rangle + b \langle \eta, \psi \rangle & \langle \psi, \phi \rangle &= \langle \phi, \psi \rangle^* \\ \langle \psi, \psi \rangle &\geq 0 & \langle \psi, \psi \rangle = 0 &\Leftrightarrow |\psi\rangle = 0 \end{aligned}$$

for $a, b \in \mathbb{C}$ and $|\phi\rangle, |\psi\rangle, |\eta\rangle \in \mathcal{H}$ is called the scalar product.

The *norm* of a state $|\psi\rangle$ is then defined as:

$$\|\psi\| := \| |\psi\rangle \| := \sqrt{\langle \psi, \psi \rangle}$$

If $\| |\psi\rangle \| = 1$, then $|\psi\rangle$ is said to be *normalized*.

- c) *Phase*: A Complex number $z \in \mathbb{C}$ with unit norm, that means $zz^* = 1$. It can be represented via a real number $\alpha \in \mathbb{R}$ as $z = e^{i\alpha}$.

The physical state associated with $|\psi\rangle$ and $e^{i\alpha} |\psi\rangle$ is THE SAME.

- d) *Complete basis*: A family of kets $(|\phi_i\rangle)_{i \in I \subseteq \mathbb{N}}$ such that for ALL states $|\psi\rangle \in \mathcal{H}$, there is a family of complex numbers $(c_i)_{i \in I \subseteq \mathbb{N}}$ (depending on $|\psi\rangle$) with:

$$|\psi\rangle = \sum_{i \in I} c_i |\phi_i\rangle$$

If $\langle \phi_i, \phi_j \rangle = \delta_{ij}$, then the basis is *complete* and *orthonormal*.

In the case of an uncountably infinite vector space the basis $(|q\rangle)_{q \in \mathbb{R}}$ can be written as a function of a real variable. The representation of $|\psi\rangle$ then is an infinite sum, that is an integral

$$|\psi\rangle = \int \psi(q) |q\rangle dq$$

and a complete and orthonormal basis is characterized by $\langle q, q' \rangle = \delta(q - q')$.

- e) *Adjoint*: For each Hilbert space \mathcal{H} there is another Hilbert space \mathcal{H}^* called *dual*, with elements $\langle f | \in \mathcal{H}^*$, which are LINEAR FUNCTIONALS acting on \mathcal{H} :

$$\begin{aligned} \langle f | : \mathcal{H} &\rightarrow \mathbb{C} \\ |\psi\rangle &\mapsto \langle f | \psi \rangle \end{aligned}$$

$\langle \psi | \phi \rangle$ is called *bracket*. Riesz theorem says, that there is a ONE TO ONE correspondence between \mathcal{H} and \mathcal{H}^* :

$$\forall_{|\psi\rangle \in \mathcal{H}} \quad \exists_{\langle \psi | \in \mathcal{H}^*} \quad \forall_{|\phi\rangle \in \mathcal{H}} : \langle \psi | \phi \rangle = \langle \psi, \phi \rangle, \quad \|\langle \psi | \| = \| |\psi\rangle \|$$

The so associated $\langle \psi |$ is the *adjoint* of $|\psi\rangle$, called *bra*, and we write:

$$\begin{aligned} \dagger : \mathcal{H} &\rightarrow \mathcal{H}^* \\ |\psi\rangle &\mapsto (|\psi\rangle)^\dagger = \langle \psi | \end{aligned}$$

The function \dagger is semilinear, that means for $a, b \in \mathbb{C}$ and $|\psi\rangle, |\phi\rangle \in \mathcal{H}$ is:

$$(a |\psi\rangle + b |\phi\rangle)^\dagger = a^* \langle \psi | + b^* \langle \phi |$$

- f) *Representation*: Assume that $(|\phi_i\rangle)_{i \in I \subseteq \mathbb{N}}$ (respectively $(|q\rangle)_{q \in \mathbb{R}}$) is a complete orthonormal basis. The ket $|\psi\rangle$ is said to “be represented” in that basis by associating:

discrete case	continous case
$ \psi\rangle \mapsto \begin{pmatrix} \langle \phi_1 \psi \rangle \\ \langle \phi_2 \psi \rangle \\ \vdots \end{pmatrix} \in \mathbb{C}^{ I }$	$ \psi\rangle \mapsto \langle q \psi \rangle =: \psi(q)$
$\langle \psi \mapsto \begin{pmatrix} \langle \phi_1 \psi \rangle^* \\ \langle \phi_2 \psi \rangle^* \\ \vdots \end{pmatrix} \in \mathbb{C}^{ I }$	$\langle \psi \mapsto \langle q \psi \rangle^* = \psi(q)^*$

This $\psi(q)$ is the wave function.

- g) *Exterior product*: The object $|\phi\rangle \langle \psi| : \mathcal{H} \rightarrow \mathcal{H}$ is a linear operator acting on $|\eta\rangle \in \mathcal{H}$ defined by

$$(|\phi\rangle \langle \psi|) |\eta\rangle := \underbrace{\langle \psi | \eta \rangle}_{\in \mathbb{C}} |\phi\rangle$$

with the adjoint:

$$(|\phi\rangle \langle \psi|)^\dagger := |\psi\rangle \langle \phi|$$

Example from linear algebra:

$$\vec{v} = (v_1, \dots, v_n)^T \qquad \vec{u} = (u_1, \dots, u_n)^T$$

$$\vec{v}^T \vec{u} = \sum_{i=1}^n v_i u_i$$

$$\vec{u} \vec{v}^T = \begin{pmatrix} u_1 v_1 & \dots & u_1 v_n \\ \vdots & \ddots & \vdots \\ u_n v_1 & \dots & u_n v_n \end{pmatrix}$$

1.2 Observables and measurements in quantum mechanics

What can be observed and how measurements affect quantum states is encoded into the following two POSTULATES:

“Observables in quantum mechanics are represented by LINEAR HERMITIAN OPERATORS on \mathcal{H} .”

“The results of a measurement of the physical quantity represented by an observable can only take values belonging to the SPECTRUM of the observable. Just after measurement, that gives one of the eigenvalues of the observable, the state belongs to corresponding eigenspace.”

- a) $\hat{A} : \mathcal{H} \rightarrow \mathcal{H}$ is a *linear* operator if and only if:

$$\hat{A}(a|\psi\rangle + b|\phi\rangle) = a\hat{A}|\psi\rangle + b\hat{A}|\phi\rangle$$

For every linear operator \hat{A} there is the identity:

$$\langle\psi|\hat{A}\phi\rangle = \langle\phi|\hat{A}^\dagger\psi\rangle^*$$

\hat{A}^\dagger is called the *adjoint* of \hat{A} . If and only if $\hat{A} = \hat{A}^\dagger$ then \hat{A} is called HERMITIAN.

- b) The *spectrum* of an operator \hat{Q} is defined by a set of numbers Q_i , called *eigenvalues*, that fulfill the equation:

$$\hat{Q}|Q_i\rangle = Q_i|Q_i\rangle$$

The $|Q_i\rangle$ are the corresponding eigenvectors. If there is more than one eigenvector for the same eigenvalue then the spectrum is called *degenerated* and the different eigenvectors are denoted by $|Q_i^{(d)}\rangle$ with $d \in D \subseteq \mathbb{N}$.

Hermitian operators have REAL eigenvalues and ORTHOGONAL eigenvectors.

That means, if $\hat{A} = \hat{A}^\dagger$, $\hat{A}|a_i\rangle = a_i|a_i\rangle$, then $a_i = a_i^*$ and $\langle a_i|a_j\rangle = 0$ if $i \neq j$.

If the eigenvectors are normalized, then we can write $\langle a_i|a_j\rangle = \delta_{ij}$.

- c) The *spectrum decomposition* of a Hermitian operator ($\hat{A} = \hat{A}^\dagger$) with

discrete case	continous case
$\hat{A} a_i\rangle = a_i a_i\rangle$	$\hat{A} a\rangle = a a\rangle$

is given by:

$$\hat{A} = \sum_{i \in I} a_i |a_i\rangle \langle a_i| \qquad \hat{A} = \int a |a\rangle \langle a| da$$

In linear algebra a symmetric matrix $M = M^T$ with eigenvalues m_i for $i \in \{1, \dots, n\}$ can be written as:

$$M = \begin{pmatrix} m_1 & 0 & \dots & 0 \\ 0 & m_2 & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \dots & 0 & m_n \end{pmatrix} = m_1 \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 0 & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \dots & 0 & 0 \end{pmatrix} + \dots + m_n \begin{pmatrix} 0 & 0 & \dots & 0 \\ 0 & \ddots & \ddots & \vdots \\ \vdots & \ddots & 0 & 0 \\ 0 & \dots & 0 & 1 \end{pmatrix}$$

- d) The *unit operator* \hat{I} defined by

$$\forall_{|\phi\rangle \in \mathcal{H}} : \hat{I}|\phi\rangle = |\phi\rangle$$

can be written as

discrete case	continous case
$\hat{I} = \sum_{i \in I} a_i\rangle \langle a_i $	$\hat{I} = \int a\rangle \langle a da$

where $(|a_i\rangle)_{i \in I \subseteq \mathbb{N}}$ (respectively $(|a\rangle)_{a \in \mathbb{R}}$) is a complete orthonormal basis.

e) The *projection* of $|\psi\rangle$ along a basis vector $|a_i\rangle$ (respectively $|a\rangle$) is:

discrete case	continous case
$(a_i\rangle \langle a_i) \psi\rangle = \langle a_i \psi \rangle a_i\rangle$	$(a\rangle \langle a) \psi\rangle = \langle a \psi \rangle a\rangle$

f) The *algebra* of the operators \hat{Q}_1 and \hat{Q}_2 is:

$$\begin{aligned} (\hat{Q}_1 \hat{Q}_2) |\psi\rangle &:= \hat{Q}_1 (\hat{Q}_2 |\psi\rangle) \neq (\hat{Q}_2 \hat{Q}_1) |\psi\rangle \\ (\hat{Q}_1 \hat{Q}_2)^\dagger &:= \hat{Q}_2^\dagger \hat{Q}_1^\dagger \end{aligned}$$

The *commutator* is defined by:

$$[\hat{Q}_1, \hat{Q}_2] := \hat{Q}_1 \hat{Q}_2 - \hat{Q}_2 \hat{Q}_1$$

It is bilinear and has the properties:

i) It is antisymmetric:

$$[\hat{Q}_1, \hat{Q}_2] = - [\hat{Q}_2, \hat{Q}_1]$$

ii) Jacobi identity:

$$[\hat{Q}_1, [\hat{Q}_2, \hat{Q}_3]] + [\hat{Q}_2, [\hat{Q}_3, \hat{Q}_1]] + [\hat{Q}_3, [\hat{Q}_1, \hat{Q}_2]] = 0$$

iii) Leibniz identity:

$$[\hat{Q}_1, \hat{Q}_2 \hat{Q}_3] = \hat{Q}_2 [\hat{Q}_1, \hat{Q}_3] + [\hat{Q}_1, \hat{Q}_2] \hat{Q}_3$$

Due to these identities the commutator is analogous to the classical Poisson bracket.

Two observables $(\hat{A} = \hat{A}^\dagger, \hat{B} = \hat{B}^\dagger)$ are called *compatible* if and only if $[\hat{A}, \hat{B}] = 0$.

Theorem: If \hat{A} and \hat{B} are compatible then there exists a basis $(|k_i\rangle)_{i \in I \subseteq \mathbb{N}}$ such that:

$$\begin{aligned} \hat{A} |k_i\rangle &= a_i |k_i\rangle \\ \hat{B} |k_i\rangle &= b_i |k_i\rangle \end{aligned}$$

Therefore the states $|k_i\rangle$ have well defined properties (a_i, b_i) .

g) i) The *matrix representation* of an operator WITH RESPECT TO THE BASIS $(|a_i\rangle)_{i \in \{1, \dots, n\}}$ (respectively $(|q\rangle)_{q \in \mathbb{R}}$) is:

discrete case	continuous case
$\hat{Q} \rightarrow \begin{pmatrix} \langle a_1 \hat{Q} a_1 \rangle & \dots & \langle a_1 \hat{Q} a_n \rangle \\ \vdots & \ddots & \vdots \\ \langle a_n \hat{Q} a_1 \rangle & \dots & \langle a_n \hat{Q} a_n \rangle \end{pmatrix}$ $= (\langle a_i \hat{Q} a_j \rangle)_{ij} =: Q_{ij}$	$\hat{Q} \rightarrow \langle q \hat{Q} q' \rangle = Q(q, q')$

ii) The *trace* of \hat{Q} is defined by

$$\begin{array}{cc} \text{discrete case} & \text{continuous case} \\ \text{Tr}(\hat{Q}) = \sum_{i=1}^n \langle a_i | \hat{Q} | a_i \rangle & \text{Tr}(\hat{Q}) = \int Q(q, q) dq \end{array}$$

ans is independent of the basis.

Therefore the equation

$$\hat{Q} |\psi\rangle = |\phi\rangle$$

can be written in this basis by:

i) Multiplying $\langle a_i |$ from the left: $\langle a_i | \hat{Q} | \phi \rangle = \langle a_i | \phi \rangle$

ii) Insert an unit operator after \hat{Q} :

$$\langle a_i | \hat{Q} \hat{I} | \phi \rangle = \left\langle a_i \left| \hat{Q} \left(\sum_{j=1}^n |a_j\rangle \langle a_j| \right) \right| \phi \right\rangle = \sum_{j=1}^n \langle a_i | \hat{Q} | a_j \rangle \langle a_j | \phi \rangle = \langle a_i | \phi \rangle$$

This is the component form of the vector equation:

$$\begin{pmatrix} Q_{11} & \cdots & Q_{1n} \\ \vdots & & \vdots \\ Q_{n1} & \cdots & Q_{nn} \end{pmatrix} \begin{pmatrix} \psi_1 \\ \vdots \\ \psi_n \end{pmatrix} = \begin{pmatrix} \phi_1 \\ \vdots \\ \phi_n \end{pmatrix}$$

The continuous version is

$$\int Q(q, q') \psi(q') dq' = \phi(q)$$

with $Q(q, q') := \langle q | \hat{Q} | q' \rangle$, $\psi(q') = \langle q' | \psi \rangle$ and $\phi(q) = \langle q | \phi \rangle$.

h) Change of basis and unitary transformations.

i) An operator \hat{U} is called *unitary* if and only if $\hat{U}^T = \hat{U}^{-1}$.

ii) Unitary transformations (“transformation” is the same as “operator”) preserve scalar products. For $|\psi\rangle, |\phi\rangle \in \mathcal{H}$ we define:

$$|\psi'\rangle := \hat{U} |\psi\rangle \qquad |\phi'\rangle := \hat{U} |\phi\rangle$$

Then the scalar product is the same:

$$\langle \psi' | \phi' \rangle = \langle \psi | \phi \rangle$$

iii) Unitary transformations represent changes of the basis.

$$\begin{aligned} |\psi\rangle &= \hat{I} |\psi\rangle = \left(\sum_i |a_i\rangle \langle a_i| \right) |\psi\rangle = \sum_{i \in I} \langle a_i | \psi \rangle |a_i\rangle \\ \langle b_i | \psi \rangle &= \sum_{j \in I} \langle b_i | a_j \rangle \langle a_j | \psi \rangle \end{aligned}$$

$$\begin{pmatrix} \psi_1^{(b)} \\ \vdots \\ \psi_N^{(b)} \end{pmatrix} = \mathbb{U} \cdot \begin{pmatrix} \psi_1^{(a)} \\ \vdots \\ \psi_N^{(a)} \end{pmatrix}$$

$$[\mathbb{U}]_{ij} = U_{ij} = \langle b_i | a_j \rangle$$

iv) For operators the transformations are:

$$\begin{aligned} Q_{ij}^{(b)} &= \langle b_i | \hat{Q} | b_j \rangle = \langle b_i | \hat{I} \hat{Q} \hat{I} | b_j \rangle = \sum_{l,m \in I} \langle b_i | a_l \rangle \langle a_l | \hat{Q} | a_m \rangle \langle a_m | b_j \rangle = \\ &= \sum_{l,m \in I} U_{il} Q_{lm}^{(a)} U_{mj} = \sum_{l,m} U_{il} Q_{lm}^{(a)} (U_{jm})^* \\ Q^{(b)} &= \mathbb{U} Q^{(a)} \mathbb{U}^\dagger \end{aligned}$$

i) One can describe functions mapping one operator to another by their Taylor series:

$$\begin{aligned} f(x) &= \sum_{n=0}^{\infty} c_n x^n \\ c_n &= \frac{1}{n!} \partial_x^n f(x) \big|_{x=0} \\ f(\hat{Q}) &:= \sum_{n=0}^{\infty} c_n \hat{Q}^n \end{aligned}$$

This is very nice. But this does not work for more than one operator, if they do not commute.

$$\begin{aligned} f(x,y) &= \sum_{n,m} c_{n,m} x^n y^m \\ f(\hat{Q}_1, \hat{Q}_2) &\neq \sum_{n,m} c_{n,m} \hat{Q}_1^n \hat{Q}_2^m \\ xy &\overset{?}{\rightarrow} \begin{cases} \hat{Q}_1 \hat{Q}_2 \\ \hat{Q}_2 \hat{Q}_1 \\ \frac{1}{2} (\hat{Q}_1 \hat{Q}_2 + \hat{Q}_2 \hat{Q}_1) \end{cases} \end{aligned}$$

In a basis of eigenvectors $|q_i\rangle$, that means $\hat{Q} |q_i\rangle = q_i |q_i\rangle$, one can write:

$$\begin{array}{cc} \text{discrete case} & \text{continuous case} \\ f(\hat{Q}) = \sum_i f(q_i) |q_i\rangle \langle q_i| & f(\hat{Q}) = \int f(q) |q\rangle \langle q| dq \end{array}$$

j) An operator can also depend on parameters:

$$\mathbb{R} \ni t \mapsto \hat{A}(t) \in \text{End}_{\mathbb{C}}(\mathcal{H})$$

It has all the nice properties, but one has to be careful not to change the ordering of the operators, if they do not commute:

$$\frac{d}{dt} (\hat{A}(t) \hat{B}(t)) = \left(\frac{d\hat{A}(t)}{dt} \right) \hat{B}(t) + \hat{A}(t) \left(\frac{d\hat{B}(t)}{dt} \right)$$

$$\frac{d\hat{A}(t)}{dt} = \hat{B}(t) \quad \Rightarrow \quad \hat{A}(t) = \int_0^t \hat{B}(t') dt' + \hat{A}_0$$

Extra property:

$$\frac{d}{dt} (\langle \psi(t) | \phi(t) \rangle) = \left(\frac{d\langle \psi(t) |}{dt} \right) | \phi(t) \rangle + \langle \psi(t) | \left(\frac{d| \phi(t) \rangle}{dt} \right)$$

As norm one usually uses:

$$\|\hat{A}\|^2 = \text{Tr}(\hat{A}\hat{A}^\dagger)$$

1.3 What does quantum actually predict?

Think about classical mechanics:

$$(x(0), p(0)) \xrightarrow{\text{evolution}} (x(t), p(t)) \rightarrow f(x(t), p(t)) \text{ is known}$$

So classical mechanics is DETERMINISTIC (the state of a system is determined by the initial conditions) and REALISTIC (the value you are going to measure already exists before the actual measurement).

Back to quantum mechanics:

The full dynamics is encoded in the postulate:

“In each quantum mechanical system there exists an observable $\hat{H}(t)$, called *Hamiltonian* which is assumed to be BOUNDED from below (and it is a nice operator). The solution of the equation

$$i\hbar \partial_t \hat{U}(t, t_0) = \hat{H}(t) \hat{U}(t, t_0) \quad \hat{U}(t_0, t_0) = \hat{I} \quad \text{for } t > t_0$$

is an unitary operator \hat{U} , called TIME EVOLUTION OPERATOR. The time evolution of an initial state $|\psi(t_0)\rangle$ is given by $|\psi(t)\rangle = \hat{U}(t, t_0) |\psi(t_0)\rangle$.”

Comments:

- a) If $\partial_t \hat{H}(t) = 0$, the one can directly solve the Schrödinger equation:

$$\hat{U}(t, t_0) = e^{-\frac{i}{\hbar} \hat{H}(t-t_0)} \quad \text{for } t > t_0 \quad (1.1)$$

- b) \hat{U} has a SEMIGROUP property:

$$\hat{U}(t, t') \hat{U}(t', t_0) = \hat{U}(t, t_0) \quad t > t' > t_0$$

Semigroup means:

$$\hat{U}(t, t_0)^{-1} = \hat{U}(t, t_0)^\dagger \neq \hat{U}(t_0, t)$$

- c) Constructing $\hat{H}(t)$ is the main step to define a quantum mechanical system, at the end it relies on experimental verification.

d) The formal solution of (1.1) is found by iteration:

$$\begin{aligned}
 \hat{U}(t, t_0) &= \hat{I} - \frac{i}{\hbar} \int_{t_0}^t \hat{H}(t') \hat{U}(t', t_0) dt' & t > t' \\
 &= \hat{I} - \frac{i}{\hbar} \int_{t_0}^t \hat{H}(t') \left(\hat{I} - \frac{i}{\hbar} \int_{t_0}^{t'} \hat{H}(t'') \hat{U}(t'', t_0) dt'' \right) dt' = \\
 &= \hat{I} - \frac{i}{\hbar} \int_{t_0}^t \hat{H}(t') dt' + \left(\frac{-i}{\hbar} \right)^2 \int_{t_0}^t \int_{t_0}^{t'} \hat{H}(t') \hat{H}(t'') \hat{U}(t'', t_0) dt'' dt' = \\
 &= \tau e^{\frac{-i}{\hbar} \int_{t_0}^t \hat{H}(t') dt'}
 \end{aligned}$$

τ is the TIME ORDERING OPERATOR.

$$\tau \left(\hat{H}(t) \hat{H}(t') \right) = \begin{cases} \hat{H}(t) \hat{H}(t') & t > t' \\ \hat{H}(t') \hat{H}(t) & t < t' \end{cases}$$

e) Stationary states: Assume $\partial_t \hat{H}(t) = 0$, then the eigenstates of \hat{H} are defined by the time-independent Schrödinger equation:

$$\hat{H} |e_n\rangle = e_n |e_n\rangle$$

Their time evolution is:

$$|e_n(t)\rangle = e^{-\frac{i}{\hbar} \hat{H} t} |e_n\rangle = \underbrace{e^{-\frac{i e_n}{\hbar} t}}_{\text{phase}} |e_n\rangle = e^{-i \omega_n t} |e_n\rangle$$

For an arbitrary initial state $|\psi(t_0)\rangle$ we can calculate the final state:

$$\begin{aligned}
 |\psi(t)\rangle &= e^{-\frac{i}{\hbar} \hat{H} \cdot (t-t_0)} |\psi(t_0)\rangle = e^{-\frac{i}{\hbar} \hat{H} \cdot (t-t_0)} \hat{I} |\psi(t_0)\rangle = \\
 &= \sum_{n=0}^{\infty} \langle e_n | \psi(t_0) \rangle e^{-i \omega_n \cdot (t-t_0)} |e_n\rangle
 \end{aligned}$$

f) Instead of finding \hat{U} we can solve the SCHRÖDINGER EQUATION directly for $|\psi(t)\rangle$:

$$\begin{aligned}
 i\hbar \partial_t |\psi(t)\rangle &= \hat{H}(t) |\psi(t)\rangle \\
 |\psi(t=t_0)\rangle &= |\psi(t_0)\rangle
 \end{aligned}$$

1.4 The probabilistic interpretation of quantum mechanics (Born's rule)

“If the states of a system at time t is given by the normalized state $|\psi(t)\rangle$, then the PROBABILITY to get the outcome a_i when the observable \hat{A} is measured is given by:”

discrete case	continuous case
$P(a_i, t) = \langle a_i \psi(t) \rangle ^2$	$P(q, t) = \langle q \psi(t) \rangle ^2 dq$

a)

discrete case

$$\sum_i P(a_i, t) = 1$$

continuous case

$$\int P(q, t) dq = 1$$

$\langle a_i | \psi(t) \rangle$ is called probability Amplitude, $P(q, t)$ is the density of probability and the probability to obtain an outcome between q and $q + dq$ is $|\langle q | \psi(t) \rangle|^2$.


b) The expectation value is given by:

$$\begin{aligned} \bar{A} &:= \sum_i a_i P(a_i, t) = \sum_i a_i |\langle a_i | \psi(t) \rangle|^2 = \sum_i a_i \langle a_i | \psi(t) \rangle \cdot \langle a_i | \psi(t) \rangle^* = \\ &= \sum_i a_i \langle \psi(t) | a_i \rangle \cdot \langle a_i | \psi(t) \rangle = \langle \psi(t) | \left(\sum_i a_i |a_i\rangle \cdot \langle a_i| \right) | \psi(t) \rangle = \\ &= \langle \psi(t) | \hat{A} | \psi(t) \rangle =: \langle \hat{A} \rangle \end{aligned}$$

$$\begin{aligned} \text{Var} \hat{A} &= \langle \hat{A}^2 \rangle - (\langle \hat{A} \rangle)^2 = \langle \psi(t) | \hat{A}^2 | \psi(t) \rangle - \langle \psi(t) | \hat{A} | \psi(t) \rangle^2 = \\ &= \langle (\hat{A} - \hat{I} \langle \hat{A} \rangle)^2 \rangle = \sum (a_i - \bar{a}_i)^2 P(a_i, t) = (\Delta \hat{A})^2 \end{aligned}$$

The only case, one gets

$$\langle (\hat{A} - \langle \hat{A} \rangle)^2 \rangle = 0$$

is, if $|\psi(t)\rangle = |a_i\rangle$.

c) Consider the eigenstates of the Hamiltonian:

$$\hat{H} |e_n\rangle = e_n |e_n\rangle$$

If at $t = 0$ the state is $|e_n\rangle$, then the time evolution is very simple:

$$\begin{aligned} |e_n(t)\rangle &= e^{-i\omega_n t} |e_n\rangle \\ \Rightarrow \langle e_n(t) | \hat{A} | e_n(t) \rangle &= \langle e_n | \hat{A} | e_n \rangle \\ \partial_t \langle \hat{A} \rangle &= 0 \end{aligned}$$

d) Define $|\eta\rangle = N(|\psi\rangle + |\phi\rangle)$. What is the probability $P_\eta(a_i)$?

$$\begin{aligned} |\langle a_i | \eta \rangle|^2 &= N^2 |\langle a_i | \psi \rangle + \langle a_i | \phi \rangle|^2 = \\ &= N^2 \left(|\langle a_i | \psi \rangle|^2 + |\langle a_i | \phi \rangle|^2 + 2\text{Re}(\langle a_i | \psi \rangle \langle a_i | \phi \rangle^*) \right) = \\ &= N^2 (P_\psi(a_i) + P_\phi(a_i) + \text{something}) \end{aligned}$$

Therefore it is not enough to know the probabilities, one also has to know the phase of the states.

1.5 “Quantization”:

Introducing the “fundamental” or basic observables

1.5.1 General approach

Historically, it all started with the Hydrogen atom. (In this system the Hilbert space \mathcal{H} is infinite-dimensional.) It is called “canonical” quantization.

- i) $\dim \mathcal{H} = \infty$ means, there can be discrete as well as continuous eigenvalues.
- ii) Every quantum mechanical observable is a function of two basic observables, the position operator \hat{q}_α and the momentum operator \hat{p}_α , satisfying:

$$[\hat{q}_\alpha, \hat{p}_\beta] = i\hbar \delta_{\alpha\beta} \hat{I} \qquad [\hat{q}_\alpha, \hat{q}_\beta] = 0 = [\hat{p}_\alpha, \hat{p}_\beta]$$

Therefore you can define an eigenbasis $|\vec{q}\rangle$ with:

$$\hat{q}_\alpha |\vec{q}\rangle = q_\alpha |\vec{q}\rangle \qquad \hat{p}_\alpha |\vec{p}\rangle = p_\alpha |\vec{p}\rangle \qquad q_\alpha, p_\alpha \in \mathbb{R}$$

$$\langle \vec{q} | \vec{q}' \rangle = \delta(\vec{q} - \vec{q}') \qquad \langle \vec{p} | \vec{p}' \rangle = \delta(\vec{p} - \vec{p}')$$

- iii) The Hamiltonian operator is given in terms of the classical Hamiltonian

$$H(\vec{q}, \vec{p}) = \left(\frac{1}{2m} \sum_\alpha p_\alpha^2 \right) + V(\vec{q})$$

as:

$$\hat{H} = H(\hat{q}_\alpha, \hat{p}_\alpha) = \left(\frac{1}{2m} \sum_\alpha \hat{p}_\alpha^2 \right) + V(\hat{q}_\alpha)$$

This works ONLY in Cartesian coordinates! In other coordinates the product pq occurs in H , but how to quantize qp ? The operators do not commute: $\hat{q}\hat{p} \neq \hat{p}\hat{q}$

For a magnetic field with vector potential $\vec{A}(\vec{q})$ the Hamiltonian is


$$H = \frac{1}{2m} \left(\vec{p} + \frac{e}{c} \vec{A}(\vec{q}) \right)^2$$

and also depends on products of \vec{p} and \vec{q} .

$$\hat{H} = \sum_n e_n |e_n\rangle \langle e_n| + \int_\Omega e(\kappa) |e(\kappa)\rangle \langle e(\kappa)| d\kappa$$

$$\begin{array}{ll} \langle e_n | e_{n'} \rangle = \delta_{n,n'} & \text{bounded} \\ \langle e(\kappa) | e(\kappa') \rangle = \delta(\kappa - \kappa') & \text{scattering} \end{array}$$

A typical effective potential is:



- iv) The representations of a state $|\psi\rangle$ in the position or momentum eigenstates are called WAVEFUNCTIONS. This means:

$$|\psi\rangle = \int \underbrace{\langle \vec{q} | \psi \rangle}_{=\psi(\vec{q})} |\vec{q}\rangle d\vec{q} \qquad |\psi\rangle = \int \underbrace{\langle \vec{p} | \psi \rangle}_{=\tilde{\psi}(\vec{p})} |\vec{p}\rangle d\vec{p}$$

Born's rule says, that $|\psi(\vec{q})|^2 d\vec{q}$ is the probability to find the particle between \vec{q} and $\vec{q} + (dq_1, dq_2, \dots)$ and $|\tilde{\psi}(\vec{p})|^2 d\vec{p}$ is the probability to find the particle between \vec{p} and $\vec{p} + (dp_1, dp_2, \dots)$.

1.5.2 Algebraic properties of \hat{q} and \hat{p}

Consider the following commutators:

$$\begin{aligned} [\hat{q}_\alpha, \hat{p}_\beta^2] &= \hat{p}_\beta [\hat{q}_\alpha, \hat{p}_\beta] + [\hat{q}_\alpha, \hat{p}_\beta] \hat{p}_\beta = 2i\hbar\delta_{\alpha\beta}\hat{p}_\beta \\ [\hat{q}_\alpha, \hat{p}_\beta^3] &= \hat{p}_\beta^2 [\hat{q}_\alpha, \hat{p}_\beta] + \hat{p}_\beta [\hat{q}_\alpha, \hat{p}_\beta] \hat{p}_\beta + [\hat{q}_\alpha, \hat{p}_\beta] \hat{p}_\beta^2 = 3i\hbar\delta_{\alpha\beta}\hat{p}_\beta^2 \\ &\vdots \\ [\hat{q}_\alpha, \hat{p}_\beta^n] &= \sum_{k=1}^n \hat{p}_\beta^{n-k} [\hat{q}_\alpha, \hat{p}_\beta] \hat{p}_\beta^{k-1} = ni\hbar\delta_{\alpha\beta}\hat{p}_\beta^{n-1} \end{aligned}$$

Since every analytic function can be represented by its Taylor series, one gets:

$$\begin{aligned} [\hat{q}_\alpha, f(\hat{p}_\beta)] &= i\hbar\delta_{\alpha\beta}\partial_{\hat{p}_\beta} f(x) \big|_{x=\hat{p}_\beta} =: i\hbar\delta_{\alpha\beta}\partial_{\hat{p}_\beta} f(\hat{p}_\beta) \\ [\hat{p}_\alpha, f(\hat{q}_\beta)] &= -i\hbar\delta_{\alpha\beta}\partial_{\hat{q}_\beta} f(\hat{q}_\beta) \end{aligned}$$

One can show:

$$\hat{q} e^{-\frac{i}{\hbar} \hat{p} q_0} |q\rangle = (q + q_0) |q + q_0\rangle$$

This follows, because $e^{-\frac{i}{\hbar} \hat{p} q_0} |q\rangle$ is an eigenstate of \hat{q} with eigenvalue $q + q_0$ and therefore:

$$e^{-\frac{i}{\hbar} \hat{p} q_0} |q\rangle = |q + q_0\rangle$$

One can also show:

$$e^{\frac{i}{\hbar} \hat{q} p_0} |p\rangle = |p + p_0\rangle$$

This is similar to:

$$e^{-\frac{i}{\hbar} \hat{H} t} |\psi(t_0)\rangle = |\psi(t_0 + t)\rangle$$

1.5.3 The Schrödinger equation in position and momentum representation

Consider the Hamiltonian:

$$\hat{H} = \frac{1}{2m} \hat{p}^2 + V(\hat{q})$$

I want to “represent”:

$$\begin{aligned} \hat{H} |e_n\rangle &= e_n |e_n\rangle \\ \hat{H} |e\rangle &= e |e\rangle \end{aligned}$$

For position representation, we apply $\langle q|$ to the equation, use the definitions $\psi_n(q) := \langle q | e_n \rangle$ and $\tilde{\psi}_n(p) := \langle p | e_n \rangle$, and multiply with the unit operator $\hat{I} = \int |p\rangle \langle p| dp$ to get:

$$\begin{aligned} \frac{1}{2m} \langle q | \hat{p}^2 \cdot \hat{I} | e_n \rangle + V(q) \langle q | e_n \rangle &= e_n \langle q | e_n \rangle \\ \frac{1}{2m} \int p^2 \langle q | p \rangle \tilde{\psi}_n(p) dp + V(q) \psi_n(q) &= e_n \psi_n(q) \end{aligned}$$

In general one gets:

$$\langle q | f(\hat{p}) | \psi \rangle = f(-\hbar \partial_q) \psi(q)$$

The recipe is:

$$\begin{array}{ll} \hat{p} \mapsto -i\hbar \partial_q & \text{one dimension} \\ \hat{p} \mapsto -i\hbar \vec{\nabla} & \text{three dimensions} \end{array}$$

In three dimensions this is:

$$\begin{array}{ll} \text{discrete case} & \text{continuous case} \\ \left(-\frac{\hbar^2}{2m} \nabla^2 + V(\vec{q}) \right) \psi_n(\vec{q}) = e_n \psi_n(\vec{q}) & \left(-\frac{\hbar^2}{2m} \nabla^2 + V(\vec{q}) \right) \psi_k(\vec{q}) = e(k) \psi_k(\vec{q}) \end{array}$$

The state is only in the discrete case normalizable:

$$\langle e_n, e_n \rangle = \int |\psi_n(\vec{q})|^2 d\vec{q} < \infty \quad \langle e(k), e(k) \rangle = \int |\psi_k(\vec{q})|^2 d\vec{q} = \infty$$

Time evolution:

$$\left(-\frac{\hbar^2}{2m}\nabla^2 + V(\vec{q})\right) \hat{U}(\vec{q}, \vec{q}', t) = i\hbar \partial_t \hat{U}(\vec{q}, \vec{q}', t) \quad \hat{U}(\vec{q}, \vec{q}', t=0) = \delta(\vec{q} - \vec{q}')$$

It is much simpler to solve the Schrödinger equation for one state:

$$\left(-\frac{\hbar^2}{2m}\nabla^2 + V(\vec{q})\right) \psi(\vec{q}, t) = i\hbar \partial_t \psi(\vec{q}, t) \quad \psi(\vec{q}, t=0) = \psi_0(\vec{q})$$

1.5.4 The Heisenberg and the Schrödinger “pictures”

Consider:

$$\begin{aligned} \langle \hat{A} \rangle_t &:= \langle \psi(t) | \hat{A} | \psi(t) \rangle \\ |\psi(t)\rangle &= \hat{U}(t, t_0) |\psi(t_0)\rangle \end{aligned}$$

This quantity admits a two-folded interpretation:

$$\begin{aligned} \langle \hat{A} \rangle_t &= \left(\langle \psi(t_0) | \hat{U}(t, t_0)^\dagger \right) \hat{A} \left(\hat{U}(t, t_0) | \psi(t_0) \rangle \right) \quad \text{the state evolves in time (Schrödinger)} \\ &= \langle \psi(t_0) | \underbrace{\left(\hat{U}(t, t_0)^\dagger \hat{A} \hat{U}(t, t_0) \right)}_{=: \hat{A}_H(t, t_0)} | \psi(t_0) \rangle \quad \text{the operator evolves in time (Heisenberg)} \end{aligned}$$

$\hat{A}_H(t, t_0)$ is the HEISENBERG OPERATOR, that is the operator \hat{A} in the Heisenberg representation. If $\hat{A} = \hat{A}^\dagger$, the also \hat{A}_H is Hermitian.

$$\hat{A}_H(t = t_0, t_0) = \hat{A}$$

An equation of motion for \hat{A} is easy to derive:

$$\begin{aligned} \partial_t \hat{A}_H(t, t_0) &= \partial_t \left(\hat{U}(t, t_0)^\dagger \hat{A}(t) \hat{U}(t, t_0) \right) = \partial_t \left(\hat{U}^\dagger \right) \hat{A} \hat{U} + \hat{U}^\dagger \partial_t \left(\hat{A} \right) \hat{U} + \hat{U}^\dagger \hat{A} \partial_t \left(\hat{U} \right) = \\ &\stackrel{\hat{H}\hat{U} = i\hbar \partial_t \hat{U}}{=} \left(-\frac{i}{\hbar} \hat{H} \hat{U} \right)^\dagger \hat{A} \hat{U} + \hat{U}^\dagger \left(\partial_t \hat{A} \right) \hat{U} + \hat{U}^\dagger \hat{A} \left(-\frac{i}{\hbar} \hat{H} \hat{U} \right) = \\ &\stackrel{\hat{U}\hat{U}^\dagger = 1}{=} -\frac{i}{\hbar} \left[\hat{A}_H(t, t_0), \hat{H}_H(t, t_0) \right] + \left(\partial_t \hat{A}(t) \right)_H \end{aligned}$$

Consider for example $\partial_t \hat{H} = 0$, $\hat{A}_1 = \hat{q}$ and $\hat{A}_2 = \hat{p}$:

$$\frac{d}{dt} \hat{q}_H = \frac{1}{m} \hat{p}_H \quad \frac{d}{dt} \hat{p}_H = -\partial_q V(q) \big|_{q=\hat{q}_H}$$

This leads to the Ehrenfest theorem:

$$\begin{aligned} m \frac{d^2}{dt^2} \hat{q}_H(t) &= -\partial_{\hat{q}_H} V(\hat{q}_H) \\ \hat{q}_H(t = t_0) &= \hat{q} \\ \hat{p}_H(t = t_0) &= \hat{p} \end{aligned}$$

For \hat{q}_H and \hat{p}_H and the usual Hamiltonian $\hat{H} = \frac{1}{2m} \hat{p}^2 + V(\hat{q}, t)$ they satisfy “Newton’s” equations. Take expectation values $\langle \psi(t_0) | \dots | \psi(t_0) \rangle$.

$$m \frac{d^2}{dt^2} \langle \hat{q} \rangle_t = -\langle \psi(t_0) | \partial_{\hat{q}_H} V(\hat{q}_H) | \psi(t_0) \rangle \stackrel{\text{in general}}{\neq} -\partial_{\langle \hat{q} \rangle_t} V(\langle \hat{q} \rangle_t)$$

But for $V(\hat{q}, t) = f_1(t) \hat{q}^2 + f_2(t) \hat{q} + f_3(t)$ the partial derivative $\partial_{\hat{q}_H} V(\hat{q}_H)$ is linear in \hat{q}_H and therefore $\langle \hat{q} \rangle_t$ describes the classical trajectory with the initial conditions $\langle \psi(t_0) | \hat{q} | \psi(t_0) \rangle$ and $\langle \psi(t_0) | \hat{p} | \psi(t_0) \rangle$.

Comment: The best compromise to do quantum mechanics in a way, that looks similar (at least for short times) to classical mechanics is GAUSSIAN wavepacket.

A wavepacket is a superposition:

$$|\psi\rangle = \int \tilde{\psi}(p) |p\rangle dp$$

A Gaussian wavepacket is:

$$\begin{aligned} |\psi_{(q_0, p_0)}\rangle &= \int \psi_{(q_0, p_0)}(q) |q\rangle dq \\ &= \int \tilde{\psi}_{(q_0, p_0)}(p) |p\rangle dp \end{aligned}$$

$$\psi_{(q_0, p_0)}(q) = \frac{1}{\sqrt[4]{2\pi}\sigma} \exp\left(-\frac{1}{4} \frac{(q - q_0)^2}{\sigma^2}\right) \exp\left(-\frac{i}{\hbar} q p_0\right) \quad (1.2)$$


For this one gets:

$$(\Delta \hat{q})^2 (\Delta \hat{p})^2 = \frac{\hbar^2}{4}$$

$$\begin{aligned} \langle \psi_{(q_0, p_0)} | \hat{q} | \psi_{(q_0, p_0)} \rangle &= q_0 \\ \langle \psi_{(q_0, p_0)} | \hat{p} | \psi_{(q_0, p_0)} \rangle &= p_0 \end{aligned}$$

2 Basic applications

2.1 Boxes




The normalized eigenstates are:

$$\psi_n(q) = \sqrt{\frac{2}{L}} \sin\left(\frac{n\pi}{L}q\right)$$

$$e_n = n^2 e_0$$

2.2 Square well



- i) Energy $E < V_0$ smaller than potential wall, leads to a bound state.

Write the equation:

$$\begin{aligned} \left(-\frac{\hbar^2}{2m} \partial_q^2 + V(q) \right) \psi(q) &= E \psi(q) \\ \partial_q^2 \psi(q) &= -k(E)^2 \psi(q) \\ k(E) &= +\sqrt{\frac{2m}{\hbar^2} (E - V(q))} \end{aligned}$$

In the three regions I, II and III one gets:

$$\begin{aligned} k_I(E) &= i |k_I(E)| \\ k_{II}(E) &= |k_{II}(E)| \\ k_{III}(E) &= i |k_{III}(E)| \end{aligned}$$

ii) Write the general solution:

$$\begin{aligned} \psi_{I(q)} &= A_I e^{|k_I(E)|q} + B_I e^{-|k_I(E)|q} \\ \psi_{II(q)} &= A_{II} e^{i|k_{II}(E)|q} + B_{II} e^{-i|k_{II}(E)|q} \\ \psi_{III(q)} &= A_{III} e^{|k_{III}(E)|q} + B_{III} e^{-|k_{III}(E)|q} \end{aligned}$$

iii) Normalization: $B_I = A_{III} = 0$

Continuity: For a finite potential ψ and $\partial_q \psi'$ are continuous.

$$\begin{aligned} \psi_I \left(-\frac{L}{2} \right) &= \psi_{II} \left(-\frac{L}{2} \right) & \partial_q \psi_I \left(-\frac{L}{2} \right) &= \partial_q \psi_{II} \left(-\frac{L}{2} \right) \\ \psi_{II} \left(\frac{L}{2} \right) &= \psi_{III} \left(\frac{L}{2} \right) & \partial_q \psi_{II} \left(\frac{L}{2} \right) &= \partial_q \psi_{III} \left(\frac{L}{2} \right) \end{aligned}$$

Now we have four equations and four variables, but one expects infinitely many solutions.

$$\mathbb{M} \begin{pmatrix} A_I \\ A_{II} \\ B_{II} \\ B_{III} \end{pmatrix} = 0 \quad \det \mathbb{M}(E) \stackrel{!}{=} 0$$

2.3 The harmonic oscillator (in one dimension)

The potential is:

$$V(\hat{q}) = \frac{m\omega_0^2}{2} \hat{q}^2$$

For all energies $e_n \in \mathbb{R}_{\geq 0}$ we have:

$$\begin{aligned} \hat{H} |e_n\rangle &= e_n |e_n\rangle \\ \langle e_n | e_n \rangle &= 1 \end{aligned}$$

i) The Schrödinger equation is:

$$\left(-\frac{\hbar^2}{2m} \partial_q^2 + \frac{m\omega_0^2}{2} q^2 \right) \psi_n(q) = e_n \psi_n(q) \quad \psi(q \rightarrow \pm\infty) = 0$$

ii) Use dimensionless variables:

$$q = \alpha x \qquad \phi(x) := \psi_n(\alpha x)$$

$$\begin{aligned} \left(-\frac{\hbar^2}{2m} \frac{1}{\alpha^2} \partial_q^2 + \frac{m\omega_0^2}{2} \alpha^2 q^2 \right) \phi(q) &= e_n \phi(x) \\ \left(\partial_x^2 - \frac{m^2 \omega_0^2}{\hbar^2} \alpha^4 x^2 \right) \phi(q) &= -\frac{2m}{\hbar^2} \alpha^2 e_n \phi(x) \end{aligned}$$

Choose:

$$\alpha := \sqrt{\frac{\hbar}{m\omega_0}} \qquad \lambda := \frac{2e_n}{\hbar\omega_0}$$

Then the problem reduces to:

$$\boxed{(\partial_x^2 + (\lambda - x^2)) \phi(x) = 0 \qquad \psi(x \rightarrow \pm\infty) = 0}$$

iii) Check the asymptotic behavior, that is, when x goes to $\pm\infty$:

$$\phi(x) \approx \exp\left(-\frac{1}{2}x^2\right)$$

Assume a solution:

$$\phi(x) = u(x) \exp\left(-\frac{1}{2}x^2\right)$$

iv) Then we get for $u(x)$ the differential equation:

$$(\partial_x^2 - 2x\partial_x + (\lambda - 1)) u(x) = 0$$

Solve this with a power series, by assuming, that $u(x)$ is analytic:

$$u(x) = \sum_{n=0}^{\infty} a_n x^n$$

This leads to the „recursion relation“:

$$a_{n+2} = \frac{(2n+1-\lambda)}{(n+1)(n+2)} a_n$$

Or equivalently:

$$a_{n-2} = \frac{(n-1)n}{(2n-3-\lambda)} a_n$$

v) Check again the asymptotic behavior, to get an idea, how $u(x)$ looks like, when $n \rightarrow \infty$:

$$\frac{a_{n+2}}{a_n} \xrightarrow{n \gg 1} \frac{2}{n}$$

But we have

$$\exp\left(-\frac{x^2}{2}\right) = \sum_{n=0}^{\infty} \underbrace{\frac{1}{n!} \left(-\frac{1}{2}\right)^n}_{=: c_n} x^{2n} = \sum_{n=0}^{\infty} c_n x^{2n}$$

with:

$$\frac{c_{n+1}}{c_n} = \frac{n!}{(n+1)!} \cdot \left(-\frac{1}{2}\right)^{n+1} \cdot (-2)^n = \frac{-1}{2(n+1)} \xrightarrow{n \gg 1} -\frac{1}{2n}$$

Therefore $u(x \rightarrow \infty)$ grows FASTER THAN $\exp\left(\frac{1}{2}x^2\right)$ and therefore:

$$\exp\left(-\frac{1}{2}x^2\right) u(x) \rightarrow \infty$$

If the series does not stop, $\phi(x)$ is *not* normalizable.

So there has to exist a „ n^* “ such that:

$$\begin{aligned} 2n^* + 1 - \lambda &= 0 \\ \lambda &= 2n^* + 1 \\ e_n &= \hbar\omega_0 \left(n^* + \frac{1}{2}\right) \end{aligned}$$

This gives:

$$\psi_n(q) = N_n \exp\left(-\frac{1}{2} \left(\frac{q}{\alpha}\right)^2\right) H_n\left(\frac{q}{\alpha}\right)$$


The polynomials $H_n(x)$ are called the HERMITE POLYNOMIALS and are explicitly:

$$H_n(x) = (-1)^n \sum_{k_1+2k_2=n} \frac{n!}{k_1! \cdot k_2!} (-1)^{k_1+k_2} (2x)^{k_1}$$

N_n is fixed by:

$$\int_{-\infty}^{\infty} |\psi_n(q)|^2 dq = 1$$

This looks like:



Then came Dirac:

2.3.1 The number representation

Introduce two operators:

$$\begin{aligned} \text{„destruction“} \quad \hat{a} &= \sqrt{\frac{m\omega_0}{2\hbar}} \left(\hat{q} + \frac{\mathbf{i}}{m\omega_0} \hat{p} \right) & \text{„creation“} \quad \hat{a}^\dagger &= \sqrt{\frac{m\omega_0}{2\hbar}} \left(\hat{q} - \frac{\mathbf{i}}{m\omega_0} \hat{p} \right) \end{aligned}$$

It is easy to calculate:

$$[\hat{a}, \hat{a}^\dagger] = 1$$

Introduce the „number“ operator:

$$\hat{n} := \hat{a}^\dagger \hat{a}$$

Then we can find easily:

$$\hat{H} = \frac{1}{2m} \hat{p}^2 + \frac{m\omega_0^2}{2} \hat{q}^2 = \hbar\omega_0 \left(\hat{n} + \frac{1}{2} \right)$$

Therefore the eigenstates of \hat{n} are automatically the eigenstates of \hat{H} .

It also satisfies:

$$[\hat{n}, \hat{a}] = -\hat{a} \quad \quad [\hat{n}, \hat{a}^\dagger] = \hat{a}^\dagger$$

We want to find $|\sigma\rangle$ such that:

$$\hat{n} |\sigma\rangle = \sigma |\sigma\rangle$$

We note:

$$\hat{n}(\hat{a}|\sigma\rangle) = (\hat{n}\hat{a})|\sigma\rangle = (\hat{a}\hat{n} - [\hat{a}, \hat{n}])|\sigma\rangle = (\hat{a}\hat{n} - \hat{a})|\sigma\rangle = (\sigma - 1)|\sigma\rangle$$

$\hat{a}|\sigma\rangle$ is an eigenstate of \hat{n} with eigenvalue $\sigma - 1$, so we get:

$$\hat{a}|\sigma\rangle = N_\sigma |\sigma - 1\rangle$$

From $\langle\sigma|\sigma\rangle = \langle\sigma - 1|\sigma - 1\rangle = 1$ follows:

$$\boxed{\hat{a}|\sigma\rangle = \sqrt{\sigma} |\sigma - 1\rangle}$$

Analogously we have:

$$\boxed{\hat{a}^\dagger|\sigma\rangle = \sqrt{\sigma + 1} |\sigma + 1\rangle}$$

Take a $\sigma > 0$ and apply \hat{a} to $|\sigma\rangle$ many times to get, that σ must be an integer, because otherwise the normalization constant $\sqrt{\sigma}$ would be imaginary.

There is a special state $|0\rangle$, the „vacuum“.

Let us call the states $|n\rangle$ for $n \in \mathbb{N}_0$.

$$\hat{n}|n\rangle = n|n\rangle \quad \hat{a}|n\rangle = \sqrt{n}|n - 1\rangle \quad \hat{a}^\dagger|n\rangle = \sqrt{n + 1}|n + 1\rangle$$

$$\hat{H}|n\rangle = e_n|n\rangle \quad e_n = \hbar\omega_0 \left(n + \frac{1}{2}\right)$$

If we know $|0\rangle$, we can construct an arbitrary $|n\rangle$:

$$|n\rangle = \frac{1}{\sqrt{n!}} (\hat{a}^\dagger)^n |0\rangle$$

To calculate $\langle q|0\rangle$ we use $\hat{a}|0\rangle = 0$ and represent.

$$\begin{aligned} \langle q|\hat{a}|0\rangle &= 0 \\ \left\langle q \left| \hat{q} + \frac{\mathbf{i}}{m\omega_0} \hat{p} \right| 0 \right\rangle &= 0 \\ \left(q + \frac{\mathbf{i}}{m\omega_0} (-i\hbar\partial_q) \right) \psi_0(q) &= 0 \\ \left(\frac{\hbar}{m\omega_0} \partial_q + q \right) \psi_0(q) &= 0 \\ \psi_0(q) &= N_0 \exp\left(-\frac{1}{2} \left(\frac{q}{\alpha}\right)^2\right) \end{aligned}$$

N_0 is fixed by normalization:

$$\int_{-\infty}^{\infty} |\psi_0(q)|^2 dq = 1$$

For example $\psi_1(q)$ is given by:

$$\begin{aligned} \psi_1(q) = \langle q|1\rangle &= \langle q|\hat{a}^\dagger|0\rangle = \sqrt{\frac{m\omega_0}{2\hbar}} \left(q - \frac{\mathbf{i}}{m\omega_0} (-i\hbar\partial_q) \right) \psi_0(q) = \\ &= N_0 \sqrt{\frac{m\omega_0}{2\hbar}} \left(q - \frac{\hbar}{m\omega_0} \partial_q \right) \exp\left(-\frac{1}{2} \left(\frac{q}{\alpha}\right)^2\right) = \\ &= N_1 \exp\left(-\frac{1}{2} \left(\frac{q}{\alpha}\right)^2\right) \cdot H_1\left(\frac{q}{\alpha}\right) \end{aligned}$$

2.3.2 The dynamics of the harmonic oscillator

The Hamiltonian is:

$$\hat{H} = \frac{1}{2m} \hat{p}^2 + \frac{m\omega_0^2}{2} \hat{q}^2$$

In principal we need:

$$\hat{U}(t) := \exp\left(-\frac{\mathbf{i}}{\hbar} \hat{H} t\right) = \exp\left(-\frac{\mathbf{i}}{\hbar} \left(\frac{1}{2m} \hat{p}^2 + \frac{m\omega_0^2}{2} \hat{q}^2\right) t\right)$$

Let's calculate in the position representation:

$$\langle q | \hat{U}(t) | q' \rangle =: U(q, q', t)$$

Remember, what \hat{U} does:

$$|\psi(t)\rangle = \hat{U}(t) |\psi(0)\rangle$$

In position representation this is:

$$\psi(q, t) = \int U(q, q', t) \psi(q', 0) dq'$$

In some books the notation for $U(q, q', t)$ is $K(q, q', t)$.

$$\begin{aligned} U(q, q', t) &= \sum_n \exp\left(-\frac{\mathbf{i}}{\hbar} E_n t\right) \langle q | n \rangle \langle n | q' \rangle = \\ &= \sum_n \exp\left(-\mathbf{i}\omega_0 \left(n + \frac{1}{2}\right) t\right) N_n \exp\left(-\frac{1}{2} \left(\frac{q}{\alpha}\right)^2\right) H_n\left(\frac{q}{\alpha}\right) \cdot \\ &\quad \cdot \left(N_n \exp\left(-\frac{1}{2} \left(\frac{q'}{\alpha}\right)^2\right) H_n\left(\frac{q'}{\alpha}\right)\right)^* = \\ &= \sum_n \exp\left(-\mathbf{i}\omega_0 \left(n + \frac{1}{2}\right) t\right) N_n^2 \exp\left(-\frac{q^2 + q'^2}{2\alpha^2}\right) H_n\left(\frac{q}{\alpha}\right) H_n\left(\frac{q'}{\alpha}\right) = \\ &= \sqrt{\frac{m\omega_0}{2\pi\mathbf{i}\hbar \sin(\omega_0 t)}} \cdot \exp\left(\frac{\mathbf{i}}{\hbar} \left(\frac{m\omega_0}{2 \sin(\omega_0 t)}\right) (q^2 + q'^2) \cos(\omega_0 t) - 2qq'\right) \end{aligned}$$

To check this, consider the limes $\omega_0 \rightarrow 0$, for which we get the free particle:

$$U(q, q', t) \xrightarrow{\omega_0 \rightarrow 0} \sqrt{\frac{m}{2\pi\mathbf{i}\hbar t}} \cdot \exp\left(\frac{\mathbf{i}}{\hbar} \frac{m}{2t} (q - q')^2\right)$$

Let's try the Heisenberg picture:

$$\begin{aligned} \frac{d}{dt} \hat{q}_H(t) &= \frac{\mathbf{i}}{\hbar} [H(\hat{q}_H, \hat{p}_H), \hat{q}_H] = \frac{1}{m} \hat{p}_H \\ \frac{d}{dt} \hat{p}_H(t) &= \frac{\mathbf{i}}{\hbar} [H(\hat{q}_H, \hat{p}_H), \hat{p}_H] = -m\omega_0 \hat{q}_H \end{aligned}$$

The initial conditions are:

$$\begin{aligned}\hat{q}_H(t=0) &= \hat{q} \\ \hat{p}_H(t=0) &= \hat{p}\end{aligned}$$

Remember the definition of an operator in the Heisenberg picture:

$$\hat{q}_H = \exp\left(\frac{\mathbf{i}}{\hbar}\hat{H}t\right) \hat{q} \exp\left(-\frac{\mathbf{i}}{\hbar}\hat{H}t\right)$$

Now we define the NESTED COMMUTATOR as

$$\begin{aligned}[\hat{B}, \hat{A}]^{(0)} &:= \hat{A} \\ [\hat{B}, \hat{A}]^{(n+1)} &:= [\hat{B}, [\hat{B}, \hat{A}]^{(n)}]\end{aligned}$$

and use the operator identity for $\lambda \in \mathbb{C}$ and arbitrary operators \hat{A}, \hat{B} :

$$\begin{aligned}\exp(\mathbf{i}\lambda\hat{B}) \hat{A} \exp(-\mathbf{i}\lambda\hat{B}) &= \sum_{n=0}^{\infty} \frac{(\mathbf{i}\lambda)^n}{n!} [\hat{B}, \hat{A}]^{(n)} \\ &\approx \hat{A} + \mathbf{i}\lambda [\hat{B}, \hat{A}] + \frac{(\mathbf{i}\lambda)^2}{2!} [\hat{B}, [\hat{B}, \hat{A}]]\end{aligned}$$

If $[\hat{A}, \hat{B}] = 0$, we get just \hat{A} .

If $[\hat{A}, [\hat{A}, \hat{B}]] = [\hat{B}, [\hat{A}, \hat{B}]] = 0$, we get $\hat{A} + \mathbf{i}\lambda [\hat{B}, \hat{A}]$.

Now we come back to the problem at hand:

$$\lambda = \frac{t}{\hbar} \qquad \hat{B} = \hat{H} \qquad \hat{A} = \hat{q}$$

We know:


$$\begin{aligned}[\hat{H}, \hat{q}] &= \frac{1}{2m} [\hat{p}^2, \hat{q}] + \underbrace{[V(\hat{q}), \hat{q}]}_{=0} = \frac{1}{2m} (\hat{p} [\hat{p}, \hat{q}] + [\hat{p}, \hat{q}] \hat{p}) = -\frac{\mathbf{i}\hbar}{m} \hat{p} \\ [\hat{H}, \hat{p}] &= \frac{1}{2m} [\hat{p}^2, \hat{p}] + \frac{m\omega_0^2}{2} [\hat{q}^2, \hat{p}] = \mathbf{i}\hbar m\omega_0^2 \hat{q}\end{aligned}$$

Together this inductively gives:

$$\begin{aligned}\hat{q}_H(t) &= \hat{q} \cos(\omega_0 t) + \frac{1}{m\omega_0} \hat{p} \sin(\omega_0 t) \\ \hat{p}_H(t) &= m\omega_0 \hat{q} \sin(\omega_0 t) + \hat{p} \cos(\omega_0 t)\end{aligned}$$

2.4 The tunneling effect

Look at the potential:



The problem is unbounded and therefore the spectrum is continuous and every energy $E \in \mathbb{R}_{>0}$ is allowed.

Imagine that we inject the particle from the left with a certain energy E .

$$k := \frac{\sqrt{2mE}}{\hbar} \qquad k_V := \frac{\sqrt{2m(V_0 - E)}}{\hbar}$$

The goal is to find R and T as functions of the Energy E , the height V_0 and width d of the potential.

Business as usual:

The boundary condition is, that A_I is know and the coefficient of e^{-ikq} in the region III is zero. We consider only $E < V_0$.

$$\begin{aligned} \psi_I(q) &= A_I e^{ikq} + R e^{-ikq} \\ \psi_{II}(q) &= A_{II} e^{k_V q} + B_{II} e^{-k_V q} \\ \psi_{III}(q) &= T e^{ikq} \end{aligned}$$

We have the four variables R, T, A_{II} and B_{II} . Now we impose the matching conditions:

$$\begin{aligned} \psi_I\left(-\frac{d}{2}\right) &= \psi_{II}\left(-\frac{d}{2}\right) & \psi_I'\left(-\frac{d}{2}\right) &= \psi_{II}'\left(-\frac{d}{2}\right) \\ \psi_I\left(\frac{d}{2}\right) &= \psi_{II}\left(\frac{d}{2}\right) & \psi_I'\left(\frac{d}{2}\right) &= \psi_{II}'\left(\frac{d}{2}\right) \end{aligned}$$

This system looks like:

$$\mathbb{M} \cdot \begin{pmatrix} B \\ A_{II} \\ B_{II} \\ T \end{pmatrix} = A_I \begin{pmatrix} -e^{-ik\frac{d}{2}} \\ -ik e^{ik\frac{d}{2}} \\ 0 \\ 0 \end{pmatrix}$$


For $k_V d \gg 1$ this roughly gives:

$$\begin{aligned} T(E) &\sim A_I e^{-\frac{k_V(E)d}{2}} \\ \frac{T(E, d)}{T(E, d')} &\sim e^{-\frac{k_V(E)}{2}(d-d')} \end{aligned}$$

2.5 Two formal consequences of the position representation of the Schrödinger equation

2.5.1 Oscillation theorem (1D, bounded)

If $\langle q | e_n \rangle$ has k -nodes, that means $\psi_n(q) = 0$ is k times, then $\langle q | e_{n+1} \rangle$ has $k + 1$ nodes.



2.5.2 The continuity equation

Let $\psi(\vec{q}, t)$ solve the Schrödinger equation for the Hamilton operator $\hat{H} = \frac{\hat{p}^2}{2m} + V(\vec{q})$. Since \hat{H} is Hermitian, $V(\vec{q})$ has to be real and solve:

$$\begin{aligned} i\hbar \partial_t \psi(\vec{q}, t) &= \left(-\frac{\hbar^2}{2m} \nabla^2 + V(\vec{q}) \right) \psi(\vec{q}, t) \\ -i\hbar \partial_t \psi^*(\vec{q}, t) &= \left(-\frac{\hbar^2}{2m} \nabla^2 + V(\vec{q}) \right) \psi^*(\vec{q}, t) \end{aligned}$$

This gives:

$$\begin{aligned} \partial_t |\psi|^2 &= \partial_t (\psi \psi^*) = (\partial_t \psi) \psi^* + \psi (\partial_t \psi^*) = \\ &= \psi^* \left(\frac{1}{i\hbar} \left(-\frac{\hbar^2}{2m} \nabla^2 + V(\vec{q}) \right) \psi \right) + \psi \left(\frac{-1}{i\hbar} \left(-\frac{\hbar^2}{2m} \nabla^2 + V(\vec{q}) \right) \psi^* \right) = \\ &= \psi^* \left(-\frac{\hbar}{2mi} \nabla^2 \psi \right) + \psi \left(\frac{\hbar}{2mi} \nabla^2 \psi^* \right) = \\ &= \frac{\hbar}{2mi} (\psi \nabla^2 \psi^* - \psi^* \nabla^2 \psi) = \\ &= \frac{\hbar}{2mi} \nabla (\psi \nabla \psi^* - \psi^* \nabla \psi) \end{aligned}$$

Therefore we have the continuity equation:

$$\partial_t |\psi|^2 - \frac{\hbar}{2mi} \nabla (\psi \nabla \psi^* - \psi^* \nabla \psi) = 0$$

Define the probability density $\varrho_\psi(\vec{q}, t) := |\psi(\vec{q}, t)|^2$ and the probability current:

$$\vec{j}_\psi(\vec{q}, t) := -\frac{i\hbar}{2m} (\psi^* \nabla \psi - \psi \nabla \psi^*)$$

They are related by

$$\partial_t \varrho_\psi(\vec{q}, t) + \nabla \cdot \vec{j}_\psi(\vec{q}, t) = 0$$

and we can interpret them as expectation values:

$$\varrho_\psi(\vec{q}, t) = |\psi(\vec{q}, t)|^2 = \left| \langle \hat{q} | \psi(t) \rangle \right|^2 = \langle \hat{q} | \underbrace{|\psi(t)\rangle \langle \psi(t)|}_{=: \hat{\varrho}} | \hat{q} \rangle = \langle \hat{q} | \hat{\varrho} | \hat{q} \rangle$$

$$\begin{aligned} \vec{j}(\vec{q}, t) &= \frac{-i\hbar}{2m} (\psi^* \nabla \psi - \psi \nabla \psi^*) = \frac{-i\hbar}{2m} (\langle \psi | \vec{q} \rangle \nabla \langle \vec{q} | \psi \rangle - \langle \vec{q} | \psi \rangle \nabla \langle \psi | \vec{q} \rangle) = \\ &= \frac{-1}{2m} \left(\langle \vec{q} | \hat{p} | \psi \rangle \langle \psi | \vec{q} \rangle - \langle \vec{q} | \psi \rangle \langle \psi | \hat{p} | \vec{q} \rangle \right) = \\ &= \frac{-1}{2m} \langle \vec{q} | \underbrace{(\hat{p} | \psi \rangle \langle \psi | - | \psi \rangle \langle \psi | \hat{p})}_{=: [\hat{p}, \hat{\varrho}]} | \vec{q} \rangle = \frac{-1}{2m} \langle \vec{q} | \underbrace{[\hat{p}, \hat{\varrho}]}_{=: -2m\vec{j}} | \vec{q} \rangle = \langle \vec{q} | \hat{j} | \vec{q} \rangle \end{aligned}$$

3 Symmetries, Rotations and Angular momentum

Remember from classical mechanics:

A symmetry is a physical operation that leaves the system invariant.

In classical mechanics this means we change \vec{q} and \vec{p} .

Examples:

- For $\mathcal{R} \in \text{SL}_n(\mathbb{R})$, that means $\mathcal{R}^{-1} = \mathcal{R}^T$ and $\det(\mathcal{R}) = 1$, we define:

$$\begin{aligned}\vec{q}' &= \mathcal{R}\vec{q} \\ \vec{p}' &= \mathcal{R}\vec{p}\end{aligned}$$

This leaves a Lagrangian of the Form $\mathcal{L}(\vec{q}, \dot{\vec{q}}) = \frac{m}{2} |\dot{\vec{q}}|^2 - V(|\vec{q}|) \Rightarrow \{H, \vec{L}\} = 0$

- The translation $\vec{q}' = \vec{q} + \vec{q}_0$ leaves the free particle with Lagrangian

$$\mathcal{L}(\vec{q}, \dot{\vec{q}}, t) = \frac{m}{2} |\dot{\vec{q}}|^2$$

invariant and therefore $\{H, \vec{p}\} = 0$.

- Emmy Noether: Symmetry \Rightarrow Existence of a constant of motion

3.1 The tensor product in quantum mechanics

Consider two degrees of freedom (for example two particles in one dimension), described by states in $\mathcal{H}^{(1)}, \mathcal{H}^{(2)}$.

A postulate of quantum mechanics says, that the composite system is described by a state in the Hilbert space $\mathcal{H} = \mathcal{H}^{(1)} \otimes \mathcal{H}^{(2)}$. This space is called TENSOR PRODUCT.

So the dimension is $\dim \mathcal{H} = \dim(\mathcal{H}^{(1)}) \cdot \dim(\mathcal{H}^{(2)})$.

If $|\psi^{(1)}\rangle \in \mathcal{H}^{(1)}$, $|\psi^{(2)}\rangle \in \mathcal{H}^{(2)}$ then $|\psi^{(1)}\rangle \otimes |\psi^{(2)}\rangle \in \mathcal{H}^{(1)} \otimes \mathcal{H}^{(2)} = \mathcal{H}$.

If $\hat{A}^{(1)}$ acts on $\mathcal{H}^{(1)}$ and $\hat{A}^{(2)}$ acts on $\mathcal{H}^{(2)}$, then $\hat{A}^{(1)} \otimes \hat{A}^{(2)}$ acts on \mathcal{H} as follows:

$$\left(\hat{A}^{(1)} \otimes \hat{A}^{(2)}\right) \left(|\psi^{(1)}\rangle \otimes |\psi^{(2)}\rangle\right) = \left(\hat{A}^{(1)} |\psi^{(1)}\rangle\right) \otimes \left(\hat{A}^{(2)} |\psi^{(2)}\rangle\right)$$

$$\hat{A}^{(1)} \mapsto \hat{A}^{(1)} \otimes \hat{I}^{(2)}$$

$$\hat{A}^{(2)} \mapsto \hat{I}^{(1)} \otimes \hat{A}^{(2)}$$

$$[\hat{A}^{(1)}, \hat{A}^{(2)}] = 0$$

For $|\psi\rangle = |\psi^{(1)}\rangle \otimes |\psi^{(2)}\rangle$ and $|\varphi\rangle = |\varphi^{(1)}\rangle \otimes |\varphi^{(2)}\rangle$ we have the scalar product:

$$\langle \varphi | \psi \rangle = \langle \varphi^{(1)} | \psi^{(1)} \rangle \langle \varphi^{(2)} | \psi^{(2)} \rangle$$

With all this together one can write any operator \hat{A} and any Ket $|\psi\rangle$:

$$\begin{aligned} \hat{A} &= \sum_{n,m} c_{n,m} \hat{A}_n^{(1)} \otimes \hat{A}_m^{(2)} \\ |\psi\rangle &= \sum_{n,m} a_{n,m} |\psi_n^{(1)}\rangle \otimes |\psi_m^{(2)}\rangle \stackrel{\text{i.A.}}{\neq} |\varphi^{(1)}\rangle \otimes |\varphi^{(2)}\rangle \end{aligned}$$

A state in the form $|\psi^{(1)}\rangle \otimes |\psi^{(2)}\rangle$ is called *separable*.

If a state is not separable, it is called *entangled*.

For example:

$$\left(\hat{A}^{(1)} \otimes \hat{I}^{(2)} + \hat{I} \otimes \hat{A}^{(2)} \right)^2 = \left(\hat{A}^{(1)} \right)^2 \otimes \hat{I}^{(2)} + 2\hat{A}^{(1)} \otimes \hat{A}^{(2)} + \hat{I} \otimes \left(\hat{A}^{(2)} \right)^2$$

Now we can understand better:

$$\hat{q}_\alpha |\vec{q}\rangle = q_\alpha |\vec{q}\rangle$$

$|\vec{q}\rangle$ is just a compact way to write:

$$|\vec{q}\rangle = |q_x, q_y, q_z\rangle := |q_x\rangle \otimes |q_y\rangle \otimes |q_z\rangle$$

3.2 Rotations and angular momentum

Come back to classical mechanics. We know, that $\vec{L} = \vec{q} \times \vec{p}$ „generates“ rotations $\mathcal{R}(\vec{n}, d\vartheta)$ around an normalized axis \vec{n} with the angle $d\vartheta$.

$$f(\vec{q}, \vec{p}) \rightarrow f(\vec{q}', \vec{p}') = f(\mathcal{R}(\vec{n}, d\vartheta) \vec{q}, \mathcal{R}(\vec{n}, d\vartheta) \vec{p}) \approx f(\vec{q}, \vec{p}) + d\vartheta \left\{ \underbrace{\vec{L} \cdot \vec{n}}_{\text{Generator}}, f(\vec{q}, \vec{p}) \right\} + O((d\vartheta)^2)$$

Another example:

$$f(\vec{q}(t), \vec{p}(t)) \rightarrow f(\vec{q}(t + \delta t), \vec{p}(t + \delta t)) \approx f(\vec{q}, \vec{p}) + dt \{H, f(\vec{q}, \vec{p})\}$$

Explicit calculation shows:

$$\{L_i, L_j\} = \varepsilon_{ijk} L_k$$

In quantum mechanics we DEFINE angular momentum \vec{j} as the generator of rotations. Define:

$$|\psi^{\mathcal{R}}\rangle = \hat{D}(\mathcal{R}) |\psi\rangle = e^{-\frac{i}{\hbar} (\vec{j} \cdot \vec{n}) \theta} |\psi\rangle$$

Check out, how an operator \hat{A} changes:

$$\hat{A}^{\mathcal{R}} = \hat{D}(\mathcal{R})^\dagger \hat{A} \hat{D}(\mathcal{R}) \xrightarrow{\theta \rightarrow d\theta} \hat{A} + \frac{i d\theta}{\hbar} [\vec{j} \cdot \vec{n}, \hat{A}] + O(d\theta^2)$$

We want, that the group properties of the \mathcal{R} 's, which are

$$\begin{aligned}\mathcal{R}(\vec{n}, 0) &= E_3 \\ \mathcal{R}(\vec{n}_1, \theta_1) \mathcal{R}(\vec{n}_2, \theta_2) &= \mathcal{R}(\vec{n}(\vec{n}_1, \vec{n}_2), \theta(\theta_1, \theta_2)) \quad (\text{cf. Goldstein Classical Mechanics}) \\ \mathcal{R}(\vec{n}, \theta)^{-1} &= \mathcal{R}(\vec{n}, -\theta) \\ (\mathcal{R}_1 \mathcal{R}_2) \mathcal{R}_3 &= \mathcal{R}_1 (\mathcal{R}_2 \mathcal{R}_3)\end{aligned}$$

go into the \hat{D} 's.

$$\begin{aligned}\hat{D}(\vec{n}, 0) &= \hat{I} \\ \hat{D}_1(\vec{n}_1, \theta_1) \hat{D}_2(\vec{n}_2, \theta_2) &= \hat{D}(\vec{n}(\vec{n}_1, \vec{n}_2), \theta(\theta_1, \theta_2)) \\ \hat{D}(\vec{n}, \theta)^{-1} &= \hat{D}(\vec{n}, -\theta) \\ (\hat{D}_1 \hat{D}_2) \hat{D}_3 &= \hat{D}_1 (\hat{D}_2 \hat{D}_3)\end{aligned}$$

In particular we want the \hat{D} 's version of:

$$\begin{aligned}\mathcal{R}(\vec{e}_x, \varepsilon) \mathcal{R}(\vec{e}_y, \varepsilon) - \mathcal{R}(\vec{e}_y, \varepsilon) \mathcal{R}(\vec{e}_x, \varepsilon) &\approx \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 - \frac{\varepsilon^2}{2} & -\varepsilon \\ 0 & \varepsilon & 1 - \frac{\varepsilon^2}{2} \end{pmatrix} \begin{pmatrix} 1 - \frac{\varepsilon^2}{2} & 0 & \varepsilon \\ 0 & 1 & 0 \\ -\varepsilon & 0 & 1 - \frac{\varepsilon^2}{2} \end{pmatrix} - \\ &- \begin{pmatrix} 1 - \frac{\varepsilon^2}{2} & 0 & \varepsilon \\ 0 & 1 & 0 \\ -\varepsilon & 0 & 1 - \frac{\varepsilon^2}{2} \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 - \frac{\varepsilon^2}{2} & -\varepsilon \\ 0 & \varepsilon & 1 - \frac{\varepsilon^2}{2} \end{pmatrix} = \\ &= \mathcal{R}(\vec{e}_z, \varepsilon^2) - E_3 = \begin{pmatrix} 0 & -\varepsilon^2 & 0 \\ \varepsilon^2 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}\end{aligned}$$

So we want:

$$[\hat{D}(\vec{e}_x, \varepsilon), \hat{D}(\vec{e}_y, \varepsilon)] = \hat{D}(\vec{e}_z, \varepsilon^2) - \hat{I} + O(\varepsilon^3)$$

Use

$$\hat{D}(\vec{e}_i, \varepsilon) = \exp\left(-\frac{\mathbf{i}}{\hbar} \hat{J}_i \varepsilon\right) \approx 1 - \frac{\mathbf{i}}{\hbar} \hat{J}_i \varepsilon + \left(-\frac{\mathbf{i}}{\hbar}\right)^2 \frac{1}{2} \hat{J}_i^2 \varepsilon^2 + \dots$$

to get:

$$[\hat{J}_i, \hat{J}_j] = \mathbf{i} \hbar \varepsilon_{ijk} \hat{J}_k$$

Funny remark: Take the spin \hat{s} and calculate:

$$\exp\left(-\frac{\mathbf{i}}{\hbar} \hat{s}_z \theta\right) |+\rangle = \exp\left(-\mathbf{i} \frac{\theta}{2}\right) |+\rangle$$

For electrons you need a rotation of 4π to come back to the same state.

3.3 Eigenstates and eigenvalues of $\hat{\vec{J}}$

First (again) notice, that since $[\hat{J}_i, \hat{J}_k] \neq 0$ for $i \neq k$ there is not any common eigenstate for the $\hat{\vec{J}}$ s, but we know:

$$\left[\left| \hat{\vec{J}} \right|^2, \hat{J}_i \right] := \left[\hat{J}^2, \hat{J}_i \right] = 0$$

Therefore we can find a basis $|a, b\rangle$ with:

$$\begin{aligned} \hat{J}^2 |a, b\rangle &= a |a, b\rangle \\ \hat{J}_z |a, b\rangle &= b |a, b\rangle \end{aligned}$$

Now we introduce:

$$\begin{aligned} \hat{J}_{\pm} &:= \hat{J}_x \pm i\hat{J}_y \\ \hat{J}_{\pm}^{\dagger} &= \hat{J}_{\mp} \end{aligned}$$

Note that this gives:

$$\left[\hat{J}^2, \hat{J}_{\pm} \right] = 0 \qquad \left[\hat{J}_+, \hat{J}_- \right] = 2\hbar \hat{J}_z \qquad \left[\hat{J}_z, \hat{J}_{\pm} \right] = \pm \hbar \hat{J}_{\pm}$$

The trick is always the same:

$$\begin{aligned} \hat{J}_z \left(\hat{J}_{\pm} |a, b\rangle \right) &= \hat{J}_z \hat{J}_{\pm} |a, b\rangle = \left(\hat{J}_{\pm} \hat{J}_z + \left[\hat{J}_z, \hat{J}_{\pm} \right] \right) |a, b\rangle = \\ &= \hat{J}_{\pm} b |a, b\rangle \pm \hbar \hat{J}_{\pm} |a, b\rangle = (b \pm \hbar) \hat{J}_{\pm} |a, b\rangle \end{aligned}$$

Therefore $\hat{J}_{\pm} |a, b\rangle$ is an eigenstate of \hat{J}_z with the eigenvalue $b \pm \hbar$:

$$\hat{J}_{\pm} |a, b\rangle = N^{\pm}(a, b) |a, b \pm \hbar\rangle$$

Now we show, that $|b|$ has an upper limit. Consider

$$\langle a, b | \hat{J}_+ \hat{J}_+^{\dagger} |a, b\rangle + \langle a, b | \hat{J}_+^{\dagger} \hat{J}_+ |a, b\rangle \in \mathbb{R}_{\geq 0}$$

and:

$$\hat{J}_+ \hat{J}_+^{\dagger} + \hat{J}_+^{\dagger} \hat{J}_+ = \hat{J}_+ \hat{J}_- + \hat{J}_- \hat{J}_+ = 2 \left(\hat{J}^2 - \hat{J}_z^2 \right)$$

Then we get:

$$\begin{aligned} 0 &\leq \langle a, b | \hat{J}^2 - \hat{J}_z^2 |a, b\rangle = \langle a, b | a - b^2 |a, b\rangle = a - b^2 \\ a &\geq b^2 \\ |b| &\leq \sqrt{a} \end{aligned}$$

This means, that there is a b_{\max} defined by:

$$\hat{J}_+ |a, b_{\max}\rangle = 0$$

From this equation we get

$$\hat{J}_- \hat{J}_+ |a, b_{\max}\rangle = 0$$

and:

$$\hat{J}^2 - \hat{J}_z^2 - \hbar \hat{J}_z = a - b_{\max}^2 - \hbar b_{\max} = 0$$

This finally leads to:

$$a = b_{\max} (b_{\max} + \hbar)$$

Analogously we get for \hat{J}_- :

$$\hat{J}_- |a, b_{\min}\rangle = 0$$

From this equation we get

$$\hat{J}_+ \hat{J}_- |a, b_{\min}\rangle = 0$$

and:

$$\hat{J}^2 - \hat{J}_z^2 + \hbar \hat{J}_z = a - b_{\min}^2 + \hbar b_{\min} = 0$$

This finally leads to:

$$a = b_{\min} (b_{\min} - \hbar)$$

A solution is $b_{\max} = -b_{\min}$ and for a $n \in \mathbb{Z}$:

$$\begin{aligned} b_{\max} &= b_{\min} + n\hbar \\ b_{\max} - b_{\min} &= 2b_{\max} = n\hbar \\ b_{\max} &= \frac{n}{2}\hbar \\ a &= \frac{n}{2}\hbar \left(\frac{n}{2}\hbar + \hbar \right) = \hbar^2 \underbrace{\frac{n}{2}}_{=:j} \left(\frac{n}{2} + 1 \right) \end{aligned}$$

Some names:

$j \in \frac{1}{2}\mathbb{Z}$ is called *the total angular momentum quantum number* and

$$m \in \{-j, \dots, -j+1, \dots, j-1, j\}$$

is called *the magnetic quantum number*:

$$b = m\hbar \qquad a = \hbar^2 j(j+1)$$

$$\hat{J}^2 |j, m\rangle = \hbar^2 j(j+1) |j, m\rangle$$

$$\hat{J}_z |j, m\rangle = \hbar m |j, m\rangle$$

With a bit of algebra we get:

$$\hat{J}_{\pm} |j, m\rangle = \hbar \sqrt{(j \mp m)(j \pm m + 1)} |j, m \pm 1\rangle$$

3.4 Irreducible representations of the rotation operators

We want to construct the matrix representations of $\hat{D}(\vec{n}, \theta)$.

In particular in the eigenbasis of \hat{J}^2 and \hat{J}_z .

Remember:

$$\hat{D}(\vec{n}, \theta) = e^{-\frac{i}{\hbar} \hat{\vec{J}} \cdot \vec{n} \theta}$$

We want:

$$\langle j, m | \hat{D}(\vec{n}, \theta) | j', m' \rangle = D_{jm}^{j'm'}(\vec{n}, \theta)$$

First note:, because $[\hat{\vec{J}}^2, \hat{\vec{J}} \cdot \vec{n}] = 0$.

$$D_{jm}^{j'm'}(\vec{n}, \theta) = D_{mm'}^{(j)}(\vec{n}, \theta) \delta_{jj'}$$

The physical reason is, that a rotation doesn't change magnitudes like j .

$j = \frac{1}{2}$ gives a 2×2 -Matrix for D , because the basis is $|\frac{1}{2}, -\frac{1}{2}\rangle$ and $|\frac{1}{2}, \frac{1}{2}\rangle$.

$$\left(\begin{array}{cc} \begin{array}{c} \xleftrightarrow{2} \\ \left(\begin{array}{c} j = \frac{1}{2} \end{array} \right) \updownarrow 2 \\ \end{array} & 0 \\ 0 & \begin{array}{c} \xleftrightarrow{3} \\ \left(\begin{array}{c} j = 1 \end{array} \right) \updownarrow 3 \\ \vdots \end{array} \end{array} \right)$$

This is the matrix representation of a rotation operator $\hat{D}(\vec{n}, \theta)$ in the basis $|j, m\rangle$ ordered as

$$\left| \frac{1}{2}, -\frac{1}{2} \right\rangle, \left| \frac{1}{2}, +\frac{1}{2} \right\rangle, |1, -1\rangle, |1, 0\rangle, |1, 1\rangle, \left| \frac{3}{2}, -\frac{3}{2} \right\rangle, \left| \frac{3}{2}, -\frac{1}{2} \right\rangle, \left| \frac{3}{2}, \frac{1}{2} \right\rangle, \left| \frac{3}{2}, \frac{3}{2} \right\rangle, \dots$$

and it is BLOCK-DIAGONAL.

3.5 Orbital angular momentum

Consider now $\hat{J}_i = \hat{L}_i = (\hat{\vec{q}} \times \hat{\vec{p}})_i$.

3.5.1 General considerations

$$\text{i) } [\hat{L}_i, \hat{L}_j] = i\hbar \varepsilon_{ijk} \hat{L}_k$$

ii) Consider:

$$\exp\left(-\frac{\mathbf{i}}{\hbar}\hat{L}_z\theta\right)|\vec{q}\rangle =: |\vec{q}\rangle^R \stackrel{?}{=} |\vec{q}^R\rangle := |\mathcal{R}(\vec{e}_z, \theta)\vec{q}\rangle$$

Do it infinitesimal:

$$\theta \rightarrow d\theta$$

$$\begin{aligned} |\vec{q}\rangle^R &\approx \left(1 - \frac{\mathbf{i}}{\hbar}\hat{L}_z d\theta\right)|\vec{q}\rangle = \left(1 - \frac{\mathbf{i}}{\hbar}d\theta(\hat{q}_x\hat{p}_y - \hat{q}_y\hat{p}_x)\right)|\vec{q}\rangle = \\ &= \left(1 - \frac{\mathbf{i}}{\hbar}d\theta(q_x\hat{p}_y - q_y\hat{p}_x)\right)|q_x\rangle \otimes |q_y\rangle \otimes |q_z\rangle \end{aligned}$$

Remember:

$$\exp\left(-\frac{\mathbf{i}}{\hbar}q^{(0)}\hat{p}_x\right)|\vec{q}\rangle = |q_x + q^{(0)}\rangle \otimes |q_y\rangle \otimes |q_z\rangle$$

Infinitesimally we get:

$$\left(1 - \frac{\mathbf{i}}{\hbar}dq^{(0)}\hat{p}_x\right)|\vec{q}\rangle = |q_x + dq^{(0)}\rangle \otimes |q_y\rangle \otimes |q_z\rangle$$

Now consider the product of two infinitesimal changes:

$$\left(1 + \frac{\mathbf{i}}{\hbar}d\theta q_y\hat{p}_x\right)\left(1 - \frac{\mathbf{i}}{\hbar}d\theta q_x\hat{p}_y\right) \approx 1 - \frac{\mathbf{i}}{\hbar}d\theta(q_x\hat{p}_y - q_y\hat{p}_x)$$

This gives:

$$\begin{aligned} |\vec{q}\rangle^R &\approx \left(1 + \frac{\mathbf{i}}{\hbar}d\theta q_y\hat{p}_x\right)\left(1 - \frac{\mathbf{i}}{\hbar}d\theta q_x\hat{p}_y\right)|q_x\rangle \otimes |q_y\rangle \otimes |q_z\rangle = \\ &= |q_x - d\theta q_y\rangle \otimes |q_y + d\theta q_x\rangle \otimes |q_z\rangle = |q_x - d\theta q_y, q_y + d\theta q_x, q_z\rangle \end{aligned}$$

Compare this with the rotated eigenstate of the position:

$$|\vec{q}^R\rangle = |\mathcal{R}(\vec{e}_z, d\theta)\vec{q}\rangle \approx \left| \begin{pmatrix} 1 & -d\theta & 0 \\ d\theta & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \vec{q} \right\rangle = |q_x - d\theta q_y, q_y + d\theta q_x, q_z\rangle \approx |\vec{q}\rangle^R$$

In general we get:

$$\exp\left(-\frac{\mathbf{i}}{\hbar}\hat{\vec{L}} \cdot \vec{n}\eta\right)|\vec{q}\rangle = |\mathcal{R}(\vec{n}, \eta)\vec{q}\rangle$$

3.5.2 Eigenstates in the position representation

We call the eigenstates $|l, m\rangle$.

We introduce:

$$\begin{aligned} \hat{r} &= \sqrt{\hat{q}_x^2 + \hat{q}_y^2 + \hat{q}_z^2} \\ \hat{q}_x &= \hat{r} \sin \hat{\theta} \cos \hat{\varphi} \\ \hat{q}_y &= \hat{r} \sin \hat{\theta} \sin \hat{\varphi} \\ \hat{q}_z &= \hat{r} \cos \theta \end{aligned}$$

$$|\vec{q}\rangle = |r\rangle \otimes |\theta, \varphi\rangle = |r\rangle \otimes |\vec{n}\rangle$$

We want to construct $\langle \theta, \varphi | l, m \rangle$. As usual we start with

$$\langle \theta, \varphi | \hat{L}_z | l, m \rangle = \hbar m \langle \theta, \varphi | l, m \rangle$$

and do it infinitesimally:

$$\langle \theta, \varphi | \hat{D}(\vec{e}_z, d\varphi) | l, m \rangle = \langle \theta, \varphi | \exp\left(-\frac{\mathbf{i}}{\hbar} \hat{L}_z d\varphi\right) | l, m \rangle = \langle \theta, \varphi - d\varphi | l, m \rangle$$

We also get for $d\varphi \rightarrow 0$:

$$\left\langle \theta, \varphi \left| \left(1 - \frac{\mathbf{i}}{\hbar} \hat{L}_z d\varphi\right) \right| l, m \right\rangle = \langle \theta, \varphi | l, m \rangle - \frac{\mathbf{i}}{\hbar} d\varphi \langle \theta, \varphi | \hat{L}_z | l, m \rangle$$

This gives:

$$\begin{aligned} m\hbar \langle \theta, \varphi | l, m \rangle &= \langle \theta, \varphi | \hat{L}_z | l, m \rangle = -\mathbf{i}\hbar \partial_\varphi \langle \theta, \varphi | l, m \rangle \\ \partial_\varphi \langle \theta, \varphi | l, m \rangle &= \mathbf{i}m \langle \theta, \varphi | l, m \rangle \end{aligned}$$

$$\langle \theta, \varphi | l, m \rangle = f_l^m(\theta) e^{\mathbf{i}m\varphi}$$

Introduce this into

$$\langle \theta, \varphi | \hat{L}^2 | l, m \rangle = \hbar^2 l(l+1) \langle \theta, \varphi | l, m \rangle$$

$$-\hbar^2 \left(\frac{1}{\sin^2 \theta} \partial_\varphi^2 + \frac{1}{\sin \theta} \partial_\theta (\sin \theta \partial_\theta) \right) \langle \theta, \varphi | l, m \rangle = \hbar^2 l(l+1) \langle \theta, \varphi | l, m \rangle$$

to get for $m, l \in \mathbb{Z}$:

$$\left(\frac{1}{\sin \theta} \frac{d}{d\theta} (\sin \theta \partial_\theta) - \frac{m^2}{\sin^2 \theta} + l(l+1) \right) f_l^m(\theta) = 0$$

Transform $x = \cos \theta$ and solve this by a power series. The solutions $P_l^m(\cos \theta)$ are the *associated Legendre polynomials*.

Together we have the *spherical harmonics*:

$$\langle \theta, \varphi | l, m \rangle = Y_l^m(\theta, \varphi) = (-1)^m \sqrt{\frac{2l+1}{4\pi} \cdot \frac{(l-m)!}{(l+m)!}} \cdot e^{\mathbf{i}m\varphi} P_l^m(\cos \theta)$$

Of course they are normalized:

$$\begin{aligned} \langle l', m' | l, m \rangle &= \delta_{ll'} \delta_{mm'} \\ \Rightarrow \int_0^{2\pi} \int_0^\pi Y_{l'}^{m'}(\theta, \varphi)^* Y_l^m(\theta, \varphi) \sin \theta d\theta d\varphi &= \delta_{ll'} \delta_{mm'} \end{aligned}$$

Therefore any function of

$$\vec{n} = \begin{pmatrix} \sin \theta \cos \varphi \\ \sin \theta \sin \varphi \\ \cos \theta \end{pmatrix}$$

can be expanded in terms of $Y_l^m(\theta, \varphi)$'s, because they form a complete basis:

$$\sum_{l=0}^{\infty} \sum_{m=-l}^l |l, m\rangle \langle l, m| = \hat{I}_{\text{angular}}$$

$$\sum_{l=0}^{\infty} \sum_{m=-l}^l Y_l^m(\theta, \varphi) Y_l^m(\theta', \varphi') = \delta(\vec{n} - \vec{n}')$$

From the general theory we also know:

$$\hat{L}_{\pm} |l, m\rangle = \hbar \sqrt{(l \mp m)(l \pm m + 1)} |l, m \pm 1\rangle$$

For an arbitrary normalized state $|\alpha\rangle$ in the angular Hilbert space one gets:

$$\langle \theta, \varphi | \hat{L}_- | \alpha \rangle = i\hbar e^{i\varphi} [i\partial_{\theta} - \cot \theta] \langle \theta, \varphi | \alpha \rangle$$

For example:

$$Y_l^{l-1}(\theta, \varphi) = -N_l^l e^{i(l-1)\varphi} (l \cot \theta - \partial_{\theta}) P_l^l(\cos \theta)$$

$$P_l^l(\cos \theta) = \sin^l(\theta)$$

$$Y_l^{-m}(\theta, \varphi) = (-1)^m (Y_l^m(\theta, \varphi))^*$$

3.5.3 Relation with the kinetic energy

Remember from classical mechanics:

$$H(\vec{q}, \vec{p}) = \frac{1}{2m} p^2 + V(\vec{q})$$

In spherical coordinates we have the canonical transformation

$$\begin{pmatrix} q_x \\ q_y \\ q_z \end{pmatrix} \rightarrow \begin{pmatrix} r \\ \theta \\ \varphi \end{pmatrix} \quad \Rightarrow \quad \begin{pmatrix} p_x \\ p_y \\ p_z \end{pmatrix} \rightarrow \begin{pmatrix} p_r \\ p_{\theta} \\ p_{\varphi} \end{pmatrix}$$

with the commutators

$$\{r, p_r\} = \{\theta, p_{\theta}\} = \{\varphi, p_{\varphi}\} = 1$$

and the other commutators are zero. The Hamiltonian is then

$$H(r, \theta, \varphi, p_r, p_{\theta}, p_{\varphi}) = \frac{1}{2m} p_r^2 + \frac{1}{2mr^2} L^2 + V(r, \theta, \varphi)$$

with:

$$L^2 = \frac{1}{\sin^2 \theta} p_{\varphi}^2 + p_{\theta}^2$$

To see this in quantum mechanics, we use

$$\hat{L}^2 = \hat{q}^2 \hat{p}^2 - (\hat{\vec{q}} \cdot \hat{\vec{p}})^2 + i\hbar \hat{\vec{q}} \cdot \hat{\vec{p}}$$

and calculate for an arbitrary state $|\alpha\rangle$:

$$\langle \vec{q} | \hat{L}^2 | \alpha \rangle = q^2 \langle \vec{q} | \hat{p}^2 | \alpha \rangle - \vec{q} \cdot (-i\hbar \vec{\nabla}) \cdot \vec{q} \cdot (-i\hbar \vec{\nabla}) \langle \vec{q} | \alpha \rangle + i\hbar \vec{q} \cdot (-i\hbar \vec{\nabla}) \langle \vec{q} | \alpha \rangle$$

$$\left\langle \vec{q} \left| \frac{1}{2m} \hat{p}^2 \right| \alpha \right\rangle = -\frac{\hbar^2}{2m} \left(\frac{1}{r} \partial_r (r \partial_r) + \frac{1}{r} \partial_r \right) \langle \vec{q} | \alpha \rangle + \frac{1}{2mr^2} \langle \vec{q} | \hat{L}^2 | \alpha \rangle =$$

$$= -\frac{\hbar^2}{2m} \left(\frac{1}{r^2} \partial_r (r^2 \partial_r) \right) \langle \vec{q} | \alpha \rangle + \frac{1}{2mr^2} \langle \vec{q} | \hat{L}^2 | \alpha \rangle$$

3.5.4 Central forces

Definition:

A central force has a potential, that depends only on the radial coordinate:

$$V(\vec{q}) = V(\hat{r})$$

Therefore it commutes with \hat{p}^2 and \hat{L}^2 and we have a set of common eigenstates:

$$\begin{aligned}\hat{H} |E, l, m\rangle &= E |E, l, m\rangle \\ \hat{L}^2 |E, l, m\rangle &= \hbar^2 l(l+1) |E, l, m\rangle \\ \hat{L}_z |E, l, m\rangle &= \hbar m |E, l, m\rangle\end{aligned}$$

We define

$$\psi_{Elm}(r, \theta, \varphi) := \langle r, \theta, \varphi | E, l, m \rangle = R_{Elm}(r) Y_l^m(\theta, \varphi)$$

and get:

$$\left(-\frac{\hbar^2}{2mr^2} \frac{d}{dr} \left(r^2 \frac{d}{dr} \right) + \frac{\hbar^2 l(l+1)}{2mr^2} + V(r) \right) R_{Elm}(r) = E R_{Elm}(r)$$

This doesn't depend on the quantum number m , so we just write $R_{Elm}(r)$. Now we transform

$$R_{El}(r) := \frac{U_{El}(r)}{r}$$

and get:

$$\left(-\frac{\hbar^2}{2m} \frac{d^2}{dr^2} + \underbrace{\left(V(r) + \frac{\hbar^2 l(l+1)}{2mr^2} \right)}_{=: V_{\text{eff}}(r)} \right) U_{El}(r) = E U_{El}(r)$$

3.6 Addition of angular momentum

Consider two particles with the same mass and momenta $\vec{p}^{(1)}$ and $\vec{p}^{(2)}$. In classical mechanics they generate translations ($k \in \{1, 2\}$):

$$f(\vec{q}^{(k)} + \delta \vec{q}^{(k)}, \vec{p}^{(k)}) \approx f(\vec{q}^{(k)}, \vec{p}^{(k)}) + \left\{ \vec{p}^{(k)}, f(\vec{q}^{(k)}) \right\} \cdot \delta \vec{q}^{(k)} + O\left(\left(\delta \vec{q}^{(k)}\right)^2\right)$$

$\vec{p} := (\vec{p}^{(1)}, \vec{p}^{(2)})$ generates translations of $\vec{q} := (\vec{q}^{(1)}, \vec{q}^{(2)})$:

$$f(\vec{q} + \delta \vec{q}, \vec{p}) \approx f(\vec{q}, \vec{p}) + \left\{ \vec{p}, f(\vec{q}) \right\} \cdot \delta \vec{q} + O\left((\delta \vec{q})^2\right)$$

What is the meaning (in the sense of generators) of:

$$\vec{P} := \vec{p}^{(1)} + \vec{p}^{(2)}$$

Remember that \vec{P} and $\vec{Q} = \frac{1}{2}(\vec{q}^{(1)} + \vec{q}^{(2)})$ are canonical, that means:

$$\{Q_i, P_j\} = \delta_{ij}$$

Therefore \vec{P} generates translations in \vec{Q} .

$$f\left(\vec{q}^{(1)} + \delta\vec{Q}, \vec{q}^{(2)} + \delta\vec{Q}, \vec{p}^{(1)}, \vec{p}^{(2)}\right) = f\left(\vec{q}^{(1)}, \vec{q}^{(2)}, \vec{p}^{(1)}, \vec{p}^{(2)}\right) + \left\{\vec{P}, f\left(\vec{q}^{(1)}, \vec{q}^{(2)}, \vec{p}^{(1)}, \vec{p}^{(2)}\right)\right\} \cdot \delta\vec{Q} + O\left(\left(\delta\vec{Q}\right)^2\right)$$

In the same way $\vec{L}^{(1)} + \vec{L}^{(2)}$ generates a RIGID rotation of the total system.

3.6.1 Formal theory (2 “particles”)

We start with two angular momenta $\hat{J}^{(1)}$ and $\hat{J}^{(2)}$ with $(n \in \{1, 2\})$:

$$\begin{aligned} \left[\hat{J}_i^{(n)}, \hat{J}_j^{(n)}\right] &= i\hbar \varepsilon_{ijk} \hat{J}_k^{(n)} \\ \left[\hat{J}_i^{(1)}, \hat{J}_j^{(2)}\right] &= 0 \end{aligned}$$

The “total” angular momentum is:

$$\hat{J}_i := \hat{J}_i^{(1)} \otimes \hat{I}^{(2)} + \hat{I}^{(1)} \otimes \hat{J}_i^{(2)} =: \hat{J}_i^{(1)} + \hat{J}_i^{(2)}$$

And a “total” rotation by \vec{n}, θ is:

$$\hat{D}(\vec{n}, \theta) := \hat{D}(\vec{n}, \theta)^{(1)} \otimes \hat{D}(\vec{n}, \theta)^{(2)} = e^{-\frac{i}{\hbar} \hat{J}^{(1)} \cdot \vec{n} \theta} \cdot e^{-\frac{i}{\hbar} \hat{J}^{(2)} \cdot \vec{n} \theta} = e^{-\frac{i}{\hbar} \hat{J} \cdot \vec{n} \theta}$$

Check whether \hat{J} is an angular momentum:

$$\left[\hat{J}_i, \hat{J}_j\right] = i\hbar \varepsilon_{ijk} \hat{J}_k$$

So we have two possible sets of commuting operators:

1. DECOUPLED BASIS: Take $\left\{\left(\hat{J}^{(1)}\right)^2, \left(\hat{J}^{(2)}\right)^2, \hat{J}_z^{(1)}, \hat{J}_z^{(2)}\right\}$ with the common eigenstates:

$$|j_1, j_2, m_1, m_2\rangle := |j_1, m_1\rangle \otimes |j_2, m_2\rangle$$

For $k \in \{1, 2\}$ we have:

$$\begin{aligned} \left(\hat{J}^{(k)}\right)^2 |j_1, j_2, m_1, m_2\rangle &= \hbar^2 (j_k + 1) |j_1, j_2, m_1, m_2\rangle \\ \hat{J}_z^{(k)} |j_1, j_2, m_1, m_2\rangle &= \hbar m_k |j_1, j_2, m_1, m_2\rangle \end{aligned}$$

2. COUPLED BASIS: Take $\left\{\left(\hat{J}^{(1)}\right)^2, \left(\hat{J}^{(2)}\right)^2, \hat{J}^2, \hat{J}_z\right\}$ – observe $\left[\hat{J}^2, \hat{J}_z^{(k)}\right] \neq 0$ – with the common eigenstates $|j_1, j_2, j, m\rangle$ with:

$$\begin{aligned} \hat{J}^2 |j_1, j_2, j, m\rangle &= \hbar^2 j(j+1) |j_1, j_2, j, m\rangle \\ \hat{J}_z |j_1, j_2, j, m\rangle &= \hbar m |j_1, j_2, j, m\rangle \end{aligned}$$

The two bases are related (for fixed j_1, j_2):

$$|j_1, j_2, j, m\rangle = \sum_{m_1=-j_1}^{j_1} \sum_{m_2=-j_2}^{j_2} \underbrace{\langle j_1, j_2, m_1, m_2 | j_1, j_2, j, m \rangle}_{\text{Lebsch-Gordan coefficients}} |j_1, j_2, m_1, m_2\rangle$$

Some properties of $\langle m_1, m_2 | j, m \rangle := \langle j_1, j_2, m_1, m_2 | j_1, j_2, j, m \rangle$:

Consider

$$\hat{J}_z - \hat{J}_z^{(1)} - \hat{J}_z^{(2)} = 0$$

and sandwich between $\langle j_1, j_2, m_1, m_2 | \dots | j_1, j_2, j, m \rangle$ to get:

$$(m - m_1 - m_2) \langle m_1, m_2 | j, m \rangle = 0$$

Therefore $\langle m_1, m_2 | j, m \rangle = 0$ for $m \neq m_1 + m_2$ and one can eliminate one of the sums above. One can also proof:

$$|j_1 - j_2| \leq j \leq j_1 + j_2$$

From unitarity of the matrix for the change of basis we get:

$$\begin{aligned} \sum_{m_1=-j_1}^{j_1} \sum_{m_2=-j_2}^{j_2} \langle m_1, m_2 | j, m \rangle \langle m_1, m_2 | j', m' \rangle^* &= \delta_{jj'} \delta_{mm'} \\ \sum_{j=|j_1-j_2|}^{j_1+j_2} \sum_{m=-j}^j \langle m_1, m_2 | j, m \rangle \langle m'_1, m'_2 | j, m \rangle^* &= \delta_{m_1 m'_1} \delta_{m_2 m'_2} \end{aligned}$$

Finally using $\hat{J}_{\pm} = \hat{J}_{\pm}^{(1)} + \hat{J}_{\pm}^{(2)}$ gives:

$$\begin{aligned} \sqrt{(j \mp m)(j \pm m + 1)} \langle m_1, m_2 | j, m \pm 1 \rangle &= \sqrt{(j_1 \pm m_1)(j_1 \mp m_1 + 1)} \langle m_1 \mp 1, m_2 | j, m \rangle + \\ &+ \sqrt{(j_2 \pm m_2)(j_2 \mp m_2 + 1)} \langle m_1, m_2 \pm 1 | j, m \rangle \end{aligned}$$

3.6.2 The formal theory for many particles

This is a REAL MESS!!

3.6.3 Example: Two spin $\frac{1}{2}$ particles

Here the total spin is:

$$\hat{S} = \hat{S}^{(1)} + \hat{S}^{(2)} \quad \hat{S}_z = \hat{S}_z^{(1)} + \hat{S}_z^{(2)} \quad \hat{S}^2 = \left(\hat{S}^{(1)}\right)^2 + \left(\hat{S}^{(2)}\right)^2 + 2\hat{S}^{(1)} \cdot \hat{S}^{(2)}$$

And the quantum numbers are $j_1 = s_1 = \frac{1}{2}$, $j_2 = s_2 = \frac{1}{2}$. The total is $m = m_1 + m_2$ and s is restricted to:

$$\begin{aligned} \left| \frac{1}{2} - \frac{1}{2} \right| &\leq s \leq \frac{1}{2} + \frac{1}{2} \\ 0 &\leq s \leq 1 \end{aligned}$$

- The decoupled basis ($|m_1, m_2\rangle$) is:

$$\left\{ \left| \frac{1}{2}, \frac{1}{2} \right\rangle, \left| \frac{1}{2}, -\frac{1}{2} \right\rangle, \left| -\frac{1}{2}, \frac{1}{2} \right\rangle, \left| -\frac{1}{2}, -\frac{1}{2} \right\rangle \right\}$$

Remember that this basis does not have a well defined \hat{S}^2 .

- The coupled basis ($|s, m\rangle$) is:

$$\begin{aligned} \{|s=0, m=0\rangle, |s=1, m=-1\rangle, |s=1, m=0\rangle, |s=1, m=1\rangle\} = \\ = \{ \underbrace{|0,0\rangle}_{\text{"Singlet"}}, \underbrace{|1, -1\rangle, |1,0\rangle, |1,1\rangle}_{\text{"Triplet" states}} \} \end{aligned}$$

Using $m = m_1 + m_2$ we get:

$$\begin{aligned} |s=1, m=1\rangle &= \left| \frac{1}{2}, \frac{1}{2} \right\rangle := \left| \frac{1}{2} \right\rangle^{(1)} \otimes \left| \frac{1}{2} \right\rangle^{(2)} \\ |s=1, m=0\rangle &= \frac{1}{\sqrt{2}} \left(\left| \frac{1}{2}, -\frac{1}{2} \right\rangle + \left| -\frac{1}{2}, \frac{1}{2} \right\rangle \right) \\ |s=1, m=-1\rangle &= \left| -\frac{1}{2}, -\frac{1}{2} \right\rangle \\ |s=0, m=0\rangle &= \frac{1}{\sqrt{2}} \left(\left| \frac{1}{2}, -\frac{1}{2} \right\rangle - \left| -\frac{1}{2}, \frac{1}{2} \right\rangle \right) \end{aligned}$$

(Take a look at: SAKURAI, Section 3.10: “*Spin Correlation measurements and Bell’s inequality*”)

Think about the Hamiltonian for $k \in \mathbb{R}_{<0}$:

$$\begin{aligned} \hat{H} = \underbrace{\hbar\omega \left(\hat{S}_z^{(1)} + \hat{S}_z^{(2)} \right)}_{\substack{\hat{S}_z^{(1)}, \hat{S}_z^{(2)} \text{ are conserved} \\ \hat{S}^2, \hat{S}_z \text{ are conserved}}} + k \hat{\vec{S}}^{(1)} \cdot \hat{\vec{S}}^{(2)} \end{aligned}$$

We use:

$$\hat{\vec{S}}^{(1)} \cdot \hat{\vec{S}}^{(2)} = \frac{1}{2} \left(\hat{S}^2 - \left(\hat{S}^{(1)} \right)^2 - \left(\hat{S}^{(2)} \right)^2 \right)$$

4 Perturbation theory (time-independent)

5 Perturbation theory (time-dependent)

6 Many-particle systems

7 Scattering theory

Appendix

Acknowledgements

My special thanks goes to Doctor Urbina, who gave this lecture and allowed me to publish this script of the lecture.

I would also like to thank all those, who found errors by careful reading and told me of them.

Andreas Völklein

GNU Free Documentation License

Version 1.3, 3 November 2008

Copyright © 2000, 2001, 2002, 2007, 2008 Free Software Foundation, Inc.

`<https://fsf.org/>`

Everyone is permitted to copy and distribute verbatim copies of this license document, but changing it is not allowed

0. PREAMBLE

The purpose of this License is to make a manual, textbook, or other functional and useful document “free” in the sense of freedom: to assure everyone the effective freedom to copy and redistribute it, with or without modifying it, either commercially or noncommercially. Secondly, this License preserves for the author and publisher a way to get credit for their work, while not being considered responsible for modifications made by others.

This License is a kind of “copyleft”, which means that derivative works of the document must themselves be free in the same sense. It complements the GNU General Public License, which is a copyleft license designed for free software.

We have designed this License in order to use it for manuals for free software, because free software needs free documentation: a free program should come with manuals providing the same freedoms that the software does. But this License is not limited to software manuals; it can be used for any textual work, regardless of subject matter or whether it is published as a printed book. We recommend this License principally for works whose purpose is instruction or reference.

1. APPLICABILITY AND DEFINITIONS

This License applies to any manual or other work, in any medium, that contains a notice placed by the copyright holder saying it can be distributed under the terms of this License. Such a notice grants a world-wide, royalty-free license, unlimited in duration, to use that work under the conditions stated herein. The “**Document**”, below, refers to any such manual or work. Any member of the public is a licensee, and is addressed as “**you**”. You accept the license if you copy, modify or distribute the work in a way requiring permission under copyright law.

A “**Modified Version**” of the Document means any work containing the Document or a portion of it, either copied verbatim, or with modifications and/or translated into another language.

A “**Secondary Section**” is a named appendix or a front-matter section of the Document that deals exclusively with the relationship of the publishers or authors of the Document to the Document’s overall subject (or to related matters) and contains nothing that could fall directly within that overall subject. (Thus, if the Document is in part a textbook of mathematics, a Secondary Section may not explain any mathematics.) The relationship could be a matter of historical connection with the subject or with related matters, or of legal, commercial, philosophical, ethical or political position regarding them.

The “**Invariant Sections**” are certain Secondary Sections whose titles are designated, as being those of Invariant Sections, in the notice that says that the Document is released under this License. If a section does not fit the above definition of Secondary then it is not allowed to be designated as Invariant. The Document may contain zero Invariant Sections. If the Document does not identify any Invariant Sections then there are none.

The “**Cover Texts**” are certain short passages of text that are listed, as Front-Cover Texts or Back-Cover Texts, in the notice that says that the Document is released under this License. A Front-Cover Text may be at most 5 words, and a Back-Cover Text may be at most 25 words.

A “**Transparent**” copy of the Document means a machine-readable copy, represented in a format whose specification is available to the general public, that is suitable for revising the document straightforwardly with generic text editors or (for images composed of pixels) generic paint programs or (for drawings) some widely available drawing editor, and that is suitable for input to text formatters or for automatic translation to a variety of formats suitable for input to text formatters. A copy made in an otherwise Transparent file format whose markup, or absence of markup, has been arranged to thwart or discourage subsequent modification by readers is not Transparent. An image format is not Transparent if used for any substantial amount of text. A copy that is not “Transparent” is called “**Opaque**”.

Examples of suitable formats for Transparent copies include plain ASCII without markup, Texinfo input format, L^AT_EX input format, SGML or XML using a publicly available DTD, and standard-conforming simple HTML, PostScript or PDF designed for human modification. Examples of transparent image formats include PNG, XCF and JPG. Opaque formats include proprietary formats that can be read and edited only by proprietary word processors, SGML or XML for which the DTD and/or processing tools are not generally available, and the machine-generated HTML, PostScript or PDF produced by some word processors for output purposes only.

The “**Title Page**” means, for a printed book, the title page itself, plus such following pages as are needed to hold, legibly, the material this License requires to appear in the title page. For works in formats which do not have any title page as such, “Title Page” means the text near the most prominent appearance of the work’s title, preceding the beginning of the body of the text.

The “**publisher**” means any person or entity that distributes copies of the Document to the public.

A section “**Entitled XYZ**” means a named subunit of the Document whose title either is precisely XYZ or contains XYZ in parentheses following text that translates XYZ in another language. (Here XYZ stands for a specific section name mentioned below, such as “**Acknowledgements**”, “**Dedications**”, “**Endorsements**”, or “**History**”.) To “**Preserve the Title**” of such a section when you modify the Document means that it remains a section “Entitled XYZ” according to this definition.

The Document may include Warranty Disclaimers next to the notice which states that this License applies to the Document. These Warranty Disclaimers are considered to be included by reference in this License, but only as regards disclaiming warranties: any other implication that

these Warranty Disclaimers may have is void and has no effect on the meaning of this License.

2. VERBATIM COPYING

You may copy and distribute the Document in any medium, either commercially or noncommercially, provided that this License, the copyright notices, and the license notice saying this License applies to the Document are reproduced in all copies, and that you add no other conditions whatsoever to those of this License. You may not use technical measures to obstruct or control the reading or further copying of the copies you make or distribute. However, you may accept compensation in exchange for copies. If you distribute a large enough number of copies you must also follow the conditions in section 3.

You may also lend copies, under the same conditions stated above, and you may publicly display copies.

3. COPYING IN QUANTITY

If you publish printed copies (or copies in media that commonly have printed covers) of the Document, numbering more than 100, and the Document's license notice requires Cover Texts, you must enclose the copies in covers that carry, clearly and legibly, all these Cover Texts: Front-Cover Texts on the front cover, and Back-Cover Texts on the back cover. Both covers must also clearly and legibly identify you as the publisher of these copies. The front cover must present the full title with all words of the title equally prominent and visible. You may add other material on the covers in addition. Copying with changes limited to the covers, as long as they preserve the title of the Document and satisfy these conditions, can be treated as verbatim copying in other respects.

If the required texts for either cover are too voluminous to fit legibly, you should put the first ones listed (as many as fit reasonably) on the actual cover, and continue the rest onto adjacent pages.

If you publish or distribute Opaque copies of the Document numbering more than 100, you must either include a machine-readable Transparent copy along with each Opaque copy, or state in or with each Opaque copy a computer-network location from which the general network-using public has access to download using public-standard network protocols a complete Transparent copy of the Document, free of added material. If you use the latter option, you must take reasonably prudent steps, when you begin distribution of Opaque copies in quantity, to ensure that this Transparent copy will remain thus accessible at the stated location until at least one year after the last time you distribute an Opaque copy (directly or through your agents or retailers) of that edition to the public.

It is requested, but not required, that you contact the authors of the Document well before redistributing any large number of copies, to give them a chance to provide you with an updated version of the Document.

4. MODIFICATIONS

You may copy and distribute a Modified Version of the Document under the conditions of sections 2 and 3 above, provided that you release the Modified Version under precisely this License, with the Modified Version filling the role of the Document, thus licensing distribution

and modification of the Modified Version to whoever possesses a copy of it. In addition, you must do these things in the Modified Version:

- A. Use in the Title Page (and on the covers, if any) a title distinct from that of the Document, and from those of previous versions (which should, if there were any, be listed in the History section of the Document). You may use the same title as a previous version if the original publisher of that version gives permission.
- B. List on the Title Page, as authors, one or more persons or entities responsible for authorship of the modifications in the Modified Version, together with at least five of the principal authors of the Document (all of its principal authors, if it has fewer than five), unless they release you from this requirement.
- C. State on the Title page the name of the publisher of the Modified Version, as the publisher.
- D. Preserve all the copyright notices of the Document.
- E. Add an appropriate copyright notice for your modifications adjacent to the other copyright notices.
- F. Include, immediately after the copyright notices, a license notice giving the public permission to use the Modified Version under the terms of this License, in the form shown in the Addendum below.
- G. Preserve in that license notice the full lists of Invariant Sections and required Cover Texts given in the Document's license notice.
- H. Include an unaltered copy of this License.
- I. Preserve the section Entitled "History", Preserve its Title, and add to it an item stating at least the title, year, new authors, and publisher of the Modified Version as given on the Title Page. If there is no section Entitled "History" in the Document, create one stating the title, year, authors, and publisher of the Document as given on its Title Page, then add an item describing the Modified Version as stated in the previous sentence.
- J. Preserve the network location, if any, given in the Document for public access to a Transparent copy of the Document, and likewise the network locations given in the Document for previous versions it was based on. These may be placed in the "History" section. You may omit a network location for a work that was published at least four years before the Document itself, or if the original publisher of the version it refers to gives permission.
- K. For any section Entitled "Acknowledgements" or "Dedications", Preserve the Title of the section, and preserve in the section all the substance and tone of each of the contributor acknowledgements and/or dedications given therein.
- L. Preserve all the Invariant Sections of the Document, unaltered in their text and in their titles. Section numbers or the equivalent are not considered part of the section titles.
- M. Delete any section Entitled "Endorsements". Such a section may not be included in the Modified Version.
- N. Do not retitle any existing section to be Entitled "Endorsements" or to conflict in title with any Invariant Section.
- O. Preserve any Warranty Disclaimers.

If the Modified Version includes new front-matter sections or appendices that qualify as Secondary Sections and contain no material copied from the Document, you may at your option designate some or all of these sections as invariant. To do this, add their titles to the list of Invariant Sections in the Modified Version's license notice. These titles must be distinct from any other section titles.

You may add a section Entitled "Endorsements", provided it contains nothing but endorsements of your Modified Version by various parties—for example, statements of peer review or that the text has been approved by an organization as the authoritative definition of a standard.

You may add a passage of up to five words as a Front-Cover Text, and a passage of up to 25 words as a Back-Cover Text, to the end of the list of Cover Texts in the Modified Version. Only one passage of Front-Cover Text and one of Back-Cover Text may be added by (or through arrangements made by) any one entity. If the Document already includes a cover text for the same cover, previously added by you or by arrangement made by the same entity you are acting on behalf of, you may not add another; but you may replace the old one, on explicit permission from the previous publisher that added the old one.

The author(s) and publisher(s) of the Document do not by this License give permission to use their names for publicity for or to assert or imply endorsement of any Modified Version.

5. COMBINING DOCUMENTS

You may combine the Document with other documents released under this License, under the terms defined in section 4 above for modified versions, provided that you include in the combination all of the Invariant Sections of all of the original documents, unmodified, and list them all as Invariant Sections of your combined work in its license notice, and that you preserve all their Warranty Disclaimers.

The combined work need only contain one copy of this License, and multiple identical Invariant Sections may be replaced with a single copy. If there are multiple Invariant Sections with the same name but different contents, make the title of each such section unique by adding at the end of it, in parentheses, the name of the original author or publisher of that section if known, or else a unique number. Make the same adjustment to the section titles in the list of Invariant Sections in the license notice of the combined work.

In the combination, you must combine any sections Entitled "History" in the various original documents, forming one section Entitled "History"; likewise combine any sections Entitled "Acknowledgements", and any sections Entitled "Dedications". You must delete all sections Entitled "Endorsements".

6. COLLECTIONS OF DOCUMENTS

You may make a collection consisting of the Document and other documents released under this License, and replace the individual copies of this License in the various documents with a single copy that is included in the collection, provided that you follow the rules of this License for verbatim copying of each of the documents in all other respects.

You may extract a single document from such a collection, and distribute it individually under this License, provided you insert a copy of this License into the extracted document, and follow this License in all other respects regarding verbatim copying of that document.

7. AGGREGATION WITH INDEPENDENT WORKS

A compilation of the Document or its derivatives with other separate and independent documents or works, in or on a volume of a storage or distribution medium, is called an “aggregate” if the copyright resulting from the compilation is not used to limit the legal rights of the compilation’s users beyond what the individual works permit. When the Document is included in an aggregate, this License does not apply to the other works in the aggregate which are not themselves derivative works of the Document.

If the Cover Text requirement of section 3 is applicable to these copies of the Document, then if the Document is less than one half of the entire aggregate, the Document’s Cover Texts may be placed on covers that bracket the Document within the aggregate, or the electronic equivalent of covers if the Document is in electronic form. Otherwise they must appear on printed covers that bracket the whole aggregate.

8. TRANSLATION

Translation is considered a kind of modification, so you may distribute translations of the Document under the terms of section 4. Replacing Invariant Sections with translations requires special permission from their copyright holders, but you may include translations of some or all Invariant Sections in addition to the original versions of these Invariant Sections. You may include a translation of this License, and all the license notices in the Document, and any Warranty Disclaimers, provided that you also include the original English version of this License and the original versions of those notices and disclaimers. In case of a disagreement between the translation and the original version of this License or a notice or disclaimer, the original version will prevail.

If a section in the Document is Entitled “Acknowledgements”, “Dedications”, or “History”, the requirement (section 4) to Preserve its Title (section 1) will typically require changing the actual title.

9. TERMINATION

You may not copy, modify, sublicense, or distribute the Document except as expressly provided under this License. Any attempt otherwise to copy, modify, sublicense, or distribute it is void, and will automatically terminate your rights under this License.

However, if you cease all violation of this License, then your license from a particular copyright holder is reinstated (a) provisionally, unless and until the copyright holder explicitly and finally terminates your license, and (b) permanently, if the copyright holder fails to notify you of the violation by some reasonable means prior to 60 days after the cessation.

Moreover, your license from a particular copyright holder is reinstated permanently if the copyright holder notifies you of the violation by some reasonable means, this is the first time you have received notice of violation of this License (for any work) from that copyright holder, and you cure the violation prior to 30 days after your receipt of the notice.

Termination of your rights under this section does not terminate the licenses of parties who have received copies or rights from you under this License. If your rights have been terminated and not permanently reinstated, receipt of a copy of some or all of the same material does not give you any rights to use it.

10. FUTURE REVISIONS OF THIS LICENSE

The Free Software Foundation may publish new, revised versions of the GNU Free Documentation License from time to time. Such new versions will be similar in spirit to the present version, but may differ in detail to address new problems or concerns. See <https://www.gnu.org/copyleft/>.

Each version of the License is given a distinguishing version number. If the Document specifies that a particular numbered version of this License "or any later version" applies to it, you have the option of following the terms and conditions either of that specified version or of any later version that has been published (not as a draft) by the Free Software Foundation. If the Document does not specify a version number of this License, you may choose any version ever published (not as a draft) by the Free Software Foundation. If the Document specifies that a proxy can decide which future versions of this License can be used, that proxy's public statement of acceptance of a version permanently authorizes you to choose that version for the Document.

11. RELICENSING

"Massive Multiauthor Collaboration Site" (or "MMC Site") means any World Wide Web server that publishes copyrightable works and also provides prominent facilities for anybody to edit those works. A public wiki that anybody can edit is an example of such a server. A "Massive Multiauthor Collaboration" (or "MMC") contained in the site means any set of copyrightable works thus published on the MMC site.

"CC-BY-SA" means the Creative Commons Attribution-Share Alike 3.0 license published by Creative Commons Corporation, a not-for-profit corporation with a principal place of business in San Francisco, California, as well as future copyleft versions of that license published by that same organization.

"Incorporate" means to publish or republish a Document, in whole or in part, as part of another Document.

An MMC is "eligible for relicensing" if it is licensed under this License, and if all works that were first published under this License somewhere other than this MMC, and subsequently incorporated in whole or in part into the MMC, (1) had no cover texts or invariant sections, and (2) were thus incorporated prior to November 1, 2008.

The operator of an MMC Site may republish an MMC contained in the site under CC-BY-SA on the same site at any time before August 1, 2009, provided the MMC is eligible for relicensing.

ADDENDUM: How to use this License for your documents

To use this License in a document you have written, include a copy of the License in the document and put the following copyright and license notices just after the title page:

Copyright © YEAR YOUR NAME.

Permission is granted to copy, distribute and/or modify this document under the terms of the GNU Free Documentation License, Version 1.3 or any later version published by the Free Software Foundation;

with no Invariant Sections, no Front-Cover Texts, and no Back-Cover Texts.

A copy of the license is included in the section entitled "GNU Free Documentation License".

If you have Invariant Sections, Front-Cover Texts and Back-Cover Texts, replace the "with ... Texts." line with this:

with the Invariant Sections being LIST THEIR TITLES, with the Front-Cover Texts being LIST, and with the Back-Cover Texts being LIST.

If you have Invariant Sections without Cover Texts, or some other combination of the three, merge those two alternatives to suit the situation.

If your document contains nontrivial examples of program code, we recommend releasing these examples in parallel under your choice of free software license, such as the GNU General Public License, to permit their use in free software.