

Data Manipulation

Contents

Data Manipulation	1
Five basic functions in data handling	1
Grouping and Combining	4
Base subsetting	5

Data Manipulation

Report exemplifying the use of dplyr in data handling on the example of dsL.

Five basic functions in data handling

For a more detailed discussion of basic verbs and operations consult the [R-Studio guide](#) or internal vignette

```
# vignette("introduction",package="dplyr")
```

The following is a brief demonstration of dplyr syntax using dsL example

`select()`

selects variables into a smaller data set

```
require(dplyr)
ds<-dsL
dim(ds)
```

```
[1] 134760      60
```

```
ds<- select(ds,id,year, byear, attend, attendF)
head(ds,13)
```

	id	year	byear	attend	attendF
1	1	1997	1981	NA	<NA>
2	1	1998	1981	NA	<NA>
3	1	1999	1981	NA	<NA>
4	1	2000	1981	1	Never
5	1	2001	1981	6	About once/week
6	1	2002	1981	2	Once or Twice
7	1	2003	1981	1	Never
8	1	2004	1981	1	Never
9	1	2005	1981	1	Never
10	1	2006	1981	1	Never
11	1	2007	1981	1	Never
12	1	2008	1981	1	Never
13	1	2009	1981	1	Never

```
dim(ds)
```

```
[1] 134760      5
```

filter()

Removes observations that do not meet criteria. The following code selects observation based on the type of sample

	sample	sampleF
1	1	Cross-Sectional
2	0	Oversample

and only between years 2000 and 2011, as only during those years the outcome of interest attend was recorded.

```
require(dplyr)
ds<- filter(dsL,sample==1, year %in% c(2000:2011))
ds<- select(ds,id, year, attend, attendF)
head(ds,13)
```

	id	year	attend	attendF
1	1	2000	1	Never
2	1	2001	6	About once/week
3	1	2002	2	Once or Twice
4	1	2003	1	Never
5	1	2004	1	Never
6	1	2005	1	Never
7	1	2006	1	Never
8	1	2007	1	Never
9	1	2008	1	Never
10	1	2009	1	Never
11	1	2010	1	Never
12	1	2011	1	Never
13	2	2000	2	Once or Twice

arrange()

Sorts observations

```
require(dplyr)
ds<- filter(dsL,sample==1, year %in% c(2000:2011))
ds<- select(ds,id, year, attend)
ds<- arrange(ds, year, desc(id))
head(ds,13)
```

	id	year	attend
1	9022	2000	1
2	9021	2000	2
3	9020	2000	2
4	9018	2000	4

5	9017	2000	6
6	9012	2000	5
7	9011	2000	6
8	9010	2000	1
9	9009	2000	2
10	9008	2000	6
11	8992	2000	NA
12	8991	2000	3
13	8987	2000	6

```
ds<- arrange(ds, id, year)
head(ds, 13)
```

	id	year	attend
1	1	2000	1
2	1	2001	6
3	1	2002	2
4	1	2003	1
5	1	2004	1
6	1	2005	1
7	1	2006	1
8	1	2007	1
9	1	2008	1
10	1	2009	1
11	1	2010	1
12	1	2011	1
13	2	2000	2

mutate()

Creates additional variables from the values of existing.

```
require(dplyr)
ds<- filter(dsL,sample==1, year %in% c(2000:2011))
ds<- select(ds,id, byear, year, attend)
ds<- mutate(ds,
            age = year-byear,
            timec = year-2000,
            linear= timec,
            quadratic= linear^2,
            cubic= linear^3)
head(ds,13)
```

	id	byear	year	attend	age	timec	linear	quadratic	cubic
1	1	1981	2000	1	19	0	0	0	0
2	1	1981	2001	6	20	1	1	1	1
3	1	1981	2002	2	21	2	2	4	8
4	1	1981	2003	1	22	3	3	9	27
5	1	1981	2004	1	23	4	4	16	64
6	1	1981	2005	1	24	5	5	25	125
7	1	1981	2006	1	25	6	6	36	216
8	1	1981	2007	1	26	7	7	49	343

9	1	1981	2008	1	27	8	8	64	512
10	1	1981	2009	1	28	9	9	81	729
11	1	1981	2010	1	29	10	10	100	1000
12	1	1981	2011	1	30	11	11	121	1331
13	2	1982	2000	2	18	0	0	0	0

```
summarize()
```

Grouping and Combining

The function `group_by` is used to identify groups in split-apply-combine (SAC) procedure: all possible interactions between the levels of supplied variables

```
require(dplyr)

ds<- filter(dsL,sample==1, year %in% c(2000:2011))
ds<- select(ds,id, year, attendF)

s <- group_by(ds, year,attendF)
s <- summarise(s, count = n())
s <- mutate(s, total = sum(count),
            percent= count/total)
head(s,10)
```

Source: local data frame [10 x 5]

Groups: year

	year	attendF	count	total	percent
1	2000	Never	1581	6748	0.234292
2	2000	Once or Twice	1304	6748	0.193242
3	2000	Less than once/month	775	6748	0.114849
4	2000	About once/month	362	6748	0.053646
5	2000	About twice/month	393	6748	0.058239
6	2000	About once/week	1101	6748	0.163159
7	2000	Several times/week	463	6748	0.068613
8	2000	Everyday	36	6748	0.005335
9	2000	NA	733	6748	0.108625
10	2001	Never	1627	6748	0.241108

The same result can be achieved with the same result use a more elegant syntax that relies on `%>%` operator, in which `x %>% f(y)` turns into `f(x, y)` :

```
ds<-dsL %>%
  filter(sample==1, year %in% c(2000:2011)) %>%
  select(id, year, attendF) %>%
  group_by(year,attendF) %>%
    summarise(count = n()) %>%
    mutate(total = sum(count),
           percent= count/total)
head(ds,10)
```

Source: local data frame [10 x 5]

Groups: year

	year	attendF	count	total	percent
1	2000	Never	1581	6748	0.234292
2	2000	Once or Twice	1304	6748	0.193242
3	2000	Less than once/month	775	6748	0.114849
4	2000	About once/month	362	6748	0.053646
5	2000	About twice/month	393	6748	0.058239
6	2000	About once/week	1101	6748	0.163159
7	2000	Several times/week	463	6748	0.068613
8	2000	Everyday	36	6748	0.005335
9	2000	NA	733	6748	0.108625
10	2001	Never	1627	6748	0.241108

verify that this is what we wanted to achieve:

```
summarize( filter(s, year==2000), should.be.one=sum(percent))
```

Source: local data frame [1 x 2]

	year	should.be.one
1	2000	1

Base subsetting

Generally we can select any desired dataset by formula

dataset[*condition for rows* , *condition for columns*], and using **dataset\$variableName** selector

```
ds<-dsL[dsL$year %in% c(2000:2011),c('id',"byear","year","attendF","ageyearF","agemon")]
print(ds[ds$id==1,])
```

	id	byear	year	attendF	ageyearF	agemon
4	1	1981	2000	Never	19	231
5	1	1981	2001	About once/week	20	243
6	1	1981	2002	Once or Twice	21	256
7	1	1981	2003	Never	22	266
8	1	1981	2004	Never	23	279
9	1	1981	2005	Never	24	290
10	1	1981	2006	Never	25	302
11	1	1981	2007	Never	26	313
12	1	1981	2008	Never	27	325
13	1	1981	2009	Never	28	337
14	1	1981	2010	Never	29	350
15	1	1981	2011	Never	29	360