Experiment M12bis: Mechanical Rutherford Scattering

Andrés Vinuesa Espinosa and Jose María Martínez Herrada
Group B2.2

Date of Experiment: April 4, 2026

Abstract

This experiment explores the mechanical analogue of Rutherford scattering using a projectile sphere interacting with a central potential. We analyze two distinct cases: a repulsive "hill" potential and an attractive "well" potential. By tracking trajectories with the software Tracker, we determine the interaction constant $k = 1.23(5) \times 10^{-3} \mathrm{kg\,m^3/s^2}$ and the offset $r_0 \approx 0$ of the center of forces. The results demonstrate a clear transition from scattering trajectories in the repulsive case to a variety of open and closed orbits in the attractive case, illustrating the effects of energy conservation and dissipative friction in a central force field.

Contents

1 Introduction and Objectives

The primary objective of this experiment is to study the mechanical scattering of particles, specifically focusing on the behavior of a projectile particle interacting with a target particle in a Rutherford-like potential. In the first part, first we represent the experimental data, then we quantitatively analyze the scattering process by measuring the trajectories of the particles under various initial conditions, determining the interaction constant of the hill-like structure, from now on the hill and the true position of the center of forces. In the second part, we will discuss the behavior of the system when the projectile particle is launched into a potential well, analyzing the resulting trajectories and comparing them with theoretical predictions.

2 Results and Data Analysis

2.1 Analysis of the Hill Scattering

2.1.1 Experimental Data



Figure 1: Set of experimental trayectories with even decimal values.



Figure 2: Experimental trajectories with odd decimal values.

The center of the axis is placed in the effective center of forces, which is the point from which the trajectories appear to be deflected. We can observe that since the initial height and thus the velocity is increasided the deflection of the trajectories is reduced, having a less pronounced curve. Although the data obtained from the experiment using the software Tracker is not perfect (i.e., it contains a noticeable amount of noise), the introduction of uncertainties in the trajectories does not significantly affect the results of the fit. For this reason, we have not included them in the visual analysis, as they would make the trajectories harder to distinguish in the plot; however, if they were to be considered, they would be approximately 5 %. We obtained the followings angles of dispersion, being 180 degrees the retrodispersion case, i.e when the ball comes back in a straight line theta$_1$4 $= np.radians(90 + 18.4) theta_1 5 = np.radians(90 + 21.4) theta_1 6 = np.radians(90 + 16) theta_1 7 = np.radians(90 + 13.9) theta_1 8 = np.radians(90 + 12.9) theta_1 9 = np.radians(90 + 12.3)$

2.1.2 Theoretical Model and Fitting Process

The experimental data was fitted to the following theoretical model:

$$b = \cos\left(\frac{\theta}{2}\right)\left[\frac{k}{mv_0^2 \sin\left(\frac{\theta}{2}\right)} - r_0\right]. \tag{1}$$

Where k is the interaction constant of the hill, m is the mass of the sphere, v_0 is the initial velocity, θ is the scattering angle, and r_0 is the true position of the center of forces, which is not necessarily the geometrical center of the hill due to inherent asymmetries. We have taken b the distance from the initial to the center of forces, as 7.5 cm in all the cases.

θ was measured as the angle between the initial trajectory of the sphere and its trajectory after the interaction with the hill. r_0 was initially estimated based on the apparent point of retrodispersion of the trajectories and then refined through the fitting process.

2.1.3 Results Interpretation

We observe that the uncertainty of r_0 is several orders of magnitude larger than the value itself. This occurs because the fit was constrained to positive values of r_0. Since the fitted value is very close to zero, we can conclude that, within our error margin, the effective center coincides with the geometrical center.

The theoretical model for the hill's curvature is given by $z = \frac{k}{r} - r_0$. Using our values for k and r_0, the predicted height of the hill at $r = 1$ cm is approximately 11 cm. This is a reasonable value considering the laboratory setup. Minor discrepancies are expected due to rolling friction and surface irregularities (e.g., small bumps), but the result confirms that k is in the correct order of magnitude.

The uncertainties for the scattering angles were estimated such that $\chi^2_{\text{reduced}} \approx 1$. Since the theoretical model is well-established, this approach ensures a reasonable estimation of the uncertainties.

Parameter	Symbol	Value
Interaction Constant	k	1.1705×10^{-3} kg m^3/s^2
Center of Forces Offset	r_0	1.8271×10^{-9} m

Table 1: Parameters obtained from the fit of the mechanical scattering model.

2.2 Potential Well Analysis (Well Scattering)

In this section, we qualitatively analyze the six trajectories obtained by launching the sphere towards the attractive potential well. Unlike the case of the hill (repulsive), here the central force tends to attract the sphere towards the origin $(0, 0)$. The recorded experimental trajectories are shown in Figure **??**.

From the observation of Figure **??**, we can individually classify the six trajectories based on the sphere's initial mechanical energy (E), ordering them from the most energetic to the least:

- **Brown Trajectory (Open):** It presents the highest initial kinetic energy ($E > 0$). The sphere enters, undergoes a deflection as it passes through the attractive well, and quickly escapes to infinity. It is the equivalent of a celestial body with a hyperbolic trajectory that visits a star only once and does not return, such as a comet.

- **Purple Trajectory (Open):** This one has an initial kinetic energy that is still large enough not to be trapped in the well and to deviate, although already making a slight curve in its turn due to the attraction of the well in question, which fails to capture it.

- **Red Trajectory (Open):** It has less energy than the purple one, bordering the critical capture limit ($E \approx 0$). It manages to complete almost a full revolution around the center of forces due to the strong attraction, but it retains just enough energy to overcome the well and end up escaping.

- **Green Trajectory (Closed):** The sphere enters with a low enough energy ($E < 0$ relative to infinity) to be trapped. It forms an elliptical orbit that makes multiple revolutions. Precession, which is the rotation of the ellipse's axis, is clearly observed, caused by the gradual loss of energy due to rolling friction in the system used in the laboratory.



Figure 3: Experimental trajectories around the attractive potential well (center of forces at $(0, 0)$). Closed orbits (blue, orange, and green) and open orbits (brown, purple, and red) are observed depending on their initial energy.

- **Orange Trajectory (Closed):** It is another bounded orbit where the sphere is strongly captured. It shows a very evident orbital decay as it loses speed and the well irremediably attracts it.
- **Blue Trajectory (Closed):** It represents one of the lowest initial energy cases. The sphere barely manages to open its orbit after the first curve and quickly collapses towards the center of the potential well after completing its spiral path.

It is worth mentioning that the transition between closed and open orbits is rather gradual, we tried do devise a completely circular one. The central forces theory tell us that at some height between h=2.05 cm and h=2cm the sphere attains an circular orbit with E=0, but in practice, due to the presence of friction and other non-idealities, this is almost impossible to achieve.

3 Discussion

The fitting of the experimental data for the hill scattering yielded an interaction constant $k = 1.1705 \times 10^{-3} \mathrm{kg\,m^3/s^2}$. This value, when used to reconstruct the hill's geometry, results in a central height of approximately 11 cm, which is consistent with the physical dimensions of the laboratory setup. The small value of r_0 indicates that the effective center of forces is well-aligned with the geometric center of the potential.

Regarding the attractive potential well, the transition from open to closed orbits was qualitatively verified. The open trajectories (brown, purple, and red) illustrate scattering at high initial energies, while the closed orbits (green, orange, and blue) represent captured states. The observed precession and orbital decay in the green trajectory are direct evidence of energy dissipation due to rolling friction, which is the primary non-ideality in this mechanical system.

4 Conclusions

In this experiment, we successfully characterized the mechanical scattering of particles in both repulsive and attractive central potentials. The quantitative analysis of the hill scattering confirmed the validity of the Rutherford-like model for this system, while the study of the potential well provided a clear demonstration of the relationship between mechanical energy and orbit topology.

Despite the noise in the experimental tracking and the dissipative effects of friction, the results are in good agreement with the theoretical framework of central forces. Future improvements could include using a high-speed camera to increase temporal resolution and a more rigorous treatment of the friction coefficients to better model the orbital decay observed in the attractive potential. Also, varying the distance from the center of forces in the hill part, could contribute to getting a better fit, since it may account for irregularities on the hill.