Dietary Reference Intakes (DRIs): Estimated Average Requirements

Food and Nutrition Board, Institute of Medicine, National Academies

Life Stage	Calcium	СНО	Protein	Vit A $(\mu g/d)^a$	Vit C (mg/d)	Vit D	Vit E (mg/d) ^b	Thiamin (mg/d)	Ribo- flavin (mg/d)	Niacin (mg/d) ^c	Vit B ₆ (mg/d)	Folate $(\mu g/d)^d$	Vit B ₁₂	Copper	Iodine	Iron (mg/d)	Magnes- ium (mg/d)	Molyb- denum (µg/d)	Phos- phorus (mg/d)	Sele- nium	Zinc (mg/d)
Group	(mg/d)	(g/d)	(g/kg/d)	(µg/u)	(IIIg/u)	(µg/d)	(IIIg/u)	(IIIg/u)	(IIIg/u)	(IIIg/u)	(IIIg/u)	(µg/u)	(µg/d)	(µg/d)	(µg/d)	(IIIg/u)	(IIIg/u)	(µg/u)	(IIIg/u)	(µg/d)	(IIIg/u)
Infants																					
0 to 6 mo																					
6 to 12																					
mo			1.0													6.9					2.5
Children																					
1–3 y	500	100	0.87	210	13	10	5	0.4	0.4	5	0.4	120	0.7	260	65	3.0	65	13	380	17	2.5
4–8 y	800	100	0.76	275	22	10	6	0.5	0.5	6	0.5	160	1.0	340	65	4.1	110	17	405	23	4.0
Males																					
9–13 y	1,100	100	0.76	445	39	10	9	0.7	0.8	9	0.8	250	1.5	540	73	5.9	200	26	1,055	35	7.0
14–18 y	1,100	100	0.73	630	63	10	12	1.0	1.1	12	1.1	330	2.0	685	95	7.7	340	33	1,055	45	8.5
19–30 y	800	100	0.66	625	75	10	12	1.0	1.1	12	1.1	320	2.0	700	95	6	330	34	580	45	9.4
31–50 y	800	100	0.66	625	75	10	12	1.0	1.1	12	1.1	320	2.0	700	95	6	350	34	580	45	9.4
51–70 y	800	100	0.66	625	75	10	12	1.0	1.1	12	1.4	320	2.0	700	95	6	350	34	580	45	9.4
> 70 y	1,000	100	0.66	625	75	10	12	1.0	1.1	12	1.4	320	2.0	700	95	6	350	34	580	45	9.4
Females																					
9–13 y	1,100	100	0.76	420	39	10	9	0.7	0.8	9	0.8	250	1.5	540	73	5.7	200	26	1,055	35	7.0
14–18 y	1,100	100	0.71	485	56	10	12	0.9	0.9	11	1.0	330	2.0	685	95	7.9	300	33	1,055	45	7.3
19–30 y	800	100	0.66	500	60	10	12	0.9	0.9	11	1.1	320	2.0	700	95	8.1	255	34	580	45	6.8
31–50 y	800	100	0.66	500	60	10	12	0.9	0.9	11	1.1	320	2.0	700	95	8.1	265	34	580	45	6.8
51–70 y	1,000	100	0.66	500	60	10	12	0.9	0.9	11	1.3	320	2.0	700	95	5	265	34	580	45	6.8
> 70 y	1,000	100	0.66	500	60	10	12	0.9	0.9	11	1.3	320	2.0	700	95	5	265	34	580	45	6.8
Pregnancy																					
14–18 y	1,000	135	0.88	530	66	10	12	1.2	1.2	14	1.6	520	2.2	785	160	23	335	40	1,055	49	10.5
19–30 y	800	135	0.88	550	70	10	12	1.2	1.2	14	1.6	520	2.2	800	160	22	290	40	580	49	9.5
31–50 y	800	135	0.88	550	70	10	12	1.2	1.2	14	1.6	520	2.2	800	160	22	300	40	580	49	9.5
Lactation																					
14–18 y	1,000	160	1.05	885	96	10	16	1.2	1.3	13	1.7	450	2.4	985	209	7	300	35	1,055	59	10.9
19–30 y	800	160	1.05	900	100	10	16	1.2	1.3	13	1.7	450	2.4	1,000	209	6.5	255	36	580	59	10.4
31–50 y	800	160	1.05	900	100	10	16	1.2	1.3	13	1.7	450	2.4	1,000	209	6.5	265	36	580	59	10.4

NOTE: An Estimated Average Requirement (EAR) is the average daily nutrient intake level estimated to meet the requirements of half of the healthy individuals in a group. EARs have not been established for vitamin K, pantothenic acid, biotin, choline, chromium, fluoride, manganese, or other nutrients not yet evaluated via the DRI process.

SOURCES: Dietary Reference Intakes for Calcium, Phosphorous, Magnesium, Vitamin D, and Fluoride (1997); Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B₆, Folate, Vitamin B₁₂, Pantothenic Acid, Biotin, and Choline (1998); Dietary Reference Intakes for Vitamin C, Vitamin E, Selenium, and Carotenoids (2000); Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc (2001); Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids (2002/2005); and Dietary Reference Intakes for Calcium and Vitamin D (2011). These reports may be accessed via www.nap.edu.

^a As retinol activity equivalents (RAEs). 1 RAE = 1 μ g retinol, 12 μ g β-carotene, 24 μ g α-carotene, or 24 μ g β-cryptoxanthin. The RAE for dietary provitamin A carotenoids is two-fold greater than retinol equivalents (RE), whereas the RAE for preformed vitamin A is the same as RE.

 $[^]b$ As α-tocopherol includes RRR-α-tocopherol, the only form of α-tocopherol that occurs naturally in foods, and the 2R-stereoisomeric forms of α-tocopherol (RRR-, RSR-, RRS-, and RSS-α-tocopherol) that occur in fortified foods and supplements. It does not include the 2S-stereoisomeric forms of α-tocopherol (SRR-, SRS-, and SSS-α-tocopherol), also found in fortified foods and supplements.

^cAs niacin equivalents (NE). 1 mg of niacin = 60 mg of tryptophan.

^dAs dietary folate equivalents (DFE). 1 DFE = 1 μg food folate = 0.6 μg of folic acid from fortified food or as a supplement consumed with food = 0.5 μg of a supplement taken on an empty stomach.