

Research Project

Mathematical Modelling of Cell Membrane Dynamics

Isak Hammer



Department of Mathematical Sciences
Norwegian University of Science and Technology

First can we detect deceases such as Alzheimer's disease, cancer cells and develop new methods and vaccines [1]. One of the primary components of the cell membranes are lipids which serve many different functions. A key function is that it is consisting of a bilayer of lipids which controls the structural rigidity and the fluidity of the membrane [2]. It also turns out that the lipids often accumulate into so-called lipid rafts which serves as a rigid platform for proteins with special properties such as intracellular trafficking of lipids and lipid-anchored proteins [3].

Modelling of lipid rafts formation can be modelled as a two-phase separation problem based on minimization of the Ginzburg-Landau energy functional [4]

$$\mathcal{E}_{ch}(\Gamma) = \int_{\Gamma} \Psi(c) + \frac{\gamma}{2} |\nabla c|^2 dx,$$

which is describing the chemical energy for a concentration $c : \Gamma \times [0, T] \mapsto [0, 1]$ over a surface membrane Γ . Several authors have solved this problem often results by deriving variants of Cahn Hilliard Equation or Allen Cahn Equation if the concentration is not conserved both standstill and evolving domains [4–8].

Assuming that the system is a single-phase system can the elastic bending energy be modelled using the Canham Helrich energy functional [5, 9]

$$\mathcal{E}_e(\Gamma) = \int_{\Gamma} c_b H^2 + c_k K dx$$

Here is $H = \frac{\kappa_1 + \kappa_2}{2}$ denoted as the mean curvature and $K = \kappa_1 \kappa_2$ as the gaussian curvature with respectively c_b and c_k as tuning parameters and κ_1 and κ_2 as principal curvatures. Using the Gauss-Bonnet theorem can it be shown that the problem above is equivalent to the so-called Willmore energy functional [10, Ch 8.5 Thm 8.29],

$$\mathcal{E}(\Gamma) = \int_{\Gamma} H^2 dx$$

This is a well known problem in the mathematical community [11, 12]. In fact, it is a mathematical tool used to study the geometry of surfaces because it can be used to study the properties of minimal surfaces, which are surfaces with the least possible area for a given boundary. This is important in many areas of mathematics, including differential geometry, topology and mathematical physics [13–15].

In this report will we establish a numerical scheme on minimization on this functional. However, we will first establish notation by including a section for definitions and important results from differential geometry. We will then derive the underlying PDE's for this equation.

Lastly we will establish the model for the problem and discretize the problem using evolutionary parametric FEM methods.

1 Differential Geometry

We may define Γ to be a smooth compact and oriented surface with no boundary in \mathbb{R}^3 . We will denote the normal unit vector outer normal vector to be $\nu \in \mathbb{R}^3$.

2 Cahn Hilliard Equation on a Closed Membrane

Let c_0 and c_1 indicate the concentration profile of the substances in a 2 -phase system such that $c_0(\mathbf{x}, t) : \Omega \times [0, \infty] \rightarrow [0, 1]$ and similarly $c_1(\mathbf{x}, t) : \Omega \times [0, \infty] \rightarrow [0, 1]$, where \mathbf{x} is a element of some surface Ω and t is time. However, in the 2 phase problem will we will restrict ourself so that $c_0(t, \mathbf{x}) + c_1(t, \mathbf{x}) = 1$ at any \mathbf{x} at time t . A property of the restriction is that we now can express c_0 using c_1 , with no loss of information. Hence, let us now define $c = c_0$ so $c(\mathbf{x}, t) : \Omega \times [0, \infty] \rightarrow [0, 1]$. It has been shown that 2 phase system if thermodynamically unstable can be evolve into a phase separation described by a evolutional differential equation [16] using a model based on chemical energy of the substances. However, further development has been done [4] to solve this equation on surfaces. Now assume model that we want to describe is a phase-separation on a closed membrane surface Γ , so that $c(\mathbf{x}, t) : \Gamma \times [0, T] \rightarrow [0, 1]$. Then is the surface Cahn Hilliard equation described such that

$$\rho \frac{\partial c}{\partial t} - \nabla_{\Gamma} (M \nabla_{\Gamma} (f'_0 - \varepsilon^2 \nabla_{\Gamma}^2 c)) = 0 \quad \text{on } \Gamma. \quad (1)$$

We define here the tangential gradient operator to be $\nabla_{\Gamma} c = \nabla c - (\mathbf{n} \cdot \nabla c) \mathbf{n}$ applied on the surface Γ restricted to $\mathbf{n} \cdot \nabla_{\Gamma} c = 0$.

Lets define ε to be the size of the layer between the substances c_1 and c_2 . The density ρ is simply defined such that $\rho = \frac{m}{S_{\Gamma}}$ is a constant based on the total mass divided by the total surface area of Γ . Here is the mobility M often derived such that is is dependent on c and is crucial for the result during a possible coarsening event [4]. However, the free energy per unit surface $f_0 = f_0(c)$ is derived based on the thermodynamical model and should according to [4] be non convex and nonlinear.

A important observation is that equation (1) is a fourth order equation which makes it more challenging to solve using conventional FEM methods. This clear when writing the equation on the equivalent weak form and second order equations arise.

3 Energy Functionals

Let $c(x, t) : \Gamma \times [0, T] \mapsto [0, 1]$. From [4] can we observe the energy functionals

$$E_1(c) = \int_{\Gamma} f(c).$$

where

$$f(c) = f_0(c) + \frac{1}{2}\varepsilon^2 |\nabla_{\Gamma} c|^2$$

and the conservation law $\rho \frac{\partial c}{\partial t} + \text{div}_{\Gamma} \mathbf{j} = 0$ for the evolution of c , derived from the Ficks Law $\mathbf{j} = -M \nabla_{\Gamma} \mu$ for the chemical potential derived by the functional derivative $\mu = \frac{\delta f}{\delta c}$. The double well function is denoted as

$$f_0(c) = \frac{\zeta}{4} c^2 (1 - c)^2$$

References

- [1] Scott A Small and Sam Gandy. “Sorting through the cell biology of Alzheimer’s disease: intracellular pathways to pathogenesis”. In: *Neuron* 52.1 (2006), pp. 15–31.
- [2] Saul L. Neidleman. “Effects of Temperature on Lipid Unsaturation”. In: *Biotechnology and Genetic Engineering Reviews* 5.1 (1987). PMID: 3314900, pp. 245–268. DOI: [10 . 1080 / 02648725 . 1987 . 10647839](https://doi.org/10.1080/02648725.1987.10647839). eprint: <https://doi.org/10.1080/02648725.1987.10647839>.
- [3] Michael Edidin. “The State of Lipid Rafts: From Model Membranes to Cells”. In: *Annual Review of Biophysics and Biomolecular Structure* 32.1 (2003). PMID: 12543707, pp. 257–283. DOI: [10 . 1146 / annurev . biophys . 32 . 110601 . 142439](https://doi.org/10.1146/annurev.biophys.32.110601.142439). eprint: <https://doi.org/10.1146/annurev.biophys.32.110601.142439>.
- [4] Vladimir Yushutin et al. “A computational study of lateral phase separation in biological membranes”. In: *International Journal for Numerical Methods in Biomedical Engineering* 35.3 (2019). e3181 cnm.3181, e3181. DOI: <https://doi.org/10.1002/cnm.3181>. eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/cnm.3181>. URL: <https://onlinelibrary.wiley.com/doi/abs/10.1002/cnm.3181>.
- [5] Udo Seifert. “Configurations of fluid membranes and vesicles”. In: *Advances in Physics* 46.1 (1997), pp. 13–137. DOI: [10 . 1080 / 00018739700101488](https://doi.org/10.1080/00018739700101488). eprint: <https://doi.org/10.1080/00018739700101488>. URL: <https://doi.org/10.1080/00018739700101488>.
- [6] Andreas Rätz. “A benchmark for the surface Cahn–Hilliard equation”. In: *Applied Mathematics Letters* 56 (2016), pp. 65–71. ISSN: 0893-9659. DOI: <https://doi.org/10.1016/j.aml.2015.12.008>. URL: <https://www.sciencedirect.com/science/article/pii/S0893965915300045>.
- [7] P. Gera and David Salac. “Cahn-Hilliard on surfaces: A numerical study”. In: *Appl. Math. Lett.* 73 (2017), pp. 56–61.
- [8] D. CAETANO and C. M. ELLIOTT. “Cahn–Hilliard equations on an evolving surface”. In: *European Journal of Applied Mathematics* 32.5 (2021), 937–1000. DOI: [10.1017/S0956792521000176](https://doi.org/10.1017/S0956792521000176).

- [9] Xiaoqiang Wang and Qiang Du. “Modelling and simulations of multi-component lipid membranes and open membranes via diffuse interface approaches”. In: *Journal of mathematical biology* 56 (Apr. 2008), pp. 347–71. DOI: [10.1007/s00285-007-0118-2](https://doi.org/10.1007/s00285-007-0118-2).
- [10] S. Montiel, A. Ros, and D.G. Babbitt. *Curves and Surfaces*. Graduate studies in mathematics. American Mathematical Society, 2009. ISBN: 9780821847633. URL: <https://books.google.no/books?id=dHDzpBDHPigC>.
- [11] Peter Topping. “Towards the Willmore conjecture”. In: *Calculus of Variations and Partial Differential Equations* 11.4 (2000), pp. 361–393.
- [12] Fernando C Marques and Andre Neves. “The willmore conjecture”. In: *Jahresbericht der Deutschen Mathematiker-Vereinigung* 116.4 (2014), pp. 201–222.
- [13] Thomas Koerber. “The area preserving Willmore flow and local maximizers of the Hawking mass in asymptotically Schwarzschild manifolds”. In: *The Journal of Geometric Analysis* 31.4 (2021), pp. 3455–3497.
- [14] Ruben Jakob. “Singularities and full convergence of the Mobius-invariant Willmore flow in the 3-sphere”. In: *arXiv preprint arXiv:2205.00604* (2022).
- [15] Fabian Rupp. “The volume-preserving Willmore flow”. In: (2020). DOI: [10.48550/ARXIV.2012.03553](https://doi.org/10.48550/ARXIV.2012.03553). URL: <https://arxiv.org/abs/2012.03553>.
- [16] John W. Cahn and John E. Hilliard. “Free Energy of a Nonuniform System. I. Interfacial Free Energy”. In: *The Journal of Chemical Physics* 28.2 (1958), pp. 258–267. DOI: [10.1063/1.1744102](https://doi.org/10.1063/1.1744102). eprint: <https://doi.org/10.1063/1.1744102>. URL: <https://doi.org/10.1063/1.1744102>.