

Project Thesis

Solving Cahn Hilliard Equation

Isak Hammer

May 31, 2022



1 Introduction

PLAN FOR REPORT

- 1) Introduction
- 2) Biharmonic Equation
 - Introduction of Biharmonic Equation
 - Weak form in $H^4(T)$
 - Weak form in $H^4(\Omega)$
- 3) Continuous Interior Penalty Method
 - CIP for Biharmonic equation
 - > Discussion of why we go for CIP method
 - > Mixed formulation
 - > B-spline basis
 - Error Estimates
 - A-priori Estimates
- 4) Numerical Experiments
 - * Manufactured solution
 - * Convergence rate
 - * Condition number (h^{-4})
- 5) Possible Extensions:
 - Compare with mixed formulation

2 Biharmonic Equation

2.1 Strong form of the Biharmonic Equation

Let $\Omega \subset \mathbb{R}^2$ be a bounded polygonal domain and $\partial\Omega$ be its corresponding boundary. Let the inhomogeneous fourth order biharmonic equation have the form,

$$\begin{aligned}\Delta^2 u + \alpha u &= f \quad \text{in } \Omega, \\ \partial_n u &= 0 \quad \text{on } \partial\Omega, \\ \partial_n \Delta u &= g(x) \quad \text{on } \partial\Omega.\end{aligned}\tag{1}$$

Here is $\Delta^2 = \Delta(\Delta)$ the biharmonic operator, also known as the bilaplacian. We will assume for the strong form that $u \in H^4(\Omega)$, α is a non-negative constant and $f \in L_2(\Omega)$. We may consider the functions $g(x)$ as time independent boundary conditions. Such problems as (1) are often associated with the Cahn-Hilliard model [1] for phase separation. However, depending on how Cahn-Hilliard model is time discretized numerically can (1) naturally arise. I refer to [2] for more information on this.

2.2 Weak Form Biharmonic Equation in $H^4(\Omega)$

We want to introduce the full weak formulation of (1). Now, let the solution space be on the form,

$$V = \{v \in H^2(\Omega) : \partial_n v = 0 \text{ on } \partial\Omega\}.$$

Let $u, v \in V$ then the derivation of the general weak form is,

$$\begin{aligned}(\Delta^2 u, v)_\Omega &= (\partial_n \Delta u, v)_{\partial\Omega} - (\nabla(\Delta u), \nabla v)_\Omega \\ &= (g, v)_{\partial\Omega} - (\nabla(\Delta u), \nabla v)_\Omega.\end{aligned}$$

In fact, the simplest formulation has the form,

$$(\nabla(\Delta u), \nabla v)_\Omega = (\Delta u, \partial_n v)_{\partial\Omega} - (\Delta u, \Delta v)_\Omega,$$

but we don't have boundary condition for Δu . Instead, we can formulate the problem so

why no b.c. for Δu ?

$$\begin{aligned}(\nabla(\Delta u), \nabla v)_\Omega &= \sum_{i=1}^d (\Delta \partial_{x_i} u, \partial_{x_i} v)_\Omega \\ &= \sum_{i=1}^d (\nabla \cdot (\nabla \partial_{x_i} u), \partial_{x_i} v)_\Omega \\ &= \sum_{i=1}^d (\partial_n \partial_{x_i} u, \nabla \partial_{x_i} v)_{\partial\Omega} - (\nabla \partial_{x_i} u, \nabla \partial_{x_i} v)_\Omega \\ &= (\partial_n \nabla u, \nabla v)_{\partial\Omega} - (D^2 u, D^2 v)_\Omega \\ &= \underbrace{(\partial_{nn} u, \partial_n v)_{\partial\Omega}}_{\partial_n v = 0 \text{ on } \partial\Omega} + (\partial_{nt} u, \partial_t v)_{\partial\Omega} - (D^2 u, D^2 v)_\Omega.\end{aligned}$$

Hence, the boundary condition of Δu is integrated into the formulation. It can be denoted that D^2 is the Hessian matrix operator such that

$$(D^2 u, D^2 v)_\Omega = \int_\Omega D^2 u : D^2 v dx,$$

where $D^2 u : D^2 v$ is the inner product and similarly for $\partial_{nn} u = n \cdot D^2 u \cdot n$.

We can now formulate the full weak formulation by solving for a $u \in V$ such that

$$a(u, v) = F(v), \quad \forall v \in V,\tag{2}$$

where

$$\begin{aligned}a(u, v)_\Omega &= (D^2 u, D^2 v)_\Omega - (\partial_{nt} u, \partial_t v)_{\partial\Omega} + \alpha(u, v)_\Omega, \\ F(v)_\Omega &= (f, v)_\Omega - (g, v)_{\partial\Omega}.\end{aligned}\tag{3}$$

In fact, the solution is unique for $\alpha > 0$. However, for $\alpha = 0$ must we assume the solvability condition,

$$\int_{\Omega} f dx = \int_{\partial\Omega} g ds.$$

This condition easily arise when using the substitution $v = 1$ in (2). To handle this, can we extended the solution space

$$V^* = \begin{cases} V & \alpha > 0 \\ \{v \in V : \int_{\Omega} v dx = 0\} & \alpha = 0, \end{cases}$$

Thus, the unique solution in $v \in V^*$ belongs to $H^3(\Omega)$ and we get the following elliptic regularity estimate [3],

$$|u|_{H^3(\Omega)} \leq C_{\Omega} \left(\|f\|_{L_2(\Omega)} + (1 + \alpha^{\frac{1}{2}}) \cdot \|w\|_{H^4(\Omega)} \right), \quad w \in H^4(\Omega). \quad (4)$$

why does it
belong to
 $H^3(\Omega)$

This regularity estimate may be important for further use cases in terms of error analysis.

3 Continuous Interior Penalty Method

3.1 Introduction

To solve (1) numerically do we want to introduce the Continuous Interior Penalty Method (CP), which is a Discontinuous Galerkin method (DG) using C^0 finite elements. There is several reasons why we want to apply C^0 instead of the often used C^1 finite elements for fourth order problems. First and foremost is the C^0 finite elements simpler than obtaining C^1 finite elements. Also, compared to other methods similar to the mixed finite element method for the problem (1), CP has in fact preserved the symmetric positive definiteness, which means the stability analysis is more straight forward. Finally and most importantly according to [2] can naive use mixed methods of splitting the boundary conditions of the problem (1) produce wrong solutions if Ω is non convex.

Write about this:

Conformal methods $V_h \subset V$ requires C^1 . Exists in a good manner in 2D, but does not exist generalization in 3D. Need reference.

Use Bspline as alternative basis. . Less flexible when generating meshes for complicated domains. Need reference.

Write in mixed formulation $\bar{w} = \Delta w$

None-conform discretization of 4th order problem using C^0 Elements. Hence CP Method

3.2 Computational Domains

Let $\mathcal{T} = \{T\}$ be a triangulation of $\Omega \subset \mathbb{R}^2$ consisting of triangles T as in figure 2. We may also define the set of all facets \mathcal{F}_h , where every facet is denoted by $F \in \mathcal{F}_h$. However, we will distinguish between the set of external facets \mathcal{F}_h^{ext} , which is all facets along $\partial\Omega$, and the interior facets \mathcal{F}_h^{int} . Let the facets be denotes as $F \in \mathcal{F}_h$, then the normal vector n is across the facets from T^+ to T^- , illustrated in figure 1.




Figure 1: Facet $F \in \mathcal{F}_h$ shared by the triangles $T^+, T^- \in \mathcal{T}_h$ and the normal unit vector n .

A parameter which is useful is the maximum diameter h of the set of triangles $\{T\}$, which we to be defined s.t.

$$\begin{aligned} h_T &= \text{diam}(T) = \max_{x_1, x_2 \in T} \text{dist}(x_1, x_2), \\ h_{min} &= \min_{T \in \mathcal{T}} h_T, \\ h_{max} &= \max_{T \in \mathcal{T}} h_T := h, \end{aligned} \tag{5}$$

where the $\text{diam}(T)$ is the largest facet for a triangle T . We will also assume mesh conform i.e., if $T_1 \neq T_2$ and $T_1 \cap T_2 \neq \emptyset$, then they share either a vertex or a facet.

Let the chunkiness parameter $c_T := h_T/r_T$, where r_T is the largest ball that be inscribed inside the element T . We can then assume that the mesh is shape-regular i.e., that $c_T \leq c$ independent of T and h . We may also assuming that the mesh is quasi-uniform only if it holds that the mesh is shape regular and $h_{max} \leq ch_{min}$.

Why is quasi-uniform important and how does shape regularity affect the numerical solution?




Figure 2: Example of a mesh of $\Omega \subset \mathbb{R}^2$ with triangulation \mathcal{T}_h .

3.3 Constructing Continuous Interior Penalty Method

Let us assume that $u, v \in H^4(T)$. Using that the results in (3) also holds for a triangle T can we write

$$(\Delta^2 u, v)_T = (D^2 u, D^2 v)_T - (\partial_{nt} u, \partial_t v)_{\partial T} - (\partial_{nn} u, \partial_n v)_{\partial T} + (\partial_n \Delta u, v)_{\partial T}. \quad (6)$$

For global continuity, let $v \in V = \{v \in H^1(\Omega) : v_T \in H^4(T), \forall T \in \mathcal{T}_h\} \cap C^0(\Omega)$ and $u \in H^4(\Omega)$ such that,

Should we require $v \in V = \{v \in H^1(\Omega) : v_T \in H^4(T) \forall T \in \mathcal{T}_h \text{ and } \partial_n v = 0 \text{ on } \partial\Omega\} \cap C^0(\Omega)$??

$$(\Delta^2 u, v)_\Omega = \sum_{T \in \mathcal{T}_h} (D^2 u, D^2 v)_T - (\partial_{nt} u, \partial_t v)_{\partial T} - (\partial_{nn} u, \partial_n v)_{\partial T} + (\partial_n \Delta u, v)_{\partial T}. \quad (7)$$

However, this expression can be written to distinguish integrating over triangles \mathcal{T}_h , integrating over exterior facets \mathcal{F}_h^{ext} and then integrate interior facets \mathcal{F}_h^{int} .

$$\begin{aligned} (\Delta^2 u, v)_\Omega &= \sum_{T \in \mathcal{T}_h} (D^2 u, D^2 v)_T + \sum_{F \in \mathcal{F}_h^{ext}} (\partial_n \Delta u, v)_F - (\partial_{nt} u, \partial_t v)_F + (\partial_{nn} u, \partial_n v)_F + \sum_{F \in \mathcal{F}_h^{int}} (\partial_{nn} u, [\partial_n v])_F \\ &= \sum_{T \in \mathcal{T}_h} (D^2 u, D^2 v)_T + \sum_{F \in \mathcal{F}_h^{ext}} (g, v)_F - (\partial_{nt} u, \partial_t v)_F + \sum_{F \in \mathcal{F}_h^{int}} (\partial_{nn} u, [\partial_n v])_F \end{aligned} \quad (8)$$

Keep in mind that any jump over a interior facet $F \subset \mathcal{F}_h^{int}$, visualized in figure 1, is defined as $\llbracket a \rrbracket = a^+ - a^-$ and likewise for the mean, $\{a\} = \frac{1}{2}(a^+ + a^-)$. The equivalence of (7) and (8) comes from the following argumentation.

$$\begin{aligned} (\Delta^2 u, v)_\Omega &= \sum_{T \in \mathcal{T}_h} (D^2 u, D^2 v)_T - (\partial_{nt} u, \partial_t v)_{\partial T} - (\partial_{nn} u, \partial_n v)_{\partial T} + (\partial_n \Delta u, v)_{\partial T} \\ &= \sum_{T \in \mathcal{T}_h} (D^2 u, D^2 v)_T \\ &\quad + \sum_{F \in \mathcal{F}_h^{ext}} \underbrace{(\partial_n \Delta u, v)_F}_{=(g, v)_F} - \underbrace{(\partial_{nt} u, \partial_t v)_F}_{=0} - \underbrace{(\partial_{nn} u, \partial_n v)_F}_{=0} \\ &\quad + \sum_{F \in \mathcal{F}_h^{int}} \underbrace{((\partial_{n^+} \Delta u^+, v^+)_F + (\partial_{n^-} \Delta u^-, v^-)_F)}_{(I)} + \underbrace{((\partial_{n^+} u^+, \partial_t v^+)_F + (\partial_{n^-} u^-, \partial_t v^-)_F)}_{(II)} + \underbrace{((\partial_{n^+} u^+, v^+)_F + (\partial_{n^-} u^-, v^-)_F)}_{(III)} \end{aligned}$$

Where integration over all interior facets $\forall F \in \mathcal{F}_h^{int}$ is computed in this way,

$$\begin{aligned}
(I) &= (\partial_{n^+} \Delta u^+, v^+)_F + (\partial_{n^-} \Delta u^+, v^-)_F = \int_F \llbracket \partial_n \Delta u \cdot v \rrbracket = \int_F \underbrace{\{\partial_n \Delta u\}}_{=0} \underbrace{\llbracket v \rrbracket}_{=0} + \underbrace{\llbracket \partial_n \Delta u \rrbracket}_{=0} \{v\} = 0 \\
(II) &= (\partial_{n^+} u^+, \partial_t v^+)_F + (\partial_{n^-} u^-, \partial_t v^-)_F = \int_F \llbracket \partial_{nt} u \cdot \partial_t v \rrbracket = \int_F \underbrace{\{\partial_{nt} u\}}_{=0} \underbrace{\llbracket \partial_t v \rrbracket}_{=0} + \underbrace{\llbracket \partial_{nt} u \rrbracket}_{=0} \{\partial_t v\} = 0 \\
(III) &= (\partial_{n^+} u^+, \partial_{nn} v^+)_F + (\partial_{n^-} u^-, \partial_{nn} v^-)_F = \int_F \llbracket \partial_{nn} u \cdot \partial_{nn} v \rrbracket = \int_F \underbrace{\{\partial_{nn} u\}}_{\neq 0} \underbrace{\llbracket \partial_{nn} v \rrbracket}_{=0} + \underbrace{\llbracket \partial_{nn} u \rrbracket}_{=0} \{\partial_{nn} v\} = (\partial_{nn} u, \llbracket \partial_{nn} v \rrbracket)_F
\end{aligned}$$

Observe that the cancellations in the term (I) appears of the continuity of $v \in V$ and $u \in H^4(\Omega)$ which makes the jumps zero. For the second term (II) does the terms become zero cancelled because the tangential derivative at the facet has no jump. However, The third term (III) is fairly interesting since the discontinuity in normal vector for $v \in V$ is a jump, while the second term is still continuous. It can also be raised that $\{\partial_{nn} u\} = \partial_{nn} u$ holds by the continuity of $H^4(\Omega)$. Anyhow, the definition of jump of should more interesting when we later weaken the continuity of u during discretization. Hence, (7) and (8) is equivalent.

3.4 Formulation of Continious Interior Penalty Method

We can finally start defining the fully discrete formulation. Let the basis be a \mathcal{P}_2 Lagrange finite element space so,

$$V_h = \{v \in C^0(\Omega) : v_T = v|_T \in P_2(T), \forall T \in \mathcal{T}_h\}$$

Again as above. Should we require $v \in V = \{v \in C^0(\Omega) : v_T = v|_T \in P_2(T), \forall T \in \mathcal{T}_h \text{ and } \partial_n v = 0 \text{ on } \partial\Omega\}$?

and

$$V_h^* = \begin{cases} V_h & \text{if } \alpha > 0 \\ \{v \in V_h : \int_{\Omega} v dx = 0\} & \text{if } \alpha = 0 \end{cases}$$

Now, if we choose $u \in V_h$ must we take account that the jump is discrete. We have now the final CP formulation. The discretized numerical problem is to solve $w_h \in V_h^*$ such that

$$\mathcal{A}(w_h, v_h) = F(v_h), \quad \forall v_h \in V_h^*. \quad (9)$$

where

$$\begin{aligned}
\mathcal{A}(w_h, v_h) &= (\alpha w_h, v_h)_{\Omega} \\
&+ \sum_{T \in \mathcal{T}_h} (D^2 w_h, D^2 v_h)_T \\
&+ \sum_{F \in \mathcal{F}_h^{int}} (\{\partial_{nn} w_h\}, \llbracket \partial_n v_h \rrbracket)_F + (\{\partial_{nn} v_h\}, \llbracket \partial_n w_h \rrbracket)_F + \frac{\gamma}{h} (\llbracket \partial_n w_h \rrbracket, \llbracket \partial_n v_h \rrbracket)_F \\
&+ \sum_{F \in \mathcal{F}_h^{ext}} -(\partial_{nt} w_h, \partial_t v_h)_F - (\partial_{nt} v_h, \partial_t w_h)_F
\end{aligned} \quad (10)$$

Is the external summation correct?

and

$$F(v_h) = (f, v_h)_{\Omega} + \sum_{F \in \mathcal{F}_h^{ext}} -(g, v_h)_F \quad (11)$$

Notice that the regulation term determined by respectively a global tuning parameter $\gamma > 0$. Another key component to the formulation in (10) after introduction of $w_h, v_h \in V_h^*$ is that we expanded $(\partial_{nn} w, \llbracket \partial_n v \rrbracket)_F \rightarrow (\{\partial_{nn} w_h\}, \llbracket \partial_n v_h \rrbracket)_F$ since we can longer not guarantee a continuous jump. For symmetric purposes we also added $(\{\partial_{nn} v_h\}, \llbracket \partial_n w_h \rrbracket)_F$. Similarly, for the summation over the external facets \mathcal{F}_h^{ext} we added the minus term $-(\partial_{nt} v_h, \partial_t v)$ for symmetry.

We may introduce the compact notation of (10).

$$\begin{aligned}
\mathcal{A}(w_h, v_h) = & (\alpha w_h, v_h)_\Omega \\
& + (D^2 w_h, D^2 v_h)_{\mathcal{T}_h} \\
& + (\{\partial_{nn} w_h\}, [\partial_n v_h])_{\mathcal{F}_h} + (\{\partial_{nn} v_h\}, [\partial_n w])_{\mathcal{F}_h} + \frac{\gamma}{h} ([\partial_n w_h], [\partial_n v_h])_{\mathcal{F}_h} \\
& - (\partial_{nt} w_h, \partial_t v_h)_{\mathcal{F}_h^{ext}} - (\partial_{nt} w_h, \partial_t v_h)_{\mathcal{F}_h^{ext}}
\end{aligned} \tag{12}$$

TODO: correct the error analysis with this \mathcal{A}_h including external facets \mathcal{F}_h^{ext}

3.5 Error and Stability Analysis of CP

To guarantee convergence and stability we may want to check coercivity and boundedness of the method.

First of all, let us now establish some important inequalities.

Cauchy-Schwarz inequality: $\|ab\| \leq \|a\| \|b\|$

Inverse inequality: $\frac{1}{h} \|\partial_{nn} v_h\|_{\mathcal{F}_h}^2 \leq C_j \|D^2 v_h\|_{\mathcal{T}_h}^2$

Youngs epsilon inequality: $2ab = 2\sqrt{\varepsilon}a \cdot \frac{b}{\sqrt{\varepsilon}} \leq \varepsilon a^2 + b^2 \frac{1}{\varepsilon}$

Let the energy norm be on the form

$$\begin{aligned}
\|v_h\|_h^2 &= \|v_h\|_{a_h}^2 = \|w_h\|_\Omega \|v_h\|_\Omega + \|D^2 v_h\|_{\mathcal{T}_h}^2 + \|h^{-\frac{1}{2}} [\partial_n v_h]\|_{\mathcal{F}_h}^2 \\
\|v\|_h^2 &= \|v\|_{a_h,*}^2 = \|v\|_{a_h}^2 + \|h^{\frac{1}{2}} \{\partial_{nn} v\}\|_F^2, \quad v \in V \oplus V_h
\end{aligned} \tag{13}$$

The method is said to be coercive if $\mathcal{A}_h(v_h, v_h) \geq C \|v_h\|_{a_h}^2$. Similarly, it is bounded if $\mathcal{A}_h(v_h, u_h) \leq C \|u_h\|_{a_h}^2 \|v_h\|_{a_h}^2$ and then, according to Lax Milgram (need reference), the solution does exist and be unique.

3.5.1 Coercivity

Suppose we have the CP problem described in (9). Then is the coercivity be computed such that

$$\begin{aligned}
\mathcal{A}(v_h, v_h) &= \alpha \|w_h v_h\|_\Omega + \|D^2 v_h\|_{\mathcal{T}_h}^2 + 2(\{\partial_{nn} v_h\}, [\partial_n v_h])_{\mathcal{F}_h} + \frac{\gamma}{h} \|[\partial_n v_h]\|_{\mathcal{F}_h}^2 \\
\text{Cauchy-Schwarz inequality} &\geq \alpha \|w_h\|_\Omega \|v_h\|_\Omega + \|D^2 v_h\|_{\mathcal{T}_h}^2 - 2\|h^{\frac{1}{2}} \{\partial_{nn} v_h\}\|_{\mathcal{F}_h} \|h^{-\frac{1}{2}} [\partial_n v_h]\|_{\mathcal{F}_h} + \gamma \|h^{-\frac{1}{2}} [\partial_n v_h]\|_{\mathcal{F}_h}^2 \\
\text{Inverse inequality} &\geq \alpha \|w_h\|_\Omega \|v_h\|_\Omega + \|D^2 v_h\|_{\mathcal{T}_h}^2 - 2C_j^{\frac{1}{2}} \|D^2 v_h\|_{\mathcal{T}_h} \|h^{-\frac{1}{2}} [\partial_n v_h]\|_{\mathcal{F}_h} + \gamma \|h^{-\frac{1}{2}} [\partial_n v_h]\|_{\mathcal{F}_h}^2 \\
\text{Youngs epsilon inequality} &\geq \alpha \|w_h\|_\Omega \|v_h\|_\Omega + \|D^2 v_h\|_{\mathcal{T}_h}^2 - \varepsilon C_j \|D^2 v_h\|_{\mathcal{T}_h}^2 - \frac{1}{\varepsilon} \|h^{\frac{1}{2}} [\partial_n v_h]\|_{\mathcal{F}_h}^2 + \gamma \|h^{-\frac{1}{2}} [\partial_n v_h]\|_{\mathcal{F}_h}^2 \\
&= \alpha \|w_h\|_\Omega \|v_h\|_\Omega + (1 - \varepsilon C_j) \|D^2 v_h\|_{\mathcal{T}_h}^2 + \left(\gamma - \frac{1}{\varepsilon}\right) \|h^{-\frac{1}{2}} [\partial_n v_h]\|_{\mathcal{F}_h}^2 \\
(\varepsilon = \frac{1}{2C_j}) \implies &= \alpha \|w_h\|_\Omega \|v_h\|_\Omega + \frac{1}{2} \|D^2 v_h\|_{\mathcal{T}_h}^2 + \underbrace{(\gamma - 2C_j)}_{\geq \frac{1}{2}} \|h^{\frac{1}{2}} [\partial_n v_h]\|_{\mathcal{F}_h}^2 \geq C \|v_h\|_{a_h}^2
\end{aligned}$$

This holds if $C = \min\{\alpha, 1/2\}$. Observe that for the first inequality is the standard **Cauchy-Schwarz inequality** such that

$$(\{\partial_{nn} v_h\}, [\partial_n v_h])_{\mathcal{F}_h} \geq -\|h^{-\frac{1}{2}} \{\partial_{nn} v_h\}\|_{\mathcal{F}_h} \|h^{\frac{1}{2}} [\partial_n v_h]\|_{\mathcal{F}_h}.$$

On the second inequality the **Inverse inequality** was applied,

$$-\|h^{\frac{1}{2}} \{\partial_{nn} v_h\}\|_{\mathcal{F}_h} \geq -C_j^{\frac{1}{2}} \|D^2 v_h\|_{\mathcal{T}_h}$$

The next step is then to use the **Youngs epsilon inequality** to be able to separate the facets and triangulation norms.

$$-2C_j^{\frac{1}{2}} \|D^2 v_h\|_{\mathcal{T}_h} \|h^{\frac{1}{2}} [\partial_n v_h]\|_{\mathcal{F}_h}^2 \geq -\varepsilon C_j \|D^2 v_h\|_{\mathcal{T}_h}^2 - \frac{1}{\varepsilon} \|h^{\frac{1}{2}} [\partial_n v_h]\|_{\mathcal{F}_h}^2$$

The last step was to choose a ε and γ as some positive constant so that the second term is restricted to be multiplied with something bigger than $\frac{1}{2}$. Thus, the term fulfils coercivity of the (13). Hence, the CP method is coercive.

3.5.2 Boundedness

We want the CP method to be bounded.

$$\begin{aligned}
\mathcal{A}(w_h, v_h) &= (\alpha w_h, v_h)_\Omega + (D^2 w_h, D^2 v_h)_{\mathcal{T}_h} + (\{\partial_{nn} w_h\}, [\partial_n v_h])_{\mathcal{F}_h} + (\{\partial_{nn} v_h\}, [\partial_n w_h])_{\mathcal{F}_h} \\
&\quad + \frac{\gamma}{h} ([\partial_n w_h], [\partial_n v_h])_{\mathcal{F}_h} \\
\text{Cauchy-Schwarz inequality} \quad &\leq \alpha \|w_h\|_\Omega \|v_h\|_\Omega + \|D^2 w_h\|_{\mathcal{T}_h} \|D^2 v_h\|_{\mathcal{T}_h} + \|h^{\frac{1}{2}} \{\partial_{nn} w_h\}\|_{\mathcal{F}_h} \|h^{-\frac{1}{2}} [\partial_n v_h]\|_{\mathcal{F}_h} \\
&\quad + \|h^{\frac{1}{2}} \{\partial_{nn} v_h\}\|_{\mathcal{F}_h} \|h^{-\frac{1}{2}} [\partial_n w_h]\|_{\mathcal{F}_h} + \gamma \|h^{-1} [\partial_n v_h]\|_{\mathcal{F}_h} \|[\partial_n w_h]\|_{\mathcal{F}_h} \\
\text{Inverse inequality} \quad &\leq \alpha \|w_h\|_\Omega \|v_h\|_\Omega + \|D^2 w_h\|_{\mathcal{T}_h} \|D^2 v_h\|_{\mathcal{T}_h} + C_j^{\frac{1}{2}} \|D^2 w_h\|_{\mathcal{T}_h} \|h^{-\frac{1}{2}} [\partial_n v_h]\|_{\mathcal{F}_h} + \\
&\quad C_j^{\frac{1}{2}} \|D^2 w_h\|_{\mathcal{T}_h} \|h^{-\frac{1}{2}} [\partial_n w_h]\|_{\mathcal{F}_h} + \gamma \|h^{-1} [\partial_n v_h]\|_{\mathcal{F}_h} \|[\partial_n w_h]\|_{\mathcal{F}_h} \\
\text{Using (14)} \quad &\leq \alpha \|w_h\|_{a_h} \|v_h\|_{a_h} + \|w_h\|_{a_h} \|v_h\|_{a_h} + 2C_j^{\frac{1}{2}} \|w_h\|_{a_h} \|v_h\|_{a_h} + \gamma \|v_h\|_{a_h} \|w_h\|_{a_h} \\
&\leq \left(\alpha + 1 + 2C_j^{\frac{1}{2}} + \gamma \right) \|v_h\|_{a_h} \|w_h\|_{a_h} \leq K \|v_h\|_{a_h} \|w_h\|_{a_h}
\end{aligned}$$

Thus, the CP method is shown to be bounded. Again, the first step was to apply the **Cauchy-Schwarz inequality** for every term. On the second inequality the **Inverse inequality** was applied so that

$$\|h^{\frac{1}{2}} \{\partial_{nn} v_h\}\|_{\mathcal{F}_h} \leq C_j^{\frac{1}{2}} \|D^2 v_h\|_{\mathcal{T}_h} \quad \text{and} \quad \|h^{\frac{1}{2}} \{\partial_{nn} w_h\}\|_{\mathcal{F}_h} \leq C_j^{\frac{1}{2}} \|D^2 w_h\|_{\mathcal{T}_h}.$$

The second step can we luckily observe that all terms individually is less than the norm such that

$$\begin{aligned}
\|w_h\|_\Omega \|v_h\|_\Omega &\leq \|w_h\|_{a_h} \|v_h\|_{a_h}, \\
\|D^2 w_h\|_{\mathcal{T}_h} \|D^2 v_h\|_{\mathcal{T}_h} &\leq \|w_h\|_{a_h} \|v_h\|_{a_h}, \\
\|D^2 w_h\|_{\mathcal{T}_h} \|h^{-\frac{1}{2}} [\partial_n v_h]\|_{\mathcal{F}_h} &\leq \|w_h\|_{a_h} \|v_h\|_{a_h}, \quad \|D^2 v_h\|_{\mathcal{T}_h} \|h^{-\frac{1}{2}} [\partial_n w_h]\|_{\mathcal{F}_h} \leq \|v_h\|_{a_h} \|w_h\|_{a_h}, \\
\text{and } \gamma \|h^{-1} [\partial_n v_h]\|_{\mathcal{F}_h} \|[\partial_n w_h]\|_{\mathcal{F}_h} &\leq \gamma \|v_h\|_{a_h} \|w_h\|_{a_h}.
\end{aligned} \tag{14}$$

Hence, the CP method is does fulfills the Lax Milgram criteria because it is both bounded and unique.

3.6 Apriori Estimates

We will now introduce the notion of an apriori estimate, which can be used the estimate the size of a solution even before we have a solution. However, we will first introduce some basic results from the theory of Galerkin methods.

3.6.1 Galerkin Orthogonality

Let us recall the weak continuous problem (15) and the weak discrete problem (16).

$$\text{Find } u \in V \quad \text{s.t. } a(u, v) = l(v) \quad \forall v \in V. \tag{15}$$

$$\text{Find } u_h \in V_h \quad \text{s.t. } a(u_h, v_h) = l(v_h) \quad \forall v_h \in V_h. \tag{16}$$

In fact, the key property of this problem formulation is that $a(u, v_h) = l(v_h) \quad \forall v_h \in V_h \subset V$. Thus, the Galerkin orthogonality property holds so that,

$$a(u, v_h) - a(u_h, v_h) = a(u - u_h, v_h) = 0, .$$

Keep in mind that we define $u - u_h \in V$

3.6.2 Céa's Lemma

Assume that the problems (15) and (16) satisfied the Lax Milgram criteria,

$$\begin{aligned}
a(u, v) &\leq C \|u\| \|v\| \quad (V_h), \quad \alpha \|u\|^2 \leq a(u, u) \quad \forall u, v \in V, \\
a(u_h, v_h) &\leq C \|u_h\| \|v_h\| \quad (V_h), \quad \alpha \|u_h\|^2 \leq a(u_h, u_h) \quad \forall u_h, v_h \in V_h.
\end{aligned}$$

Then the Céa's lemma says that this is satisfied,

$$\|u - u_h\| \leq \frac{C}{\alpha} \inf_{v_h \in V_h} \|v_h - v\|. \tag{17}$$

The lemma naturally arise using the following argumentation,

$$\begin{aligned}
\alpha \|u - u_h\|^2 &\leq a(u - u_h, u - u_h) \\
&= a(u - u_h, u - v_h) - a(u - u_h, v_h - u_h) \\
&\leq C \|u - u_h\| \|u - v_h\| \\
\implies \alpha \|u - u_h\| &\leq C \|v_h - u\| \leq C \inf_{v_h \in V_h} \|v_h - u\|.
\end{aligned} \tag{18}$$

3.6.3 Céa's Lemma for CP Method,

Since we have discrete coercivity, then $V_h \not\subset V$, thus the standard method does not work. Firstly, we want to use the results from 3.5. We have shown that

$$\begin{aligned}
\text{Discrete coercivity} \quad \hat{\alpha} \|u_h\|_{a_h}^2 &\leq \mathcal{A}(u_h, u_h) \\
\text{Boundedness (semi-discrete)} \quad \mathcal{A}(v, w_h) &\leq \tilde{C} \|v\|_{a_{h,*}} \|w_h\|_{a_h} \quad \forall v \in V_h \oplus H^4(\Omega). \\
\text{Boundedness (fully discrete)} \quad \mathcal{A}(v_h, w_h) &\leq \bar{C} \|v\|_{a_{h,*}} \|w_h\|_{a_h} \quad \forall v_h, w_h \in V_h
\end{aligned}$$

Let the difference have the form $u - u_h = (u - v_h) + (v_h - u_h)$ and the define identity

$$\|w_h\|_{a_{h,*}} \leq D \|w_h\|_{a_h}, \forall w_h \in V_h.$$

Thus, the norm can now be computed such that

$$\|u - u_h\|_{a_{h,*}} \leq \|u - v_h\|_{a_{h,*}} + \|v_h - u_h\|_{a_{h,*}} \leq \|u - v_h\|_{a_{h,*}} + D \|u_h - v_h\|_{a_h}.$$

Finally, following the same procedure as in (18) we get

$$\begin{aligned}
\|u_h - v_h\|_{a_h}^2 \hat{\alpha} &\leq \mathcal{A}(u_h - v_h, u_h - v_h) \\
&= \mathcal{A}_h(u_h - u, u_h - v_h) + \mathcal{A}(u - v_h, u_h - v_h) \\
&\leq \mathcal{A}(u - v_h, u_h - v_h) \\
&\leq \tilde{C} \|u - v_h\|_{a_{h,*}} \|u - v_h\|_{a_h}
\end{aligned}$$

Observe that we now have $\|u_h - v_h\|_{a_h} \leq \frac{\tilde{C}}{\hat{\alpha}} \|u - v_h\|_{a_{h,*}}$ and $\|u - u_h\|_{a_{h,*}} \leq \left(1 + D\tilde{C}/\hat{\alpha}\right) \cdot \|u - v_h\|_{a_{h,*}}$. Hence, we have derived a equivalent Céa's lemma for the CP method.

$$\begin{aligned}
\|u_h - v_h\|_{a_h} &\leq \frac{\tilde{C}}{\hat{\alpha}} \inf_{v_h \in V_h} \|v_h - u\|_{a_{h,*}} \\
\|u - u_h\|_{a_{h,*}} &\leq \left(1 + D\tilde{C}/\hat{\alpha}\right) \cdot \inf_{v_h \in V_h} \|u - v_h\|_{a_{h,*}}
\end{aligned}$$

4 Numerical Results

4.1 Manufactured Solution

4.2 Convergence Rate

4.3 Condition Number

5 HCP Method copied from NGSolve

We consider the Kirchhoff plate equation: Find $w \in H^2$, such that

$$\int \nabla^2 w : \nabla^2 v = \int f v$$

A conforming method requires C^1 continuous finite elements. But there is no good option available, and thus there is no H^2 conforming finite element space in NGSolve.

$$\sum_T \nabla^2 w : \nabla^2 v - \int_E \{\nabla^2 w\}_{nn} [\partial_n v] - \int_E \{\nabla^2 v\}_{nn} [\partial_n w] + \alpha \int_E [\partial_n w] [\partial_n v]$$

[Baker 77, Brenner Gudi Sung, 2010]

We consider its hybrid DG version, where the normal derivative is a new, facet-based variable:

$$\sum_T \nabla^2 w : \nabla^2 v - \int_{\partial T} (\nabla^2 w)_{nn} (\partial_n v - \widehat{v}_n) - \int_{\partial T} (\nabla^2 v)_{nn} (\partial_n w - \widehat{w}_n) + \alpha \int_E (\partial_n v - \widehat{v}_n)(\partial_n w - \widehat{w}_n)$$

6 Cahn Hilliard Equation on a Closed Membrane

Let c_0 and c_1 indicate the concentration profile of the substances in a 2 -phase system such that $c_0(\mathbf{x}, t) : \Omega \times [0, \infty] \rightarrow [0, 1]$ and similarly $c_1(\mathbf{x}, t) : \Omega \times [0, \infty] \rightarrow [0, 1]$, where \mathbf{x} is a element of some surface Ω and t is time. However, in the 2 phase problem will we will restrict ourself so that $c_0(t, \mathbf{x}) + c_1(t, \mathbf{x}) = 1$ at any \mathbf{x} at time t . A property of the restriction is that we now can express c_0 using c_1 , with no loss of information. Hence, let us now define $c = c_0$ so $c(\mathbf{x}, t) : \Omega \times [0, \infty] \rightarrow [0, 1]$. It has been shown that 2 phase system if thermodynamically unstable can be evolve into a phase separation described by a evolutional differential equation [1] using a model based on chemical energy of the substances. However, further development has been done [4] to solve this equation on surfaces. Now assume model that we want to describe is a phase-separation on a closed membrane surface Γ , so that $c(\mathbf{x}, t) : \Gamma \times [0, T] \rightarrow [0, 1]$. Then is the surface Cahn Hilliard equation described such that

$$\rho \frac{\partial c}{\partial t} - \nabla_\Gamma (M \nabla_\Gamma (f'_0 - \varepsilon^2 \nabla_\Gamma^2 c)) = 0 \quad \text{on } \Gamma. \quad (19)$$

We define here the tangential gradient operator to be $\nabla_\Gamma c = \nabla c - (\mathbf{n} \nabla c) \mathbf{n}$ applied on the surface Γ restricted to $\mathbf{n} \cdot \nabla_\Gamma c = 0$.

Lets define ε to be the size of the layer between the substances c_1 and c_2 . The density ρ is simply defined such that $\rho = \frac{m}{S_\Gamma}$ is a constant based on the total mass divided by the total surface area of Γ . Here is the mobility M often derived such that is dependent on c and is crucial for the result during a possible coarsening event [4]. However, the free energy per unit surface $f_0 = f_0(c)$ is derived based on the thermodynamical model and should according to [4] be non convex and nonlinear.

A important observation is that equation (19) is a fourth order equation which makes it more challenging to solve using conventional FEM methods. This clear when writing the equation on the equivalent weak form and second order equations arise.

7 Appendix

7.1 The Space $L^2(\Omega)$

Using the definition from [5] and we let Ω be a an open set in \mathbb{R}^d and $p \in \mathbb{R}$ such that $p \geq 1$. Then we denote $L^p(\Omega)$ to be the set of measurable function $u : \Omega \rightarrow \mathbb{R}$ such that it is equipped in a finite Banach space

$$\|u\|_{L^p(\Omega)} = \left(\int_{\Omega} |u|^p \right)^{\frac{1}{p}}.$$

Now let $u, v : \Omega \rightarrow \mathbb{R}$. Then is $L^2(\Omega)$ a Hilbert space when the inner product is finite such that this exists

$$(u, v)_{L^2(\Omega)} = \int_{\Omega} uv.$$

If the integral is finite do we say that $u, v \in L^p(\Omega)$.

7.2 The Space $H^m(\Omega)$, $m > 1$

Again using the definition from [5]. Let $\alpha = (\alpha_1, \dots, \alpha_d)$, $\alpha \geq 0$, such that $|\alpha| = \sum_{i=1}^d \alpha_i$. Now we define the space

$$H^m(\Omega) = \{u \in L^2(\Omega) : D^{\alpha}u \in L^2(\Omega) \quad \forall \alpha : |\alpha| \leq m\}.$$

Suppose that u, v is measurable functions. We can now define $u \in H^m(\Omega)$ the Banach space is finite .

$$\|u\|_{H^m(\Omega)} = \left(\|u\|_{L^2(\Omega)}^2 + \sum_{k=1}^m \|u\|_{H^k(\Omega)}^2 \right)^{\frac{1}{2}}, \quad \|u\|_{H^k(\Omega)} = \sqrt{\sum_{|\alpha|=k} \|D^{\alpha}u\|_{L^2(\Omega)}^2}$$

Similarly for the finite Hilbert space

$$(u, v)_{H^m(\Omega)} = \sum_{|\alpha| \leq m} \int_{\Omega} D^{\alpha}u D^{\alpha}v$$

References

- [1] John W. Cahn and John E. Hilliard. “Free Energy of a Nonuniform System. I. Interfacial Free Energy”. In: *The Journal of Chemical Physics* 28.2 (1958), pp. 258–267. DOI: [10.1063/1.1744102](https://doi.org/10.1063/1.1744102). eprint: <https://doi.org/10.1063/1.1744102>. URL: <https://doi.org/10.1063/1.1744102>.
- [2] Susanne C Brenner et al. “A Quadratic C^0 Interior Penalty Method for Linear Fourth Order Boundary Value Problems with Boundary Conditions of the Cahn–Hilliard Type”. In: *SIAM Journal on Numerical Analysis* 50.4 (2012), pp. 2088–2110.
- [3] S. Gu and La.). Department of Mathematics Louisiana State University (Baton Rouge. *C0 Interior Penalty Methods for Cahn-Hilliard Equations*. Dissertation (Louisiana State University (Baton Rouge, La.)) Louisiana State University, 2012. URL: <https://books.google.no/books?id=eKP1xQEACAAJ>.
- [4] Vladimir Yushutin et al. “A computational study of lateral phase separation in biological membranes”. In: *International Journal for Numerical Methods in Biomedical Engineering* 35.3 (2019). e3181 cnm.3181, e3181. DOI: <https://doi.org/10.1002/cnm.3181>. eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/cnm.3181>. URL: <https://onlinelibrary.wiley.com/doi/abs/10.1002/cnm.3181>.
- [5] A. Manzoni, A. Quarteroni, and S. Salsa. *Optimal Control of Partial Differential Equations: Analysis, Approximation, and Applications*. Applied Mathematical Sciences. Springer International Publishing, 2021. ISBN: 9783030772253. URL: <https://books.google.no/books?id=V3NpzgEACAAJ>.