

# Mixed finite elements for port-Hamiltonian models of von Kármán beams

7th IFAC Conference on Lagrangian and Hamiltonian method for non linear control

Andrea Brugnoli<sup>1</sup>   Ramy Rashad<sup>1</sup>   Federico Califano<sup>1</sup>   Stefano Stramigioli<sup>1</sup>   Denis Matignon<sup>2</sup>

<sup>1</sup>University of Twente, Enschede (NL)

<sup>2</sup>ISAE-SUPAERO, Toulouse (FR)

11-13 October, 2021



# Overview

---

- 1 On the von-Karman assumption

# Outline

---

- 1 On the von-Karman assumption

# The von-Karman

---

This theory has puzzled mathematicians and physicists as the derivation of the model rely on some not well justified assumptions.

# Membrane-bending problem for thin beams

---

For small deformations the membrane and bending behavior are uncoupled:

$$\frac{\partial}{\partial t} \begin{pmatrix} \alpha_u \\ \alpha_\varepsilon \\ \alpha_w \\ \alpha_\kappa \end{pmatrix} = \begin{bmatrix} 0 & \partial_x & 0 & 0 \\ \partial_x & 0 & 0 & 0 \\ 0 & 0 & 0 & -\partial_{xx} \\ 0 & 0 & \partial_{xx} & 0 \end{bmatrix} \begin{pmatrix} e_u \\ e_\varepsilon \\ e_w \\ e_\kappa \end{pmatrix}$$

If the material is isotropic

# References I

---