| | Farm economy | | | Farm functioning | | | Environment | |
|---|---|---|---|---|---|---|---|---|
| | Productivity | Profitability | Labour demand | Soil health | Inputs efficiency | Resilience to pests | Biodiversity | Climate mitigation |
| **a) Cereals** | | | | | | | | |
| **Biomass management, compared with no residues or mulch** | | | | | | | | |
| Organic mulch | **73%** n = 278 | 38% n = 24 | | **76%** n = 319 | **79%** n = 63 | 69% n = 13 | 0% n = 4 | |
| Residue incorporation | **67%** n = 263 | 18% n = 11 | | **61%** n = 92 | | | | 48% n = 40 |
| Residue retention | **85%** n = 78 | 42% n = 52 | | 77% n = 26 | | | 83% n = 6 | 0% n = 4 |
| **Nutrient management, compared with no fertilizer applied** | | | | | | | | |
| Organic fertilizer | **66%** n = 151 | 59% n = 27 | | **90%** n = 71 | | | 75% n = 108 | 50% n = 4 |
| Mixed fertilizer | **84%** n = 218 | 65% n = 20 | | **70%** n = 67 | 100% n = 4 | | | 43% n = 23 |
| Inoculation | 48% n = 80 | | | | | 33% n = 24 | | |
| Biochar amendment | **63%** n = 67 | | | 0% n = 3 | | | | 47% n = 15 |
| **Nutrient management, compared with inorganic fertilizer** | | | | | | | | |
| Mixed fertilizer | 100% n = 17 | | | | | | | |
| Organic fertilizer | 100% n = 5 | | | | | | | |
| **Diversification, compared with monoculture** | | | | | | | | |
| Intercropping | **80%** n = 505 | 50% n = 12 | | | | **96%** n = 72 | 71% n = 24 | 17% n = 6 |
| Agroforestry | **48%** n = 229 | 50% n = 22 | | | | 41% n = 44 | 0% n = 5 | |
| Crop rotation | 92% n = 100 | | | | | | 75% n = 4 | |
| Natural Fallow | 34% n = 80 | | | | | 29% n = 24 | | |
| Improved fallow | 79% n = 42 | | | | | | | |
| **Diversification, compared with intercropping** | | | | | | | | |
| Agroforestry | | 71% n = 14 | | | | | | |
| **Soil management, compared with conventional tillage** | | | | | | | | |
| Reduced tillage | **56%** n = 124 | 70% n = 53 | 100% n = 1 | 42% n = 19 | | | | |
| **Soil management, compared with no liming** | | | | | | | | |
| Liming | 80% n = 5 | | | | | | | |
| **Variety management, compared with improved variety** | | | | | | | | |
| Local variety | 25% n = 4 | | | | | | | |
| **Pest management, compared with no chemical applied** | | | | | | | | |
| Mechanical weeding | | | | | | | 100% n = 1 | |
| **b) Pulses and legumes** | | | | | | | | |
| **Diversification, compared with monoculture** | | | | | | | | |
| Intercropping | **27%** n = 260 | 69% n = 32 | | | | 92% n = 13 | 50% n = 20 | 17% n = 6 |
| Agroforestry | 27% n = 15 | 50% n = 22 | | | | | | |
| Crop rotation | 75% n = 4 | | | | | | 75% n = 4 | |
| **Diversification, compared with intercropping** | | | | | | | | |
| Agroforestry | | 71% n = 14 | | | | | | |
| **Biomass management, compared with no residues or mulch** | | | | | | | | |
| Residue incorporation | **31%** n = 152 | 18% n = 11 | | 0% n = 18 | | | | |
| Residue retention | 30% n = 10 | | | | | | 83% n = 6 | |
| Organic mulch | 0% n = 3 | | | | | | | |
| **Nutrient management, compared with no fertilizer applied** | | | | | | | | |
| Organic fertilizer | 79% n = 66 | 82% n = 11 | | 48% n = 64 | | | | |
| Biochar amendment | 88% n = 24 | | | | | | | |
| Mixed fertilizer | 100% n = 24 | | | | | | | |
| **Variety management, compared with improved variety** | | | | | | | | |
| Local variety | 22% n = 36 | | | 50% n = 24 | | | | |
| **Water management, compared with rainfed agriculture** | | | | | | | | |
| Supplemental irrigation | 100% n = 12 | | | | 11% n = 18 | | | |
| **Soil management, compared with conventional tillage** | | | | | | | | |
| Zero tillage | 0% n = 12 | | | | | | | |
| Reduced tillage | 0% n = 6 | | | | | | | |
| **c) Vegetables** | | | | | | | | |
| **Water management, compared with rainfed agriculture** | | | | | | | | |
| Supplemental irrigation | 66% n = 402 | 50% n = 12 | | | | | | |
| **Diversification, compared with monoculture** | | | | | | | | |
| Intercropping | 28% n = 32 | | | | | 62% n = 16 | | |
| **Variety management, compared with improved variety** | | | | | | | | |
| Local variety | 42% n = 36 | | | | | | | |
| **Nutrient management, compared with no fertilizer applied** | | | | | | | | |
| Mixed fertilizer | 100% n = 2 | 75% n = 4 | | | | | | 0% n = 2 |
| Organic fertilizer | 100% n = 2 | 75% n = 4 | | | | | | 50% n = 2 |
| **d) Roots and tubers** | | | | | | | | |
| **Diversification, compared with monoculture** | | | | | | | | |
| Intercropping | 8% n = 48 | 69% n = 32 | | | | 100% n = 4 | 50% n = 8 | |
| **Nutrient management, compared with no fertilizer applied** | | | | | | | | |
| Mixed fertilizer | 0% n = 6 | | | | | | | |
| **e) Nuts and seeds** | | | | | | | | |
| **Diversification, compared with monoculture** | | | | | | | | |
| Intercropping | | | | | | 100% n = 6 | 50% n = 12 | |

**% positive comparisons**

| | Negative (≤35%) | Neutral (36–64%) | Positive (≥65%) |
|---|---|---|---|
| Strong evidence (n ≥ 50, ≥ 3 studies) | | | |
| Limited evidence (fewer comparisons or studies) | | | |

Empty cells: not measured