

List of papers included in the dataset

Title

A global dataset of experimental intercropping and agroforestry studies in horticulture

<https://doi.org/10.57745/HV33V1>

1. Abbes, Z., Trabelsi, I., Kharrat, M., & Amri, M. (2019). Intercropping with fenugreek (*Trigonella foenum-graecum*) enhanced seed yield and reduced *Orobanche foetida* infestation in faba bean (*Vicia faba*). *Biological Agriculture & Horticulture*, 35(4), 238–247. <https://doi.org/10.1080/01448765.2019.1616614>
2. Abd El-Gai, M. A., Al-Dokeshy, M. H., & M.T. Nasse, D. (2014). Effects of Intercropping System of Tomato and Common Bean on Growth, Yield Components and Land Equivalent Ratio in New Valley Governorate. *Asian Journal of Crop Science*, 6(3), 254–261. <https://doi.org/10.3923/ajcs.2014.254.261>
3. Abdel-Wahab, S. I., & Abdel-Wahab, E. I. (2021). Cropping Systems of Fenugreek with Faba Bean to Reduce Broomrape Infestation. *Legume Research, Of*. <https://doi.org/10.18805/LR-621>
4. Abidi, I., Daoui, K., Abouabdillah, A., Belqadi, L., Mahyou, H., Bazile, D., Douaik, A., Gaboun, F., Hassane Sidikou, A. A., & Alaoui, S. B. (2024). Quinoa–Olive Agroforestry System Assessment in Semi-Arid Environments: Performance of an Innovative System. *Agronomy*, 14(3), 495. <https://doi.org/10.3390/agronomy14030495>
5. Abou-Husse, S. D., Salman, S. R., Abdel-Mawgoud, A. M. R., & Ghoname, A. A. (2005). Productivity, Quality and Profit of Sole or Intercropped Green Bean (*Phaseolus vulgaris* L.) Crop. *Journal of Agronomy*, 4(2), 151–155. <https://doi.org/10.3923/ja.2005.151.155>
6. Adeniyi, O. R. (2011). Economic aspects of intercropping systems of vegetables (okra, tomato and cowpea). *African Journal of Plant Science*, 5(11), 648–655.
7. Adeoti, A. O., & Pitan, O. R. (2023). Influence of time of introduction and proportion on the efficacy of grain amaranth trap crop against *Aulacophora foveicollis* and *Epilachna chrysomelina* on cucumber (*Cucumis sativus* L.). *Archives of Phytopathology and Plant Protection*, 56(14), 1062–1075. <https://doi.org/10.1080/03235408.2023.2256648>
8. Adhikary, S., Bagchi, D. K., Ghosal, P., Banerjee, R. N., & Chatterjee, B. N. (1991). Studies on Maize-Legume Intercropping and their Residual Effects on Soil Fertility

Status and Succeeding Crop in Upland Situation. *Journal of Agronomy and Crop Science*, 167(5), 289–293. <https://doi.org/10.1111/j.1439-037X.1991.tb00959.x>

9. Ahmed, M., Ishtiaque, S., Sarker, M., Khan, A. M., Choudhury, A., Hasan, M., Hossain, F., Paul, S., & Islam, M. (2016). Hybrid Maize and Chilli Intercropping in the Hilly Areas of Bandarban. *Bangladesh Agronomy Journal*, 19(1), 45–48. <https://doi.org/10.3329/baj.v19i1.29869>
10. Albuquerque, J. D. A. A. D., Sediya, T., Alves, J. M. A., Silva, A. A. D., & Uchôa, S. C. P. (2012). Cultivo de mandioca e feijão em sistemas consorciados realizado em Coimbra, Minas Gerais, Brasil. *Revista Ciência Agronômica*, 43(3), 532–538. <https://doi.org/10.1590/S1806-66902012000300016>
11. Al-Dalain, S. A. (2009). Effect of Intercropping of Zea Maize with Potato Solanum tuberosum, L. on Potato Growth and on the Productivity and Land Equivalent Ratio of Potato and Zea Maize. *Agricultural Journal*, 4(3), 164–170.
12. Álvarez-Solís, J. D., Mendoza-Núñez, J. A., León-Martínez, N. S., Castellanos-Albores, J., & Federico A, G.-M. (2016). Effect of Bokashi and Vermicompost Leachate on Yield and Quality of Pepper (capsicum Annuum) and Onion (allium Cepa) Under Monoculture and Intercropping Cultures. *Ciencia e Investigación Agraria*, 43(2), 7–7. <https://doi.org/10.4067/S0718-16202016000200007>
13. Alves, S. M. C., Abboud, A. C. D. S., Ribeiro, R. D. L. D., & Almeida, D. L. D. (2004). Balanço do nitrogênio e fósforo em solo com cultivo orgânico de hortaliças após a incorporação de biomassa de guandu. *Pesquisa Agropecuária Brasileira*, 39(11), 1111–1117. <https://doi.org/10.1590/S0100-204X2004001100009>
14. Amassaghrou, A., Barkaoui, K., Bouaziz, A., Alaoui, S. B., Fatemi, Z. E. A., & Daoui, K. (2023). Yield and related traits of three legume crops grown in olive-based agroforestry under an intense drought in the South Mediterranean. *Saudi Journal of Biological Sciences*, 30(4), 103597. <https://doi.org/10.1016/j.sjbs.2023.103597>
15. Ambrosano, E. J., Salgado, G. C., Otsuk, I. P., Prati, P., Henrique, C. M., & Melo, P. C. T. D. (2018). Organic cherry tomato yield and quality as affect by intercropping green manure. *Acta Scientiarum. Agronomy*, 40(1), 36530. <https://doi.org/10.4025/actasciagron.v40i1.36530>
16. Amiriyan Chelan, Z., Amini, R., & Dabbagh Mohammadi Nasab, A. (2023). Essential oil yield and compositions of Dracocephalum moldavica L. in intercropping with fenugreek, inoculation with mycorrhizal fungi and bacteria. *Scientific Reports*, 13(1), 8039. <https://doi.org/10.1038/s41598-023-35156-x>
17. Amiriyan Chelan, Z., Amini, R., & Dabbagh Mohammadi Nasab, A. (2024). Optimizing fenugreek (Trigonella foenum-graecum L.) oil yield and compositions in

intercropping through growth-promoting bacteria and mycorrhiza. *Frontiers in Agronomy*, 6, 1422236. <https://doi.org/10.3389/fagro.2024.1422236>

18. An, R., Yu, R.-P., Xing, Y., Zhang, J.-D., Bao, X.-G., Lambers, H., & Li, L. (2023). Enhanced phosphorus-fertilizer-use efficiency and sustainable phosphorus management with intercropping. *Agronomy for Sustainable Development*, 43(5), 57. <https://doi.org/10.1007/s13593-023-00916-6>
19. Anjum, M. A., Qasim, S. A., Ahmad, S., & Hussain, S. (2015). Assessment of Advantages of Pea and Non-Legume Winter Vegetable Intercropping Systems Through Competition and Economic Indices. *Experimental Agriculture*, 51(3), 327–343. <https://doi.org/10.1017/S0014479714000337>
20. Auler, P. A. M., Fidalski, J., Pavan, M. A., & Neves, C. S. V. J. (2008). Produção de laranja “Pêra” em sistemas de preparo de solo e manejo nas entrelinhas. *Revista Brasileira de Ciência do Solo*, 32(1), 363–374. <https://doi.org/10.1590/S0100-06832008000100034>
21. Azandémè-Hounmalon, G. Y., Logbo, J., Gbèblonoudo Dassou, A., Lokossi, L., Akpla, E., Fiaboe, K. K. M., & Tamò, M. (2023). Investigation of amaranth production constraints and pest infestation reduction by basil intercropping. *Journal of Agriculture and Food Research*, 12, 100627. <https://doi.org/10.1016/j.jafr.2023.100627>
22. Azevedo, F. A. D., Rossetto, M. P., Schinor, E. H., Martelli, I. B., & Pacheco, C. D. A. (2012). Influência do manejo da entrelinha do pomar na produtividade da laranjeira-’Pera’. *Revista Brasileira de Fruticultura*, 34(1), 134–142. <https://doi.org/10.1590/S0100-29452012000100019>
23. Badawy, S., & Shalaby, G. (2015). Effect of Intercropping of Sugar Beet with Onion and Garlic on Insect Infestation, Sugar Beet Yield and Economics. *Journal of Plant Production*, 6(6), 903–914. <https://doi.org/10.21608/jpp.2015.49788>
24. Bai, W., Sun, Z., Zheng, J., Du, G., Feng, L., Cai, Q., Yang, N., Feng, C., Zhang, Z., Evers, J. B., Van Der Werf, W., & Zhang, L. (2016). Mixing trees and crops increases land and water use efficiencies in a semi-arid area. *Agricultural Water Management*, 178, 281–290. <https://doi.org/10.1016/j.agwat.2016.10.007>
25. Banik, P., & Sharma, R. C. (2009). Yield and Resource Utilization Efficiency in Baby Corn—Legume-Intercropping System in the Eastern Plateau of India. *Journal of Sustainable Agriculture*, 33(4), 379–395. <https://doi.org/10.1080/10440040902834970>
26. Bantie, Y. B. (2015). Determination of Effective Spatial Arrangement for Intercropping of Maize (*Zea mays* L.) and Potato (*Solanum tuberosum* L.) Using Competition

Indices Ethiopia. *Journal of Horticulture*, 02(02). <https://doi.org/10.4172/2376-0354.1000137>

27. Baumann, D. T., Bastiaans, L., & Kropff, M. J. (2001). Competition and Crop Performance in a Leek–Celery Intercropping System. *Crop Science*, 41(3), 764–774. <https://doi.org/10.2135/cropsci2001.413764x>
28. Bavec, M., Zuljan, M., Robačar, M., & Bavec, F. (2012). White Cabbage Productivity in Intercropping Production Systems. *Acta Horticulturae*, 933, 343–346. <https://doi.org/10.17660/ActaHortic.2012.933.44>
29. Begam, A., Pramanick, M., Dutta, S., Paramanik, B., Dutta, G., Patra, P. S., Kundu, A., & Biswas, A. (2024). Inter-cropping patterns and nutrient management effects on maize growth, yield and quality. *Field Crops Research*, 310, 109363. <https://doi.org/10.1016/j.fcr.2024.109363>
30. Begam, A., Pramanick, M., Dutta, S., Ray, M., & Sengupta, K. (2023). Growth and Yield Responses of Cowpea (*Vigna unguiculata* L.) as Influenced by Crop Geometry and Nutrient Management Practices. *Legume Research*, 46(9), 1184–1191. <https://doi.org/10.18805/LR-4839>
31. Begum, S., Zaman, M., & Khan, A. (2015). Intercropping of root crops with chilli in charlands of Mymensingh. *Progressive Agriculture*, 26(2), 109–114. <https://doi.org/10.3329/pa.v26i2.25964>
32. Benalia, H., Fatima, L.-L., Fatma-Zohra, B., Omar, K., Bahia, Z., Mounir, S., Fouzi, G., Merwa, F., & Mourad, L. (2024). Comparative assessment of agroforestry performances between pea and barley intercropping with olive tree under Mediterranean conditions. *Agroforestry Systems*, 98(3), 619–635. <https://doi.org/10.1007/s10457-023-00933-0>
33. Bezerra Neto, F., Silva, M. L. D., Lima, J. S. S. D., Barros Júnior, A. P., Silva, I. N., & Chaves, A. P. (2019). Productive Viability and Profitability of Carrot-Cowpea Intercropping Using Different Amounts of Calotropis Procera. *Revista Caatinga*, 32(1), 62–71. <https://doi.org/10.1590/1983-21252019v32n107rc>
34. Bhagat, R., Walia, S. S., Dheri, G. S., Singh, G., & Sharma, K. (2024). Pear (*Pyrus communis*)-based agroecosystem improves soil nutrient dynamics, microbial biomass, enzymatic activity, farm productivity and profitability. *Scientia Horticulturae*, 336, 113398. <https://doi.org/10.1016/j.scienta.2024.113398>
35. Bhardwaj, A., Kaur, N., Dhat, A. S., & Gill, R. I. S. (2021). Optimization of onion planting time and variety under *Populus deltoides*-based agroforestry system in North-Western India. *Agroforestry Systems*, 95(3), 533–546. <https://doi.org/10.1007/s10457-021-00600-2>

36. Bhardwaj, R., Kushwaha, H. S., Birla, J., Turkar, Y., & Patidar, K. (2022). Studies on Nutrient Management in Pigeon Pea [*Cajanus cajan* (L.) Millsp.] based Inter Cropping System. *Legume Research, Of*. <https://doi.org/10.18805/LR-3947>
37. Bhat, R., Wani, W. M., Sharma, M. K., & Ashraf, N. (2014). Studies on Intercropping with Leguminous and Non-leguminous Crops on Yield, Leaf Nutrient Status and Relative Economic Yield of Apple cv. Red Delicious. *International Journal of Horticulture*. <https://doi.org/10.5376/ijh.2014.04.0005>
38. Biswas, B., Chakraborty, D., Timsina, J., Bhowmick, U. R., Dhara, P. K., Ghosh (Lkn), D. K., Sarkar, A., Mondal, M., Adhikary, S., Kanthal, S., Patra, K., Sarkar, S., Parsad, R., & Ray, B. R. (2022). Agroforestry offers multiple ecosystem services in degraded lateritic soils. *Journal of Cleaner Production*, 365, 132768. <https://doi.org/10.1016/j.jclepro.2022.132768>
39. Błażewicz-Woźniak, M., & Wach, D. (2011). The Effect of Intercropping on Yielding of Root Vegetables of Apiaceae Family. *Acta Scientiarum Polonorum*, 10(4).
40. Bomford, M. K. (2009). Do Tomatoes Love Basil but Hate Brussels Sprouts? Competition and Land-Use Efficiency of Popularly Recommended and Discouraged Crop Mixtures in Biointensive Agriculture Systems. *Journal of Sustainable Agriculture*, 33(4), 396–417. <https://doi.org/10.1080/10440040902835001>
41. Bordin, I., Neves, C. S. V. J., Francio Filho, P., Preti, E. A., & Cardoso, C. (2008). Crescimento de milho e guandu, desempenho de plantas cítricas e propriedades físicas do solo escarificado em um pomar. *Revista Brasileira de Ciência do Solo*, 32(4), 1409–1418. <https://doi.org/10.1590/S0100-06832008000400005>
42. Brintha, I., & Seran, T. H. (2012). Effect of intercropping chili (*Capsicum annuum* L.) with onion (*Allium cepa* L.) in sandy regosol. *Bangladesh Journal of Agricultural Research*, 37(3), 547–550. <https://doi.org/10.3329/bjar.v37i3.12132>
43. Bustamante-González, C. A., Ferrás-Negrín, Y., Hernández-Forte, I., & Rivera-Espinosa, R. (2022). Dr Beneficios del intercalamiento de canavalia inoculada con hongos micorrízicos y *Rhizobium* en *Coffea canephora*. *Agronomía Mesoamericana*, 46288. <https://doi.org/10.15517/am.v33i2.46288>
44. Caetano, L. C. S., Ferreira, J. M., & Araújo, M. L. D. (1999). Produtividade de cenoura e alface em sistema de consorciação. *Horticultura Brasileira*, 17(2), 143–146. <https://doi.org/10.1590/S0102-05361999000200013>
45. Carlos, T. J., Cecílio Filho, A. B., Dos Reis Cardoso Passos, D., & Dos Santos Reis, I. (2021). Collard greens and chicory intercropping efficiency as a function of

chicory (*Cichorium intybus*) transplant time. *Revista de La Facultad de Ciencias Agrarias UNCuyo*, 53(2), 91–99. <https://doi.org/10.48162/rev.39.043>

46. Carrubba, A., Torre, R., Saiano, F., & Aiello, P. (2008). Sustainable production of fennel and dill by intercropping. *Agronomy for Sustainable Development*, 28(2), 247–256. <https://doi.org/10.1051/agro:2007040>
47. Carvalho, A. J. D., Andrade, M. J. B. D., Guimarães, R. J., & Moraes, A. R. D. (2010). Sistemas de produção de feijão intercalado com cafeeiro adensado em período de formação ou após recepa. *Revista Ceres*, 57(3), 383–392. <https://doi.org/10.1590/S0034-737X2010000300015>
48. Carvalho, L. M. D., Nunes, M. U. C., Oliveira, I. R. D., & Leal, M. D. L. D. S. (2009). Produtividade do tomateiro em cultivo solteiro e consorciado com espécies aromáticas e medicinais. *Horticultura Brasileira*, 27(4), 458–464. <https://doi.org/10.1590/S0102-05362009000400010>
49. Castro, C. M. D., Alves, B. J. R., Almeida, D. L. D., & Ribeiro, R. D. L. D. (2004). Adubação verde como fonte de nitrogênio para a cultura da berinjela em sistema orgânico. *Pesquisa Agropecuária Brasileira*, 39(8), 779–785. <https://doi.org/10.1590/S0100-204X2004000800008>
50. Cecílio Filho, A. B., Alves, A. U., Galati, V. C., Bezerra Neto, F., Barbosa, J. C., & Machado, B. Q. V. (2022). Intercropping of Eggplant and Tomato as Function of Times of Transplant and Cropping Season. *Revista Caatinga*, 35(2), 276–287. <https://doi.org/10.1590/1983-21252022v35n204rc>
51. Cecílio Filho, A. B., Machado, B. Q., Alves, A. U., Pereira, B. D. J., Guerra, N. M., & Bezerra Neto, F. (2022). Bio-agronomic efficiency indices of eggplant and tomato intercropping. *Horticultura Brasileira*, 40(2), 181–189. <https://doi.org/10.1590/s0102-0536-20220207>
52. Cecílio Filho, A. B., Medelo, M. J. Y., Pontes, S. C. D., & Nascimento, C. S. (2021). Chicory and Arugula in Intercropping with Collard Greens. *Revista Caatinga*, 34(4), 772–779. <https://doi.org/10.1590/1983-21252021v34n404rc>
53. Cecílio Filho, A. B., Rezende, B. L. A., Barbosa, J. C., Feltrim, A. L., Silva, G. S. D., & Grangeiro, L. C. (2008). Interação entre alface e tomateiro consorciados em ambiente protegido, em diferentes épocas. *Horticultura Brasileira*, 26(2), 158–164. <https://doi.org/10.1590/S0102-05362008000200006>
54. Cecílio Filho, A. B., Rezende, B. L. A., Barbosa, J. C., & Grangeiro, L. C. (2011). Agronomic efficiency of intercropping tomato and lettuce. *Anais Da Academia Brasileira de Ciências*, 83(3), 1109–1119. <https://doi.org/10.1590/S0001-37652011000300029>

55. Cecílio Filho, A. B., Rezende, B. L. A., & Canato, G. H. D. (2007). Produtividade de alface e rabanete em cultivo consorciado estabelecido em diferentes épocas e espaçamentos entre linhas. *Horticultura Brasileira*, 25(1), 15–19. <https://doi.org/10.1590/S0102-05362007000100004>
56. Cecílio Filho, A. B., Rezende, B. L. A., & Costa, C. C. (2010). Economic analysis of the intercropping of lettuce and tomato in different seasons under protected cultivation. *Horticultura Brasileira*, 28(3), 326–336. <https://doi.org/10.1590/S0102-05362010000300015>
57. Cecílio Filho, A. B., Rezende, B. L. A., & Dutra, A. F. (2019). Yield of Intercropped Lettuce and Cucumber as a Function of Population Density and Cropping Season. *Revista Caatinga*, 32(4), 943–951. <https://doi.org/10.1590/1983-21252019v32n410rc>
58. Cenpukdee, U., & Fukai, S. (1992). Cassava/legume intercropping with contrasting cassava cultivars. 1. Competition between component crops under three intercropping conditions. *Field Crops Research*, 29(2), 113–133. [https://doi.org/10.1016/0378-4290\(92\)90082-K](https://doi.org/10.1016/0378-4290(92)90082-K)
59. Cesar, M. N. Z., Ribeiro, R. D. L. D., Paula, P. D. D., Polidoro, J. C., Manera, T. D. C., & Guerra, J. G. M. (2007). Desempenho do pimentão em cultivo orgânico, submetido ao desbaste e consórcio. *Horticultura Brasileira*, 25(3), 322–326. <https://doi.org/10.1590/S0102-05362007000300002>
60. Chabi-Olaye, A., Nolte, C., Schulthess, F., & Borgemeister, C. (2006). Relationships of soil fertility and stem borers damage to yield in maize-based cropping system in Cameroon. *Annales de La Société Entomologique de France (N.S.)*, 42(3–4), 471–479. <https://doi.org/10.1080/00379271.2006.10697481>
61. Chaudhari, D. T., Vekariya, P. D., Talpada, M. M., & Sutaria, G. S. (2016). Enhancing productivity of groundnut based intercropping systems under rainfed conditions of Gujarat. *Legume Research*. <https://doi.org/10.18805/lr.v0i0.7849>
62. Chaves, A. P., Bezerra Neto, F., Lima, J. S. S., Silva, J. N., Santos, E. C., Nunes, R. L. C., & Lino, V. A. D. S. (2022). Bio-agroeconomic returns in beet-cowpea intercropping by optimization of population densities and spatial arrangements. *Acta Scientiarum. Agronomy*, 44, e55146. <https://doi.org/10.4025/actasciagron.v44i1.55146>
63. Choudhary, V. K., Dixit, A., & Chauhan, B. S. (2016). Resource-use maximisation through legume intercropping with maize in the eastern Himalayan region of India. *Crop and Pasture Science*, 67(5), 508. <https://doi.org/10.1071/CP15233>

64. Choudhary, V. K., Dixit, A., Suresh Kumar, P., & Chauhan, B. S. (2014). Productivity, Weed Dynamics, Nutrient Mining, and Monetary Advantage of Maize-Legume Intercropping in the Eastern Himalayan Region of India. *Plant Production Science*, 17(4), 342–352. <https://doi.org/10.1626/pps.17.342>
65. Chowdhury, M. K., & Rosario, E. L. (1992). Utilization efficiency of applied nitrogen as related to yield advantage in maize/mungbean intercropping. *Field Crops Research*, 30(1–2), 41–51. [https://doi.org/10.1016/0378-4290\(92\)90055-E](https://doi.org/10.1016/0378-4290(92)90055-E)
66. Ciftci, V., . N. T., . Y. T., & . Y. D. (2005). The Effects of Intercropping Sowing Systems with Dry Bean and Maize on Yield and Some Yield Components. *Journal of Agronomy*, 5(1), 53–56. <https://doi.org/10.3923/ja.2006.53.56>
67. Costa, A. P. D., Bezerra Neto, F., Silva, M. L. D., Lima, J. S. S. D., Barros Júnior, A. P., & Porto, V. C. N. (2017). Intercropping of Carrot X Cowpea-Vegetables: Evaluation of Cultivar Combinations Fertilized with Roostertree. *Revista Caatinga*, 30(3), 633–641. <https://doi.org/10.1590/1983-21252017v30n311rc>
68. Coutinho, P. W. R., Oliveira, P. S. R. D., Echer, M. D. M., Cadorin, D. A., & Vanelli, J. (2017). Establishment of intercropping of beet and chicory depending on soil management. *Revista Ciência Agronômica*, 48(4). <https://doi.org/10.5935/1806-6690.20170078>
69. Cuartero, J., Pascual, J. A., Vivo, J.-M., Özbolat, O., Sánchez-Navarro, V., Egea-Cortines, M., Zornoza, R., Mena, M. M., Garcia, E., & Ros, M. (2022). A first-year melon/cowpea intercropping system improves soil nutrients and changes the soil microbial community. *Agriculture, Ecosystems & Environment*, 328, 107856. <https://doi.org/10.1016/j.agee.2022.107856>
70. De Costa, W. A. J. M., & Perera, M. K. K. W. (1998). Effects of Bean Population and Row Arrangement on the Productivity of Chilli/Dwarf Bean (*Capsicum annuum/Phaseolus vulgaris* L.) Intercropping in Sri Lanka. *Journal of Agronomy and Crop Science*, 180(1), 53–58. <https://doi.org/10.1111/j.1439-037X.1998.tb00369.x>
71. De Lima, J. S. S., Bezerra Neto, F., Gomes, E. G., De Negreiros, M. Z., Pontes, F. S. T., De Medeiros, M. A., & Barros Júnior, A. P. (2014). Agroeconomic evaluation of intercropping rocket and carrot by uni- and multivariate analyses in a semi-arid region of Brazil. *Ecological Indicators*, 41, 109–114. <https://doi.org/10.1016/j.ecolind.2014.01.031>
72. Deemak, C., Lerslerwong, L., Nampila, S., Meetha, S., & Isarangkool Na Ayutthaya, S. (2023). Evaluation of Intercropped Papaya with Banana on Growth, Fruit, and Nutrient Status. *The Horticulture Journal*, 92(2), 142–150. <https://doi.org/10.2503/hortj.QH-011>

73. Demir, H., & Polat, E. (2011). Effects of broccoli-crispy salad intercropping on yield and quality under greenhouse conditions. *African Journal of Agricultural Research*, 6(17), 4116–4121.
74. Devides, A. C. P., Ribeiro, R. D. L. D., Valle, T. L., Almeida, D. L. D., Castro, C. M. D., & Feltran, J. C. (2009). Produtividade de raízes de mandioca consorciada com milho e caupi em sistema orgânico. *Bragantia*, 68(1), 145–153. <https://doi.org/10.1590/S0006-87052009000100016>
75. Dolijanović, Ž., Momirović, N., Oljača, S., Simić, M., Oljača, M., & Janošević, B. (2015). Productivity of Intercropping Maize (zea Mays L.) and Pumpkins (cucurbita Maxima Duch.) Under Conventional Vs. Conservation Farming System. *Turkish Journal Of Field Crops*, 20(1). <https://doi.org/10.17557/1.75885>
76. Dua, V. K., Kumar, S., & Jatav, M. K. (2017). Effect of nitrogen application to intercrops on yield, competition, nutrient use efficiency and economics in potato (Solanum Tuberosum L.) + French bean (Phaseolus Vulgaris L.) system in north-western hills of India. *Legume Research*, 40(4), 698–703. <https://doi.org/10.18805/lr.v0i0.7841>
77. Fathi, S. A. A. (2023). Eggplant-garlic intercrops reduce the density of Tetranychus urticae on eggplant and improve crop yield. *Experimental and Applied Acarology*, 91(1), 43–55. <https://doi.org/10.1007/s10493-023-00821-3>
78. Favacho, F. S., Lima, J. S. S. D., Bezerra Neto, F., Silva, J. N. D., & Barros Júnior, A. P. (2017). Productive and economic efficiency of carrot intercropped with cowpeavegetable resulting from green manure and different spatial arrangements. *Revista Ciência Agronômica*, 48(2). <https://doi.org/10.5935/1806-6690.20170039>
79. Fayyaz, F. A., Anwar, A., & Yousaf, G. (2024). On-Farm Crop Productivity and Economic Assessment of Mungbean-Sorghum Intercropping System Under Integrated Nutrient Management. *Journal of Agricultural Sciences – Sri Lanka*, 19(2), 233–249. <https://doi.org/10.4038/jas.v19i2.9988>
80. Fonseca, M. C. M., Sediya, M. A. N., Bonfim, F. P. G., Dores, R. G. R. D., Gonçalves, M. G., Prado, A. L. D., & Lopes, I. P. D. C. (2016). Lettuce and marigold intercropping: Crops productivity and marigold's flavonoid content. *Ciência Rural*, 46(9), 1553–1558. <https://doi.org/10.1590/0103-8478cr20150712>
81. Fu, Z., Chen, P., Zhang, X., Du, Q., Zheng, B., Yang, H., Luo, K., Lin, P., Li, Y., Pu, T., Wu, Y., Wang, X., Yang, F., Liu, W., Song, C., Yang, W., & Yong, T. (2023). Maize-legume intercropping achieves yield advantages by improving leaf functions and dry matter partition. *BMC Plant Biology*, 23(1), 438. <https://doi.org/10.1186/s12870-023-04408-3>

82. Furtado, G. D. F., Souza, A. D. S., Sousa Júnior, J. R. D., Sousa, J. R. M. D., Lacerda, R. R. D. A., & Silva, S. S. D. (2014). Rendimento e correlações da mamoneira consorciada com feijão-caupi e gergelim no semiárido paraibano. *Revista Brasileira de Engenharia Agrícola e Ambiental*, 18(9), 892–898. <https://doi.org/10.1590/1807-1929/agriambi.v18n09p892-898>
83. Gao, H., Meng, W., Zhang, C., Van Der Werf, W., Zhang, Z., Wan, S., & Zhang, F. (2020). Yield and nitrogen uptake of sole and intercropped maize and peanut in response to N fertilizer input. *Food and Energy Security*, 9(1), e187. <https://doi.org/10.1002/fes3.187>
84. Gao, J., & Xie, H. (2023). Daylily intercropping: Effects on soil nutrients, enzyme activities, and microbial community structure. *Frontiers in Plant Science*, 14, 1107690. <https://doi.org/10.3389/fpls.2023.1107690>
85. Gao, L., Xu, H., Bi, H., Xi, W., Bao, B., Wang, X., Bi, C., & Chang, Y. (2013). Intercropping Competition between Apple Trees and Crops in Agroforestry Systems on the Loess Plateau of China. *PLoS ONE*, 8(7), e70739. <https://doi.org/10.1371/journal.pone.0070739>
86. Gawade, M. H., Patil, J. D., & Kakade, D. S. (2003). Studies on Effect of Intercrops on Yield and Monetary Returns of Cauliflower. *Agricultural Science Digest*, 23(1), 73–74.
87. Gebru, H. (2015). Evaluation of Tomato (*Lycopersicon esculentum* Mill) and Maize (*Zea mays* L.) Intercropping System for Profitability of the Crops in Wolaita Zone, Southern Ethiopia. *Journal of Biology*.
88. Ghanbari, A., Dahmardeh, M., Siahsar, B. A., & Ramroudi, M. (2010). Effect of maize (*Zea mays* L.)—Cowpea (*Vigna unguiculata* L.) intercropping on light distribution, soil temperature and soil moisture in arid environment. *Journal of Food, Agriculture & Environment*, 8(1), 102–108.
89. Gitari, H. I., Nyawade, S. O., Kamau, S., Karanja, N. N., Gachene, C. K. K., Raza, M. A., Maitra, S., & Schulte-Geldermann, E. (2020). Revisiting intercropping indices with respect to potato-legume intercropping systems. *Field Crops Research*, 258, 107957. <https://doi.org/10.1016/j.fcr.2020.107957>
90. Grangeiro, L. C., Santos, A. P., Freitas, F. C. L., Simão, L. M. C., & Bezerra Neto, F. (2011). Avaliação agroeconômica das culturas da beterraba e coentro em função da época de estabelecimento do consórcio. *Revista Ciência Agronômica*, 42(1), 242–248. <https://doi.org/10.1590/S1806-66902011000100030>

91. Grewal, S. S., Mittal, S. P., Dyal, S., & Agnihotri, Y. (1992). Agroforestry systems for soil and water conservation and sustainable production from foothill areas of north India. *Agroforestry Systems*, 17(3), 183–191. <https://doi.org/10.1007/BF00054146>
92. Guimarães, M. D. A., Lemos Neto, H. D. S., Lima Neto, B. P., Hendges, A. R. A., Viana, C. D. S., Rabelo, J. D. S., & Oliveira, F. R. A. D. (2022). Respostas fisiológicas da rúcula consorciada com nirá sob diferentes arranjos e épocas de cultivo. *Horticultura Brasileira*, 40(3), 255–260. <https://doi.org/10.1590/s0102-0536-20220302>
93. Guldán, S. J., Martin, C. A., & Falk, C. L. (1998). Interseeding Snap Pea into Stands of Chile Pepper Reduces Yield of Pea More Than That of Chile. *HortScience*, 33(4), 660–662. <https://doi.org/10.21273/HORTSCI.33.4.660>
94. Güvenç, I., & Yıldırım, E. (2005). Intercropping with eggplant for proper utilisation of interspace under greenhouse conditions. *European Journal of Horticultural Science*, 300–302. <https://doi.org/10.1079/ejhs.2005/36481>
95. Guvenc, I., & Yildirim, E. (2006). Increasing Productivity with Intercropping Systems in Cabbage Production. *Journal of Sustainable Agriculture*, 28(4), 29–44. https://doi.org/10.1300/J064v28n04_04
96. Hadidi, N., Sharaiha, R., & Debei, H. A.-. (2011). Effect of Intercropping on the Performance of Some Summer Vegetable Crops Grown Under Different Row Arrangements. *Lucrări Științifice*, 54(2).
97. Haile, M. A., Karanja, N. N., Nyawade, S. O., Gitari, H. I., Cheruto, G., Nyawira, L., Raza, M. A., & Kamau, S. (2024). Lupin and Lima Beans Diminish Potatoes' N and P Uptake, Uptake Efficiency and Use Efficiency. *Potato Research*, 67(1), 37–53. <https://doi.org/10.1007/s11540-023-09625-9>
98. Hamdani, J. S., & Suradinata, Y. R. (2015). Effects of Row Intercropping System of Corn and Potato and Row Spacing of Corn on the Growth and Yields of Atlantic Potato Cultivar Planted in Medium Altitude. *Asian Journal of Agricultural Research*, 9(3), 104–112. <https://doi.org/10.3923/ajar.2015.104.112>
99. Hammad, S. A., Bahnasy, M. I., Alzamel, N. M., Hussein, M. F. A., Mahmoud, A. A. A., & Loutfy, N. (2024). Agro-Ecological Practice for Sustaining Higher Productivity of Fennel Plant Using Alley Cropping System and Endophytic Fungi. *Sustainability*, 16(12), 5167. <https://doi.org/10.3390/su16125167>
100. Haque, M. E., . M. R., . M. A. R., . A. K. R., & . B. S. (2003). Lablab Bean Based Intercropping System in Northwest Region of Bangladesh. *Pakistan Journal of Biological Sciences*, 6(10), 948–951. <https://doi.org/10.3923/pjbs.2003.948.951>

101. Härdter, R., & Horst, W. J. (1991). Nitrogen and phosphorus use in maize sole cropping and maize/cowpea mixed cropping systems on an Alfisol in the northern Guinea Savanna of Ghana. *Biology and Fertility of Soils*, 10(4), 267–275. <https://doi.org/10.1007/BF00337377>
102. Hendges, A. R. A. D. A., Guimarães, M. D. A., Silva, J. C. D., & Lima Neto, B. P. (2019). Agronomic Performance and Biological Efficiency of Kale Intercropped with Spice Species. *Revista Caatinga*, 32(1), 7–15. <https://doi.org/10.1590/1983-21252019v32n102rc>
103. Hu, F., Gan, Y., Chai, Q., Feng, F., Zhao, C., Yu, A., Mu, Y., & Zhang, Y. (2016). Boosting system productivity through the improved coordination of interspecific competition in maize/pea strip intercropping. *Field Crops Research*, 198, 50–60. <https://doi.org/10.1016/j.fcr.2016.08.022>
104. Hu, F., Tan, Y., Yu, A., Zhao, C., Fan, Z., Yin, W., Chai, Q., Coulter, J. A., & Cao, W. (2020). Optimizing the split of N fertilizer application over time increases grain yield of maize-pea intercropping in arid areas. *European Journal of Agronomy*, 119, 126117. <https://doi.org/10.1016/j.eja.2020.126117>
105. Huang, C., Liu, Q., Li, H., Li, X., Zhang, C., & Zhang, F. (2018). Optimised sowing date enhances crop resilience towards size-asymmetric competition and reduces the yield difference between intercropped and sole maize. *Field Crops Research*, 217, 125–133. <https://doi.org/10.1016/j.fcr.2017.12.010>
106. Hugar, H. Y., & Palled, Y. B. (2008). Studies on Maize-Vegetable Intercropping Systems. *Karnataka Journal of Agricultural Sciences*, 21(2), 162–164.
107. Ilakiya, T., Swarnapriya, R., Pugalandhi, L., Geethalakshmi, V., Laxmanan, A., & Sathees Kumar, K. (2023). Land Use and Weed Control Efficiency of Intercropping in Elephant Foot Yam. *Applied Ecology and Environmental Research*, 21(5), 4005–4018. https://doi.org/10.15666/aeer/2105_40054018
108. Islam, M. N., Akhteruzzaman, M., Alom, M. S., & Salim, M. (2014). Hybrid Maize and Sweet Potato Intercropping: A Technology to Increase Productivity and Profitability for Poor Hill Farmers in Bangladesh. *Saarc Journal of Agriculture*, 12(2), 101–111.
109. Jiang, Y., Khan, M. U., Lin, X., Lin, Z., Lin, S., & Lin, W. (2022). Evaluation of maize/peanut intercropping effects on microbial assembly, root exudates and peanut nitrogen uptake. *Plant Physiology and Biochemistry*, 171, 75–83. <https://doi.org/10.1016/j.plaphy.2021.12.024>
110. Jinger, D., Kakade, V., Bhatnagar, P. R., Paramesh, V., Dinesh, D., Singh, G., N, N. K., Kaushal, R., Singhal, V., Rathore, A. C., Tomar, J. M. S., Singh, C., Yadav, L. P., Jat, R. A., Kaledhonkar, M. J., & Madhu, M. (2024). Enhancing productivity and

sustainability of ravine lands through horti-silviculture and soil moisture conservation: A pathway to land degradation neutrality. *Journal of Environmental Management*, 364, 121425. <https://doi.org/10.1016/j.jenvman.2024.121425>

111. Jørgensen, V., & Møller, E. (2000). Intercropping of Different Secondary Crops in Maize. *Acta Agriculturae Scandinavica, Section B - Soil & Plant Science*, 50(2), 82–88. <https://doi.org/10.1080/09064710050505026>
112. Kabiraj, J., Das, R., Das, S. P., & Mandal, A. R. (2017). A Study on Cauliflower (*Brassica oleracea* var. Botrytis) Based Intercropping System. *International Journal of Current Microbiology and Applied Sciences*, 6(7), 2595–2602. <https://doi.org/10.20546/ijcmas.2017.607.306>
113. Kabura, B. H., Musa, B., & Odo, P. E. (2007). Evaluation of the Yield Components and Yield of Onion (*Allium cepa* L.)-Pepper (*Capsicum annuum* L.) Intercrop in the Sudan Savanna. *Journal of Agronomy*, 7(1), 88–92. <https://doi.org/10.3923/ja.2008.88.92>
114. Kang, B. T., Caveness, F. E., Tian, G., & Kolawole, G. O. (1999). Longterm alley cropping with four hedgerow species on an Alfisol in southwestern Nigeria – effect on crop performance, soil chemical properties and nematode population. *Nutrient Cycling in Agroecosystems*, 54, 145–155.
115. Karlidag, H., & Yildirim, E. (2007). The Effects of Nitrogen Fertilization on Intercropped Strawberry and Broad Bean. *Journal of Sustainable Agriculture*, 29(4), 61–74. https://doi.org/10.1300/J064v29n04_06
116. Karlidag, H., & Yildirim, E. (2009). Strawberry Intercropping with Vegetables for Proper Utilization of Space and Resources. *Journal of Sustainable Agriculture*, 33(1), 107–116. <https://doi.org/10.1080/10440040802587462>
117. Kermah, M., Franke, A. C., Adjei-Nsiah, S., Ahiabor, B. D. K., Abaidoo, R. C., & Giller, K. E. (2017). Maize-grain legume intercropping for enhanced resource use efficiency and crop productivity in the Guinea savanna of northern Ghana. *Field Crops Research*, 213, 38–50. <https://doi.org/10.1016/j.fcr.2017.07.008>
118. Khongruang, P., Rison, K., Disyatat, N. R., & Chaisuekul, C. (2025). Flowering zinnia as companion plant increases predatory arthropod abundance and improves yard-long bean yield. *BioControl*, 70(1), 19–31. <https://doi.org/10.1007/s10526-024-10275-0>
119. Kidane, B. Z., Hailu, M. H., & Haile, H. T. (2017). Maize and Potato Intercropping: A Technology to Increase Productivity and Profitability in Tigray. *Open Agriculture*, 2(1), 411–416. <https://doi.org/10.1515/opag-2017-0044>

120. Köneke, A., & Böckmann, E. (2024). Wheat companion plants reduced aphid and flea beetle infestations in Brassica vegetable crops in a multiple-year field study in central Germany. *Crop Protection*, 182, 106732.
<https://doi.org/10.1016/j.cropro.2024.106732>
121. Köneke, A., Uesugi, R., Herz, A., Tabuchi, K., Yoshimura, H., Shimoda, T., Nagasaka, K., & Böckmann, E. (2023). Effects of wheat undersowing and sweet alyssum intercropping on aphid and flea beetle infestation in white cabbage in Germany and Japan. *Journal of Plant Diseases and Protection*, 130(3), 619–631.
<https://doi.org/10.1007/s41348-023-00730-y>
122. Kour, M., Thakur, N. P., Kumar, P., & Charak, A. S. (2016). Productivity and profitability of maize (*Zea mays*) as influenced by intercropping of rajmash (*Phaseolus vulgaris*) and nutrient management techniques under sub-alpine conditions of Jammu, India. *Legume Research*, 39(6), 970–975.
<https://doi.org/10.18805/lr.v0iOF.11042>
123. Kushwah, A. S., Rawat, G. S., Gupta, S., Patil, D., & Prajapati, N. (2017). Production and profitability assessment of clusterbean (*Cyamopsis tetragonoloba* L. Taub.) based intercropping systems under different row arrangement. *Legume Research*, 40(5), 916–919. <https://doi.org/10.18805/LR-3809>
124. Labigaline, I., Sala, F. C., Gomes, T. M., Corsini, I., & Rossi, F. (2020). Green manure, *Trichoderma asperellum* and homeopathy in cultivating the biquinho pepper. *Revista Ciência Agronômica*, 51(3). <https://doi.org/10.5935/1806-6690.20200054>
125. Latati, M., Bargaz, A., Belarbi, B., Lazali, M., Benlahrech, S., Tellah, S., Kaci, G., Drevon, J. J., & Ounane, S. M. (2016). The intercropping common bean with maize improves the rhizobial efficiency, resource use and grain yield under low phosphorus availability. *European Journal of Agronomy*, 72, 80–90.
<https://doi.org/10.1016/j.eja.2015.09.015>
126. Latati, M., Blavet, D., Alkama, N., Laoufi, H., Drevon, J. J., Gérard, F., Pansu, M., & Ounane, S. M. (2014). The intercropping cowpea-maize improves soil phosphorus availability and maize yields in an alkaline soil. *Plant and Soil*, 385(1–2), 181–191.
<https://doi.org/10.1007/s11104-014-2214-6>
127. Legodi, K. D., & Ogola, J. B. O. (2020). Cassava-legume intercrop: I. Effects of relative planting dates of legumes on cassava productivity. *Acta Agriculturae Scandinavica, Section B — Soil & Plant Science*, 70(2), 150–157.
<https://doi.org/10.1080/09064710.2019.1682185>

128. Leishangthem, K., Patel, B., Prajapati, D. D., & Pandya, H. (2016). Intercropping of Tuber Crops in Young Orchard of Sapota Cv. Kalipatti. *Journal of Tree Sciences*, 35(2), 85. <https://doi.org/10.5958/2455-7129.2016.00014.5>
129. Leite, A. A. L., Ferraz Junior, A. S. D. L., Moura, E. G. D., & Aguiar, A. D. C. F. (2008). Comportamento de dois genótipos de milho cultivados em sistema de aléias preestabelecido com diferentes leguminosas arbóreas. *Bragantia*, 67(4), 875–882. <https://doi.org/10.1590/S0006-87052008000400009>
130. Lenka, S., Swain, S. K., & Pradhan, K. C. (2023). Performance of Different Groundnut (*Arachis hypogaea* L.) based Intercropping Systems with Millets under Rainfed Condition of Odisha. *Legume Research*, 46(2), 154–159. <https://doi.org/10.18805/LR-4690>
131. Li, P., Liu, H., Chen, Y., Zhang, X., Cao, N., Sun, Y., Jia, M., Wu, M., Tong, X., Jiang, X., Yu, X., & Cheng, Y. (2023). Effect of Dandelion (*Taraxacum mongolicum* Hand.-Mazz.) Intercropping with Different Plant Spacing on Blight and Growth of Pepper (*Capsicum annuum* L.). *Phyton*, 92(8), 2227–2244. <https://doi.org/10.32604/phyton.2023.027392>
132. Liang, Q., Zhang, T., Liu, Z., Gao, W., Cheng, Y., & Feng, H. (2023). Effects of Different Green Manure Crops on Soil Water, Nitrogen, and Yield: Preliminary Results in an Apple Orchard on the Loess Plateau, China. *Agronomy*, 13(8), 2009. <https://doi.org/10.3390/agronomy13082009>
133. Lima, A. D. A., Caldas, R. C., Borges, A. L., Ritzinger, C. H. S. P., Trindade, A. V., Pires, M. D. M., Midlej, M. M. B. C., Mata, H. T. D. C., & Souza, J. D. S. (2002). Cultivos intercalares e controle de plantas daninhas em plantios de maracujá-amarelo. *Revista Brasileira de Fruticultura*, 24(3), 711–713. <https://doi.org/10.1590/S0100-29452002000300036>
134. Lima, J. S. S., Bezerra-Neto, F., Negreiro, M. Z., Cardoso-Ribeiro, M. C., & Barros-Júnior, A. P. (2010). Productive performance of carrot and rocket cultivars in strip-intercropping system and sole crops. *Agrociencia*, 44, 561–574.
135. Lima, M. E. D., Carvalho, D. F. D., Souza, A. P. D., Rocha, H. S., & Guerra, J. G. M. (2012). Desempenho do cultivo da berinjela em plantio direto submetida a diferentes lâminas de irrigação. *Revista Brasileira de Engenharia Agrícola e Ambiental*, 16(6), 604–610. <https://doi.org/10.1590/S1415-43662012000600003>
136. Liu, L., Wang, Y., Yan, X., Li, J., Jiao, N., & Hu, S. (2017). Biochar amendments increase the yield advantage of legume-based intercropping systems over monoculture. *Agriculture, Ecosystems & Environment*, 237, 16–23. <https://doi.org/10.1016/j.agee.2016.12.026>

137. Liu, Y., Huang, L., Liu, Q., Li, Z., Liu, C., Yuan, J., Liao, J., Luo, L., Yu, C., & Feng, Y. (2023). Effects of tomato-Sedum alfredii Hance intercropping on crop production and Cd remediation as affected by soil types. *Environmental Science and Pollution Research*, 31(3), 3696–3706. <https://doi.org/10.1007/s11356-023-31318-6>
138. Liu, Y., Zhang, Y., Xiao, T., Wu, Y., Li, Y., He, J., Xiang, Y., & Yao, B. (2023). Effect of Intercropping on Fruit Yield and Financial Benefits of Rosa roxburghii Tratt Orchard in Southwest China. *Agronomy*, 13(12), 2953. <https://doi.org/10.3390/agronomy13122953>
139. Lu, P., Zhao, C., Yin, W., Hu, F., Fan, Z., Yu, A., & Fan, H. (2023). Microbial Community Shifts with Soil Properties and Enzyme Activities in Inter-/Mono-Cropping Systems in Response to Tillage. *Agronomy*, 13(11), 2707. <https://doi.org/10.3390/agronomy13112707>
140. Mahant, H. D., Patil, S. J., Bhalerao, P. P., & Gaikwad, S. S. (2012). Economics and Land Equivalent Ratio of Different Intercrops in Banana (musa Paradisiaca L.) Cv. Grand Naine Under Drip Irrigation. *The Asian Journal of Horticulture*, 7(2), 330–332.
141. Marcos-Pérez, M., Sánchez-Navarro, V., Martínez-Martínez, S., Martínez-Mena, M., García, E., & Zornoza, R. (2023). Intercropping organic melon and cowpea combined with return of crop residues increases yields and soil fertility. *Agronomy for Sustainable Development*, 43(4), 53. <https://doi.org/10.1007/s13593-023-00902-y>
142. Marcos-Pérez, M., Sánchez-Navarro, V., & Zornoza, R. (2023). Intercropping systems between broccoli and fava bean can enhance overall crop production and improve soil fertility. *Scientia Horticulturae*, 312, 111834. <https://doi.org/10.1016/j.scienta.2023.111834>
143. Marouelli, W. A., Lage, D. A. D. C., Gravina, C. S., Michereff Filho, M., & Souza, R. B. D. (2013). Sprinkler and drip irrigation in the organic tomato for single crops and when intercropped with coriander. *Revista Ciência Agronômica*, 44(4), 825–833. <https://doi.org/10.1590/S1806-66902013000400020>
144. Mauro Shindoi, E.L. Avico, P.C. Sarco, & D.E. Monteros Solito. (2019). Evaluación de la intersiembra de rúcula (Eruca sativa Mill.) y lechuga (Lactuca sativa L.) en Colonia Benítez, Chaco. *Revista de Investigaciones Agropecuarias*, 45(2), 252–258.
145. Mbah, E. U. U., & Ogbodo, E. N. N. (2013). Assessment of intercropped sweet corn (Zea mays var. Saccharata) and vegetable cowpea (Vigna unguiculata (L.) Walp) using competitive indices in the derived savannah of south-eastern Nigeria. *Journal of Biology, Agriculture and Healthcare*, 3(3), 84–93.

146. Md. Hafiz All Amin, Sumaia Sumi, Warangkana Jutidamrongphan, & Kua-anan Techato. (2022). Economic potentiality of *Colocasia esculenta* L. under multipurpose tree-based agroforestry systems. *Asian Journal of Agriculture and Biology*, 3. <https://doi.org/10.35495/ajab.2021.01.059>
147. Mehr-un-Nisa, Shafiq, F., Anwar, S., Mahmood, A., Iqbal, M., Ullah, K., Zulqarnain, M., Haider, I., Ashraf, M., & Zhang, L. (2023). Physiological effects of some engineered nanomaterials on radish (*Raphanus sativus* L.) intercropped with pea (*Pisum sativum* L.). *Environmental Science and Pollution Research*, 30(32), 78353–78366. <https://doi.org/10.1007/s11356-023-27400-8>
148. Menezes, C. S. L., Rezende, R., Terassi, D. D. S., Hachmann, T. L., & Saath, R. (2022). Lettuce and Radish Grown in Single Crop and Intercropping Systems Under Different Irrigation Water Depths in a Protected Environment. *Revista Caatinga*, 35(3), 658–666. <https://doi.org/10.1590/1983-21252022v35n317rc>
149. Miah, Md. G., Islam, M. M., Rahman, Md. A., Ahamed, T., Islam, Md. R., & Jose, S. (2018). Transformation of jackfruit (*Artocarpus heterophyllus* Lam.) orchard into multistory agroforestry increases system productivity. *Agroforestry Systems*, 92(6), 1687–1697. <https://doi.org/10.1007/s10457-017-0118-1>
150. Midega, C. A. O., Salifu, D., Bruce, T. J., Pittchar, J., Pickett, J. A., & Khan, Z. R. (2014). Cumulative effects and economic benefits of intercropping maize with food legumes on *Striga hermonthica* infestation. *Field Crops Research*, 155, 144–152. <https://doi.org/10.1016/j.fcr.2013.09.012>
151. Mndzebele, B., Ncube, B., Fessehazion, M., Mabhaudhi, T., & Modi, A. T. (2023). Assessment of the effects of NPK fertilizer on edible yield and agro-biological parameters in a cowpea–amaranth intercrop. *Frontiers in Sustainable Food Systems*, 7, 1210984. <https://doi.org/10.3389/fsufs.2023.1210984>
152. Molina-Anzures, M. F., Chávez-Servia, J. L., Gil-Muñoz, A., López, P. A., Hernández-Romero, E., & Ortiz-Torres, E. (2016). Productive efficiencies in corn, bean and squash (*Cucurbita pepo* L.) associations, intercropped with rows of fruit trees. *Phyton*, 85(1), 36–50. <https://doi.org/10.32604/phyton.2016.85.036>
153. Mondal, M., Begum, F., & Alam, M. (2012). Study on Intercropping Carrot with Groundnut Under Different Row Arrangements. *Bangladesh Journal of Agricultural Research*, 37(2), 215–223. <https://doi.org/10.3329/bjar.v37i2.11223>
154. Morais, H., Caramori, P. H., Ribeiro, A. M. D. A., Gomes, J. C., & Koguishi, M. S. (2006). Microclimatic characterization and productivity of coffee plants grown under shade of pigeon pea in Southern Brazil. *Pesquisa Agropecuária Brasileira*, 41(5), 763–770. <https://doi.org/10.1590/S0100-204X2006000500007>

155. Morgado, L. B., & Willey, R. W. (2003). Effects of plant population and nitrogen fertilizer on yield and efficiency of maize-bean intercropping. *Pesquisa Agropecuária Brasileira*, 38(11), 1257–1264. <https://doi.org/10.1590/S0100-204X2003001100002>
156. Morgado, L. B., & Willey, R. W. (2008). Optimum plant population for maize-bean intercropping system in the Brazilian semi-arid region. *Scientia Agricola*, 65(5), 474–480. <https://doi.org/10.1590/S0103-90162008000500005>
157. Morkeliūnė, A., Rasiukevičiūtė, N., Dėnė, L., Dambrauskienė, E., Duchovskienė, L., & Valiuškaitė, A. (2024). Effect of White Cabbage Intercropping with Aromatic Plant on Yield, Mineral and Biochemical Composition. *Plants*, 13(13), 1870. <https://doi.org/10.3390/plants13131870>
158. Motagally, F. M. F. A., & Metwally, A. K. (2014). Maximizing Productivity by Intercropping Onion on Sugar Beet. *Asian Journal of Crop Science*, 6(3), 226–235. <https://doi.org/10.3923/ajcs.2014.226.235>
159. Mulugeta, T., Mulatu, B., Tekie, H., Yesuf, M., Andreasson, E., & Alexandersson, E. (2019). Phosphite alters the behavioral response of potato tuber moth (*Phthorimaea operculella*) to field-grown potato. *Pest Management Science*, 75(3), 616–621. <https://doi.org/10.1002/ps.5152>
160. Mutiga, S. K., S. Gohole, L., & O. Auma, Elmada. (2011). Agronomic Performance of Collards under Two Intercrops and Varying Nitrogen Application Levels as Assessed Using Land Equivalent Ratios. *Journal of Agricultural Science*, 3(1), p22. <https://doi.org/10.5539/jas.v3n1p22>
161. Ndengu, G., Mponela, P., Chataika, B., Desta, L. T., Chirwa, R., & Sileshi, G. G. (2022). Effect of combining organic manure and inorganic fertilisers on maize–bush bean intercropping. *Experimental Agriculture*, 58, e29. <https://doi.org/10.1017/S0014479722000102>
162. Newman, S. M. (1986). A Pear and Vegetable Interculture System: Land Equivalent Ratio, Light Use Efficiency and Productivity. *Experimental Agriculture*, 22(4), 383–392. <https://doi.org/10.1017/S0014479700014630>
163. Ngwenya, Z. D., Mohammed, M., & Dakora, F. D. (2024). Monocropping and Intercropping of Maize with Six Food Legumes at Malkerns in Eswatini: Their Effects on Plant Growth, Grain Yield and N₂ Fixation, Measured using the ¹⁵N Natural Abundance and Ureide Techniques. *Symbiosis*, 92(2), 257–269. <https://doi.org/10.1007/s13199-024-00971-x>
164. Ngwira, A. R., Aune, J. B., & Mkwinda, S. (2012). On-farm evaluation of yield and economic benefit of short term maize legume intercropping systems under

conservation agriculture in Malawi. *Field Crops Research*, 132, 149–157.
<https://doi.org/10.1016/j.fcr.2011.12.014>

165. Nissen, T. M., Midmore, D. J., & Cabrera, M. L. (1991). Aboveground and belowground competition between intercropped cabbage and young *Eucalyptus torelliana*. *Agricultural Systems*, 46, 83–93.
166. Njira, K. O. W., Semu, E., Mrema, J. P., & Nalivata, P. C. (2021). Productivity of pigeon pea, cowpea and maize under sole cropping, legume–legume and legume–cereal intercrops on Alfisols in Central Malawi. *Agroforestry Systems*, 95(2), 279–291. <https://doi.org/10.1007/s10457-020-00589-0>
167. Nkongho, R. N., Ibeku, N. E., Ndam, L. M., Agbor, D. T., & Yinda, G. S. (2024). A comparative analysis of the performance of common cropping systems used in the production of maize and selected grain legumes in Cameroon. *Scientific African*, 24, e02131. <https://doi.org/10.1016/j.sciaf.2024.e02131>
168. Noda-Leyva, Y., Martín-Martín, G. J., Martín-Alonso, G. M., & Perez Vazquez, A. (2024). Effect of Different Spatial Arrangements of *Jatropha Curcas* L. on Food Crops. *Agrociencia*. <https://doi.org/10.47163/agrociencia.v58i2.3019>
169. Nthabiseng, T. R., Irvine, K. M., & Moshibudi, P. M. (2015). Response of a maize or dry bean intercrop to maize density and dry bean arrangement under rainfed conditions. *International Journal of Agronomy and Agricultural Research*, 6(6), 18–29.
170. Nwofia, G. E., Jiwuba, F., Okpara, D. A., & Mbah, E. U. (2017). Yield and Productivity of Eggplant Genotypes Intercropped with Vegetable Cowpea in the Humid Tropics. *International Journal of Vegetable Science*, 23(5), 400–410.
<https://doi.org/10.1080/19315260.2017.1310779>
171. Nyawade, S. O., Gitari, H. I., Karanja, N. N., Gachene, C. K. K., Schulte-Geldermann, E., & Parker, M. L. (2021). Yield and evapotranspiration characteristics of potato-legume intercropping simulated using a dual coefficient approach in a tropical highland. *Field Crops Research*, 274, 108327.
<https://doi.org/10.1016/j.fcr.2021.108327>
172. Obiagwu, C. J. (1995). Estimated yield and nutrient contributions of legume cover crops intercropped with yam, cassava, and maize in the Benue River basins of Nigeria. *Journal of Plant Nutrition*, 18(12), 2775–2782.
<https://doi.org/10.1080/01904169509365099>
173. Ocimati, W., Ntamwira, J., Groot, J. C. J., Taulya, G., Tittonell, P., Dhed'a, B., Van Asten, P., Vanlauwe, B., Ruhigwa, B., & Blomme, G. (2019). Banana leaf pruning to facilitate annual legume intercropping as an intensification strategy in the East

African highlands. *European Journal of Agronomy*, 110, 125923.
<https://doi.org/10.1016/j.eja.2019.125923>

174. Octavia, D., Wijayanto, N., Budi, S. W., Batubara, I., & Suharti, S. (2023). The Potential of Cardamom Leaf in the Agroforestry System: Essential Oil Yield and 1.8-Cineol Content. *Indonesian Journal of Forestry Research*, 11(1), 17–32.
<https://doi.org/10.59465/ijfr.2024.11.1.17-32>
175. Ofori, K., & Gamedoagbao, D. K. (2005). Yield of scarlet eggplant (*Solanum aethiopicum* L.) as influenced by planting date of companion cowpea. *Scientia Horticulturae*, 105(3), 305–312. <https://doi.org/10.1016/j.scienta.2005.02.003>
176. Ogbuehi, C. R. A., & Orzolek, M. D. (1987). Intercropping carrot and sweetcorn in a multiple cropping system. *Scientia Horticulturae*, 31(1–2), 17–24.
[https://doi.org/10.1016/0304-4238\(87\)90102-6](https://doi.org/10.1016/0304-4238(87)90102-6)
177. Ohse, S., Alves Rezende, B. L., Sleutjes Silveira, L., Fernandes Otto, R., & Gonçalves Cortez, M. (2012). Viabilidade agrônômica de consórcios de brócolis e alface estabelecidos em diferentes épocas. *Idesia (Arica)*, 30(2), 29–37.
<https://doi.org/10.4067/S0718-34292012000200004>
178. Olasantan, F. O. (1991). Response of tomato and okra to nitrogen fertilizer in sole cropping and intercropping with cowpea. *Journal of Horticultural Science*, 66(2), 191–199. <https://doi.org/10.1080/00221589.1991.11516144>
179. Olasantan, F. O. (1998). Effects of preceding maize (*Zea mays*) and cowpea (*Vigna unguiculata*) in sole cropping and intercropping on growth, yield and nitrogen requirement of okra (*Abelmoschus esculentus*). *The Journal of Agricultural Science*, 131(3), 293–298. <https://doi.org/10.1017/S0021859698005711>
180. Oliveira, F. L. D., Ribas, R. G. T., Junqueira, R. M., Padovan, M. P., Guerra, J. G. M., Almeida, D. L. D., & Ribeiro, R. D. L. D. (2005). Desempenho do consórcio entre repolho e rabanete com pré-cultivo de crotalária, sob manejo orgânico. *Horticultura Brasileira*, 23(2), 184–188. <https://doi.org/10.1590/S0102-05362005000200004>
181. Osonubi, O., Atayese, M. O., & Mulongoy, K. (1995). The effect of vesicular-arbuscular mycorrhizal inoculation on nutrient uptake and yield of alley-cropped cassava in a degraded Alfisol of southwestern Nigeria. *Biology and Fertility of Soils*, 20(1), 70–76. <https://doi.org/10.1007/BF00307844>
182. Ouma, G., & Jeruto, P. (2010). Sustainable horticultural crop production through intercropping: The case of fruits and vegetable crops: A review. *Agriculture and Biology Journal of North America*, 1(5), 1098–1105.
<https://doi.org/10.5251/abjna.2010.1.5.1098.1105>

183. Panozzo, A., Bernazeau, B., & Desclaux, D. (2020). Durum wheat in organic olive orchard: Good deal for the farmers? *Agroforestry Systems*, 94(3), 707–717. <https://doi.org/10.1007/s10457-019-00441-0>
184. Paoletti, A., Benincasa, P., Famiani, F., & Rosati, A. (2023). Spear yield and quality of wild asparagus (*Asparagus acutifolius* L.) as an understory crop in two olive systems. *Agroforestry Systems*, 97(8), 1361–1373. <https://doi.org/10.1007/s10457-023-00860-0>
185. Paslawar, A. N., Mhashakhetri, S., Shingrup, P. V., Chirde, P. N., & Darekar, N. K. (2023). Productivity Dynamics of Blackgram with Intercropping of Minor Millets on BBF under Organic Cultivation. *Legume Research*. <https://doi.org/10.18805/LR-5031>
186. Paula, P. D. D., Guerra, J. G. M., Ribeiro, R. D. L., Cesar, M. N. Z., Guedes, R. E., & Polidoro, J. C. (2009). Viabilidade agrônômica de consórcios entre cebola e alface no sistema orgânico de produção. *Horticultura Brasileira*, 27(2), 202–206. <https://doi.org/10.1590/S0102-05362009000200014>
187. Paulo, E. M., Berton, R. S., Cavichioli, J. C., Bulisani, E. A., & Kasai, F. S. (2006). Produtividade do cafeeiro Mundo Novo enxertado e submetido à adubação verde antes e após recepa da lavoura. *Bragantia*, 65(1), 115–120. <https://doi.org/10.1590/S0006-87052006000100015>
188. Paz, L. B. D., Gallo, A. D. S., Souza, R. D. L., Oliveira, L. V. N. D., Cunha, C. D., & Silva, R. F. D. (2019). Desempenho e produtividade do milho safrinha em consórcio com leguminosas em sistema orgânico. *Revista de Ciências Agrárias*, 788-794 Páginas. <https://doi.org/10.19084/RCA16240>
189. Perrin, B., Leroy, C., Parès, L., Pradere, P., Goude, M., Salvador, B., Marrec, T., Comes, L., Huot-Marchand, R., Guillot, E., & Lefèvre, A. (2023). Experimental dataset of the impact assessment of vegetable intercropping on agroeconomic performances, pests and beneficials, and soil resources. *Data in Brief*, 50, 109607. <https://doi.org/10.1016/j.dib.2023.109607>
190. Phiri, A., Njira, K., & Dixon, A. (2024). Comparative effects of legume-based intercropping systems involving pigeon pea and cowpea under deep-bed and conventional tillage systems in Malawi. *Agrosystems, Geosciences & Environment*, 7(2), e20503. <https://doi.org/10.1002/agg2.20503>
191. Polthanee, A., & Trelo-ges, V. (2003). Growth, Yield and Land Use Efficiency of Corn and Legumes Grown under Intercropping Systems. *Plant Production Science*, 6(2), 139–146. <https://doi.org/10.1626/pps.6.139>

192. Powers, L. E., McSorley, R., Dunn, R. A., & Montes, A. (1994). The agroecology of a cucurbit-based intercropping system in the Yeguaré Valley of Honduras. *Agriculture, Ecosystems & Environment*, 48(2), 139–147. [https://doi.org/10.1016/0167-8809\(94\)90084-1](https://doi.org/10.1016/0167-8809(94)90084-1)
193. Puiatti, M., Fávero, C., Finger, F. L., & Gomes, J. M. (2000). Crescimento e produtividade de inhame e de milho doce em cultivo associado. *Horticultura Brasileira*, 18(1), 24–30. <https://doi.org/10.1590/S0102-05362000000100006>
194. Qasim, S. A., Anjum, M. A., Hussain, S., & Ahmad, S. (2013). Effect of Pea Intercropping on Biological Efficiencies and Economics of Some Non-Legume Winter Vegetables. *Pakistan Journal of Agricultural Sciences*, 50(3), 399–406.
195. Qiang, X., Sun, Z., Li, X., Li, S., Yu, Z., He, J., Li, Q., Han, L., & He, L. (2024). The Impacts of Planting Patterns Combined with Irrigation Management Practices on Watermelon Growth, Photosynthesis, and Yield. *Plants*, 13(10), 1402. <https://doi.org/10.3390/plants13101402>
196. Quiroz, A. I., & Marín, D. (2003). Rendimiento En Granos Y Eficiencia De Una Asociación Maíz (zea Mays) Y Quinchoncho con o sin fertilización. *Bioagro*, 15(2), 121–128.
197. Rahman, M. M., Rahman, M. A., Ali, M. A., Gaber, A., Skalicky, M., & Hossain, A. (2024). The prospects of intercropping of brinjal (*Solanum melongena* L.) with other vegetables on sorjan system in vulnerable southern coastal saline and non-saline areas of Bangladesh. *Journal of Agriculture and Food Research*, 16, 101192. <https://doi.org/10.1016/j.jafr.2024.101192>
198. Rajput, R. L., & Kushwaha, B. B. (2020). Yield Analysis of Chickpea (*Cicer arietinum*) with Mustard (*Brassica juncea*) Intercropping System as Influenced by Weed Management Practices. *Legume Research*. <https://doi.org/10.18805/LR-3842>
199. Rajput, R. L., & Rawat, G. S. (2019). Yield and Yield Attributes as Influenced by Nutrient Management Practices in Pigeonpea and Clusterbean Intercropping System. *Legume Research*, 48(6), 834–837. <https://doi.org/10.18805/LR-3841>
200. Ranjan, R., Sharma, N. K., Kumar, A., Pramanik, M., Mehta, H., Ojasvi, P. R., & Yadav, R. S. (2021). Evaluation of maize-based intercropping on runoff, soil loss, and yield in foothills of the Indian sub-Himalayas. *Experimental Agriculture*, 57(2), 69–84. <https://doi.org/10.1017/S0014479721000053>
201. Rao, M. R., & Singh, M. (1990). Productivity and risk evaluation in contrasting intercropping systems. *Field Crops Research*, 23(3–4), 279–293. [https://doi.org/10.1016/0378-4290\(90\)90060-O](https://doi.org/10.1016/0378-4290(90)90060-O)

202. Rapholo, E., Odhiambo, J. J. O., Nelson, W. C. D., Rötter, R. P., Ayisi, K., Koch, M., & Hoffmann, M. P. (2020). Maize–lablab intercropping is promising in supporting the sustainable intensification of smallholder cropping systems under high climate risk in southern Africa. *Experimental Agriculture*, 56(1), 104–117. <https://doi.org/10.1017/S0014479719000206>
203. Resende, A. L. S., Viana, A. J. D. S., Oliveira, R. J., Aguiar-Menezes, E. D. L., Ribeiro, R. D. L., Ricci, M. D. S., & Guerra, J. G. M. (2010). Consórcio couve-coentro em cultivo orgânico e sua influência nas populações de joaninhas. *Horticultura Brasileira*, 28(1), 41–46. <https://doi.org/10.1590/S0102-05362010000100008>
204. Reynafarje, X., Siura, S., & Pérez, K. (2016). Mixed cropping of vegetables to improve organic tomato (*Solanum lycopersicum* L.) production in small farmer systems. *Acta Horticulturae*, 1128, 299–304. <https://doi.org/10.17660/ActaHortic.2016.1128.45>
205. Rezaei-Chiyaneh, E., Amani Machiani, M., Javanmard, A., Mahdavia, H., Maggi, F., & Morshedloo, M. R. (2021). Vermicompost Application in Different Intercropping Patterns Improves the Mineral Nutrient Uptake and Essential Oil Compositions of Sweet Basil (*Ocimum basilicum* L.). *Journal of Soil Science and Plant Nutrition*, 21(1), 450–466. <https://doi.org/10.1007/s42729-020-00373-0>
206. Rezapour Kavishahi, T., Sayfzadeh, S., Mostafavi Rad, M., Valadabady, A. R., & Hadidi Masouleh, E. (2023). Quantitative and Qualitative Yield of Groundnut (*Arachis Hypogaea* L.) in Response to Intercropping Pattern and Integrated Application of Chemical and Biological Phosphorus Fertilizers. *Communications in Soil Science and Plant Analysis*, 54(14), 1955–1968. <https://doi.org/10.1080/00103624.2023.2211109>
207. Rezende, B. L. A., Canato, G. H. D., & Cecílio Filho, A. B. (2005). Influência das épocas de cultivo e do estabelecimento do consórcio na produção de tomate e alface consorciados. *Ciência e Agrotecnologia*, 29(1), 77–83. <https://doi.org/10.1590/S1413-70542005000100009>
208. Rezende, G. D. S. P., & Ramalho, M. A. P. (1994). Competitive ability of maize and common bean (*Phaseolus vulgaris*) cultivars intercropped in different environments. *The Journal of Agricultural Science*, 123(2), 185–190. <https://doi.org/10.1017/S0021859600068441>
209. Ribas, R. G. T., Cecílio Filho, A. B., Dutra, A. F., Barbosa, J. C., & Rolim, G. D. S. (2020). Land Equivalent Ratio in the Intercropping of Cucumber with Lettuce as a Function of Cucumber Population Density. *Agriculture*, 10(3), 88. <https://doi.org/10.3390/agriculture10030088>

210. Ribeiro, G. M., Bezerra Neto, F., Lima, J. S. S. D., Silva, M. L. D., Barros Júnior, A. P., & Santos, E. C. D. (2017). Agro-Economic Efficiency of the Intercropping of Carrot X Cowpea-Vegetable Under Different Spatial Arrangements and Population Densities. *Revista Caatinga*, 30(4), 847–854. <https://doi.org/10.1590/1983-21252017v30n404rc>
211. Ribera, L. M., Cecilio Filho, A. B., Peres, N. D., Santana, D. C., & Silva, M. L. D. (2023). Lettuce and arugula production in intercropping and organic fertilization. *Revista Caatinga*, 36(4), 794–801. <https://doi.org/10.1590/1983-21252023v36n407rc>
212. Rodrigo, V. H. L., Stirling, C. M., Teklehaimanot, Z., & Nugawela, A. (1997). The effect of planting density on growth and development of component crops in rubber/banana intercropping systems. *Field Crops Research*, 52(1–2), 95–108. [https://doi.org/10.1016/S0378-4290\(96\)01069-6](https://doi.org/10.1016/S0378-4290(96)01069-6)
213. Roltsch, W. J., & Gage, S. H. (1990). Influence of Bean–Tomato Intercropping on Population Dynamics of the Potato Leafhopper (Homoptera: Cicadellidae). *Environmental Entomology*, 19(3), 534–543. <https://doi.org/10.1093/ee/19.3.534>
214. Romero Fernández, A. D. J., González-Chávez, Ma. D. C. Á., Herrera Cabrera, B. E., Corona Sánchez, J. E., & Carrillo González, R. (2023). The coffee-mango association promotes favorable soil conditions for better-nourished and higher-yielding plants. *Agriculture, Ecosystems & Environment*, 354, 108589. <https://doi.org/10.1016/j.agee.2023.108589>
215. Ronner, E., Thuijsman, E., Ebanyat, P., Descheemaeker, K., & Giller, K. E. (2021). Intercropping of climbing bean (*Phaseolus vulgaris* , L.) and East African highland banana (*Musa spp.*) in the Ugandan highlands. *Experimental Agriculture*, 57(1), 1–14. <https://doi.org/10.1017/S0014479720000411>
216. Saidi, M., & Itulya, F. M. (2006). Effect of intercropping collard with beans or onions on aphid populations and yields of collard under high altitude conditions in Kenya. *Tanzania Journal of Agricultural Sciences*, 7(2), 57–66.
217. Sairam, M., G, S. P., Maitra, S., & Shankar, T. (2023). Effect of the summer pearl millet-groundnut intercropping system on the growth, productivity and competitive ability of crops under south Odisha conditions. *Plant Science Today*. <https://doi.org/10.14719/pst.2627>
218. Salgado, G. C., Ambrosano, E. J., Rossi, F., Otsuk, I. P., Trivelin, P. C. O., Muraoka, T., Ambrosano, G. M. B., Dias, F. L. F., Tavares, S., & Melo, P. C. T. D. (2020). Nitrogen transfer from green manure to organic cherry tomato in a greenhouse intercropping system. *Journal of Plant Nutrition*, 43(8), 1119–1135. <https://doi.org/10.1080/01904167.2020.1724304>

219. Sampaio, L. R., Araújo, J. R., Sousa, E. H., Ferraz Júnior, A. S., & Araujo, A. M. (2015). Cultivo de abóbora, suplementada com biofertilizante, em aléias de leguminosas arbóreas. *Horticultura Brasileira*, 33(1), 40–44. <https://doi.org/10.1590/S0102-053620150000100007>
220. Sánchez, A. A., Krishnamurthy, L., & Graniel, J. A. B. (2009). Leguminosas De Cobertera Para Mejorar Y Sostener La Productividad De Maíz En El Sur De Yucatán. *Terra Latinoamericana*, 27(1), 63–69.
221. Sánchez-Navarro, V., Marcos-Pérez, M., Contreras, J., & Zornoza, R. (2024). Biological nitrogen fixation and intra- and interspecific competition in two vegetable/legume intercropping systems and their relationship to crop yield. *Scientia Horticulturae*, 337, 113502. <https://doi.org/10.1016/j.scienta.2024.113502>
222. Santillano-Cázares, J., Ruiz-Alvarado, C., García-López, A. M., Escobosa-García, I., Cárdenas-Salazar, V., Morales-Maza, A., & Núñez-Ramírez, F. (2018). Assessment of Intercropping and Plastic Mulch as Tools to Manage Heat Stress, Productivity and Quality of Jalapeño Pepper. *Agronomy*, 8(12), 307. <https://doi.org/10.3390/agronomy8120307>
223. Santin, M., Parichanon, P., Sciampagna, M. C., Ranieri, A., & Castagna, A. (2024). Enhancing Tomato Productivity and Quality in Moderately Saline Soils through Salicornia-Assisted Cultivation Methods: A Comparative Study. *Horticulturae*, 10(6), 655. <https://doi.org/10.3390/horticulturae10060655>
224. Sarkar, R. K., & Shit, D. (1993). Effect of Intercropping Cereal, Pulse and Oilseed Crops in Redgram on Yield, Competition and Advantage. *Journal of Agronomy and Crop Science*, 170(3), 171–176. <https://doi.org/10.1111/j.1439-037X.1993.tb01072.x>
225. Sarker, P. K., Rahman, M. M., & Das, B. C. (2007). Effect of Intercropping of Mustard with Onion and Garlic on Aphid Population and Yield. *Journal of Bio-Science*, 15, 35–40.
226. Schultz, B., McGuinness, H., Horwith, B., Vandermeer, J., Phillips, C., Perfecto, I., Rosset, P., Ambrose, R., & Hansen, M. (1987). Effects of Planting Densities, Irrigation, and Hornworm Larvae on Yields in Experimental Intercrops of Tomatoes and Cucumbers. *Journal of the American Society for Horticultural Science*, 112(5), 747–755. <https://doi.org/10.21273/JASHS.112.5.747>
227. Schultz, B., Phillips, C., Rosset, P., & Vandermeer, J. (1982). An experiment in intercropping cucumbers and tomatoes in Southern Michigan, U.S.A. *Scientia Horticulturae*, 18(1), 1–8. [https://doi.org/10.1016/0304-4238\(82\)90096-6](https://doi.org/10.1016/0304-4238(82)90096-6)

228. Schwerdtner, U., & Spohn, M. (2021). Interspecific root interactions increase maize yields in intercropping with different companion crops. *Journal of Plant Nutrition and Soil Science*, 184(5), 596–606. <https://doi.org/10.1002/jpln.202000527>
229. Sekamatte, B. M., Ogenga-Latigo, M., & Russell-Smith, A. (2003). Effects of maize–legume intercrops on termite damage to maize, activity of predatory ants and maize yields in Uganda. *Crop Protection*, 22(1), 87–93. [https://doi.org/10.1016/S0261-2194\(02\)00115-1](https://doi.org/10.1016/S0261-2194(02)00115-1)
230. Senaratne, R., Liyanage, N. D. L., & Ratnasinghe, D. S. (1993). Effect of K on nitrogen fixation of intercrop groundnut and the competition between intercrop groundnut and maize. *Fertilizer Research*, 34(1), 9–14. <https://doi.org/10.1007/BF00749954>
231. Sharaiha, R., & Gliessman, S. (1992). The Effects of Crop Combination and Row Arrangement in the Intercropping of Lettuce, Favabean and Pea on Weed Biomass and Diversity and on Crop Yields. *Biological Agriculture & Horticulture*, 9(1), 1–13. <https://doi.org/10.1080/01448765.1992.9754615>
232. Sharma, O. P., & Gupta, A. K. (2001). Comparing the Feasibilities of Pearlmillet-Based Intercropping Systems Supplied with Varying Levels of Nitrogen and Phosphorus. *Journal of Agronomy and Crop Science*, 186(2), 91–95. <https://doi.org/10.1046/j.1439-037X.2001.00449.x>
233. Sharma, R. C., & Banik, P. (2015). Baby Corn-Legumes Intercropping Systems: I. Yields, Resource Utilization Efficiency, and Soil Health. *Agroecology and Sustainable Food Systems*, 39(1), 41–61. <https://doi.org/10.1080/21683565.2014.942764>
234. Sheha, A. M., El-Mehy, A. A., Mohamed, A. S., & Saleh, S. A. (2022). Different wheat intercropping systems with tomato to alleviate chilling stress, increase yield and profitability. *Annals of Agricultural Sciences*, 67(1), 136–145. <https://doi.org/10.1016/j.aoas.2022.06.005>
235. Shen, L., Wang, X., Liu, T., Wei, W., Zhang, S., Zhu, Y., Tuerti, T., Li, L., & Zhang, W. (2023). Apple–Soybean Mixed Stand Increased Fine Root Distribution and Soil Water Content with Reduced Soil Nitrate Nitrogen. *Agronomy*, 13(2), 548. <https://doi.org/10.3390/agronomy13020548>
236. Silue, B. K., Koné, A. W., Kotaix, A. J. A., Coulibaly, K., Aidara, S., Chapuis-Lardy, L., & Masse, D. (2024). Effects of shade tree legumes on cacao biomass and bean yields after 20 years of intercropping in Ivory Coast. *Experimental Agriculture*, 60, e16. <https://doi.org/10.1017/S0014479724000097>

237. Silva, Í. N., Bezerra Neto, F., Barros Júnior, A. P., Lima, J. S. S. D., Batista, T. M. D. V., & Lins, H. A. (2018). Green manure and spatial arrangement in the sustainability improvement of lettuce-beet intercrops. *Revista Brasileira de Engenharia Agrícola e Ambiental*, 22(7), 451–457. <https://doi.org/10.1590/1807-1929/agriambi.v22n7p451-457>
238. Silva, P. S. L. (2001). Consorciação milho e feijão caupí para produção de espigas verdes e grãos verdes. *Horticultura Brasileira*, 19(1), 4–9. <https://doi.org/10.1590/S0102-05362001000100002>
239. Singh, A. P., Bijalwan, A., Bisht, T. S., Singh, B., Kumar, S., & Tariyal, N. (2023). Evaluation of growth, yield, economics and soil properties of agri-horticulture systems in mid-hill situations of Himalayas. *Agroforestry Systems*, 97(6), 1113–1130. <https://doi.org/10.1007/s10457-023-00851-1>
240. Singh, A., Weisser, W. W., Hanna, R., Houmgny, R., & Zytynska, S. E. (2017). Reduce pests, enhance production: Benefits of intercropping at high densities for okra farmers in Cameroon. *Pest Management Science*, 73(10), 2017–2027. <https://doi.org/10.1002/ps.4636>
241. Singh, M., Singh, A., Singh, S., Tripathi, R. S., & Patra, D. D. (2011). Production potential and economics of safed musli (*Chlorophytum borivillianum*) under intercropping system. *Archives of Agronomy and Soil Science*, 57(6), 669–678. <https://doi.org/10.1080/03650341003785768>
242. Souza Dos Santos, N., Arcanjo Alves, J. M., Pereira Uchoa, S. C., Alves Albuquerque, J. A., Oliveira Da Silva, D. C., & Molina Alvarez, R. A. (2024). Calidad de fruta de la piña y evaluación del consorcio del piñal, yuca y caupí. *Bioagro*, 36(2), 163–174. <https://doi.org/10.51372/bioagro362.4>
243. Spagnollo, E., Bayer, C., Wildner, L. P., Ernani, P. R., Albuquerque, J. A., & Proença, M. M. (2002). Leguminosas estivais intercalares como fonte de nitrogênio para o milho, no sul do Brasil. *Revista Brasileira de Ciência do Solo*, 26(2), 417–423. <https://doi.org/10.1590/S0100-06832002000200015>
244. Sridhar, H. S., & Salakinkop, S. R. (2021). Competitive functions, pest dynamics and bio-economic analysis in traditional maize and legumes intercropping systems under rainfed situation of South India. *Indian Journal of Traditional Knowledge*, 20(3), 827–837. <https://doi.org/10.56042/ijtk.v20i3.30361>
245. Srivastava, A. K., Huchche, A. D., Ram, L., & Singh, S. (2007). Yield prediction in intercropped versus monocropped citrus orchards. *Scientia Horticulturae*, 114(1), 67–70. <https://doi.org/10.1016/j.scienta.2007.05.005>

246. Stavridou, E., Kristensen, H. L., Krumbein, A., Schreiner, M., & Thorup-Kristensen, K. (2012). Effect of Differential N and S Competition in Inter- and Sole Cropping of *Brassica* Species and Lettuce on Glucosinolate Concentration. *Journal of Agricultural and Food Chemistry*, 60(25), 6268–6278. <https://doi.org/10.1021/jf300091e>
247. Suresha, B. A., Allolli, T. B., Patil, M. G., Desai, B. K., & Hussain, S. (2007). Yield and Economics of Chilli Based Intercropping System. *Journal of Agricultural Science*, 20(4), 807–809.
248. Syafruddin & Suwardi. (2020). Intercropping of maize-mungbean to increase the farmer's income. *IOP Conference Series: Earth and Environmental Science*, 484(1), 012054. <https://doi.org/10.1088/1755-1315/484/1/012054>
249. Tarafder, I. H., . M. S. R., . A. K. M. M. H., . J. A. S., & . M. M. R. (2003). Economic Returns and Yield of Chilli as Intercropped with Varying Onion Population. *Pakistan Journal of Biological Sciences*, 6(2), 149–152. <https://doi.org/10.3923/pjbs.2003.149.152>
250. Tchapga, F. J. N., Chotangui, A. H., Fouegag, M. T., & Mubeteneh, T. C. (2023). Effects of potato (*Solanum tuberosum* L.)– *Mucuna pruriens* intercropping pattern on the agronomic performances of potato and the soil physicochemical properties of the western highlands of Cameroon. *Open Agriculture*, 8(1), 20220142. <https://doi.org/10.1515/opag-2022-0142>
251. Teixeira, M. L. F., & Franco, A. A. (2007). Infestação por larvas de *Ceratomyxa arcuata* (Olivier) (Coleoptera: Chrysomelidae) em nódulos de feijoeiro em cultivo com cobertura morta ou em consórcio com milho ou com caupi. *Ciência Rural*, 37(6), 1529–1535. <https://doi.org/10.1590/S0103-84782007000600004>
252. Telles, C. C., Junqueira, A. M. R., & Fukushi, Y. K. D. M. (2020). Technical feasibility of conventional and nonconventional vegetables' intercropping under organic fertilization. *Bioscience Journal*, 36, 1–13. <https://doi.org/10.14393/BJ-v36n0a2020-48219>
253. Tembakazi Silwana, T., & Lucas, E. O. (2002). The effect of planting combinations and weeding on the growth and yield of component crops of maize/bean and maize/pumpkin intercrops. *The Journal of Agricultural Science*, 138(2), 193–200. <https://doi.org/10.1017/S0021859601001861>
254. Temesgen, A., Fukai, S., & Rodriguez, D. (2015). As the level of crop productivity increases: Is there a role for intercropping in smallholder agriculture. *Field Crops Research*, 180, 155–166. <https://doi.org/10.1016/j.fcr.2015.06.003>

255. Thierfelder, C., Cheesman, S., & Rusinamhodzi, L. (2012). A comparative analysis of conservation agriculture systems: Benefits and challenges of rotations and intercropping in Zimbabwe. *Field Crops Research*, 137, 237–250. <https://doi.org/10.1016/j.fcr.2012.08.017>
256. Tinoco-Jaramillo, L., Vargas-Tierras, Y., Habibi, N., Caicedo, C., Chanaluisa, A., Paredes-Arcos, F., Viera, W., Almeida, M., & Vásquez-Castillo, W. (2024). Agroforestry Systems of Cocoa (*Theobroma cacao* L.) in the Ecuadorian Amazon. *Forests*, 15(1), 195. <https://doi.org/10.3390/f15010195>
257. Tivelli, S. W., Kano, C., Purquerio, L. F. V., Wutke, E. B., & Ishimura, I. (2013). Desempenho do quiabeiro consorciado com adubos verdes eretos de porte baixo em dois sistemas de cultivo. *Horticultura Brasileira*, 31(3), 483–488. <https://doi.org/10.1590/S0102-05362013000300023>
258. Tripathi, A. K., & Singh, A. K. (2014). Productivity, economic viability and energy efficiency of intercropping winter maize (*Zea mays*) and rajmash bean (*Phaseolus vulgaris*) in potato (*Solanum tuberosum*) with border ridge technique. *Current Advances in Agricultural Sciences*, 6(1), 10–15.
259. Tripathi, P., Shah, S., Kashyap, S. D., & Tripathi, A. (2019). Fruit yield and quality characteristics of high density *Prunus persica* (L.) Batsch plantation intercropped with medicinal and aromatic plants in the Indian Western Himalayas. *Agroforestry Systems*, 93(5), 1717–1728. <https://doi.org/10.1007/s10457-018-0276-9>
260. Ugrinovic, M., Mijatovic, M., Zdravkovic, J., Girek, Z., Kuzmanovic, D., Rasulic, N., & Josic, D. (2014). Intercropped red beet and radish with green bean affected microbial communities and nodulation by indigenous rhizobia. *Agricultural and Food Science*, 23(3), 173–185. <https://doi.org/10.23986/afsci.41385>
261. Van Asten, P. J. A., Wairegi, L. W. I., Mukasa, D., & Uringi, N. O. (2011). Agronomic and economic benefits of coffee–banana intercropping in Uganda’s smallholder farming systems. *Agricultural Systems*, 104(4), 326–334. <https://doi.org/10.1016/j.agsy.2010.12.004>
262. Varghese, L. (2000). Indicators of Production Sustainability in Intercropped Vegetable Farming on Montmorillonitic Soils in India. *Journal of Sustainable Agriculture*, 16(4), 5–17. https://doi.org/10.1300/J064v16n04_03
263. Veeramani, P., Ravichandran, V., Saravanan, P. A., Manickam, S., Venkatachalam, S. R., Arutchenthil, P., & Velmurugan, M. (2024). Effect of Different Tillage Practises on Castor (*ricinus Communis* L.) Cultivation in Legume-Based Intercropping System. *Applied Ecology and Environmental Research*, 22(2), 1139–1148. https://doi.org/10.15666/aeer/2202_11391148

264. Verma, K., Singh, A. K., & Singh, S. (2023). Assessing the suitability of traditional food crops and resource use efficiency in kalmegh [*Andrographis paniculata* (Burm. F.) Wall ex. Nees] based intercropping systems. *Industrial Crops and Products*, 203, 117203. <https://doi.org/10.1016/j.indcrop.2023.117203>
265. Vieira, J. C. B., Puiatti, M., Cecon, P. R., Bhering, A. D. S., Silva, G. D. C. C. D., & Colombo, J. N. (2014). Viabilidade agroeconômica da consorciação do taro com feijão-vagem indeterminado em razão da época de plantio. *Revista Ceres*, 61(2), 226–233. <https://doi.org/10.1590/S0034-737X2014000200010>
266. Vilhekar, S. C., Badole, K. K., & Ghanbahadur, M. R. (2014). Performance and economics of sweet corn as influenced by leafy vegetables intercropping system under rainfed condition. *Advance Research Journal of Crop Improvement*, 5(2), 118–121. <https://doi.org/10.15740/HAS/ARJCI/5.2/118-121>
267. Visscher, A. M., Chavez, E., Caicedo, C., Tinoco, L., & Pulleman, M. (2024). Biological soil health indicators are sensitive to shade tree management in a young cacao (*Theobroma cacao* L.) production system. *Geoderma Regional*, 37, e00772. <https://doi.org/10.1016/j.geodrs.2024.e00772>
268. Walia, S., & Kumar, R. (2021). Elucidating the yield and quality response of *Tagetes minuta* L. intercropped with *Zea mays* L. under different spacing in the western Himalayas. *Industrial Crops and Products*, 171, 113850. <https://doi.org/10.1016/j.indcrop.2021.113850>
269. Wang, X., Cao, B., Zou, J., Xu, A., & Feng, X. (2022). Intercropping Gramineae Herbage in Semiarid Jujube Cultivar ‘LingwuChangzao’ (*Ziziphus jujuba* Mill. Cv. LingwuChangzao) Orchard Improves Productivity, Plant Nutritional Quality, and Soil Quality. *Horticulturae*, 8(9), 834. <https://doi.org/10.3390/horticulturae8090834>
270. Wangiyana, W., Jaya, I. K. D., & Suheri, H. (2021). Application of Mycorrhiza-based biofertilizer to increase yields of several varieties of small chili intercropped with peanut or shallot. *IOP Conference Series: Earth and Environmental Science*, 648(1), 012178. <https://doi.org/10.1088/1755-1315/648/1/012178>
271. Warren Raffa, D., Migliore, M., Campanelli, G., Leteo, F., & Trinchera, A. (2022). Effects of Faba Bean Strip Cropping in an Outdoor Organic Tomato System on Soil Nutrient Availability, Production, and N Budget under Different Fertilizations. *Agronomy*, 12(6), 1372. <https://doi.org/10.3390/agronomy12061372>
272. Waweru, B. W., Rukundo, P., Kilalo, D. C., Miano, D. W., & Kimenju, J. W. (2021). Effect of border crops and intercropping on aphid infestation and the associated viral diseases in hot pepper (*Capsicum* sp.). *Crop Protection*, 145, 105623. <https://doi.org/10.1016/j.cropro.2021.105623>

273. Wei, W., Liu, T., Zhang, S., Shen, L., Wang, X., Li, L., Zhu, Y., & Zhang, W. (2024). Root spatial distribution and belowground competition in an apple/ryegrass agroforestry system. *Agricultural Systems*, 215, 103869. <https://doi.org/10.1016/j.agsy.2024.103869>
274. Workayehu, T., & Wortmann, C. S. (2011). Maize–Bean Intercrop Weed Suppression and Profitability in Southern Ethiopia. *Agronomy Journal*, 103(4), 1058–1063. <https://doi.org/10.2134/agronj2010.0493>
275. Xiao, X., Cheng, Z., Meng, H., Khan, M. A., & Li, H. (2012). Intercropping with garlic alleviated continuous cropping obstacle of cucumber in plastic tunnel. *Acta Agriculturae Scandinavica, Section B — Soil & Plant Science*, 62(8), 696–705. <https://doi.org/10.1080/09064710.2012.697571>
276. Xie, Y., & Kristensen, H. L. (2017). Intercropping leek (*Allium porrum* L.) with dyer's woad (*Isatis tinctoria* L.) increases rooted zone and agro-ecosystem retention of nitrogen. *European Journal of Agronomy*, 82, 21–32. <https://doi.org/10.1016/j.eja.2016.09.017>
277. Xu, H., Bi, H., Gao, L., & Yun, L. (2019). Alley Cropping Increases Land Use Efficiency and Economic Profitability Across the Combination Cultivation Period. *Agronomy*, 9(1), 34. <https://doi.org/10.3390/agronomy9010034>
278. Yan, Z., Wang, J., Liu, Y., You, Z., Zhang, J., Guo, F., Gao, H., Li, L., & Wan, S. (2023). Maize/Peanut Intercropping Reduces Carbon Footprint Size and Improves Net Ecosystem Economic Benefits in the Huang-Huai-Hai Region: A Four-Year Study. *Agronomy*, 13(5), 1343. <https://doi.org/10.3390/agronomy13051343>
279. Yang, L., Luo, Y., Lu, B., Zhou, G., Chang, D., Gao, S., Zhang, J., Che, Z., & Cao, W. (2023). Long-term maize and pea intercropping improved subsoil carbon storage while reduced greenhouse gas emissions. *Agriculture, Ecosystems & Environment*, 349, 108444. <https://doi.org/10.1016/j.agee.2023.108444>
280. Yang, Z., Zhang, Y., Wang, Y., Zhang, H., Zhu, Q., Yan, B., Fei, J., Xiangmin, R., Peng, J., & Luo, G. (2022). Intercropping regulation of soil phosphorus composition and microbially-driven dynamics facilitates maize phosphorus uptake and productivity improvement. *Field Crops Research*, 287, 108666. <https://doi.org/10.1016/j.fcr.2022.108666>
281. Yildirim, E., & Guvenc, I. (2005a). Intercropping based on cauliflower: More productive, profitable and highly sustainable. *European Journal of Agronomy*, 22(1), 11–18. <https://doi.org/10.1016/j.eja.2003.11.003>

282. Yap, V. Y., Xaphokhame, P., De Neergaard, A., & Bech Bruun, T. (2019). Barriers to Agro-Ecological Intensification of Smallholder Upland Farming Systems in Lao PDR. *Agronomy*, 9(7), 375. <https://doi.org/10.3390/agronomy9070375>
283. Yildirim, E., & Guvenc, I. (2005b). Intercropping in cucumber (*Cucumis sativus*) under greenhouse conditions. *The Indian Journal of Agricultural Sciences*, 74(12), 663–664.
284. Yimer, T., Abera, G., Beyene, S., & Rasche, F. (2022). Optimizing maize–bean cropping systems for sustainable intensification in southern Ethiopia. *Agronomy Journal*, 114(6), 3283–3296. <https://doi.org/10.1002/agj2.21143>
285. Žalac, H., Herman, G., Ergović, L., Jović, J., Zebec, V., Bubalo, A., & Ivezić, V. (2022). Ecological and Agronomic Benefits of Intercropping Maize in a Walnut Orchard—A Case Study. *Agronomy*, 13(1), 77. <https://doi.org/10.3390/agronomy13010077>
286. Zhang, W., Wang, B. J., Gan, Y. W., Duan, Z. P., Hao, X. D., Xu, W. L., & Li, L. H. (2019). Different tree age affects light competition and yield in wheat grown as a companion crop in jujube-wheat agroforestry. *Agroforestry Systems*, 93(2), 653–664. <https://doi.org/10.1007/s10457-017-0160-z>
287. Zhang, W., Wang, B. J., Gan, Y. W., Duan, Z. P., Hao, X. D., Xu, W. L., Lv, X., & Li, L. H. (2017). Competitive interaction in a jujube tree/wheat agroforestry system in northwest China's Xinjiang Province. *Agroforestry Systems*, 91(5), 881–893. <https://doi.org/10.1007/s10457-016-9962-7>
288. Zhang, W., Xu, W. L., Ahanbieke, P., Wang, B. J., Hao, X. D., Zhu, Y., & Li, L. H. (2019). Competition for ¹⁵N-labeled nitrogen in a jujube tree (*Zizyphus jujuba* Mill.)/wheat (*Triticum aestivum* L.) agroforestry system in northwestern China. *Agroforestry Systems*, 93(6), 2097–2110. <https://doi.org/10.1007/s10457-018-0332-5>
289. Zhao, J., Bedoussac, L., Sun, J., Chen, W., Li, W., Bao, X., Van Der Werf, W., & Li, L. (2023). Competition-recovery and overyielding of maize in intercropping depend on species temporal complementarity and nitrogen supply. *Field Crops Research*, 292, 108820. <https://doi.org/10.1016/j.fcr.2023.108820>
290. Zhu, L., Li, X., He, J., Zhou, X., Wang, F., Zhao, Y., Liang, X., Nan, X., Li, Y., Qin, K., & Cao, Y. (2023). Development of *Lycium barbarum*–Forage Intercropping Patterns. *Agronomy*, 13(5), 1365. <https://doi.org/10.3390/agronomy13051365>
291. Ziaie-Juybari, H., Pirdashti, H., Abo-Elyousr, K. A. M., & Mottaghian, A. (2021). Abiotic benefits of intercropping legumes and maize to reduce pests. *Archives of*

Phytopathology and Plant Protection, 54(17–18), 1539–1552.
<https://doi.org/10.1080/03235408.2021.1919592>

292. Zougmore, R., Kambou, F. N., Ouattara, K., & Guillobez, S. (2000). Sorghum-cowpea Intercropping: An Effective Technique Against Runoff and Soil Erosion in the Sahel (Saria, Burkina Faso). *Arid Soil Research and Rehabilitation*, 14(4), 329–342.
<https://doi.org/10.1080/08903060050136441>