	Farm economy			Farm functioning			Environment	
	Productivity	Profitability	Labour demand	Soil health	Inputs efficiency	Resilience to pests	Biodiversity	Climate mitigation
a) Cereals								
Biomass management, compared with no residues or mulch								
Organic mulch	**69%** n = 196	58% n = 12		**76%** n = 270	**78%** n = 54	69% n = 13	0% n = 4	
Residue incorporation	**67%** n = 202	18% n = 11		**67%** n = 82				47% n = 30
Residue retention	86% n = 42	42% n = 26		88% n = 16			80% n = 5	0% n = 2
Nutrient management, compared with no fertilizer applied								
Organic fertilizer	**66%** n = 151	59% n = 27		**90%** n = 71			75% n = 108	50% n = 4
Mixed fertilizer	**84%** n = 215	65% n = 20		**70%** n = 67	100% n = 4			43% n = 23
Biochar amendment	**63%** n = 67			0% n = 3				47% n = 15
Inoculation	60% n = 40					42% n = 12		
Nutrient management, compared with inorganic fertilizer								
Mixed fertilizer	100% n = 17							
Organic fertilizer	100% n = 5							
Diversification, compared with monoculture								
Intercropping	**76%** n = 564	50% n = 12				**96%** n = 72	71% n = 24	0% n = 3
Agroforestry	**52%** n = 175	50% n = 22				47% n = 32	0% n = 5	
Crop rotation	**49%** n = 90						50% n = 2	
Improved fallow	60% n = 40					42% n = 12		
Natural Fallow	42% n = 40					50% n = 12		
Diversification, compared with intercropping								
Agroforestry		71% n = 14						
Soil management, compared with conventional tillage								
Reduced tillage	**55%** n = 82	73% n = 33	100% n = 1	40% n = 15				
Variety management, compared with improved variety								
Local variety	25% n = 4							
Pest management, compared with no chemical applied								
Mechanical weeding							100% n = 1	
b) Pulses and legumes								
Diversification, compared with monoculture								
Intercropping	**28%** n = 248	62% n = 16				100% n = 12	50% n = 20	0% n = 3
Agroforestry	27% n = 15	50% n = 22						
Crop rotation	12% n = 8						50% n = 2	
Diversification, compared with intercropping								
Agroforestry		71% n = 14						
Biomass management, compared with no residues or mulch								
Residue incorporation	**31%** n = 152	18% n = 11		0% n = 18				
Residue retention	20% n = 5						100% n = 3	
Organic mulch	0% n = 3							
Nutrient management, compared with no fertilizer applied								
Organic fertilizer	79% n = 66	82% n = 11		48% n = 64				
Biochar amendment	88% n = 24							
Mixed fertilizer	100% n = 24							
Variety management, compared with improved variety								
Local variety	22% n = 36			50% n = 24				
Water management, compared with rainfed agriculture								
Supplemental irrigation	100% n = 12				11% n = 18			
Soil management, compared with conventional tillage								
Zero tillage	0% n = 12							
Reduced tillage	0% n = 6							
c) Vegetables								
Water management, compared with rainfed agriculture								
Supplemental irrigation	66% n = 402	50% n = 12						
Diversification, compared with monoculture								
Intercropping	32% n = 28					71% n = 14		
Variety management, compared with improved variety								
Local variety	42% n = 36							
Nutrient management, compared with no fertilizer applied								
Mixed fertilizer	100% n = 2	75% n = 4						0% n = 2
Organic fertilizer	100% n = 2	75% n = 4						50% n = 2
d) Roots and tubers								
Diversification, compared with monoculture								
Intercropping	7% n = 28	62% n = 16				100% n = 4	50% n = 8	
Nutrient management, compared with no fertilizer applied								
Mixed fertilizer	0% n = 6							
e) Nuts and seeds								
Diversification, compared with monoculture								
Intercropping						100% n = 6	50% n = 12	

% positive comparisons

	Negative (≤35%)	Neutral (36–64%)	Positive (≥65%)
Strong evidence (n ≥ 50, ≥ 3 studies)			
Limited evidence (fewer comparisons or studies)			

Empty cells: not measured