

Contents

1	Hurwitz Numbers	1
1.1	Maps of Riemann surfaces and Hurwitz numbers	1
1.2	Maps of Riemann surfaces as ramified covers	2
1.3	Interlude: representation theory of S_d	4
1.4	Burnside’s formula	7
	Bibliography	9

Chapter 1

Hurwitz Numbers

1.1 Maps of Riemann surfaces and Hurwitz numbers

In the following we will assume the Riemann surfaces that we will consider to be compact and connected. Let $f: X \rightarrow Y$ be a non-constant holomorphic map of Riemann surfaces (hence surjective¹)

Check where connectedness is needed.

Definition 1.1. • The *ramification index* of f at $x \in X$ is the integer $k_x \in \mathbb{Z}_{\geq 1}$ s.t. f is locally of the form z_x^k , where z is a local coordinate centered in x .

- The *differential length* of f at $x \in X$ is $\nu_x = k_x - 1$.
- A point $x \in X$ s.t. $\nu_x > 0$ is called a *ramification point*. The *ramification locus* of f is the set of its ramification points. If $\nu_x = 0$ we say that f is *unramified* at x . If $\nu_x = 1$ we say that f has *simple ramification* at x .
- If $x \in X$ is a ramification point, then $f(x) \in Y$ is called a *branch point*. The *branch locus* B_f of f is the set of its branch points.

The ramification locus and branch locus are finite sets of X and Y respectively.² We assume f non-constant. For any $y, y' \in Y \setminus B_f$ we have $|f^{-1}(y)| = |f^{-1}(y')|$.³ We call *degree* of f the integer $\deg f := |f^{-1}(y)|$ for any $y \in Y \setminus B_f$. Then:

Theorem 1.2 (Riemann-Hurwitz [CM16, Thm. 4.4.1]). *Let g_X and g_Y denote the genus of X and Y respectively. Then*

$$(1.3) \quad \underbrace{2g_X - 2}_{\chi(X)} = \underbrace{(2g_Y - 2)}_{\chi(Y)} \deg f + \sum_{x \in X} \nu_x$$

We also have that for any $y \in Y$

$$(1.4) \quad \deg f = \sum_i k_{x_i}$$

where i labels the elements of $f^{-1}(y) = \{x_1, \dots, x_n\}$. Denote $d := \deg f$.

¹[CM16, Thm. 4.3.1]

²[CM16, Lemma 4.2.5.]

³[CM16, Thm. 4.3.3]

Definition 1.5. Let $y \in Y$, $f^{-1} = \{x_1, \dots, x_n\}$. We call *ramification profile* of f at y the partition of d given by $(k_{x_1}, \dots, k_{x_n})$. We say that f is *unramified* / *simply ramified* / *fully ramified* if its ramification profile is $(1, \dots, 1)$ / $(2, 1, \dots, 1)$ / (d) respectively.

Definition 1.6. Two holomorphic maps of Riemann surfaces $f: X \rightarrow Y$ and $\tilde{X} \rightarrow Y$ are called isomorphic if there is an *isomorphism* of Riemann surfaces $\phi: X \rightarrow \tilde{X}$ s.t. $f = g \circ \phi$.

An *automorphism* of $f: X \rightarrow Y$ is an isomorphism $\psi: X \rightarrow X$ s.t. $f = f \circ \psi$. The group of automorphisms of f is denoted $\text{Aut}(f)$.

Definition 1.7. Let Y be a connected compact Riemann surface of genus g , $B = \{b_1, \dots, b_n\} \subset Y$ finite subset, $d \in \mathbb{Z}_{\geq 1}$, $\eta_1, \dots, \eta_n \vdash d$. We define the *(degree d) connected Hurwitz number* to be

$$(1.8) \quad H_{h \xrightarrow{d} g}^{\circ}(\eta_1, \dots, \eta_n) = \sum_{[f]} \frac{1}{|\text{Aut}(f)|}$$

and the *(degree d) Hurwitz number* to be

$$(1.9) \quad H_{h \xrightarrow{d} g}^{\bullet}(\eta_1, \dots, \eta_n) = \sum_{[f]} \frac{1}{|\text{Aut}(f)|}$$

where both sums runs over isomorphism classes of holomorphic maps $f: X \rightarrow Y$ s.t.

- X is a compact Riemann surface of genus h ⁴
- the branch locus of f is B
- the ramification profile of f at b_i is η_i

In the case of connected Hurwitz numbers we further require X to be connected. Numbers (1.9) are also called *disconnected Hurwitz number* since X is allowed to be disconnected.

Note that for any $f: X \rightarrow Y$ as in the definition above we have

$$(1.10) \quad \sum_{x \in X} \nu_x = \sum_{i=1}^n (d - \ell(\eta_i)) = nd - \sum_{i=1}^n \ell(\eta_i)$$

where $\ell(\eta_i)$ denotes the length of the partition η_i (i.e. the number of cycles in η_i), so due to Riemann-Hurwitz theorem we have

$$(1.11) \quad 2h - 2 = (2g - 2)d + nd - \sum_{i=1}^n \ell(\eta_i)$$

In particular the ramification profile of $f: X \rightarrow Y$ and the genus of Y uniquely determines the genus of X .

1.2 Maps of Riemann surfaces as ramified covers

Maps of Riemann surfaces as above can be regarded as ramified covers:

⁴For X disconnected, its genus is defined by $\chi = 2g - 2$, where the Euler characteristic χ is naturally additive under disjoint unions. The Riemann-Hurwitz formula then applies identically in also for X disconnected.

Definition 1.12. A *ramified cover* is a continuous function between compact topological surfaces $f: X \rightarrow Y$ s.t. there is a finite set $B \subset Y$ and

- $f^{-1}(B)$ is finite,
- $p: X \setminus f^{-1}(B) \rightarrow Y \setminus B$ is a covering.

Vice-versa, we have

Theorem 1.13 (Riemann's Existence Thm. [CM16, Thm. 6.2.2]). *Let Y compact Riemann surface, X^0 topological surface, $\{b_1, \dots, b_n\} \subset Y$ finite subset, $f^0: X^0 \rightarrow Y \setminus \{b_1, \dots, b_n\}$ covering of finite degree. Then there exists a unique (up to isomorphisms) compact Riemann surface X s.t.*

- X^0 is a dense subset of X ,
- f^0 extends to a holomorphic map of Riemann surfaces $f: X \rightarrow Y$.

Definition 1.14. Let $y_0 \in Y \setminus B_f$ and fix a bijection $f^{-1}(y_0) \cong \{1, \dots, d\}$. The ramified cover $f: X \rightarrow Y$ determines a group homomorphism

$$(1.15) \quad \Phi: \pi_1(Y \setminus B_f, y_0) \rightarrow S_d, \quad \gamma \mapsto \sigma_\gamma$$

called *monodromy representation*.

Now consider $b \in B_f$ and $\gamma \in \pi_1(Y \setminus B_f, y_0)$ simple loop winding once around b (and with zero winding number around the other branch points). If the ramification profile of f at b is $\eta = (k_1, \dots, k_l)$, then σ_γ has cycle type η (to see this recall that the local expression of f around ramification points is z^k and consider a circle around y_0 of unit radius in the chart).

Definition 1.16. Let Y be a connected Riemann surface of genus g , $y_0, b_1, \dots, b_n \in Y$ points, $d \in \mathbb{Z}_{\geq 1}$ and $\eta_1, \dots, \eta_n \vdash d$. A *monodromy representation of type $(g, d, \eta_1, \dots, \eta_n)$* is a group homomorphism $\Phi: \pi_1(Y \setminus \{b_1, \dots, b_n\}, y_0) \rightarrow S_d$ s.t. if γ_k is a small loop around b_k then $\Phi(\gamma_k)$ has cycle type η_k .

If moreover the subgroup $\text{im } \Phi \subset S_d$ acts transitively on $\{1, 2, \dots, d\}$ we say that Φ is a *connected monodromy representation*.

We obtained that a degree d map $f: X \rightarrow Y$ between compact connected Riemann surfaces s.t. the ramification profile over each branch point is η_i gives rise to a connected monodromy representation Φ of type $(g_Y, d, \eta_1, \dots, \eta_n)$. If we let X to be non connected, then the monodromy representation may not be connected anymore, more precisely: the monodromy representation is connected if and only if X is connected.⁵ We also have that isomorphic maps give the same monodromy representation.

Conversely:

Theorem 1.17 ([CM16, Thm. 7.2.2]). *Let Y be a Riemann surface of genus g , Φ a monodromy representation of type $(g, d, \eta_1, \dots, \eta_n)$, $B = \{b_1, \dots, b_n\} \subset Y$ a finite subset. Then exists a holomorphic map of Riemann surface covering Y with branch locus B whose associated monodromy is Φ . Such map is unique up to isomorphisms.*

Our discussion leads to a bijection between isomorphisms classes of holomorphic maps with a given ramification profile and monodromy representations. More precisely:

⁵For further details: [CM16, §§7.1].

Theorem 1.18 ([CM16, Thm. 7.3.1, Thm. 7.3.2]). *Let M° (resp M^\bullet) be the set of connected monodromy representations (resp. monodromy representations) of type $(g, d, \eta_1, \dots, \eta_n)$. Then*

$$(1.19) \quad H_{h \rightarrow g}^\circ(\eta_1, \dots, \eta_n) = \frac{|M^\circ|}{d!}$$

and

$$(1.20) \quad H_{h \rightarrow g}^\bullet(\eta_1, \dots, \eta_n) = \frac{|M^\bullet|}{d!}$$

where h is determined by (1.11).

Although the information carried by connected Hurwitz numbers is usually more interesting for geometrical purposes, it turns out that it is easier to compute the (possibly disconnected) Hurwitz numbers. We will see later how it is possible to recover the connected Hurwitz numbers from the disconnected ones.

We mention that using some “degeneration formulas” (which heuristically correspond to shrink the Riemann surface Y producing nodal curves) all disconnected degree d Hurwitz numbers are determined in terms of Hurwitz numbers of the form $H_{h \rightarrow 0}^\bullet(\eta_1, \eta_2, \eta_3)$.⁶ For this (and other) reason we later restrict our discussion to the case $g = 0$.

1.3 Interlude: representation theory of S_d

Using theorem 1.18 the problem of computing degree d Hurwitz numbers can be translated into a problem in representation theory of the symmetric group S_d . In order to show this we need some facts about representation theory (see [FH04, Part I] and [CM16, §8] for more details).

Definition 1.21. The *group algebra* of the symmetric group S_d is the complex algebra generated by the elements of S_d , that is

$$(1.22) \quad \mathbb{C}[S_d] := \left\{ \sum_{\sigma \in S_d} a_\sigma \sigma \mid a_\sigma \in \mathbb{C} \right\}$$

with operations

$$(1.23) \quad \sum_{\sigma \in S_d} a_\sigma \sigma + \sum_{\sigma \in S_d} b_\sigma \sigma = \sum_{\sigma \in S_d} (a_\sigma + b_\sigma) \sigma$$

$$(1.24) \quad \left(\sum_{\sigma \in S_d} a_\sigma \sigma \right) \cdot \left(\sum_{\sigma \in S_d} b_\sigma \sigma \right) = \sum_{\sigma \in S_d} \sum_{\sigma' \in S_d} a_\sigma b_{\sigma'} (\sigma \cdot \sigma')$$

$$(1.25) \quad t \cdot \left(\sum_{\sigma \in S_d} a_\sigma \sigma \right) = \sum_{\sigma \in S_d} (ta_\sigma) \sigma$$

⁶[CM16, Thm. 7.5.3]

where $t \in \mathbb{C}$. Expression in the r.h.s. of (1.24) (before multiplying σ and σ') is called formal expansion of the product.

We define *class algebra* of S_d the center of the group algebra

$$(1.26) \quad \mathcal{Z}\mathbb{C}[S_d] = \{x \in \mathbb{C}[S_d] \mid yx = xy \text{ for all } y \in \mathbb{C}[S_d]\}$$

The following functions are very important for our discussion:

Definition 1.27. A *class function* on S_d is a map $\alpha: S_d \rightarrow \mathbb{C}$ which is constant on conjugacy classes, i.e. $\forall h \in S_d$ we have $\alpha(h^{-1}gh) = \alpha(h)$. Let $\mathbb{C}_{\text{class}}$ denote the vector space of class functions on S_d . We define on $\mathbb{C}_{\text{class}}$ the following Hermitian inner product

$$(1.28) \quad (\alpha, \beta) := \frac{1}{d!} \sum_{\sigma \in S_d} \alpha(\sigma) \overline{\beta(\sigma)} = \frac{1}{d!} \sum_{C \subset S_d} |C| \alpha(C) \overline{\beta(C)}$$

for any $\alpha, \beta \in \mathbb{C}_{\text{class}}$, where \sum_C runs over the conjugacy classes of S_d .

We have the following

Lemma 1.29 ([FH04, §§3.4]).

$$(1.30) \quad \mathcal{Z}\mathbb{C}[S_d] = \left\{ \sum_{\sigma \in S_d} \alpha(\sigma) \sigma \mid \alpha \in \mathbb{C}_{\text{class}} \right\}$$

For $\eta \vdash d$ denote by $C_\eta \subset S_d$ the conjugacy class corresponding to all elements of cycle type η . Let $\alpha_\eta: S_d \rightarrow \mathbb{C}$ the class function which takes value 1 on elements of C_η and 0 otherwise. It is clear that the set $\{\alpha_\eta \mid \eta \vdash d\}$ gives a basis of $\mathbb{C}_{\text{class}}$. Let $c_\eta \in \mathcal{Z}\mathbb{C}[S_d]$ denote the corresponding element in the center of the group algebra, that is

$$(1.31) \quad c_\eta = \sum_{\sigma \in S_d} \alpha_\eta(\sigma) \sigma = \sum_{\sigma \in C_\eta} \sigma$$

We get that $\{c_\eta \mid \eta \vdash d\}$ form a basis of $\mathcal{Z}\mathbb{C}[S_d]$ as a complex vector space, called *conjugacy class basis*

$$(1.32) \quad \mathcal{Z}\mathbb{C}[S_d] = \bigoplus_{\eta \vdash d} \langle c_\eta \rangle_{\mathbb{C}}$$

Note that the identity element c_e in $\mathcal{Z}\mathbb{C}[S_d]$ corresponds to the partition $e = (1, \dots, 1)$.

A (complex) representation of S_d is a homomorphism $\rho: S_d \rightarrow \text{Aut}(V_\rho)$ making the finite dimensional complex vector space V_ρ into a S_d -module. We define the dimension of the representation to be $\dim \rho := \dim_{\mathbb{C}} V_\rho$. Given any representation ρ , notice that this extends to a homomorphism $\mathbb{C}[S_d] \rightarrow \text{End}(V_\rho)$, making V_ρ into a $\mathbb{C}[S_d]$ -module.

Representations of S_d are described by their characters, which are defined as follows:

Definition 1.33. Let ρ be a representation of S_d . The *character* of ρ is the class function $\chi_\rho \in \mathbb{C}_{\text{class}}$ defined by

$$(1.34) \quad \chi_\rho(\sigma) := \text{tr}(\rho(\sigma))$$

The fact that χ_ρ is a class function follows from the cyclicity of the trace. From the definition it follows that χ_ρ does not depend on the choice of basis for V_ρ (due to the corresponding property of the trace) and that $\chi_\rho(e) = \dim \rho$ (since $\rho(e) = \text{id}$). It is also easy to see that $\chi_{\rho_1 \oplus \rho_2} = \chi_{\rho_1} + \chi_{\rho_2}$.

A representation ρ is *irreducible* if V_ρ does not contain any nontrivial S_d -submodules. For complex representation we have the following fundamental result

Theorem 1.35 ([FH04, Thm. 2.12]). *In terms of the inner product (1.28) the characters of the irreducible representations of S_d are orthonormal:*

$$(1.36) \quad (\chi_{\rho_1}, \chi_{\rho_2}) = \begin{cases} 1 & \text{if } \rho_1 \cong \rho_2 \\ 0 & \text{if } \rho_1 \not\cong \rho_2 \end{cases}$$

where ρ_1, ρ_2 are irreducible representations.

Any complex representation decomposes uniquely into the direct sum of irreducible representations. More precisely, we have

$$(1.37) \quad R = \bigoplus_{\rho} V_{\rho}^{\oplus \dim V_{\rho}}$$

where R is the obvious representation of S_d on $\mathbb{C}[S_d]$, called *regular representation*, while the (big) direct sum runs over all the irreducible representations of S_d . This also implies the following isomorphism of algebras

$$(1.38) \quad \mathbb{C}[S_d] \cong \bigoplus_{\rho} \text{End}(V_{\rho})$$

which is defined extending $S_d \rightarrow \bigoplus_{\rho} \text{End}(V_{\rho})$ by linearity.

Theorem 1.39 ([FH04, Thm. 4.3]). *To each partition $\lambda \vdash d$ corresponds a unique irreducible representation V_{λ} of S_d . The corresponding character, denoted χ^{λ} , is given by Frobenius formula [FH04, Eq. 4.10]. In particular χ^{λ} is a real class function.*

By dimensional arguments, we get that characters of irreducible representations form another basis of $\mathbb{C}_{\text{class}}$. Moreover, denoting

$$(1.40) \quad e_{\lambda} := \sum_{\sigma \in S_d} \chi^{\lambda}(\sigma) \sigma$$

we get that $\{e_{\lambda} \mid \lambda \vdash d\}$ gives another basis of $\mathbb{Z}\mathbb{C}[S_d]$

$$(1.41) \quad \mathbb{Z}\mathbb{C}[S_d] = \bigoplus_{\lambda \vdash d} \langle e_{\lambda} \rangle \mathbb{C}$$

which will be called *character basis* or *idempotent basis*. Indeed from characters orthogonality (and the fact that $\overline{\chi^{\lambda}} = \chi^{\lambda}$) we get

$$(1.42) \quad e_{\lambda_i} \cdot e_{\lambda_j} = \begin{cases} e_{\lambda_i} & \text{if } e_{\lambda_i} = e_{\lambda_j} \\ 0 & \text{otherwise} \end{cases}$$

Since $\mathcal{Z}\mathbb{C}[S_d]$ admits a basis of idempotent elements, we say that it is a *semisimple algebra*. The formulas for the change of basis are given by the characters themselves

$$(1.43) \quad e_\lambda = \frac{\dim \lambda}{d!} \sum_{\eta \vdash d} \chi^\lambda(C_\eta) c_\eta \quad \text{and} \quad c_\eta = |C_\eta| \sum_{\lambda \vdash d} \frac{\chi^\lambda(C_\eta)}{\dim \lambda} e_\lambda$$

where $\dim \lambda := \dim V_\lambda$. In order to simplify the notation, set also $\chi_\eta^\lambda := \chi^\lambda(C_\eta)$.

1.4 Burnside's formula

Now we are ready to use the notions that we introduced in order to rewrite the expression of Hurwitz numbers in terms of characters of the irreducible representations of S_d . Existence of a idempotent basis for $\mathcal{Z}\mathbb{C}[S_d]$ and formulas (1.43) will be crucial.

Recall that

$$(1.44) \quad H_{h \rightarrow g}^\bullet(\eta_1, \dots, \eta_n) = \frac{|M^\bullet|}{d!}$$

where M^\bullet is the set of monodromy representations of type $(g, d, \eta_1, \dots, \eta_r)$ where η_1, \dots, η_n are partitions of d .

In order to account for the case $g \neq 0$ one needs

Definition 1.45. Let $d \in \mathbb{Z}_{\geq 1}$, $\eta \vdash d$. Denote by $\xi(\eta)$ the centralizer of C_η . We define the *kommutator* to be the element

$$(1.46) \quad \mathfrak{K} := \sum_{\eta \vdash d} |\xi(\eta)| c_\eta^2 \in \mathcal{Z}\mathbb{C}[S_d]$$

Then we have the following

Proposition 1.47 ([CM16, Prop. 9.2.3]).

$$(1.48) \quad H_{h \rightarrow g}^\bullet(\eta_1, \dots, \eta_n) = \frac{1}{d!} [c_e] \mathfrak{K}^g c_{\eta_n} \cdots c_{\eta_2} c_{\eta_1}$$

where $[c_e] \mathfrak{K}^g c_{\eta_n} \cdots c_{\eta_2} c_{\eta_1}$ denotes the coefficient of c_e after writing the product $\mathfrak{K}^g c_{\eta_n} \cdots c_{\eta_2} c_{\eta_1}$ as a linear combination of the basis elements $c_\eta \in \mathcal{Z}\mathbb{C}[S_d]$. As usual h is determined by Riemann-Hurwitz formula.

By changing basis from the conjugacy basis to the idempotent basis we get

Theorem 1.49 (Burnside's Character Formula [CM16, Thm. 9.3.1]).

$$(1.50) \quad H_{h \rightarrow g}^\bullet(\eta_1, \dots, \eta_n) = \sum_{\lambda \vdash d} \left(\frac{\dim \lambda}{d!} \right)^{2-2g} \prod_{j=1}^n f_{C_j}(\lambda)$$

where

$$(1.51) \quad f_{C_j}(\lambda) := |C_j| \frac{\chi^\lambda(C_j)}{\dim \lambda}$$

and $C_j := C_{\eta_j}$.

Maybe add proof for $g = 0$.

Recall that from the change of basis formula,

$$(1.52) \quad c_\eta = \sum_{\lambda \vdash d} f_{C_\eta}(\lambda) e_\lambda$$

From Bournside's formula we see that such coefficients for the change of basis correspond to the contribution of the ramification profile η to the disconnected Hurwitz numbers.

1.5 The generating function

In the following we will restrict ourself to the case of $g = 0$. Recall that there are some degeneration formulas which allows to express all the Hurwitz numbers in terms of those for $g = 0$. For the same discussion in more generality see [CM16, §10]. We will follow instead [Oko00, §§2.2, 2.3].

Rather than considering the different Hurwitz numbers separately, it worth to collect them together into generating functions.

Bibliography

- [CM16] Renzo Cavalieri and Eric Miles. *Riemann Surfaces and Algebraic Curves: A First Course in Hurwitz Theory*. London Mathematical Society Student Texts. Cambridge University Press, 2016. DOI: [10.1017/CB09781316569252](https://doi.org/10.1017/CB09781316569252).
- [FH04] William Fulton and Joe Harris. *Representation Theory*. Springer New York, 2004. DOI: [10.1007/978-1-4612-0979-9](https://doi.org/10.1007/978-1-4612-0979-9).
- [Joh15] Paul Johnson. “Double Hurwitz numbers via the infinite wedge”. In: *Transactions of the American Mathematical Society* 367.9 (Apr. 2015), pp. 6415–6440. DOI: [10.1090/s0002-9947-2015-06238-2](https://doi.org/10.1090/s0002-9947-2015-06238-2).
- [MJD00] T. Miwa, M. Jimbo, and E. Date. *Solitons: Differential Equations, Symmetries and Infinite Dimensional Algebras*. Ed. by M. Reid, W. Fulton, A. Katok, F. Kirwan, B. Bollobas, P. Sarnak, et al. Cambridge Tracts in Mathematics. Cambridge University Press, 2000. ISBN: 9780521561617.
- [Oko01] A. Okounkov. “Infinite wedge and random partitions”. In: *Selecta Mathematica* 7.1 (Apr. 2001), pp. 57–81. DOI: [10.1007/pl00001398](https://doi.org/10.1007/pl00001398).
- [Oko00] Andrei Okounkov. “Toda equations for Hurwitz numbers”. In: *Mathematical Research Letters* 7.4 (2000), pp. 447–453. DOI: [10.4310/mrl.2000.v7.n4.a10](https://doi.org/10.4310/mrl.2000.v7.n4.a10).
- [OP06a] Andrei Okounkov and Rahul Pandharipande. “Gromov–Witten theory, Hurwitz theory, and completed cycles”. In: *Annals of Mathematics* 163.2 (Mar. 2006), pp. 517–560. DOI: [10.4007/annals.2006.163.517](https://doi.org/10.4007/annals.2006.163.517).
- [OP06b] Andrei Okounkov and Rahul Pandharipande. “The equivariant Gromov–Witten theory of P^1 ”. In: *Annals of Mathematics* 163.2 (Mar. 2006), pp. 561–605. DOI: [10.4007/annals.2006.163.561](https://doi.org/10.4007/annals.2006.163.561).