

Abstract

In this document we are going to prove **De Rham** theorem, which states that De Rham cohomology and singular cohomology, over a *paracompact* manifold M are isomorphic. Moreover we will prove the explicit form of this isomorphism, given by the integration of forms on singular simplices. This result is going to be proved using F -injective resolutions and tools from sheaf theory.

1 Useful results

Let M be a *paracompact smooth real manifold*.

Lemma 1.1. *Every **fine** sheaf on a M is **soft**.*

Lemma 1.2. *The full subcategory of **soft sheaves** is $\Gamma(M, -)$ -injective.*

2 Singular Cohomology

For this section consider K to be a PID, M a *paracompact manifold* and $U \subset M$ an open subset of M . Let us define the *continuous singular simplices*:

Definition 2.1: Standard p -simplex.

Let $\mathbb{N} \ni p \geq 1$. We define the *standard p -simplex*

$$\Delta^p := \left\{ (a_1, \dots, a_p) \in \mathbb{R}^p \mid \sum_{i=1}^p a_i \leq 1 \text{ and, } \forall i \in \{1, \dots, p\}, a_i \geq 0 \right\}. \quad (2.1)$$

For $p = 0$ we set $\Delta^0 := \{0\}$ the 1-point space, and we call Δ^0 the *standard 0-simplex*.

Definition 2.2: Continuous/differentiable singular p -simplex.

We define a *continuous singular simplex* σ in U to be a *continuous* map $\sigma : \Delta^p \rightarrow U$.

If $p \geq 1$ we say that σ is a *differentiable singular p -simplex* in U iff it can be extended to a differentiable (\mathcal{C}^∞) map of a neighborhood of Δ^p in \mathbb{R}^p into U .

Remark 1 The theory can be developed also for *differentiable* singular simplices, but it requires some care we are not willing to give. In general it exactly the same as for *continuous* ones, apart from where elsewhere noted. In case you want to develop such theory, please refer to [War83].

From now on we are going to deal only with *continuous* singular simplices, hence we will stop indicating their continuity.

Definition 2.3: Singular p -chains with integer coefficients.

Fixed $U \overset{\text{open}}{\subset} M$, we define $S_p(U)$ to be the *free abelian group* (equivalently the *free \mathbb{Z} -module*) generated by the singular p -simplices in U . Its elements are called *singular p -chains with integer coefficients*, and can be written as finite formal sums, with integer coefficients, of singular simplices, such as:

$$c = \sum_{j=1}^n n_j \sigma_j, \quad (2.2)$$

where $n_j \in \mathbb{Z} \setminus \{0\}$ and σ_j a singular p -simplex for every $1 \leq j \leq n$.

Definition 2.4: Boundary.

We define, for each $p \geq 0$ and $0 \leq i \leq p+1$, the collection of maps $k_i^p : \Delta^p \rightarrow \Delta^{p+1}$:

$$\begin{aligned} \text{for } p = 0, \quad & \begin{cases} k_0^0(0) := 1 \\ k_1^0(0) := 0 \end{cases} \\ \text{for } p \geq 1, \quad & \begin{cases} k_0^p(a_1, \dots, a_p) := \left(1 - \sum_{j=1}^p a_j, a_1, \dots, a_p\right) \\ k_i^p(a_1, \dots, a_p) := (a_1, \dots, a_{i-1}, 0, a_{i+1}, \dots, a_p) \quad \text{for } 1 \leq i \leq p \end{cases} \end{aligned} \quad (2.3)$$

Given a singular p -simplex σ in $U \overset{\text{open}}{\subset} M$, we define its i th *face*, for $0 \leq i \leq p$, to be the singular $(p-1)$ -simplex

$$\sigma^i := \sigma \circ k_i^{p-1}. \quad (2.4)$$

We will use the superscript index to denote the face of a singular simplex. Finally we define the *boundary* of σ to be the singular $(p-1)$ -chain

$$\partial\sigma := \sum_{j=0}^p (-1)^j \sigma^j \in S_{p-1}(U), \quad (2.5)$$

in which, as stated before, σ^j denotes the j th face of σ .

Extending the *boundary operator* ∂ by linearity, we obtain a *homomorphism*

$$\partial : S_p(U) \rightarrow S_{p-1}(U). \quad (2.6)$$

More explicitly, the boundary operator acts as follows on a singular p -chain:

$$\partial \left(\sum_{j=1}^n a_j \sigma_j \right) := \sum_{j=1}^n a_j \partial \sigma_j = \sum_{j=1}^n \sum_{i=0}^p (-1)^i a_j \sigma_j^i. \quad (2.7)$$

Lemma 2.5. $k_i^{p+1} \circ k_j^p = k_{j+1}^{p+1} \circ k_i^p$ for any $p \geq 0$ and $i \leq j$.

Proof. If $p = 0$ it can be checked directly (there are only 3 cases). For $p \geq 1$ it will be computed directly. The first term acts as

$$k_i^{p+1} \circ k_j^p(a_1, \dots, a_p) = \begin{cases} (a_1, \dots, a_{i-1}, 0, a_i, \dots, a_{j-1}, 0, a_j, \dots, a_p) & \text{if } 1 \leq i < j \\ (a_1, \dots, a_{i-1}, 0, 0, a_i, \dots, a_p) & \text{if } 1 \leq i = j \\ \left(1 - \sum_{j=1}^p a_j, \dots, a_{j-1}, 0, a_j, \dots, a_p\right) & \text{if } 0 = i < j \\ \left(0, \sum_{j=1}^p a_j, a_1, \dots, a_p\right) & \text{if } 0 = i = j \end{cases} \quad (2.8)$$

Analogously we can compute that the second term acts as

$$k_{j+1}^{p+1} \circ k_i^p(a_1, \dots, a_p) = \begin{cases} (a_1, \dots, a_{i-1}, 0, a_i, \dots, a_{j-1}, 0, a_j, \dots, a_p) & \text{if } 1 \leq i < j \\ (a_1, \dots, a_{i-1}, 0, 0, a_i, \dots, a_p) & \text{if } 1 \leq i = j \\ \left(1 - \sum_{j=1}^p a_j, \dots, a_{j-1}, 0, a_j, \dots, a_p\right) & \text{if } 0 = i < j \\ \left(0, \sum_{j=1}^p a_j, a_1, \dots, a_p\right) & \text{if } 0 = i = j \end{cases} \quad (2.9)$$

■

Lemma 2.6. $\partial \circ \partial = 0$.

Proof. We are gonna prove this result only for singular p simplices, then by linearity this result will hold for arbitrary singular p -chains. If $p = 0, 1$ the result is trivial. Let, now, $p \geq 2$ and σ be a singular p -simplex. By definition ∂ acts on σ as

$$\partial\sigma = \sum_{i=0}^p (-1)^i \left(\sigma \circ k_i^{p-1} \right). \quad (2.10)$$

Hence we can compute the double *boundary* as

$$\partial \circ \partial\sigma = \sum_{i=0}^p \sum_{j=0}^{p-1} (-1)^{i+j} \left(\sigma \circ k_i^{p-1} \circ k_j^{p-2} \right). \quad (2.11)$$

We can divide the sum for $i \leq j$ and $i > j$, as

$$\partial \circ \partial\sigma = \sum_{j=0}^{p-1} \sum_{i=j+1}^p (-1)^{i+j} \left(\sigma \circ k_i^{p-1} \circ k_j^{p-2} \right) + \sum_{\tilde{j}=0}^{p-1} \sum_{\tilde{i}=0}^{\tilde{j}} (-1)^{\tilde{i}+\tilde{j}} \left(\sigma \circ k_{\tilde{i}}^{p-1} \circ k_{\tilde{j}}^{p-2} \right) \quad (2.12)$$

$$= \sum_{j=0}^{p-1} \sum_{i=j+1}^p (-1)^{i+j} \left(\sigma \circ k_i^{p-1} \circ k_j^{p-2} \right) + \sum_{i=1}^p \sum_{j=0}^{i-1} (-1)^{i+j+1} \left(\sigma \circ k_i^{p-1} \circ k_j^{p-2} \right) \quad (2.13)$$

$$= 0, \quad (2.14)$$

where, in the second sum we put $i := \tilde{j} + 1$ and $j := \tilde{i}$, and we concluded with the last line since the two sums are over all i, j s.t. $i > j$ and they only differ by a sign. ■

Definition 2.7: Singular p -cochain on U .

Let $S^p(U, K)$, with $U \overset{\text{open}}{\subset} M$ and K a PID as usual, be the set of functions that map a singular p -simplex in U into an element of K . An element of $S^p(U, K)$ is called *singular p -cochain* on U .

The set $S^p(U, K)$ can be made into a K -module by defining the following addition and scalar multiplication:

$$(kf)(\sigma) := k \cdot f(\sigma) \quad (2.15)$$

$$(f + g)(\sigma) := f(\sigma) + g(\sigma). \quad (2.16)$$

Also note that each singular p -cochain can be extended into a *homomorphism* of $S_p(U)$ into K by linearity. This actually determines an isomorphism of $S^p(U, K)$ into the K -module of morphisms of $S_p(U)$ into K . We will, hence, identify elements of $S^p(U, K)$ with the corresponding morphism.

Definition 2.8: Presheaf of singular p -cochains.

We can define the functor $\tilde{S}_K^p : \text{Op}_M^{\text{op}} \rightarrow \text{Mod}(K)$ as the functor which acts

- on the objects: $\tilde{S}_K^p(U) := S^p(U, K)$, for any $U \in \text{Op}_M$,
- on the morphisms: $V \overset{\text{open}}{\subset} U$ iff we have $U \rightarrow V$ in Op_M^{op} . \tilde{S}_K^p maps this morphism to $\rho_{V,U} : S^p(U, K) \rightarrow S^p(V, K)$, the function which maps any $f \in S^p(U, K)$ to its restriction to singular p -simplices on V .

Clearly, for any $p \geq 1$ these are *presheaves*.

Note that these *presheaves* satisfy **S2**, but not **S1**:

S2: Given $\mathcal{U} := \{U_i\}_{i \in I}$ an open covering of $U \stackrel{\text{open}}{\subset} X$ and p -cochains $f_i \in \tilde{S}_K^p(U_i)$ s.t. $f_i|_{U_i \cap U_j} = f_j|_{U_i \cap U_j}$ for any $i, j \in I$ we can construct $f \in \tilde{S}_K^p(U)$ s.t. $f|_{U_i} = f_i$ for every $i \in I$. In fact $f|_{U_i}$ is f acting only on singular p -simplices in U_i . This, combined with the fact that we are not imposing any continuity conditions on f means that we can take f to be define as f_i when computed on any singular p -simplex in U_i and any arbitrary value – e.g. 0 – when computed on a singular p -simplex with range not contained in any U_i .

S1: The last sentence should make clear the fact that the extension is not unique – i.e. we can map the last singular simplices to any value in K – giving at least two different extensions for any set of local cochains. We then have that equality cannot be checked locally.

This shortcoming hints to the fact that we will have to consider the *sheafification* of this presheaf in order to prove an isomorphism of cohomologies (spoiler alert: we are going to use resolutions).

Let's now lay the ground for the resolution we will construct:

Definition 2.9: Coboundary homomorphism.

One can define the *coboundary homomorphism* $d(U) : S^p(U, K) \rightarrow S^{p+1}(U, K)$ setting

$$d(U)f(\sigma) := f(\partial\sigma), \quad (2.17)$$

for any $f \in S^p(U, K)$ and σ a singular $(p+1)$ -simplex with range in U .

Remark 2 Note that, since $\partial \circ \partial = 0$, also for the coboundary homomorphism, fixed $U \stackrel{\text{open}}{\subset} M$, we have $d(U) \circ d(U) = 0$. Moreover d yields a *morphism of presheaves*:

$$d : \tilde{S}_K^p \rightarrow \tilde{S}_K^{p+1}. \quad (2.18)$$

In order to check it we have to show that, for any $V \stackrel{\text{open}}{\subset} U \in \mathbf{Op}_M$, the following diagram commutes:

$$\begin{array}{ccc} S^p(U, K) & \xrightarrow{d(U)} & S^{p+1}(U, K) \\ \rho_{U,V} \downarrow & & \downarrow \rho_{U,V} \\ S^p(V, K) & \xrightarrow{d(V)} & S^{p+1}(V, K) \end{array}, \quad (2.19)$$

where, as defined above, $d(U)$ represents the *coboundary* homomorphism from $S^p(U, K)$ and $d(V)$ the one from $S^p(V, K)$. The diagram clearly commutes, since d commutes with restriction (the boundary operator is not affected: the range of σ has to be in the restricted domain).

Definition 2.10: Complex of presheaves of singular cochains.

Consider the following chain of presheaves of *singular q -cochains* (consider them to be 0 for every $q < 0$):

$$\dots \rightarrow 0 \rightarrow \tilde{S}_K^0 \xrightarrow{d} \tilde{S}_K^1 \xrightarrow{d} \tilde{S}_K^2 \xrightarrow{d} \dots \quad (2.20)$$

This is a complex, since we have proved that the coboundary morphism satisfies

$$d \circ d = 0. \quad (2.21)$$

We denote it with \tilde{S}_K^\bullet and call it the *complex of presheaves of singular cochains*.

Definition 2.11: Singular cohomology.

We associate, to the global sections of the complex of presheaves of singular cochains, the classical singular cohomology, i.e.

$$H_\Delta^q(M) := H^q(\tilde{S}_K^\bullet(M)). \quad (2.22)$$

Let's now create the corresponding *complex of sheaves*:

Definition 2.12: Sheaf of singular p -cochains.

We define the *sheaf of p -cochains* $S_K^p := (\tilde{S}_K^p)^a$ the *sheafification* of the presheaf of p -cochains.

With the aim of obtaining a $\Gamma(M; -)$ -injective resolution for the constant sheaf in K , we want to prove that the just defined sheaf is fine:

Lemma 2.13. *For any p the sheaf S_K^p is **fine**.*

Proof. Recall that M is a *paracompact* manifold. Hence it admits a locally finite open cover $\{U_i\}_{i \in I}$, with associated *partition of unity* $\{\varphi_i\}_{i \in I}$. As proved by Warner (see [War83] section 5.22) we can choose φ_i that only assume the values 0 and 1, with $\text{supp } \varphi_i = \overline{V}_i \subset U_i$.

We aim to exhibit a partition of unity for the sheaf S_K^p . Our starting block will be the family $\{\tilde{l}_i\}_{i \in I}$ of endomorphisms of the presheaf \tilde{S}_K^p . Fixed $U \xrightarrow{\text{open}} M$, $f \in \tilde{S}_K^p(U)$ and σ a singular p -simplex, then these morphisms act as:

$$\tilde{l}_i(U)(f)(\sigma) := \varphi_i(\sigma(0))f(\sigma). \quad (2.23)$$

The fact that these morphisms commute with restriction is trivially true, hence $\{\tilde{l}_i\}_{i \in I}$ is really a family of morphisms of *presheaves*.

We can, now, consider $\{l_i\}_{i \in I}$, the family of morphisms of *sheaves* associated by sheafification. We want to prove that this is a *partition of unity* for the sheaf S_K^p :

- By definition $\text{supp } l_i := \overline{\{m \in M \mid (l_i)_m \neq 0\}}$. We also know that the stalk of the sheafification is isomorphic to the stalk of the original presheaf, so we will check the support of l_i by checking the behaviour of \tilde{l}_i at the level of stalks. If $m \notin \overline{V}_i$, then $\exists U_m$ an open neighborhood of m s.t. $U_m \cap \overline{V}_i = \emptyset$. Since \overline{V}_i is the support of φ_i we deduce that $(l_i)_m f_m = 0$ for any $f_m \in (\tilde{S}_K^p)_m$. It immediately follows that $\text{supp } l_i \subset \overline{V}_i \subset U_i$;
- The morphism $\sum_{i \in I} l_i(U)$ sends $s \in S_K^p(U)$ to

$$\sum_{i \in I} l_i(U)(s) : m \mapsto \sum_{i \in I} (l_i)_m s(m). \quad (2.24)$$

Clearly the last term is exactly $s(m)$ by definition of $\{\tilde{l}_i\}_{i \in I}$. This shows that

$$\sum_{i \in I} l_i(U) = \text{id}_{S_K^p}, \quad (2.25)$$

hence that $\{l_i\}_{i \in I}$ is a *partition of unity* for S_K^p . ■

Definition 2.14: Complex of sheaves of singular cochains.

Let us denote (with a little abuse of notation) by d the image of the *coboundary morphism* by the sheafification functor, d^a . Then, associated with 2.10, we have the *complex of sheaves of singular cochains*

$$\dots \rightarrow 0 \rightarrow S_K^0 \xrightarrow{d} S_K^1 \xrightarrow{d} S_K^2 \xrightarrow{d} \dots \quad (2.26)$$

Remark 3 This is going to be the starting block for the resolution of the constant sheaf K_M . Our aim in the next few lemmas is, in fact, to prove that

$$0 \rightarrow K_M \rightarrow S_K^0 \xrightarrow{d} S_K^1 \xrightarrow{d} S_K^2 \xrightarrow{d} \dots, \quad (2.27)$$

is exact.

Remark 4 Since the functor $(-)^a$ is exact we can check exactness of (2.27) at K_M and at S_K^0 by checking the exactness of

$$0 \rightarrow K_M \rightarrow \tilde{S}_K^0 \xrightarrow{d} \tilde{S}_K^1. \quad (2.28)$$

- It is exact at K_M iff the morphism from the constant sheaf with values in K in \tilde{S}_K^0 is mono. It is mono iff for any U open in M the associated morphism of modules is injective. This map just sends a locally constant function $f \in K_M(U)$ to its associated locally constant singular 0-cochain (recall that a singular 0-simplex is just the data of a point in the manifold), hence it's clearly injective.
- It is exact at \tilde{S}_K^0 iff $\ker d$ contains only the locally constant 0-cochains. This is the case, since by definition

$$df(\sigma) := f(\partial\sigma) = f(1) - f(0). \quad (2.29)$$

Since any singular 1-chain is by definition a continuous path we desume that $f \in \ker d$ iff f is constant on every connected subset of M , i.e. iff it is locally constant.

Remark 5 For $p \geq 1$ we are going to check exactness at the level of stalks. Let's fix p and $m \in M$, we want to check exactness of

$$(S_K^{p-1})_m \xrightarrow{d_m} (S_K^p)_m \xrightarrow{d_m} (S_K^{p+1})_m \quad (2.30)$$

at $(S_K^p)_m$. We have already proved that $d \circ d = 0$, hence also $d_m \circ d_m = 0$. This means we only have to prove that, given $f \in (S_K^p)_m$ s.t. $d_m f = 0$, there exist $g \in (S_K^{p-1})_m$ for which $f = dg$.

Since we are at the level of stalks we can consider the stalks of the presheaves: $(\tilde{S}_K^p)_m$. Here $d_m f = 0$ iff there exists a small enough neighborhood of m s.t. $d(f|_U) = 0$. Similarly $f = dg$ iff there exists a small enough neighborhood of m on which $d(g|_U) = f|_U$.

Remark 6 Moved by the above remark we are going to concentrate on an arbitrarily small neighborhood U of m . Since we are in a manifold M , which is locally *euchidean*, we can assume U to be the open unit ball in \mathbb{R}^d , where d is the dimension of M .

We are not going to prove exactness, but the stronger fact that the following complex

$$0 \rightarrow S^0(U, K) \xrightarrow{d} S^1(U, K) \xrightarrow{d} S^2(U, K) \xrightarrow{d} \dots \quad (2.31)$$

is homotopic to zero. In other words we have to construct a family of morphisms

$$h_p : S^p(U, K) \rightarrow S^{p-1}(U, K), \quad (2.32)$$

for all $p \geq 1$, s.t.

$$d \circ h_p + h_{p+1} \circ d = id_{S^p(U, K)}. \quad (2.33)$$

In fact, if we take $f \in S^p(U, K)$ s.t. $df = 0$ then, by the above formula, we get

$$f = id(f) = (d \circ h_p + h_{p+1} \circ d)f = d \circ h_p(f). \quad (2.34)$$

If we call $g := h_p(f)$, then we obtain the desired result: the complex is exact.

As just stated, for the following theorems,

$$U := \mathbb{B} = \{x \in \mathbb{R}^d \mid \|x\| < 1\}, \quad (2.35)$$

where d is the dimension of M , i.e. the generic simply connected open neighborhood in the manifold M .

Definition 2.15 Let $f \in S_K^p(U)$ and σ a singular $(p-1)$ -simplex in U . We define $h_p : S^p(U, K) \rightarrow S^{p-1}(U, K)$ by

$$h_p(f)(\sigma) := f(\tilde{h}_p(\sigma)), \quad (2.36)$$

where $\tilde{h}_p(\sigma)$ is the p -simplex that maps the origin, in Δ^p , to the origin, in U , and, for any $(a_1, \dots, a_p) \neq 0$, defined by

$$\tilde{h}_p(\sigma)(a_1, \dots, a_p) := \left(\sum_{j=1}^p a_j \right) \sigma \left(\frac{a_2}{\sum_{j=1}^p a_j}, \dots, \frac{a_p}{\sum_{j=1}^p a_j} \right). \quad (2.37)$$

In what follows we will consider \tilde{h}_p to be the linear extension of the above definition to the p -cochains, i.e. to $\tilde{h}_p : S_{p-1}(U) \rightarrow S_p(U)$.

Remark 7 With this definition we are only granting *continuity* to $\tilde{h}_p(\sigma)$, but not *differentiability*. This is one of the few points in which the theories for continuous and differentiable simplices diverge.

Lemma 2.16. $\partial \circ \tilde{h}_{p+1} + \tilde{h}_p \circ \partial = id$.

Proof. It is a simple – and tedious – computation: let σ be a singular p -simplex.

$$\begin{aligned} \left(\partial \circ \tilde{h}_{p+1} + \tilde{h}_p \circ \partial \right) (\sigma) &= \sum_{i=0}^{p+1} (-1)^i \left(\tilde{h}_{p+1}(\sigma) \right)^i + \sum_{i=0}^p (-1)^i \tilde{h}_p(\sigma^i) \\ &= \left(\tilde{h}_{p+1}(\sigma) \right)^0 + \sum_{i=1}^{p+1} (-1)^i \left\{ \left(\tilde{h}_{p+1}(\sigma) \right)^i - \tilde{h}_p(\sigma^{i-1}) \right\}. \end{aligned} \quad (2.38)$$

(2.39)

Let's now compute the individual terms.

Let $i > 1$, by definition the first summand is

$$\left(\tilde{h}_{p+1}(\sigma)\right)^i(a_1, \dots, a_p) = \tilde{h}_{p+1}(\sigma) \circ k_i^p(a_1, \dots, a_p) \quad (2.40)$$

$$= \tilde{h}_{p+1}(\sigma)(a_1, \dots, a_{i-1}, 0, a_{i+1}, \dots, a_p) \quad (2.41)$$

$$= \left(\sum_{j=1}^p a_j\right) \sigma \left(\frac{a_2}{\sum_{j=1}^p a_j}, \dots, \frac{a_{i-1}}{\sum_{j=1}^p a_j}, 0, \frac{a_{i+1}}{\sum_{j=1}^p a_j}, \dots, \frac{a_p}{\sum_{j=1}^p a_j} \right). \quad (2.42)$$

Again, by definition, the second summand is

$$\left(\tilde{h}_p(\sigma^{i-1})\right)(a_1, \dots, a_p) = \tilde{h}_p(\sigma \circ k_{i-1}^{p-1})(a_1, \dots, a_p) \quad (2.43)$$

$$= \left(\sum_{j=1}^p a_j\right) \left(\sigma \circ k_{i-1}^{p-1}\right) \left(\frac{a_2}{\sum_{j=1}^p a_j}, \dots, \frac{a_p}{\sum_{j=1}^p a_j} \right) \quad (2.44)$$

$$= \left(\sum_{j=1}^p a_j\right) \sigma \left(\frac{a_2}{\sum_{j=1}^p a_j}, \dots, \frac{a_{i-1}}{\sum_{j=1}^p a_j}, 0, \frac{a_{i+1}}{\sum_{j=1}^p a_j}, \dots, \frac{a_p}{\sum_{j=1}^p a_j} \right). \quad (2.45)$$

Let, now, $i = 1$. Still by definition, the first summand is

$$\left(\tilde{h}_{p+1}(\sigma)\right)^1(a_1, \dots, a_p) = \tilde{h}_{p+1}(\sigma) \circ k_1^p(a_1, \dots, a_p) \quad (2.46)$$

$$= \tilde{h}_{p+1}(\sigma)(0, a_1, \dots, a_p) \quad (2.47)$$

$$= \left(\sum_{j=1}^p a_j\right) \sigma \left(\frac{a_1}{\sum_{j=1}^p a_j}, \dots, \frac{a_p}{\sum_{j=1}^p a_j} \right). \quad (2.48)$$

The second summand, instead, is

$$\left(\tilde{h}_p(\sigma^0)\right)(a_1, \dots, a_p) = \tilde{h}_p(\sigma \circ k_0^{p-1})(a_1, \dots, a_p) \quad (2.49)$$

$$= \left(\sum_{j=1}^p a_j\right) \left(\sigma \circ k_0^{p-1}\right) \left(\frac{a_2}{\sum_{j=1}^p a_j}, \dots, \frac{a_p}{\sum_{j=1}^p a_j} \right) \quad (2.50)$$

$$= \left(\sum_{j=1}^p a_j\right) \sigma \left(\frac{a_1}{\sum_{j=1}^p a_j}, \dots, \frac{a_p}{\sum_{j=1}^p a_j} \right), \quad (2.51)$$

where, in the last equality, we used

$$1 - \frac{\sum_{j=2}^p a_j}{\sum_{j=1}^p a_j} = \frac{\sum_{j=1}^p a_j - \sum_{j=2}^p a_j}{\sum_{j=1}^p a_j} = \frac{a_1}{\sum_{j=1}^p a_j}. \quad (2.52)$$

All of the above terms are equal and, in (2.39), they appear with different sign. This allows us to simplify the first equality to:

$$\left(\partial \circ \tilde{h}_{p+1} + \tilde{h}_p \circ \partial\right)(\sigma) = \left(\tilde{h}_{p+1}(\sigma)\right)^0. \quad (2.53)$$

Let's compute this last term:

$$\left(\tilde{h}_{p+1}(\sigma)\right)^0(a_1, \dots, a_p) = \tilde{h}_{p+1}(\sigma) \circ k_0^p(a_1, \dots, a_p) \quad (2.54)$$

$$= \tilde{h}_{p+1}(\sigma) \left(1 - \sum_{j=1}^p a_j, a_1, \dots, a_p\right) \quad (2.55)$$

$$= \left(1 - \sum_{j=1}^p a_j + \sum_{j=1}^p a_j\right) \sigma(a_1, \dots, a_p) \quad (2.56)$$

$$= \sigma(a_1, \dots, a_p). \quad (2.57)$$

We, then, have the desired result! ■

Lemma 2.17. $d \circ h_p + h_{p+1} \circ d = id.$

Proof. From lemma 2.16 we know that

$$\partial \circ \tilde{h}_{p+1} + \tilde{h}_p \circ \partial = id. \quad (2.58)$$

In order to use this result to prove our statement we are gonna compute the action of $d \circ h_p(f) + h_{p+1} \circ d(f)$, for $f \in S^p(U, M)$, on an arbitrary p -chain σ :

$$(d \circ h_p + h_{p+1} \circ d) f(\sigma) = d \circ h_p f(\sigma) + h_{p+1} \circ df(\sigma) \quad (2.59)$$

$$= h_p f(\partial \sigma) + df(\tilde{h}_{p+1} \sigma) \quad (2.60)$$

$$= f\left(\tilde{h}_p(\partial \sigma)\right) + f\left(\partial(\tilde{h}_{p+1} \sigma)\right) \quad (2.61)$$

$$= f\left(\tilde{h}_p(\partial \sigma) + \partial(\tilde{h}_{p+1} \sigma)\right) \quad (2.62)$$

$$= f(\sigma). \quad (2.63)$$

■

Remark 8 As of this point we have checked everything we needed to in order to say that

$$0 \rightarrow K_M \rightarrow S_K^0 \xrightarrow{d} S_K^1 \xrightarrow{d} S_K^2 \xrightarrow{d} \dots \quad (2.64)$$

is a $\Gamma(M; -)$ -injective resolution for K_M . In fact it is exact, by the above lemmas, and it is $\Gamma(M; -)$ -injective, by 1.2, since each term is *fine*.

Our next task is to prove that the cohomology of the global sections of this resolution coincides with the singular cohomology on our manifold. This will be a lengthy construction, for which we'll outline the most interesting parts, leaving the lengthy computations to [War83].

Proposition 2.18. *Let $P \in \text{PSh}(k_M)$ be a presheaf that satisfies **S2**. Let $S := P^a$ its **sheafification** and $(P(M))_0$ the K -module defined by*

$$(P(M))_0 := \{s \in P(M) \mid s_m = 0 \ \forall m \in M\}. \quad (2.65)$$

Then the sequence of modules

$$0 \rightarrow (P(M))_0 \rightarrow P(M) \xrightarrow{\theta} S(M) \rightarrow 0 \quad (2.66)$$

is exact.

Proof. We need to check exactness at each point in the sequence.

- At $(P(M))_0$ is clear, since it is a submodule of $P(M)$ and the inclusion map is injective.
- At $P(M)$ we need to check that $\ker \theta = (P(M))_0$. Let $s \in P(M)$, then

$$s \mapsto \left\{ M \xrightarrow{\theta(s)} \prod_{m \in M} P_m \right\}, \quad (2.67)$$

where $\theta(s)(m) = s_m \in P_m$. It is clear that $s \in \ker \theta$ iff $s_m = 0$ for every $m \in M$, i.e. iff $s \in (P(M))_0$.

- At $S(M)$ we need to check that θ is surjective. We will do it explicitly: given $t \in S(M)$ we will construct $s \in P(M)$ s.t. $t = \theta(s)$. Recall that an element t in $S(M)$ is of the form

$$M \xrightarrow{t} \prod_{m \in M} P_m \text{ s.t. } t(m) \in P_m \forall m \in M \quad (2.68)$$

satisfying $\forall m \in M \exists V_m \overset{\text{open}}{\subset} M$ with $m \in V_m$ and $\exists s \in P(V_m)$ s.t. $s_x = t(x) \forall x \in V_m$. $\{V_m\}_{m \in M}$ clearly is an open cover of M , which is *paracompact*. We can extract a *locally finite* open refinement $\{U_\alpha\}_{\alpha \in \mathcal{A}}$ in which, for every $\alpha \in \mathcal{A}$, there exists $s_\alpha \in P(U_\alpha)$ s.t. $\theta(s_\alpha) = t|_{U_\alpha}$. Let, now, $\{V_\alpha\}_{\alpha \in \mathcal{A}}$ be a refinement s.t. $\bar{V}_\alpha \subset U_\alpha$. Let I_m be the collection of indices α for which $m \in \bar{V}_\alpha$ and W_m a neighborhood of m that satisfies:

- $W_m \cap \bar{V}_\alpha = \emptyset$ if $\alpha \notin I_m$,
- $W_m \subset \cap_{\alpha \in I_m} U_\alpha$, which is open and nonempty,
- $s_\alpha|_{W_m} = s_\beta|_{W_m}$ for any $\alpha, \beta \in I_m$, possible by the definition of stalk and sheafification.

Let $s_m \in W_m$ be the common image of the third point.

Consider, now, $n, m \in U$ s.t. $W_m \cap W_n \neq \emptyset$ and $p \in W_m \cap W_n$. From the first condition we know that $I_p \subset I_n \cap I_m$. Let $\alpha \in I_p$. By the third condition we have

$$s_m = s_\alpha|_{W_m} \quad \text{and} \quad s_\alpha|_{W_n} = s_n. \quad (2.69)$$

This implies

$$s_m|_{W_m \cap W_n} = s_\alpha|_{W_m \cap W_n} = s_n|_{W_m \cap W_n}. \quad (2.70)$$

We have just constructed a family $\{s_m\}_{m \in U}$ of sections of the presheaf that satisfy condition **S2**. We can hence patch them together to obtain an element $s \in P(M)$ s.t.

$$s|_{W_m} = s_m \quad (2.71)$$

for any m . Then, by definition, $\theta(s) = t$. ■

Theorem 2.19. *Let M be a **paracompact** manifold, K a PID, then for every $q \geq 0$*

$$H_\Delta^q(M) \simeq H^q(S_K^\bullet(M)). \quad (2.72)$$

Proof. Recall that

$$H_{\Delta}^q(M) = H^q(\tilde{S}_K^{\bullet}(M)). \quad (2.73)$$

This means that we have to prove

$$H^q(\tilde{S}_K^{\bullet}(M)) \simeq H^q(S_K^{\bullet}(M)). \quad (2.74)$$

By proposition 2.18, applied to the *presheaf* \tilde{S}_K^p , we have the following short exact sequence of K -modules

$$0 \rightarrow (\tilde{S}_K^p(M))_0 \rightarrow \tilde{S}_K^p(M) \xrightarrow{\theta} S_K^p(M), \quad (2.75)$$

for any $p \in \mathbb{N}$. It immediately follows that the associated sequence of complexes

$$0 \rightarrow (\tilde{S}_K^{\bullet}(M))_0 \rightarrow \tilde{S}_K^{\bullet}(M) \xrightarrow{\theta^{\bullet}} S_K^{\bullet}(M) \quad (2.76)$$

is exact. As usual, to this exact sequence of complexes we can associate the long cohomology sequence

$$\cdots \rightarrow H^q((\tilde{S}_K^{\bullet}(M))_0) \rightarrow H^q(\tilde{S}_K^{\bullet}(M)) \xrightarrow{\theta^q} H^q(S_K^{\bullet}(M)) \xrightarrow{\delta^q} H^{q+1}((\tilde{S}_K^{\bullet}(M))_0) \rightarrow \cdots \quad (2.77)$$

If we managed to prove that, for every q , $H^q((\tilde{S}_K^{\bullet}(M))_0) = 0$, then θ^q would be both mono and epi, hence (we are working in the category of K -modules) an iso.

Let's now check it explicitly:

q < 0: For $q < 0$ $(\tilde{S}_K^q(M))_0 = 0$, then also the associated cohomology module is,

q = 0: Note that, for $q = 0$, \tilde{S}_K^0 is actually a sheaf. In fact singular 0-simplices in M are just the assignment of a point in M . It follows that a singular 0-cochain on M is just a function from M to K . If it is determined in an open cover of M , then it is uniquely determined, recall that \tilde{S}_K^0 satisfies **S1**. This means that any element of $(\tilde{S}_K^0(M))_0$ is the zero element of $\tilde{S}_K^0(M)$.

q > 0: We won't explicitly give the whole construction for this case, but we will reference a technical lemma, found in [War83]. Let $\mathfrak{U} = \{U_i\}_{i \in I}$ be an open cover of M . We define $S_{\mathfrak{U}}^{\bullet}(M, K)$ to be the set of singular cochains f with values in K , defined only on \mathfrak{U} -small singular simplices. We say that a singular simplex is \mathfrak{U} -small iff its range is contained in U_i for one $U_i \in \mathfrak{U}$. Clearly to any element $f \in S^p(M, K)$ one can associate an element in $S_{\mathfrak{U}}^p(M, K)$ given by the restriction of f to \mathfrak{U} -small singular p -simplices only. This association gives rise to a surjective morphism of cochains

$$j_{\mathfrak{U}} : S^{\bullet}(M, K) \rightarrow S_{\mathfrak{U}}^{\bullet}(M, K). \quad (2.78)$$

Denoting with $K_{\mathfrak{U}}^{\bullet}$ the kernel of this morphism, we get a short exact sequence of complexes

$$0 \rightarrow K_{\mathfrak{U}}^{\bullet} \rightarrow S^{\bullet}(M, K) \xrightarrow{j_{\mathfrak{U}}} S_{\mathfrak{U}}^{\bullet}(M, K) \rightarrow 0. \quad (2.79)$$

From this we obtain a corresponding long exact cohomology sequence

$$\cdots \rightarrow H^q(K_{\mathfrak{U}}^{\bullet}) \xrightarrow{i_{\mathfrak{U}}^q} H^q(S^{\bullet}(M, K)) \xrightarrow{j_{\mathfrak{U}}^q} H^q(S_{\mathfrak{U}}^{\bullet}(M, K)) \xrightarrow{\delta^q} H^{q+1}(K_{\mathfrak{U}}^{\bullet}) \rightarrow \cdots \quad (2.80)$$

We want to prove that the map $j_{\mathfrak{U}}^q$ induced in the cohomology sequence is an iso for any q . If it were the case, it would follow that, for every q

$$H^q(K_{\mathfrak{U}}^{\bullet}) = 0. \quad (2.81)$$

In fact, since $j_{\mathfrak{U}}^q$ induces an iso, it means that $\ker \delta^q = H^q(S^{\bullet}(M, K))$ and $\operatorname{im} \delta^q = 0$. But $\operatorname{im} \delta^q = \ker i_{\mathfrak{U}}^q$, i.e. this last map is a mono. Analogously it can be proved that $\operatorname{im} i_{\mathfrak{U}}^q = 0$, hence that $H^q(K_{\mathfrak{U}}^{\bullet}) = 0$.

Following [War83] this statement is proved using the results of lemma 2.20, below. In fact, by (2.84), $j_{\mathfrak{U}}$ induces surjections of the cohomology modules. By (2.85) $k \circ j_{\mathfrak{U}}$ induces the identity on cohomology, which means that $j_{\mathfrak{U}}$ must induce injections. Putting it all together $j_{\mathfrak{U}}$ induces isomorphisms.

With this fact we want to prove that $H^q((\tilde{S}_K^{\bullet}(M))_0) = 0$. Consider $f \in (\tilde{S}_K^q(M))_0$ s.t. $df = 0$. By definition of $(\tilde{S}_K^q(M))_0$ we know that $f_m = 0$ for every $m \in M$, hence that for any m there exist an open neighborhood U_m of m s.t. f maps every singular q -chain with range in U_m to 0. This means that there exists an open cover \mathfrak{U} of U consisting of sufficiently small sets, for which $f \in K_{\mathfrak{U}}^q$. From (2.81) we know that $\exists g \in K_{\mathfrak{U}}^{q-1} \subset (\tilde{S}_K^{q-1}(M))_0$ s.t. $f = dg$, i.e.

$$H^q((\tilde{S}_K^{\bullet}(M))_0) = 0. \quad (2.82)$$

■

Lemma 2.20. *Let M be a **paracompact** manifold,*

$$j_{\mathfrak{U}} : S^{\bullet}(m, K) \rightarrow S_{\mathfrak{U}}^{\bullet}(M, K) \quad (2.83)$$

be the inclusion map defined in the above proof. Then there exists a map

$$k : S_{\mathfrak{U}}^{\bullet}(M, K) \rightarrow S^{\bullet}(M, K) \quad (2.84)$$

s.t. $j_{\mathfrak{U}} \circ k = \operatorname{id}$ and homotopy operators $h^p : S^p(M, K) \rightarrow S^{p-1}(M, K)$ s.t.

$$h^{p+1} \circ d + d \circ h^p = \operatorname{id} - k^p \circ j^p. \quad (2.85)$$

The proof for this lemma consists in explicitly constructing these maps. It is rather long and boring. Also we have used almost the same notation as [War83], so it can be checked there – in section 5.32 – without any issue.

Now, before moving onto the main result of this work, let's define a last concept:

Definition 2.21: Integration of forms over singular simplices.

Let M be a *differentiable manifold*, σ a *differentiable* singular p -simplex in U and ω a *continuous* p -form also defined on $U \stackrel{\text{open}}{\subset} M$.

- If $p = 0$ σ is just the data of a point, $\sigma(0)$, in U and ω just a continuous (differentiable) function. We define the integral of ω over σ to be

$$\int_{\sigma} \omega := \omega(\sigma(0)). \quad (2.86)$$

- If $p \geq 1$ σ extends to a smooth map from a neighborhood of Δ^p into U . This implies that the pullback of ω is defined in a neighborhood of Δ^p . We can then compute its integral on the p -simplex. From this we define the integral of ω over σ to be

$$\int_{\sigma} \omega := \int_{\Delta^p} \sigma^*(\omega). \quad (2.87)$$

We linearly extend those definitions to chains: let $c = \sum_i a_i \sigma_i$, then

$$\int_c \omega := \sum_i a_i \int_{\sigma_i} \omega. \quad (2.88)$$

Let's now state (without proving it) a famous theorem that links the *exterior differential* for forms, with the *boundary operator* for simplices.

Theorem 2.22 (Stokes' theorem). *Let M be a differentiable manifold, $U \subset^{open} M$, c a p -chain in U , with $p \geq 1$, and ω be a smooth $(p-1)$ -form, still defined on U . Then*

$$\int_{\partial c} \omega = \int_c d\omega. \quad (2.89)$$

If you are interested in the proof you can find it in [War83], section 4.7.

Remark 9 Let us define, for $p \geq 0$, the following homomorphism

$$k^p : \Omega^p(U) \rightarrow S^p(U, \mathbb{R}) \quad (2.90)$$

by setting, for $\omega \in \Omega^p(U)$ and σ a differentiable singular p -simplex,

$$k^p(\omega)(\sigma) := \int_{\sigma} \omega. \quad (2.91)$$

Then *Stokes' theorem* makes this a morphism of complexes

$$k^\bullet : \Omega^\bullet(U) \rightarrow \tilde{S}_\mathbb{R}^\bullet(U). \quad (2.92)$$

We, in fact, need to check the commutativity of the following diagram

$$\begin{array}{ccc} \Omega^p(U) & \xrightarrow{d} & \Omega^{p+1}(U) \\ k^p \downarrow & & \downarrow k^{p+1} \\ S^p(U, \mathbb{R}) & \xrightarrow{d} & S^{p+1}(U, \mathbb{R}) \end{array}, \quad (2.93)$$

i.e. that $k^{p+1} \circ d = d \circ k^p$. Let σ and ω be as above, then

$$(d \circ k^p)(\omega)(\sigma) = k^p(\omega)(\partial\sigma) = \int_{\partial\sigma} \omega \quad (2.94)$$

$$= \int_{\sigma} d\omega = k^{p+1}(d\omega)(\sigma) = (k^{p+1} \circ d)(\omega)(\sigma). \quad (2.95)$$

This, in turn, gives a morphism of the respective cohomologies:

$$k^q : H^q(\Omega^\bullet(U)) \rightarrow H^q(S^\bullet(U, \mathbb{R})), \quad (2.96)$$

called the *de Rham homomorphism*. Moreover, since this family of morphisms clearly commutes with restrictions, it also induces a family of morphisms of *presheaves*:

$$k^p : \Omega^p \rightarrow \tilde{S}_\mathbb{R}^p. \quad (2.97)$$

Which, as before, gives a morphism of complexes of presheaves

$$k^\bullet : \Omega^\bullet \rightarrow \tilde{S}_\mathbb{R}^\bullet. \quad (2.98)$$

We are finally ready to state and prove our final result:

Theorem 2.23 (De Rham). *Let M be a smooth paracompact manifold. Then, for every q , there exists an isomorphism $H_{DR}^q(M) \simeq H_{\Delta}^q(M)$. Moreover this isomorphism is given by integration of forms on singular chains.*

Proof. When taking the PID $K = \mathbb{R}$ to be the field of real numbers, we have constructed two $\Gamma(M, -)$ -injective resolutions for the sheaf of constant functions in \mathbb{R} :

$$0 \rightarrow \mathbb{R}_M \rightarrow \Omega^0 \xrightarrow{d} \Omega^1 \xrightarrow{d} \Omega^2 \xrightarrow{d} \dots, \quad (2.99)$$

and

$$0 \rightarrow \mathbb{R}_M \rightarrow S_{\mathbb{R}}^0 \xrightarrow{d} S_{\mathbb{R}}^1 \xrightarrow{d} S_{\mathbb{R}}^2 \xrightarrow{d} \dots \quad (2.100)$$

Moreover we know that $\mathbf{Mod}(\mathbb{R}_M)$ has enough *injectives* and $\Gamma(M, -)$ is left exact. It follows that we can compute the q -th right derived functor of $\Gamma(M, -)$ by computing the $(q - 1)$ -th cohomology of these two sequences. As a consequence we have the following isomorphism

$$H_{DR}^q(M) \simeq H^q(S_{\mathbb{R}}^{\bullet}(M)). \quad (2.101)$$

By theorem 2.19 we know that $H_{\Delta}^q(M) \simeq H^q(S_{\mathbb{R}}^{\bullet}(M))$, which gives the desired isomorphism

$$H_{DR}^q(M) \simeq H_{\Delta}^q(M). \quad (2.102)$$

We are only left to prove that this isomorphism is the above defined *de Rham isomorphism*. In fact it gives rise to the following morphism of complexes

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{R}_M & \longrightarrow & \Omega^0 & \xrightarrow{d} & \Omega^1 \xrightarrow{d} \dots \\ & & \downarrow id & & \downarrow k^0 & & \downarrow k^1 \\ 0 & \longrightarrow & \mathbb{R}_M & \longrightarrow & \tilde{S}_{\mathbb{R}}^0 & \xrightarrow{d} & \tilde{S}_{\mathbb{R}}^1 \xrightarrow{d} \dots \end{array} \quad (2.103)$$

This, in turn, by *sheafification* becomes

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{R}_M & \longrightarrow & \Omega^{\bullet} & & \\ & & \downarrow id & & \downarrow (k^{\bullet})^a & & \\ 0 & \longrightarrow & \mathbb{R}_M & \longrightarrow & S_{\mathbb{R}}^{\bullet} & & \end{array} \quad (2.104)$$

Since both of those are $\Gamma(M, -)$ -injective resolutions, $(k^{\bullet})^a$ induces a quasi isomorphism, i.e. isomorphisms at the level of cohomologies, between the following complexes

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{R}_M & \longrightarrow & \Omega^0(M) & \xrightarrow{d} & \Omega^1(M) \xrightarrow{d} \dots \\ & & \downarrow id & & \downarrow (k^0)^a & & \downarrow (k^1)^a \\ 0 & \longrightarrow & \mathbb{R}_M & \longrightarrow & S_{\mathbb{R}}^0(M) & \xrightarrow{d} & S_{\mathbb{R}}^1(M) \xrightarrow{d} \dots \end{array} \quad (2.105)$$

Moreover we have proved that F -injective resolutions differ by a unique isomorphism, at the level of cohomology, hence the canonical isomorphism in equation (2.101) corresponds with the one induced by $(k^{\bullet})^a$. This, in turn, means that the following commutative diagram

$$\begin{array}{ccc} \Omega^{\bullet}(M) & & \\ k^{\bullet} \downarrow & \searrow (k^{\bullet})^a & \\ \tilde{S}_{\mathbb{R}}^{\bullet}(M) & \xrightarrow{\theta} & S_{\mathbb{R}}^{\bullet}(M) \end{array}, \quad (2.106)$$

where θ is the natural morphism $F \xrightarrow{\theta} F^a$, gives rise to the following commutative diagram of cohomologies

$$\begin{array}{ccc} H_{DR}^\bullet(M) & & \\ \downarrow k^\bullet & \searrow \sim & \\ H_\Delta^\bullet(M) & \xrightarrow{\sim} & H^\bullet(S_{\mathbb{R}}^\bullet(M)) \end{array} . \quad (2.107)$$

Since the other two are isomorphisms, also k^\bullet is. Moreover it is the canonical (and unique) one, we have found before. \blacksquare

References

- [War83] F.W. Warner. *Foundations of Differentiable Manifolds and Lie Groups*. Graduate Texts in Mathematics. Springer, 1983.