

A review of computational and mathematical modeling contributions to our understanding of *Mycobacterium tuberculosis* within-host infection and treatment

Denise Kirschner¹, Elsje Pienaar^{1,2}, Simeone Marino¹ and Jennifer J. Linderman²

Abstract

Tuberculosis (TB) is an ancient and deadly disease characterized by complex host-pathogen dynamics playing out over multiple time and length scales and physiological compartments. Mathematical and computational modeling can be used to integrate various types of experimental data and suggest new hypotheses, mechanisms, and therapeutic approaches to TB. Here, we offer a first-time comprehensive review of work on within-host TB models that describe the immune response to infection, including the formation of lung granulomas. The models include systems of ordinary and partial differential equations and agent-based models as well as hybrid and multi-scale models that are combinations of these. Many aspects of *Mycobacterium tuberculosis* infection, including host dynamics in the lung (typical site of infection for TB), granuloma formation, roles of cytokine and chemokine dynamics, and bacterial nutrient availability have been explored. Finally, we survey applications of these within-host models to TB therapy and prevention and suggest future directions to impact this global disease.

Addresses

¹ Department of Microbiology and Immunology, University of Michigan Medical School, Ann Arbor, MI, USA

² Department of Chemical Engineering, University of Michigan, Ann Arbor, MI, USA

Corresponding authors: Kirschner, Denise; Linderman, Jennifer J (kirschne@umich.edu), (linderman@umich.edu)

Current Opinion in Systems Biology 2017, 3:170–185

This review comes from a themed issue on **Mathematical modelling (2017)**

Edited by **Martin Falcke**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 22 May 2017

<http://dx.doi.org/10.1016/j.coisb.2017.05.014>

2452-3100/© 2017 Published by Elsevier Ltd.

Keywords

Systems biology, Simulations, Granuloma, Antibiotic treatment, Sensitivity analysis, Agent-based model.

Introduction

Tuberculosis (TB) is a disease caused by infection with *Mycobacterium tuberculosis* (Mtb). An estimated one fourth of the world population has latent Mtb infection, and in

2014 TB was responsible for 1.5 million deaths [1,2]. There is no broadly efficacious vaccine, and drug treatment is complicated by a necessarily long course of multiple antibiotics (at least 6 months of chemotherapy) in part to prevent development of resistance. This long time course contributes to toxicity and patient non-compliance, which can lead to relapse. The infection process typically occurs after long and repeated exposure to infected individuals via inhalation of bacteria into lungs [3]. Once infection is established in the lung, macrophages take up bacteria and a complex and dynamic process follows that leads to granuloma formation. Granulomas are spherical collections of cells, bacteria and necrotic tissue that serve to both immunologically contain and physically constrain Mtb. However, granulomas also provide an environment that allows bacteria to persist, in some cases for the lifetime of the host. As granulomas are the battleground for host-pathogen interactions during infection, understanding these structures is key to developing intervention strategies [4–6].

A long-standing view of TB describes it as latent, asymptomatic infection or active TB disease. Patients with latent infection are known to sometimes progress to active TB disease, in some cases decades after the initial infection [7]. The challenge with this dichotomous view is that it does not help identify which patients are at highest risk of progression to active TB disease [8]. However, Mtb infection is now understood to result in a spectrum of patient outcomes ranging from natural cure by innate immune mechanisms to active TB disease [9]. Patients can move along this spectrum throughout their lives. Overall, approximately 10% of individuals infected with Mtb progress to active TB at some point in their lifetime, driven by a number of factors including aging, co-morbidities or immune-suppression [10]. If left untreated, active TB leads to death in about half of those infected in an average of 3 years [11]. There are ongoing efforts to find biomarkers that can identify patients at highest risk of developing active TB disease for preventive therapy [12–14].

Systems biology approaches can be critical to advance our understanding and treatment of TB [15–22]. There is a need to integrate data across multiple time and

length scales, incorporating, for example, knowledge about how bacteria can alter infected host cells, how trafficking of immune cells to lungs influences granuloma formation, how infection can spread to lymph nodes, and how vaccines and antibiotics impact disease progression [23–29]. Mathematical and computational models can be used to integrate these different types of data, as well as to bridge between experimental measurements, better understand hypothesized mechanisms, run virtual experiments (e.g. virtual clinical trials, virtual deletions/depletions) when animal experiments are too expensive or difficult, interpret data, and offer new explanations for observed phenomena.

There have been three main foci of computational models relevant to TB. First, early and continuing extensive mathematical modeling centered on the epidemiology of TB. A comprehensive review of this literature by the TB Modelling Alliance Consortium can be found at [<http://tb-mac.org/Resources/Resource/4>]. These models describe the population-level dynamics of TB under different scenarios. Second, models were developed to explore mycobacterium growth, drug-resistance development, metabolism, and adaptive response to stress conditions such as hypoxia, nutrient starvation and intracellular survival [18–22,30–33]. Finally, we and others have performed extensive modeling that describes within-host interactions in Mtb infection. This is the first review on this topic, and thus it focuses solely on this latter body of literature, exploring within-host dynamics of TB infection (Table 1). Much of this work was born out of the authors' studies, so the majority of references represent our efforts. In addition, the search protocol for all other references and relevant authors working in this area were found by PubMed and ScienceDirect search term “(mathematical model OR computational OR *in silico* OR systems biology) AND mycobacterium tuberculosis AND within host AND host-pathogen NOT epidemiology NOT metabolic”. The resulting list of papers was manually filtered to exclude experimental, -omics (big

data), epidemiology and structural biology papers. The goal of this review is to stimulate more work in this area and provide a comprehensive resource for both education and research purposes. Efforts at this within-host level, which include necessary biological detail and a range of relevant length (molecular to host) and time (minutes to lifetime) scales, may be especially useful in understanding both immune mechanisms and interventions to treat disease [34,35].

Multiple physiological compartments are relevant to TB (Figure 1). Antigen-presenting cells from the lungs travel to lymph nodes (LNs), ultimately resulting in recruitment of T cells to lungs where formation of granulomas occurs. Granulomas have a characteristic composition, typically with a central core of caseous necrosis surrounded by a ring of macrophages, neutrophils, giant cells, and an outer ring of lymphocytes including T and B cells. Bacteria are either trapped inside the caseous center (non-replicating bacteria), within macrophages (intracellular bacteria) or in extracellular spaces. Multiple granulomas form in the lungs following infection in adults and granulomas have independent trajectories leading to outcomes ranging from sterilizing to controlled bacteria growth without clearance to uncontrolled bacteria growth [36]. The collective outcome of all granulomas likely determines whether the host's disease trajectory (e.g. active disease).


Despite much experimental research on TB over the past century (Leeuwenhoek identified Mtb using the first microscope in the 1700s), there are many features of infection that are not well understood and we still lack a broadly efficacious vaccine [28]. Antibiotics are an effective treatment for drug sensitive disease, but drug-resistant TB is more difficult, and in some cases not possible, to treat [90]. Mathematical and computational modeling can assist in better understanding this complex system, and a systems biology approach affords the pairing of multiple modalities, such as *in vitro* and *in vivo*

Table 1

Mathematical and computational models describing aspects of within-host infection dynamics during Mtb infection. Models can be classified by physiological scales (rows) and applications (columns).

	Antibiotic treatment	Vaccines	Biomarker discovery	Drug target identification	Molecular mechanisms	Cellular mechanisms	Pathogen mechanisms	Other immune mechanisms	Virtual clinical trials
Whole host models	[37]		[38]		[39]	[39,40]		[41]	
Whole lung models	[42–44]				[45]	[44,46–50]	[45,50]	[41,51]	
Granuloma models	[52–55]			[56,57]	[56,58–63]	[64–71]	[57,72,73]	[74]	[63]
Lung/Lymph node models			[38]		[75,76]	[76–81]	[77]		
Lymph node/Blood models		[82]							
Single cell models					[83–87]		[88]	[89]	

Figure 1



Physiological compartments involved in *Mtb* infection dynamics. Infection begins when *Mtb* are inhaled into the lung, taken up by macrophages, and begin multiplying. Antigen-presenting cells (including dendritic cells, DCs) travel through afferent lymphatics (AL) to lung-draining lymph nodes (LNs) where priming of naïve T cells occurs. This process initiates the *Mtb*-specific immune response generating both CD4+ and CD8+ T cells. These T cells have phenotypes that include activated, effector, effector memory (EM), and central memory (CM). Effector and memory T cells flow out of the LN into blood to traffic to the lung to participate in granuloma formation and function. Multiple cytokines also play a role in granuloma formation and infection control. Granulomas are dynamic, can last the lifetime of a host, and fall into major categories of solid cellular, caseous (a central core of necrotic material), fibrotic (a cuff of fibrosis surrounds the granuloma) and disseminating (increasing in both size and bacteria). Orally administered antibiotics are absorbed into the blood and distribute to lungs (pharmacokinetics). Most human data reflects the blood compartment, as it is the most easily sampled compartment, in the hope that blood data may be useful for predicting disease progression.

studies together with modeling and computation. Many of the modeling concepts, approaches and tools are summarized in [BOX](#).

Overview of in-host computational models

In-host computational models for *Mtb* infection focus on events occurring in specific host locations – the lung, lymph nodes, and blood. At first, data were available only at higher levels of resolution (e.g. health of host, total bacteria and immune cells in the lung), and early models reflected that. More detailed data, including lung PET/CT, cytokine concentrations, and activation of signaling networks, are now increasingly available. [Table 2](#) gives an overview of the types of data now available to build, validate, and calibrate in-host computational models. The data presented are not exhaustive, but emphasize the types and quality that are available. We only show data derived for the lung compartment, but other datasets in TB from both blood (humans and non-human primates-NHPs) and lymph nodes (from NHPs) are available as well [12,13,38,91–96]. Here we give a brief overview of model evolution, including

approaches that were developed to capture dynamics at the molecular, cellular/multicellular, and organ/tissue scales. Model formulations include continuous models, discrete models, and hybrid models that combine both ([BOX](#)). These models provide the platforms on which to examine both disease mechanisms but also, ultimately, disease therapy and prevention.

Models that capture the immune response to *Mtb* infection in the whole lung

Modeling host-immune infection dynamics of TB began with models bridging the rich literature of modeling HIV-1 infection to co-infection with *Mtb* [44,49]. TB is a primarily lung-localized infection, whereas HIV-1 is a systemic infection. The first host immune response models of TB were ODE-based virtual human models of the lung that qualitatively and quantitatively characterized the cellular and cytokine network known to occur during TB infection [46,48,50]. It was appropriate to start with ODEs ([BOX](#)) as the calibration data were derived from humans and also whole lungs isolated from mice, as no granuloma spatial data were yet available.

BOX. Computational model concepts, approaches and tools.

Agent-based model (ABM): A class of computational models that allows simulation of stochastic actions and interactions of discrete agents, with a view to identify system behavior. The TB ABMs described here incorporate spatial and temporal aspects of granuloma formation and function. Incorporation of stochastic elements in TB models allows for variability in outcomes, as seen in biology.

Continuous models: Mathematical models, including ordinary differential equations (ODEs) and partial differential equations (PDEs), formulated to capture average behaviors of collections of items (e.g. populations) over time. In TB, overall lung dynamics and receptor-ligand binding and signaling have been described with ODEs, while antibiotic diffusion within a granuloma has been described by PDEs.

Diffusion: Random motion of molecules. In TB models, cytokines and antibiotics are examples of molecules that diffuse in tissue.

Discrete models: In contrast to continuous models, discrete models track the behavior of individual elements over time. Model types that capture these include agent-based models and Markov models. In TB, ABMs are used to track granuloma formation in lungs.

Emergent behavior: Interactions of entities may produce a system-wide behavior that is not readily predictable. The formation of a granuloma from immune cell and bacterial interactions in lungs is an emergent phenomenon.

Hybrid model: Mathematical or computational model that is composed of at least two model formulations, e.g. differential equations and difference equations. Many of the TB models incorporate both differential equations to track diffusion of molecules and ABM components to track cellular interactions.

Model calibration: The process of using data to identify parameter values in a model to yield outputs that align with biological data. For TB, model outputs may be compared to, for example, mouse, non-human primate and human data to calibrate a model.

Model validation: Performing studies to test whether model outputs align with unique biological data or known behaviors. Data used to calibrate and validate models must be distinct. In TB, model validation may test whether model output agrees with data from primate systems, for example.

Multi-compartment model: A model that is comprised of linked sub-models representing different physiological compartments. TB models may include lung, lymph node, lymphatics and blood compartments.

Multi-scale model: A model that is comprised of linked sub-models representing different physiological scales in space or time. TB models may include molecular, cellular, and tissue level interactions across minutes to years.

Parameter estimation: Identifying model parameter values using a variety of techniques, including deriving values directly from data, using uncertainty analysis, or using model calibration to estimate them.

Pharmacokinetic/pharmacodynamic (PK/PD) modeling: PK refers to quantifying the movement of drugs throughout body compartments. PD refers to the actions of drugs in the body. TB models that describe antibiotic delivery and action include PK/PD equations.

Uncertainty and sensitivity analysis: Uncertainty analysis (UA) is performed to investigate uncertainty in model outputs that is generated from uncertainty in parameter inputs. Sensitivity analysis (SA) assesses how variations in model outputs can be apportioned, qualitatively or quantitatively, to different input sources. In TB modeling, these techniques guide parameter estimation, identification of critical mechanisms and drug targets, and model fine- and coarse-graining.

Tuneable resolution: Tuneable resolution proposes that multi-scale models should be built with multiple levels of resolution, so that a fine- or coarse-grained version can be employed with user discretion, allowing adjustment of the level(s) of resolution specific to a question, an experiment, a particular use, or a change in perspective or scale of interest. In our TB models, for example, the level of detail around tumor-necrosis factor (TNF) interactions can be adjusted according the question being asked.

Virtual clinical trial (VCT): *In silico* simulation of a clinical trial, including patient variability as well as treatment protocols. VCTs in TB have included anti-TNF treatment of a population to predict outcomes.

Virtual Deletion/Depletions: *In silico* simulation of a gene knockout or system-wide depletion of a molecule or cell. Multiple simultaneous virtual deletions or depletions, although difficult to perform in animal studies, are easily performed *in silico*.

Table 2

A sampling of experimental datasets on immune responses to Mtb in lungs are listed. Key components of the TB mathematical models have been determined in animal models (mouse, guinea pigs, rabbits, zebrafish, non-human primates) and with human data where available. The type of data derived for each component is listed. Experiments in mouse (knockouts), monkey and human (depletion by antibody or natural mutations) are indicated and can be performed *in silico* for model validation. Experimental techniques used to generate and validate data on each component are listed.

Immune factor	Animal model (mouse, guinea pig, rabbit, zebrafish) data	Monkey data (and human where available)	Knockouts or depletions	References and techniques used in experiments
CD4 T cells, CD8 T cells	Numbers Phenotypes Mtb specific Dynamics	Numbers Phenotypes Mtb specific Dynamics Spatial	Mouse Monkey	[97–114] Flow cytometry, ELISPOT, IHC, ISH, <i>in vivo</i> and <i>in vitro</i> cytotoxicity assays
Macrophages	Numbers Phenotypes Spatial	Numbers Phenotypes	Mouse, Monkey	[99,112,115–118] Flow cytometry, IHC, ISH
Dendritic cells	Numbers	Numbers	Mouse, Monkey	[119–122] Flow cytometry, ISH
Regulatory T cells	Numbers	Numbers Dynamics	Mouse Monkey Human	[123–127] Flow cytometry, IHC
Cytokines, e.g. IFN- γ , TNF, IL-12, TGF-B, IL-10	Cellular source Levels	Cellular source Levels	Mouse Monkey	[97,99,113–115,128–132] Flow cytometry, ELISPOT, Luminex, IHC, ISH
Chemokines	Cellular source Levels	Cellular source Levels Spatial expression	Mouse, Monkey	[99,115,116,133–140] Flow cytometry, IHC, ISH, Luminex
Mtb bacilli	Numbers Spatial Dynamics	Numbers Sites	Mouse Monkey Human	[99] Culture, IHC, AFB, ISH
Granuloma	Morphology Composition Formation (ZF) Dynamics	Morphology Composition Dynamics		[4,99,112,135,141–145] Histology, IHC, ISH, flow cytometry, live imaging, PET/CT imaging

The known dynamics of human immune responses in lungs were included. For example, in mice TH2 cells make IL-10, but in humans many cell types produce it and this was included in the model. These original models predicted specific cell and cytokine factors that determine the ability of a host to control infection. Interestingly, the model indicated that even if infection is controlled a balance of pro- and anti-inflammatory signals is required to prevent excessive tissue damage. These models did not include the role of CD8+ T cells; however, the models predicted that a cell type that added to cytotoxicity of infected cells and was a producer of mediators was missing. Later data showed that indeed CD8+ T cells are important in TB, and a number of models emerged to understand their role [47,48].

Computational models of granulomas in Mtb-infected lungs

Although most early datasets were derived from whole lungs of mice, cellular and molecular interactions within individual lung granulomas became increasingly understood as key to understanding TB infection. Granulomas are structures that evolve in both space and time, and so

ODEs would not suffice to represent them. Concomitantly, new spatial data on granulomas were beginning to emerge from an NHP system, allowing us to visualize 2-D slices of granulomas at disease endpoints (see Table 2). We tried multiple mathematical formulations to capture the spatial evolution of granulomas in lungs. The first spatial model of granulomas developed was an agent-based model (ABM) [64,69]; formation of the granuloma was an emergent behavior in the model. PDE models [58,68,71] and other formulations such as meta-population models [70] were also used and compared [146]. One model examined early events in mice, including neutrophils and the role they play in granuloma formation [66], while others examined the role of the bacterial burst size on granulomas [67]. The influence of lung structure also indicated an important role for neutrophils [74].

Receptor-ligand dynamics (described by ODEs) [147] and molecular diffusion (PDEs) were next integrated into the ABM framework to form a hybrid single granuloma model, and thus the granuloma models became explicitly multi-scale (BOX) [59,61]. The general

computational framework is reviewed in Ref. [148] and may sometimes rely on development of individual level models, e.g. single cell models [58,83,86,87,149], for incorporation into a multi-scale framework. These models were supported and spurred by novel NHP data, including data on individual granulomas, as well as questions from the experimental literature surrounding the roles of specific pro- and anti-inflammatory factors. In particular, TNF and IL-10 dynamics were captured by including cell secretion, diffusion within the granuloma, receptor/ligand binding and internalization, and signaling leading to particular cellular outcomes [59–62]. The observation that some anti-TNF therapies (for unrelated autoimmune disease) were more likely to cause TB reactivation was interpreted as the result of differences in TNF/drug binding kinetics and permeabilities, both in an ODE setting [63] and in an ABM platform [56,150]. IL-10 was predicted to play an important role in controlling the early immune response to *Mtb* at the granuloma scale, limiting host-inflammation-induced caseation and preventing granuloma sterilization [61].

Two studies [65,151] address regulation of macrophage microenvironment-specific polarization (as pro- or anti-inflammatory, often termed M1 and M2 respectively) and effects on granuloma outcome. A new metric, the macrophage polarization ratio, captures how cytokine signaling translates into polarization of individual macrophages and collectively drives the emergence of an inflammation phenotype at the granuloma scale. Results suggest that the NF- κ B signaling pathway is a viable therapeutic target to promote M1 polarization early during infection and to improve outcome. Dynamics of M1 and M2 macrophages have also been modeled using ODEs to understand how switching time (e.g. from an M2 to an M1-dominated environment) can impact infection outcome (measured as total CFU in the whole lung) [75,81]. Finally, a PDE model exploring host-directed therapies in the context of M1/M2 was recently explored [71]. Taken together, these more detailed models further support the view that there is a complex balance of pro- and anti-inflammatory mechanisms that attempt to “tune” each granuloma to contain infection but prevent its elimination.

Mtb utilize oxygen and carbon sources such as lipids. Nutrient availability can be predicted within single granuloma models, with various levels of complexity in how bacteria are described [57,72,73]. These models predict a continuous fluctuation in bacterial numbers, bacterial phenotypes and host responses throughout years of persistent infection [57]. They also address the contribution of granuloma structure to hypoxia [73] and bacterial phenotypic changes [72]. The bacterial focus of these models illuminates the constant feedback between bacterial adaptation and host immune responses that result in persistent infection.

Two compartment (lung and lymph node) models

Because cells move between lymph nodes and the lung, models also connect these compartments mathematically. Several 2-compartment ODE models capture cellular activation and priming in the lung-draining LN as well as trafficking between lungs and LN [76–78,80]. Additional models examined the link between lung granulomas and LNs [75,79]. These studies identified that mechanisms relevant to CD4+ T cell priming and trafficking might be manipulated to prevent establishment of *Mtb* infection or to clear bacteria from an established granuloma.

Attempting to scale in-host models to the whole host

Modeling an entire host can help bridge understanding from data and models derived at a smaller scale (e.g. single granulomas) to data derived from the host scale, including disease manifestations. Such models may also be critical to identifying new therapeutic interventions and predicting results of clinical trials. These models can be challenging from both the perspective of computational cost and capturing all relevant mechanisms. Approaches vary using either a whole lung scaling of granulomas to a whole host level model. A Boolean network model representing a whole host integrated multiple cellular and molecular mechanisms. They identified the importance of specific cytokines and phagosome functions [39]. In other work, multiple independently-evolving granulomas were linked together [38,152], created virtual NHPs that matched NHP datasets. A 3-compartment model (lung, LN and blood) captured formation of independent granulomas in lungs as well as T cell profiles in blood, and potential biomarkers that can predict infection outcome were identified. Rather than match datasets, multiple granulomas were tracked and an improved representation of DC trafficking was implemented [40]. Early events after *Mtb* infection were shown to be critical to establishing a timely and effective response. Finally, in a coarse-grained model, granulomas (population I, infected) were considered as variables that could spread to multiple lymph nodes (population S, susceptible) leading to infection there, with each tissue having the chance to resolve and clear infection (population R, recovered) using an SIR formalism [41]. This work showed that in early phases of infection granulomas suppress infection by providing additional antigens to the site of immune priming (in LNs); however, this induces a more rapid reactivation at later stages by disrupting immune responses.

Computational and mathematical modeling applications to TB

The mathematical and computational models described above have helped us better understand the bacterial and immune dynamics relevant to TB. They have begun to elucidate a critical aspect of infection, namely why some individuals develop clinically active disease and others latent disease. Confirmed by further wetlab

studies, we now understand containment of infection with a granuloma from the perspective of an appropriate balance of pro- and anti-inflammatory factors. Models can also help improve and direct how to prevent, treat, and monitor (via biomarkers) TB.

Within-host antibiotic treatment dynamics of TB

Sufficient exposure of bacteria to antibiotics is required for effective TB treatment [42,153]. Many studies investigate the pharmacokinetics (PK) and pharmacodynamics (PD) of anti-bacterial drugs for TB, focusing on blood and patient status (c.f. [25,154–161]). The spatial complexity of the granuloma, with *Mtb* within and external to cells that are located in this structure, complicates treatment, as antibiotic penetration into granulomas is often spatially heterogeneous [162–165] and bacteria develop phenotypic tolerance to antibiotics inside granulomas. To assess antibiotic efficacy within a granuloma, a single granuloma model (described above) is integrated with two additional elements [52,54]. First, ODE and PDE models are used to describe plasma and tissue pharmacokinetics (PK) following oral antibiotic administration, capturing antibiotic concentrations available to the granuloma as a function of time. Second, drug pharmacodynamics (PD) is described by ODEs that capture the mechanisms by which extracellular, intracellular replicating, and non-replicating bacteria are killed. Direct efficacy comparisons between drugs and dosing regimens (e.g. daily vs. weekly dosing) are possible, as one can examine the same simulated granulomas under different conditions. Thus far, two first line antibiotics, isoniazid (INH) and rifampin (RIF), have been examined within granulomas, with more antibiotics currently being incorporated [54,166]. This work demonstrates that antibiotic penetration into a granuloma limits treatment efficacy, particularly for intermittent dosing regimens as compared to daily regimens. It also demonstrates that intracellular bacteria present the main reservoir of bacteria at risk for developing resistance to RIF, one of the main first-line anti-TB antibiotics. In other work, the possibility of inhaled or orally dosed nanoparticle antibiotic delivery formulations was tested [53,55]. Both models highlight that the PK of released antibiotics influence the feasibility of nanoparticle delivery systems. While one drug (INH) was deemed suitable for inhaled delivery, the other (RIF) appeared less promising [53]. Overall, these *in silico* systems pharmacology approaches to examining the design space for antibiotic treatment may facilitate the identification of new treatment regimens for TB.

Within host-immune response modeling to guide vaccine development and testing

To date there is no broadly effective vaccine for TB, although many are in various stages of clinical development [167]. These clinical trials are difficult, expensive, and time-consuming, providing incentives to take a computational systems biology approach. Computational

efforts in this area are limited but hold promise. Computational models have been applied to the study of other vaccines such as those for influenza and LCMV [168–170] but have yet to be applied in the context of TB. To begin to understand the generation of vaccine immunity in lymph nodes, an ABM representing the dynamics occurring during priming in a LN was developed. This model was used to simulate the generation of effector and memory T cells in response to a vaccine [24,82,171]. In one study, a “memory design space” was postulated and explored in which various ratios of effector and memory T cells were generated and their ability to generate an appropriate response determined. Given a set of desired characteristics for antigen-specific memory populations, the model can be used as a tool to predict vaccine formulations that will generate memory cell populations as well as their effect on granuloma-scale outcomes [24,82]. Here, virtual clinical trials that allow for host-to-host variation in a population will be important to perform.

Computational modeling in biomarker discovery in TB

Biomarkers associated with the observed spectrum of TB infection have yet to be identified [9]. This gap is particularly relevant when designing and testing new treatments and vaccines and defining the correlates of protection [172–175]. Recent work partly addresses this lack of understanding in a natural immunity setting [38]. In this work, the authors generated a unique large dataset of cell and cytokine markers measured in blood samples from *Mtb*-infected NHPs up to 6 months post infection. The NHPs were classified as either latent or active TB cases. The experimental data were used to build and calibrate a multi-organ computational model (lung at a single granuloma scale, with *in silico* blood level readouts of T cell memory markers). The analysis of *in silico* blood measures identifies *Mtb*-specific frequencies of effector T cell phenotypes at various time points post infection as promising indicators of infection outcome. The next challenge will be to experimentally validate such biomarkers; however, more reliable experimental techniques are needed to label *Mtb*-specific epitopes, which is not yet possible in primates. Another direction of research will be to address the same questions but in vaccine-induced immunity scenarios.

Tools for guiding new drug formulations for TB treatment

Computational models and systems biology approaches may also suggest modifications of existing drugs to improve efficacy as well as new drug targets. Application of uncertainty and sensitivity analyses to computational models can help with this identification. For example, since antibiotic properties are parameters in the computational models, analysis of these models can identify the appropriate antibiotic properties to modify

by systematically tuning each parameter and evaluating effects on treatment outcomes in the context of complex host immunity, PK and PD [54]. Identifying good bacterial targets for new or repurposed antibiotics is challenging since growth inhibition in liquid culture or *in vitro* mammalian cell infection does not necessarily translate into infection inhibition *in vivo*. Computational models that include host immunity and bacterial metabolism are able to identify bacterial genes that will have an effect on infection progression when targeted mid-infection in an established granuloma, a realistic approximation of when antibiotics would be administered in patients [57]. Further, single granuloma computational model analyses have suggested NF κ -B-associated signaling and TNF internalization pathways as well as a variety of Mtb metabolic enzymes as promising targets [56,58,59].

Successes, limitations and future challenges

Tuberculosis is a centuries old disease that is in desperate need of new and improved strategies for treatment and prevention. Copious amounts of data are being generated on many aspects of infection and disease, and we argue that a systems biology approach can help integrate and make sense of it. Collaborations between experimental and theoretical scientists can lead to advancements in the field. Over the past 15 years as described above, mathematical and computational models of within-host dynamics of TB infection have been developed and have provided insights to advance the study of this disease. For example, initially the anti-inflammatory cytokine IL-10 was shown to have no importance in mouse models of TB, but our computational models indicated that this negative regulator was a key determinant of infection outcome [46,61,62]. Later and ongoing studies in NHPs driven by our findings have supported this role and suggest that it can potentially be harnessed for immunotherapy [75,176] and unpublished data]. It is this close connection between theory and experiment that may have the largest impacts on therapy development as well.

As described in Table 2, datasets to build, validate, and test in-host models are available. Different and disparate types of data are available from unique animal systems, at both different length and time scales and from different types of experiments and in different physiological compartments, including cell numbers in blood and lung, PET/CT images from NHP granulomas, as well as *in vitro* datasets (i.e. antibiotic microbicidal activity), and patient disease status and blood protein levels, etc. The computational approaches reviewed here are an excellent way to integrate various types of data into a more cohesive picture, providing better understanding of the immune factors and bacterial properties that drive and control infection.

Current and forthcoming data, together with the types of models described herein, position computational biologists to take on new challenges in TB. The first challenge is to better understand the factors that limit the efficacy of current and (forthcoming) antibiotics. The four front-line anti-TB drugs, INH, RIF, pyrazinamide, and ethambutol, as well as second-line agents used against drug-resistant strains, were implemented decades ago. More recently, broader spectrum antibiotics — aminoglycosides, fluoroquinolones and oxazolidinones — have been repurposed to treat drug-resistant TB. In addition, two representatives of new drug classes were approved for use against drug-resistant TB: Bedaquiline and Delamanid. More than a dozen drugs or drug candidates from these various classes are currently in clinical development or repositioning [177–179]. While the potential of new classes of antibiotics is promising, the “design space” for treatment regimens (combinations of antibiotics, doses, and dose frequencies) is huge. There is no standard approach to identifying treatment regimens and drugs are currently assigned in an ad hoc fashion, e.g. based on local guidelines and cost rather than comparative clinical data, which are largely missing. Exploring the design space in animal models or human clinical trials is extremely costly, lengthy, and difficult to assess. Integrating multi-scale computational models with relevant animal model data could allow rapid *in silico* testing of antibiotic therapies to predict more effective treatment regimens and accelerate clinical research.

We also need to better understand the development of antibiotic resistance. While some mathematical and computational approaches have been applied at the bacterial scale [31,180–184], including additional scales may be necessary in the multi-scale approaches described earlier. The genetic scale can capture resistance that arises due to point mutations or other genetic changes, up to the population scale, which captures the spread of drug resistance. The power of granuloma level models can be harnessed to understand how resistance can develop in the presence of antibiotics, esp. inadequate concentrations of antibiotics. These models may also require additional computational and modeling approaches to efficiently capture rare events in the context of what are computationally intensive simulations.

Challenges in the modeling approaches to study TB remain. For example, the single granuloma models described above capture in great detail the dynamics of immune cells, cytokines and chemokines, and Mtb; however the model simulations are time-consuming and costly, particularly given the large numbers of replications needed to gain meaningful statistics. In addition, on average, there are 42 granulomas in NHP lungs during TB infection [4–6]. Modeling techniques need to be developed to track the evolution of multiple granulomas. Early approaches assumed a well-mixed lung compartment; current approaches improve on this

but are still fairly simplified [38]. Further, we need computational approaches that mechanistically link to blood datasets (the most easily sampled compartment in humans) with the simultaneous evolving infection in the host lung. Such studies will advance the identification and interpretation of biomarkers of infection and correlates of protection for vaccine studies. However, given the computational demands of multi-scale modeling, especially ABMs, models should be developed with an appropriate level of complexity; techniques such as tuneable resolution (BOX) [185] and other coarse-graining tools can assist in analysis [186–188]. Developing models capturing within-host and population-scale dynamics is computationally difficult but necessary to explore the population-scale impacts of host-level factors [189–191].

While there are few applications of multi-scale modeling to infectious disease, the approaches here build on and contribute to the broader application of multi-scale modeling in biology, including many focused in cancer modeling and blood-related studies (e.g. Refs. [192–196]). In addition, there are other granulomatous diseases such as Schistosomiasis and Leishmania that would benefit from modeling, and lessons there could be applicable to TB and vice versa [197]. To advance infectious disease research further, it will likely take scientists and modelers collaborating across fields as different as HIV and TB, promoting cross-fertilization of ideas, methods, therapies and analysis techniques to gain a better understanding of how to combat and prevent infections.

Acknowledgement

This research was supported by: R01 AI123093, R01 EB012579, R01 HL110811, HL1U01HL131072. We also thank our fantastic experimental collaborators and students without whom our work would not have been possible.

References

1. WHO: *Global tuberculosis report 2016*. World Health Organization; 2016.
2. Houben RM, Dodd PJ: **The global burden of latent tuberculosis infection: a re-estimation using mathematical modelling**. *PLoS Med* 2016, **13**:e1002152. <http://dx.doi.org/10.1371/journal.pmed.1002152>. PubMed PMID: 27780211; PubMed Central PMCID: PMC5079585.
3. Feng Z, Castillo-Chavez C, Capurro AF: **A model for tuberculosis with exogenous reinfection**. *Theor Popul Biol* 2000, **57**: 235–247. <http://dx.doi.org/10.1006/tpbi.2000.1451>. Epub 2000/06/01, S0040–5809(00)91451-5 [pii]. PubMed PMID: 10828216.
4. Via LE, Lin PL, Ray SM, Carrillo J, Allen SS, Eum SY, Taylor K, Klein E, Manjunatha U, Gonzales J, Lee EG, Park SK, Raleigh JA, Cho SN, McMurray DN, Flynn JL, Barry 3rd CE: **Tuberculous granulomas are hypoxic in guinea pigs, rabbits, and nonhuman primates**. *Infect Immun* 2008, **76**:2333–2340. <http://dx.doi.org/10.1128/IAI.01515-07>. Epub 2008/03/19, [pii] 10.1128/IAI.01515-07. PubMed PMID: 18347040; PubMed Central PMCID: PMC2423064.
5. Lin PL, Flynn JL: **Understanding latent tuberculosis: a moving target**. *J Immunol* 2010, **185**:15–22. <http://dx.doi.org/10.4049/jimmunol.0903856>. Epub 2010/06/22, [pii] 10.4049/jimmunol.0903856. PubMed PMID: 20562268.
6. Ford CB, Lin PL, Chase MR, Shah RR, Iartchouk O, Galagan J, Mohaideen N, Ioerger TR, Sacchetti JC, Lipsitch M, Flynn JL, Fortune SM: **Use of whole genome sequencing to estimate the mutation rate of *Mycobacterium tuberculosis* during latent infection**. *Nat Genet* 2011, **43**:482–486. <http://dx.doi.org/10.1038/ng.811>. Epub 2011/04/26, PubMed PMID: 21516081; PubMed Central PMCID: PMC3101871.
7. Lillebaek T, Dirksen A, Baess I, Strunge B, Thomsen VO, Andersen AB: **Molecular evidence of endogenous reactivation of *Mycobacterium tuberculosis* after 33 years of latent infection**. *J Infect Dis* 2002, **185**:401–404. PubMed PMID: 11807725.
8. Esmail H, Barry 3rd CE, Young DB, Wilkinson RJ: **The ongoing challenge of latent tuberculosis**. *Philos Trans R Soc Lond B Biol Sci* 2014, **369**:20130437. <http://dx.doi.org/10.1098/rstb.2013.0437>. Epub 2014/05/14, rstb.2013.0437 [pii]. PubMed PMID: 24821923; PubMed Central PMCID: PMC4024230.
9. Barry 3rd CE, Boshoff HI, Dartois V, Dick T, Ehrt S, Flynn J, Schnappinger D, Wilkinson RJ, Young D: **The spectrum of latent tuberculosis: rethinking the biology and intervention strategies**. *Nat Rev Microbiol* 2009, **7**:845–855. <http://dx.doi.org/10.1038/nrmicro2236>. Epub 2009/10/27, [pii] 10.1038/nrmicro2236. PubMed PMID: 19855401.
10. Sadikot RT: **Identifying patients at high risk of tuberculosis recurrence**. *Int J Mycobacteriol* 2016, **5**(Suppl. 1):S66. <http://dx.doi.org/10.1016/j.ijmyco.2016.08.017>. PubMed PMID: 28043619.
11. Tiemersma EW, van der Werf MJ, Borgdorff MW, Williams BG, Nagelkerke NJ: **Natural history of tuberculosis: duration and fatality of untreated pulmonary tuberculosis in HIV negative patients: a systematic review**. *PLoS One* 2011, **6**:e17601. <http://dx.doi.org/10.1371/journal.pone.0017601>. PubMed PMID: 21483732; PubMed Central PMCID: PMC3070694.
12. Petruccioli E, Scriba TJ, Petrone L, Hatherill M, Cirillo DM, Joosten SA, Ottenhoff TH, Denkinger CM, Goletti D: **Correlates of tuberculosis risk: predictive biomarkers for progression to active tuberculosis**. *Eur Respir J* 2016, **48**:1751–1763. <http://dx.doi.org/10.1183/13993003.01012-2016>. PubMed PMID: 27836953.
13. Zak DE, Penn-Nicholson A, Scriba TJ, Thompson E, Suliman S, Amon LM, Mahomed H, Erasmus M, Whatney W, Hussey GD, Abrahams D, Kafaar F, Hawkrigge T, Verver S, Hughes EJ, Ota M, Sutherland J, Howe R, Dockrell HM, Boom WH, Thiel B, Ottenhoff TH, Mayanja-Kizza H, Crampin AC, Downing K, Hatherill M, Valvo J, Shankar S, Parida SK, Kaufmann SH, Walzl G, Aderem A, Hanekom WA, Acs, groups GCcs: **A blood RNA signature for tuberculosis disease risk: a prospective cohort study**. *Lancet* 2016, **387**:2312–2322. [http://dx.doi.org/10.1016/S0140-6736\(15\)01316-1](http://dx.doi.org/10.1016/S0140-6736(15)01316-1). PubMed PMID: 27017310.
14. Dodd CE, Schlesinger LS: **New concepts in understanding latent tuberculosis**. *Curr Opin Infect Dis* 2017. <http://dx.doi.org/10.1097/QCO.0000000000000367>. PubMed PMID: 28177961.
15. Young D, Stark J, Kirschner D: **Systems biology of persistent infection: tuberculosis as a case study**. *Nat Rev Microbiol* 2008, **6**:520–528. <http://dx.doi.org/10.1038/nrmicro1919>. Epub 2008/06/10, PubMed PMID: 18536727; PubMed Central PMCID: PMCn/a.
16. Fallahi-Sichani M, Marino S, Flynn JL, Linderman JJ, Kirschner DE: **A systems biology approach for understanding granuloma formation and function in tuberculosis**. In *Systems biology of tuberculosis*. Edited by McFadden J, Beste DJV, Kierzek AM, Springer; 2013.
17. Walpole J, Papin JA, Peirce SM: **Multiscale computational models of complex biological systems**. *Annu Rev Biomed Eng* 2013, **15**:137–154. <http://dx.doi.org/10.1146/annurev-bioeng-071811-150104>. Epub 2013/05/07, PubMed PMID: 23642247.
18. Jamshidi N, Palsson BO: **Investigating the metabolic capabilities of *Mycobacterium tuberculosis* H37Rv using the in silico strain iNJ661 and proposing alternative drug targets**. *BMC Syst Biol* 2007, **1**:26. <http://dx.doi.org/10.1186/1752-0509-1-26>. Epub 2007/06/09, PubMed PMID: 17555602; PubMed Central PMCID: PMC1925256.

19. Bordbar A, Lewis NE, Schellenberger J, Palsson BO, Jamshidi N: **Insight into human alveolar macrophage and *M. tuberculosis* interactions via metabolic reconstructions.** *Mol Syst Biol* 2010, **6**:422, <http://dx.doi.org/10.1038/msb.2010.68>. PubMed PMID: 20959820; PubMed Central PMCID: PMC2990636.
20. Shi L, Sohaskey CD, Pfeiffer C, Datta P, Parks M, McFadden J, North RJ, Gennaro ML: **Carbon flux rerouting during *Mycobacterium tuberculosis* growth arrest.** *Mol Microbiol* 2010, **78**: 1199–1215, <http://dx.doi.org/10.1111/j.1365-2958.2010.07399.x>. Epub 2010/11/26, PubMed PMID: 21091505; PubMed Central PMCID: PMC3072047.
21. Beste DJ, Noh K, Niedenfuhr S, Mendum TA, Hawkins ND, Ward JL, Beale MH, Wiechert W, McFadden J: **(13)C-flux spectral analysis of host-pathogen metabolism reveals a mixed diet for intracellular *Mycobacterium tuberculosis*.** *Chem Biol* 2013, **20**:1012–1021, <http://dx.doi.org/10.1016/j.chembiol.2013.06.012>. Epub 2013/08/06, PubMed PMID: 23911587; PubMed Central PMCID: PMC3752972.
22. Rienksma RA, Suarez-Diez M, Spina L, Schaap PJ, Martins Dos Santos VA: **Systems-level modeling of mycobacterial metabolism for the identification of new (multi-)drug targets.** *Semin Immunol* 2014, **26**:610–622, <http://dx.doi.org/10.1016/j.smim.2014.09.013>. PubMed PMID: 25453232.
23. Kirschner DE, Young D, Flynn JL: **Tuberculosis: global approaches to a global disease.** *Curr Opin Biotechnol* 2010, **21**: 524–531, <http://dx.doi.org/10.1016/j.copbio.2010.06.002>. Epub 2010/07/20, PubMed PMID: 20637596; PubMed Central PMCID: PMC2943033.
24. Linderman JJ, Cilfone NA, Pienaar E, Gong C, Kirschner DE: **A multi-scale approach to designing therapeutics for tuberculosis.** *Integr Biol (Camb)* 2015, **7**:591–609, <http://dx.doi.org/10.1039/c4ib00295d>. PubMed PMID: 25924949; PubMed Central PMCID: PMC4436084.
25. Dooley KE, Luetkemeyer AF, Park JG, Allen R, Cramer Y, Murray S, Sutherland D, Aweeka F, Koletar SL, Marzan F, Bao J, Savic R, Haas DW, Team ACTGAS: **Phase I safety, pharmacokinetics, and pharmacogenetics study of the antituberculosis drug PA-824 with concomitant lopinavir-ritonavir, efavirenz, or rifampin.** *Antimicrob Agents Chemother* 2014, **58**: 5245–5252, <http://dx.doi.org/10.1128/AAC.03332-14>. PubMed PMID: 24957823; PubMed Central PMCID: PMC4135849.
26. Via LE, Savic R, Weiner DM, Zimmerman MD, Pradeaux B, Irwin SM, Lyon E, O'Brien P, Gopal P, Eum S, Lee M, Lanoix JP, Dutta NK, Shim T, Cho JS, Kim W, Karakousis PC, Lenaerts A, Nuernberger E, Barry 3rd CE, Dartois V: **Host-mediated bio-activation of pyrazinamide: implications for efficacy, resistance, and therapeutic alternatives.** *ACS Infect Dis* 2015, **1**: 203–214, <http://dx.doi.org/10.1021/id500028m>. PubMed PMID: 26086040; PubMed Central PMCID: PMC4467917.
27. WHO: **Target regimen profiles for TB treatment: candidates: rifampicin-susceptible, rifampicin resistant and pan-TB treatment regimens.** 2016.
28. Andersen P, Woodworth JS: **Tuberculosis vaccines—rethinking the current paradigm.** *Trends Immunol* 2014, **35**: 387–395, <http://dx.doi.org/10.1016/j.it.2014.04.006>. PubMed PMID: 24875637.
29. Wong EA, Agger EM, Andersen P, Flynn JL: **Vaccination route has an impact on level of protection of non-human primates from tuberculosis.** *J Immunol* 2016, **196**(1 Suppl.):146.21.
30. Lipsitch M, Levin BR: **Population dynamics of tuberculosis treatment: mathematical models of the roles of non-compliance and bacterial heterogeneity in the evolution of drug resistance.** *Int J Tuberc Lung Dis* 1998, **2**:187–199. Epub 1998/04/04. PubMed PMID: 9526190.
31. Ankomah P, Levin BR: **Two-drug antimicrobial chemotherapy: a mathematical model and experiments with *Mycobacterium marinum*.** *PLoS Pathog* 2012, **8**:e1002487, <http://dx.doi.org/10.1371/journal.ppat.1002487>. Epub 2012/01/19, PubMed PMID: 22253599; PubMed Central PMCID: PMC3257304.
32. Ghosh S, Sureka K, Ghosh B, Bose I, Basu J, Kundu M: **Phenotypic heterogeneity in mycobacterial stringent response.** *BMC Syst Biol* 2011, **5**:18, <http://dx.doi.org/10.1186/1752-0509-5-18>. PubMed PMID: 21272295; PubMed Central PMCID: PMC3045321.
33. Alavez-Ramirez J, Castellanos JR, Esteva L, Flores JA, Fuentes-Allen JL, Garcia-Ramos G, Gomez G, Lopez-Estrada J: **Within-host population dynamics of antibiotic-resistant *M. tuberculosis*.** *Math Med Biol A J IMA* 2007, **24**:35–56, <http://dx.doi.org/10.1093/imammb/dql026>. PubMed PMID: 17060491.
34. Marino S, Linderman JJ, Kirschner DE: **A multifaceted approach to modeling the immune response in tuberculosis.** *Wiley Interdiscip Rev Syst Biol Med* 2011, **3**:479–489, <http://dx.doi.org/10.1002/wsbm.131>. Epub 2011/01/05, PubMed PMID: 21197656; PubMed Central PMCID: PMC3110521.
35. Kirschner D: **The multi-scale immune response to pathogens: *M. tuberculosis* as an example.** *In silico Immunol* 2007: 289–311. PubMed PMID: exempt.
36. Lin PL, Ford CB, Coleman MT, Myers AJ, Gawande R, Ioerger T, Sacchetti J, Fortune SM, Flynn JL: **Sterilization of granulomas is common in active and latent tuberculosis despite within-host variability in bacterial killing.** *Nat Med* 2014, **20**:75–79, <http://dx.doi.org/10.1038/nm.3412>. Epub 2013/12/18, PubMed PMID: 24336248.
37. Lalande L, Bourguignon L, Bihari S, Maire P, Neely M, Jelliffe R, Goutelle S: **Population modeling and simulation study of the pharmacokinetics and antituberculosis pharmacodynamics of isoniazid in lungs.** *Antimicrob Agents Chemother* 2015, **59**: 5181–5189, <http://dx.doi.org/10.1128/AAC.00462-15>. PubMed PMID: 26077251; PubMed Central PMCID: PMC4538517.
38. Marino S, Gideon HP, Gong C, Mankad S, McCrone JT, Lin PL, Linderman JJ, Flynn JL, Kirschner DE: **Computational and empirical studies predict mycobacterium tuberculosis-specific T cells as a biomarker for infection outcome.** *PLoS Comput Biol* 2016, **12**:e1004804, <http://dx.doi.org/10.1371/journal.pcbi.1004804>. PubMed PMID: 27065304; PubMed Central PMCID: PMC4827839.
39. Raman K, Bhat AG, Chandra N: **A systems perspective of host-pathogen interactions: predicting disease outcome in tuberculosis.** *Mol Biosyst* 2010, **6**:516–530, <http://dx.doi.org/10.1039/b912129c>. PubMed PMID: 20174680.
40. Marino S, Kirschner DE: **A multi-compartment hybrid computational model predicts key roles for dendritic cells in tuberculosis infection.** *Computation* 2016, **4**. In Press.
41. Gong C, Linderman JJ, Kirschner D: **A population model capturing dynamics of tuberculosis granulomas predicts host infection outcomes.** *Math Biosci Eng* 2015, **12**:625–642, <http://dx.doi.org/10.3934/mbe.2015.12.625>. PubMed PMID: 25811559; PubMed Central PMCID: PMC4447319.
42. Magombedze G, Garira W, Mwenje E: **Mathematical modeling of chemotherapy of human TB infection.** *J Biol Syst* 2006, **14**: 509–553, <http://dx.doi.org/10.1142/s0218339006001945>.
43. de Espindola AL, Girardi D, Penna TJP, Bauch CT, Troca Cabella BC, Martinez AS: **An antibiotic protocol to minimize emergence of drug-resistant tuberculosis.** *Phys A Stat Mech its Appl* 2014, **400**:80–92, <http://dx.doi.org/10.1016/j.physa.2013.12.039>.
44. Kirschner D: **Dynamics of co-infection with *M. Tuberculosis* and HIV-1.** *Theor Popul Biol* 1999, **55**:94–109, <http://dx.doi.org/10.1006/tpbi.1998.1382>. Epub 1999/02/02, PubMed PMID: 9925811.
45. Magombedze G, Mulder N: **A mathematical representation of the development of *Mycobacterium tuberculosis* active, latent and dormant stages.** *J Theor Biol* 2012, **292**:44–59, <http://dx.doi.org/10.1016/j.jtbi.2011.09.025>. Epub 2011/10/05, PubMed PMID: 21968442.
46. Wigginton JE, Kirschner D: **A model to predict cell-mediated immune regulatory mechanisms during human infection with *Mycobacterium tuberculosis*.** *J Immunol* 2001, **166**: 1951–1967. Epub 2001/02/13. PubMed PMID: 11160244; PubMed Central PMCID: PMCExempt.
47. Sud D, Bigbee C, Flynn JL, Kirschner DE: **Contribution of CD8+ T cells to control of *Mycobacterium tuberculosis* infection.**

- J Immunol* 2006, **176**:4296–4314. PubMed PMID: 16547267; PubMed Central PMCID: PMCexempt.
48. Magombedze G, Garira W, Mwenje E: **Modelling the human immune response mechanisms to mycobacterium tuberculosis infection in the lungs.** *Math Biosci Eng* 2006, **3**:661–682. Epub 2006/10/01. PubMed PMID: 20361838.
 49. Bauer AL, Hogue IB, Marino S, Kirschner DE: **The effects of HIV Infection on latent tuberculosis.** *Math Model Nat Phenom* 2008, **3**:229–266. PubMed PMID: n/a; PubMed Central PMCID: PMCn/a.
 50. Antia R, Koella JC, Perrot V: **Models of the within-host dynamics of persistent mycobacterial infections.** *Proc Biol Sci* 1996, **263**:257–263. <http://dx.doi.org/10.1098/rspb.1996.0040>. Epub 1996/03/22. PubMed PMID: 8920248.
 51. Prats C, Vilaplana C, Valls J, Marzo E, Cardona PJ, Lopez D: **Local inflammation, dissemination and coalescence of lesions are key for the progression toward active tuberculosis: the bubble model.** *Front Microbiol* 2016, **7**:33. <http://dx.doi.org/10.3389/fmicb.2016.00033>. PubMed PMID: 26870005; PubMed Central PMCID: PMC4736263.
 52. Pienaar E, Cilfone NA, Lin PL, Dartois V, Mattila JT, Butler JR, Flynn JL, Kirschner DE, Linderman JJ: **A computational tool integrating host immunity with antibiotic dynamics to study tuberculosis treatment.** *J Theor Biol* 2015, **367**:166–179. <http://dx.doi.org/10.1016/j.jtbi.2014.11.021>. Epub 2014/12/17. PubMed PMID: 25497475; PubMed Central PMCID: PMC4332617.
 53. Cilfone NA, Pienaar E, Thurber GM, Kirschner DE, Linderman JJ: **Systems pharmacology approach toward the design of inhaled formulations of rifampicin and isoniazid for treatment of tuberculosis.** *CPT Pharmacomet Syst Pharmacol* 2015, **4**: 193–203. e00022. <http://dx.doi.org/10.1002/psp4.22>. PubMed PMID: 26225241; PubMed Central PMCID: PMC4394619.
 54. Pienaar E, Dartois V, Linderman JJ, Kirschner DE: **In silico evaluation and exploration of antibiotic tuberculosis treatment regimens.** *BMC Syst Biol* 2015, **9**:79. <http://dx.doi.org/10.1186/s12918-015-0221-8>. PubMed PMID: 26578235; PubMed Central PMCID: PMC4650854.
 55. Grobler A, Perez Sierra Z, Viljoen HJ: **Modeling nanoparticle delivery of TB drugs to granulomas.** *J Theor Biol* 2016, **388**: 85–95. <http://dx.doi.org/10.1016/j.jtbi.2015.10.004>. PubMed PMID: 26493361.
 56. Fallahi-Sichani M, Kirschner DE, Linderman JJ: **NF-kappaB signaling dynamics play a key role in infection control in tuberculosis.** *Front Physiol* 2012, **3**:170. <http://dx.doi.org/10.3389/fphys.2012.00170>. Epub 2012/06/12. PubMed PMID: 22685435; PubMed Central PMCID: PMC3368390.
 57. Pienaar E, Matern WM, Linderman JJ, Bader JS, Kirschner DE: **Multiscale model of Mycobacterium tuberculosis infection maps metabolite and gene perturbations to granuloma sterilization predictions.** *Infect Immun* 2016, **84**:1650–1669. <http://dx.doi.org/10.1128/IAI.01438-15>. PubMed PMID: 26975995; PubMed Central PMCID: PMC4862722.
 58. Fallahi-Sichani M, Schaller MA, Kirschner DE, Kunkel SL, Linderman JJ: **Identification of key processes that control tumor necrosis factor availability in a tuberculosis granuloma.** *PLoS Comput Biol* 2010, **6**:e1000778. <http://dx.doi.org/10.1371/journal.pcbi.1000778>. Epub 2010/05/14. PubMed PMID: 20463877; PubMed Central PMCID: PMC2865521.
 59. Fallahi-Sichani M, El-Kebir M, Marino S, Kirschner DE, Linderman JJ: **Multiscale computational modeling reveals a critical role for TNF-alpha receptor 1 dynamics in tuberculosis granuloma formation.** *J Immunol* 2011, **186**:3472–3483. <http://dx.doi.org/10.4049/jimmunol.1003299>. Epub 2011/02/16. PubMed PMID: 21321109; PubMed Central PMCID: PMC3127549.
 60. Fallahi-Sichani M, Flynn JL, Linderman JJ, Kirschner DE: **Differential risk of tuberculosis reactivation among anti-TNF therapies is due to drug binding kinetics and permeability.** *J Immunol* 2012, **188**:3169–3178. <http://dx.doi.org/10.4049/jimmunol.1103298>. Epub 2012/03/02. PubMed PMID: 22379032; PubMed Central PMCID: PMC3311778.
 61. Cilfone NA, Perry CR, Kirschner DE, Linderman JJ: **Multi-scale modeling predicts a balance of tumor necrosis factor-alpha and interleukin-10 controls the granuloma environment during Mycobacterium tuberculosis infection.** *PLoS One* 2013, **8**:e68680. <http://dx.doi.org/10.1371/journal.pone.0068680>. Epub 2013/07/23. PubMed PMID: 23869227; PubMed Central PMCID: PMC3711807.
 62. Cilfone NA, Ford CB, Marino S, Mattila JT, Gideon HP, Flynn JL, Kirschner DE, Linderman JJ: **Computational modeling predicts IL-10 control of lesion sterilization by balancing early host immunity-mediated antimicrobial responses with caseation during mycobacterium tuberculosis infection.** *J Immunol* 2015, **194**:664–677. <http://dx.doi.org/10.4049/jimmunol.1400734>. Epub 2014/12/17. PubMed PMID: 25512604; PubMed Central PMCID: PMC4283220.
 63. Marino S, Sud D, Plessner H, Lin PL, Chan J, Flynn JL, Kirschner DE: **Differences in reactivation of tuberculosis induced from anti-TNF treatments are based on bioavailability in granulomatous tissue.** *PLoS Comput Biol* 2007, **3**: 1909–1924. PubMed PMID: 17953477; PubMed Central PMCID: PMCn/a.
 64. Ray JC, Flynn JL, Kirschner DE: **Synergy between individual TNF-dependent functions determines granuloma performance for controlling Mycobacterium tuberculosis infection.** *J Immunol* 2009, **182**:3706–3717. <http://dx.doi.org/10.4049/jimmunol.0802297>. Epub 2009/03/07. PubMed PMID: 19265149; PubMed Central PMCID: PMC3182770.
 65. Marino S, Cilfone NA, Mattila JT, Linderman JJ, Flynn JL, Kirschner DE: **Macrophage polarization drives granuloma outcome during Mycobacterium tuberculosis infection.** *Infect Immun* 2015, **83**:324–338. <http://dx.doi.org/10.1128/IAI.02494-14>. Epub 2014/11/05. PubMed PMID: 25368116; PubMed Central PMCID: PMC4288886.
 66. Bru A, Cardona PJ: **Mathematical modeling of tuberculosis bacillary counts and cellular populations in the organs of infected mice.** *PLoS One* 2010, **5**:e12985. <http://dx.doi.org/10.1371/journal.pone.0012985>. Epub 2010/10/05. PubMed PMID: 20886087; PubMed Central PMCID: PMC2944881.
 67. Repasy T, Lee J, Marino S, Martinez N, Kirschner DE, Hendricks G, Baker S, Wilson AA, Kotton DN, Kornfeld H: **Intracellular bacillary burden reflects a burst size for Mycobacterium tuberculosis in vivo.** *PLoS Pathog* 2013, **9**: e1003190. <http://dx.doi.org/10.1371/journal.ppat.1003190>. Epub 2013/02/26. PubMed PMID: 23436998; PubMed Central PMCID: PMC3578792.
 68. Gammack D, Doering CR, Kirschner DE: **Macrophage response to Mycobacterium tuberculosis infection.** *J Math Biol* 2004, **48**: 218–242. <http://dx.doi.org/10.1007/s00285-003-0232-8>. Epub 2004/01/28. PubMed PMID: 14745511; PubMed Central PMCID: PMCexempt.
 69. Segovia-Juarez JL, Ganguli S, Kirschner D: **Identifying control mechanisms of granuloma formation during M. tuberculosis infection using an agent-based model.** *J Theor Biol* 2004, **231**: 357–376. <http://dx.doi.org/10.1016/j.jtbi.2004.06.031>. Epub 2004/10/27. PubMed PMID: 15501468; PubMed Central PMCID: PMCexempt.
 70. Ganguli S, Gammack D, Kirschner DE: **A metapopulation model of granuloma formation in the lung during infection with mycobacterium tuberculosis.** *Math Biosci Eng* 2005, **2**: 535–560. <http://dx.doi.org/10.1267/s00285-005-0232-8>. Epub 2005/07/01. PubMed PMID: 20369939.
 71. Hao W, Schlesinger LS, Friedman A: **Modeling granulomas in response to infection in the lung.** *PLoS One* 2016, **11**: e0148738. <http://dx.doi.org/10.1371/journal.pone.0148738>. PubMed PMID: 26986986; PubMed Central PMCID: PMC4795641.
 72. Serksen CL, Plimpton SJ, May EE: **Oxygen modulates the effectiveness of granuloma mediated host response to Mycobacterium tuberculosis: a multiscale computational biology approach.** *Front Cell Infect Microbiol* 2016, **6**:6. <http://dx.doi.org/10.3389/fcimb.2016.00006>. PubMed PMID: 26913242; PubMed Central PMCID: PMC4753379.
 73. Datta M, Via LE, Chen W, Baish JW, Xu L, Barry 3rd CE, Jain RK: **Mathematical model of oxygen transport in tuberculosis granulomas.** *Ann Biomed Eng* 2016, **44**:863–872. <http://dx.doi.org/10.1007/s10439-015-1415-3>. PubMed PMID: 26253038; PubMed Central PMCID: PMC4795989.

74. Cardona PJ, Prats C: **The small breathing amplitude at the upper lobes favors the attraction of polymorphonuclear neutrophils to *Mycobacterium tuberculosis* lesions and helps to understand the evolution toward active disease in an individual-based model.** *Front Microbiol* 2016, **7**:354, <http://dx.doi.org/10.3389/fmicb.2016.00354>. PubMed PMID: 27065951; PubMed Central PMCID: PMC4810076.
75. Marino S, Myers A, Flynn JL, Kirschner DE: **TNF and IL-10 are major factors in modulation of the phagocytic cell environment in lung and lymph node in tuberculosis: a next-generation two-compartmental model.** *J Theor Biol* 2010, **265**: 586–598, <http://dx.doi.org/10.1016/j.jtbi.2010.05.012>. Epub 2010/06/01, PubMed PMID: 20510249; PubMed Central PMCID: PMC3150786.
76. Marino S, Kirschner DE: **The human immune response to *Mycobacterium tuberculosis* in lung and lymph node.** *J Theor Biol* 2004, **227**:463–486. PubMed PMID: 15038983; PubMed Central PMCID: PMCexempt.
77. Myers AJ, Marino S, Kirschner DE, Flynn JL: **Inoculation dose of *Mycobacterium tuberculosis* does not influence priming of T cell responses in lymph nodes.** *J Immunol* 2013, **190**: 4707–4716, <http://dx.doi.org/10.4049/jimmunol.1203465>. Epub 2013/04/03, PubMed PMID: 23547119; PubMed Central PMCID: PMC3674545.
78. Marino S, Pawar S, Fuller CL, Reinhart TA, Flynn JL, Kirschner DE: **Dendritic cell trafficking and antigen presentation in the human immune response to *Mycobacterium tuberculosis*.** *J Immunol* 2004, **173**:494–506. PubMed PMID: 15210810; PubMed Central PMCID: PMCexempt.
79. Marino S, El-Kebir M, Kirschner D: **A hybrid multi-compartment model of granuloma formation and T cell priming in tuberculosis.** *J Theor Biol* 2011, **280**:50–62, <http://dx.doi.org/10.1016/j.jtbi.2011.03.022>. Epub 2011/03/30, PubMed PMID: 21443879; PubMed Central PMCID: PMC3740747.
80. Magombedze G, Garira W, Mwenje E: **The role of dendritic cells and other immune mechanisms during human infection with *Mycobacterium tuberculosis*.** *Int J Biomath* 2009, **02**:69–105, <http://dx.doi.org/10.1142/S1793524509000534>.
81. Day J, Friedman A, Schlesinger LS: **Modeling the immune rheostat of macrophages in the lung in response to infection.** *Proc Natl Acad Sci U S A* 2009, **106**:11246–11251, <http://dx.doi.org/10.1073/pnas.0904846106>. Epub 2009/06/25, [pii] 10.1073/pnas.0904846106. PubMed PMID: 19549875; PubMed Central PMCID: PMC2708732.
82. Ziraldo C, Gong C, Kirschner DE, Linderman JJ: **Strategic priming with multiple antigens can yield memory cell phenotypes optimized for infection with *Mycobacterium tuberculosis*: a computational study.** *Front Microbiol* 2015, **6**:1477, <http://dx.doi.org/10.3389/fmicb.2015.01477>. PubMed PMID: 26779136; PubMed Central PMCID: PMC4701940.
83. Chang ST, Linderman JJ, Kirschner DE: **Multiple mechanisms allow *Mycobacterium tuberculosis* to continuously inhibit MHC class II-mediated antigen presentation by macrophages.** *Proc Natl Acad Sci U S A* 2005, **102**:4530–4535, <http://dx.doi.org/10.1073/pnas.0500362102>. Epub 2005/03/16, PubMed PMID: 15767567; PubMed Central PMCID: PMC55518.
84. Chang ST, Ghosh D, Kirschner DE, Linderman JJ: **Peptide length-based prediction of peptide-MHC class II binding.** *Bioinformatics* 2006, **22**:2761–2767, <http://dx.doi.org/10.1093/bioinformatics/btl479>. Epub 2006/09/27, PubMed PMID: 17000752.
85. Chang ST, Linderman JJ, Kirschner DE: **Effect of multiple genetic polymorphisms on antigen presentation and susceptibility to *Mycobacterium tuberculosis* infection.** *Infect Immun* 2008, **76**:3221–3232, <http://dx.doi.org/10.1128/IAI.01677-07>. Epub 2008/04/30, PubMed PMID: 18443099; PubMed Central PMCID: PMC2446740.
86. Ray JC, Wang J, Chan J, Kirschner DE: **The timing of TNF and IFN-gamma signaling affects macrophage activation strategies during *Mycobacterium tuberculosis* infection.** *J Theor Biol* 2008, **252**:24–38, <http://dx.doi.org/10.1016/j.jtbi.2008.01.010>. Epub 2008/03/07, PubMed PMID: 18321531; PubMed Central PMCID: PMC2459258.
87. Pedruzzi G, Rao KV, Chatterjee S: **Mathematical model of mycobacterium-host interaction describes physiology of persistence.** *J Theor Biol* 2015, **376**:105–117, <http://dx.doi.org/10.1016/j.jtbi.2015.03.031>. PubMed PMID: 25865521.
88. Rapanoel HA, Mazandu GK, Mulder NJ: **Predicting and analyzing interactions between *Mycobacterium tuberculosis* and its human host.** *PLoS One* 2013, **8**:e67472, <http://dx.doi.org/10.1371/journal.pone.0067472>. PubMed PMID: 23844013; PubMed Central PMCID: PMC3699628.
89. Gough M, May E: **An in silico model of the effects of vitamin D3 on mycobacterium infected macrophage.** In *Conference proceedings: annual international conference of the IEEE engineering in medicine and biology society IEEE engineering in medicine and biology society annual conference*; 2016: 1443–1446, <http://dx.doi.org/10.1109/EMBC.2016.7590980>. PubMed PMID: 28268597.
90. Gunther G: **Multidrug-resistant and extensively drug-resistant tuberculosis: a review of current concepts and future challenges.** *Clin Med* 2014, **14**:279–285, <http://dx.doi.org/10.7861/clinmedicine.14-3-279>. PubMed PMID: 24889573.
91. Scriba TJ, Tameris M, Mansoor N, Smit E, van der Merwe L, Isaacs F, Keyser A, Moyo S, Brittain N, Lawrie A, Gelderbloem S, Veldsman A, Hatherill M, Hawkrige A, Hill AV, Hussey GD, Mahomed H, McShane H, Hanekom WA: **Modified vaccinia Ankara-expressing Ag85A, a novel tuberculosis vaccine, is safe in adolescents and children, and induces polyfunctional CD4+ T cells.** *Eur J Immunol* 2010, **40**:279–290, <http://dx.doi.org/10.1002/eji.200939754>. PubMed PMID: 20017188; PubMed Central PMCID: PMC3044835.
92. Flynn JL, Gideon HP, Mattila JT, Lin PL: **Immunology studies in non-human primate models of tuberculosis.** *Immunol Rev* 2015, **264**:60–73, <http://dx.doi.org/10.1111/imr.12258>. Epub 2015/02/24, PubMed PMID: 25703552; PubMed Central PMCID: PMC4339213.
93. Gideon HP, Phuah J, Myers AJ, Bryson BD, Rodgers MA, Coleman MT, Maiello P, Rutledge T, Marino S, Fortune SM, Kirschner DE, Lin PL, Flynn JL: **Variability in tuberculosis granuloma T cell responses exists, but a balance of pro- and anti-inflammatory cytokines is associated with sterilization.** *PLoS Pathog* 2015, **11**:e1004603, <http://dx.doi.org/10.1371/journal.ppat.1004603>. Epub 2015/01/23, PubMed PMID: 25611466; PubMed Central PMCID: PMC4303275.
94. Andersen P, Doherty TM: **The success and failure of BCG – implications for a novel tuberculosis vaccine.** *Nat Rev Microbiol* 2005, **3**:656–662, <http://dx.doi.org/10.1038/nrmi-cro1211>. PubMed PMID: 16012514.
95. Ginsberg AM, Ruhwald M, Mearns H, McShane H: **TB vaccines in clinical development.** *Tuberc (Edinb)* 2016, **99**(Suppl. 1): S16–S20, <http://dx.doi.org/10.1016/j.tube.2016.05.013>. PubMed PMID: 27470538.
96. Luabeya AK, Kagina BM, Tameris MD, Geldenhuys H, Hoff ST, Shi Z, Kromann I, Hatherill M, Mahomed H, Hanekom WA, Andersen P, Scriba TJ, Group HTS, Schoeman E, Krohn C, Day CL, Africa H, Makhethle L, Smit E, Brown Y, Suliman S, Hughes EJ, Bang P, Snowden MA, McClain B, Hussey GD: **First-in-human trial of the post-exposure tuberculosis vaccine H56: IC31 in *Mycobacterium tuberculosis* infected and non-infected healthy adults.** *Vaccine* 2015, **33**:4130–4140, <http://dx.doi.org/10.1016/j.vaccine.2015.06.051>. PubMed PMID: 26095509.
97. Lazarevic V, Myers AJ, Scanga CA, Flynn JL: **CD40, but not CD40L, is required for the optimal priming of T cells and control of aerosol *M. tuberculosis* infection.** *Immunity* 2003, **19**:823–835. PubMed PMID: 14670300.
98. Caruso AM, Serbina N, Klein E, Triebold K, Bloom BR, Flynn JL: **Mice deficient in CD4 T cells have only transiently diminished levels of IFN-gamma, yet succumb to tuberculosis.** *J Immunol* 1999, **162**:5407–5416. Epub 1999/05/05. PubMed PMID: 10228018.
99. Lin PL, Rodgers M, Smith L, Bigbee M, Myers A, Bigbee C, Chioseia I, Capuano SV, Fuhrman C, Klein E, Flynn JL: **Quantitative comparison of active and latent tuberculosis in the cynomolgus macaque model.** *Infect Immun* 2009, **77**:4631–4642, <http://dx.doi.org/10.1128/IAI.00592-09>. Epub 2009/07/22, [pii]

- 10.1128/IAI.00592-09. PubMed PMID: [19620341](#); PubMed Central PMCID: [PMC2747916](#).
100. Saunders BM, Frank AA, Orme IM, Cooper AM: **CD4 is required for the development of a protective granulomatous response to pulmonary tuberculosis.** *Cell Immunol* 2002, **216**:65–72. PubMed PMID: [12381351](#).
 101. Khader SA, Bell GK, Pearl JE, Fountain JJ, Rangel-Moreno J, Cilley GE, Shen F, Eaton SM, Gaffen SL, Swain SL, Locksley RM, Haynes L, Randall TD, Cooper AM: **IL-23 and IL-17 in the establishment of protective pulmonary CD4+ T cell responses after vaccination and during Mycobacterium tuberculosis challenge.** *Nat Immunol* 2007, **8**:369–377. PubMed PMID: [17351619](#).
 102. Scanga CA, Mohan VP, Yu K, Joseph H, Tanaka K, Chan J, Flynn JL: **Depletion of CD4(+) T cells causes reactivation of murine persistent tuberculosis despite continued expression of interferon gamma and nitric oxide synthase 2.** *J Exp Med* 2000, **192**:347–358. Epub 2000/08/10. PubMed PMID: [10934223](#); PubMed Central PMCID: [PMC2193220](#).
 103. Winslow GM, Cooper A, Reiley W, Chatterjee M, Woodland DL: **Early T-cell responses in tuberculosis immunity.** *Immunol Rev* 2008, **225**:284–299. <http://dx.doi.org/10.1111/j.1600-065X.2008.00693.x>. Epub 2008/10/08. [pii] 10.1111/j.1600-065X.2008.00693.x. PubMed PMID: [18837789](#).
 104. Ordway DJ, Shanley CA, Caraway ML, Orme EA, Bucy DS, Hascall-Dove L, Henao-Tamayo M, Harton MR, Shang S, Ackart D, Kraft SL, Lenaerts AJ, Basaraba RJ, Orme IM: **Evaluation of standard chemotherapy in the guinea pig model of tuberculosis.** *Antimicrob Agents Chemother* 2010, **54**:1820–1833. <http://dx.doi.org/10.1128/AAC.01521-09>. Epub 2010/02/18. [pii] 10.1128/AAC.01521-09. PubMed PMID: [20160055](#); PubMed Central PMCID: [PMC2863665](#).
 105. Demangel C, Bean AG, Martin E, Feng CG, Kamath AT, Britton WJ: **Protection against aerosol Mycobacterium tuberculosis infection using Mycobacterium bovis Bacillus Calmette Guérin-infected dendritic cells.** *Eur J Immunol* 1999, **29**:1972–1979. [http://dx.doi.org/10.1002/\(SICI\)1521-4141\(199906\)29:06<1972::AID-IMMU1972>3.0.CO;2-1](http://dx.doi.org/10.1002/(SICI)1521-4141(199906)29:06<1972::AID-IMMU1972>3.0.CO;2-1). Epub 1999/06/26. [pii] 10.1002/(SICI)1521-4141(199906)29:06<1972::AID-IMMU1972>3.0.CO;2-1. PubMed PMID: [10382760](#).
 106. Serbina NV, Lazarevic V, Flynn JL: **CD4(+) T cells are required for the development of cytotoxic CD8(+) T cells during Mycobacterium tuberculosis infection.** *J Immunol* 2001, **167**:6991–7000. Epub 2001/12/12. PubMed PMID: [11739519](#); PubMed Central PMCID: [PMCexempt](#).
 107. Behar SM, Dascher CC, Grusby MJ, Wang CR, Brenner MB: **Susceptibility of mice deficient in CD1D or TAP1 to infection with Mycobacterium tuberculosis.** *J Exp Med* 1999, **189**:1973–1980. Epub 1999/06/22. PubMed PMID: [10377193](#); PubMed Central PMCID: [PMC2192974](#).
 108. Kamath A, Woodworth JS, Behar SM: **Antigen-specific CD8+ T cells and the development of central memory during mycobacterium tuberculosis infection.** *J Immunol* 2006, **177**:6361–6369. PubMed PMID: [17056567](#).
 109. Kamath AB, Woodworth J, Xiong X, Taylor C, Weng Y, Behar SM: **Cytolytic CD8+ T cells recognizing CFP10 are recruited to the lung after Mycobacterium tuberculosis infection.** *J Exp Med* 2004, **200**:1479–1489. PubMed PMID: [15557351](#).
 110. Chen CY, Huang D, Wang RC, Shen L, Zeng G, Yao S, Shen Y, Halliday L, Fortman J, McAllister M, Estep J, Hunt R, Vasconcelos D, Du G, Porcelli SA, Larsen MH, Jacobs Jr WR, Haynes BF, Letvin NL, Chen ZW: **A critical role for CD8 T cells in a nonhuman primate model of tuberculosis.** *PLoS Pathog* 2009, **5**:e1000392. <http://dx.doi.org/10.1371/journal.ppat.1000392>. Epub 2009/04/22. PubMed PMID: [19381260](#); PubMed Central PMCID: [PMC2663842](#).
 111. Chen J, Liu XS: **Development and function of IL-10 IFN-gamma-secreting CD4(+) T cells.** *J Leukoc Biol* 2009, **86**:1305–1310. <http://dx.doi.org/10.1189/jlb.0609406>. Epub 2009/09/11. [pii] 10.1189/jlb.0609406. PubMed PMID: [19741156](#).
 112. Lin PL, Pawar S, Myers A, Pegu A, Fuhrman C, Reinhart TA, Capuano SV, Klein E, Flynn JL: **Early events in Mycobacterium tuberculosis infection in cynomolgus macaques.** *Infect Immun* 2006, **74**:3790–3803. <http://dx.doi.org/10.1128/IAI.00064-06>. Epub 2006/06/23. [pii] 10.1128/IAI.00064-06. PubMed PMID: [16790751](#); PubMed Central PMCID: [PMC1489679](#).
 113. Flynn JL, Chan J, Triebold KJ, Dalton DK, Stewart TA, Bloom BR: **An essential role for interferon gamma in resistance to Mycobacterium tuberculosis infection.** *J Exp Med* 1993, **178**:2249–2254. Epub 1993/12/01. PubMed PMID: [7504064](#); PubMed Central PMCID: [PMC2191274](#).
 114. Einarsdottir T, Lockhart E, Flynn JL: **Cytotoxicity and secretion of gamma interferon are carried out by distinct CD8 T cells during Mycobacterium tuberculosis infection.** *Infect Immun* 2009, **77**:4621–4630. <http://dx.doi.org/10.1128/IAI.00415-09>. Epub 2009/08/12. [pii] 10.1128/IAI.00415-09. PubMed PMID: [19667047](#); PubMed Central PMCID: [PMC2747936](#).
 115. Fuller CL, Flynn JL, Reinhart TA: **In situ study of abundant expression of proinflammatory chemokines and cytokines in pulmonary granulomas that develop in cynomolgus macaques experimentally infected with Mycobacterium tuberculosis.** *Infect Immun* 2003, **71**:7023–7034. Epub 2003/11/26. PubMed PMID: [14638792](#); PubMed Central PMCID: [PMC308896](#).
 116. Algood HM, Lin PL, Yankura D, Jones A, Chan J, Flynn JL: **TNF influences chemokine expression of macrophages in vitro and that of CD11b+ cells in vivo during Mycobacterium tuberculosis infection.** *J Immunol* 2004, **172**:6846–6857. Epub 2004/05/22. PubMed PMID: [15153503](#); PubMed Central PMCID: [PMCn/a](#).
 117. Clay H, Davis JM, Beery D, Huttenlocher A, Lyons SE, Ramakrishnan L: **Dichotomous role of the macrophage in early Mycobacterium marinum infection of the zebrafish.** *Cell Host Microbe* 2007, **2**:29–39. <http://dx.doi.org/10.1016/j.chom.2007.06.004>. Epub 2007/11/17. [pii] S1931-3128(07)00130-8. PubMed PMID: [18005715](#).
 118. Cosma CL, Humbert O, Ramakrishnan L: **Superinfecting mycobacteria home to established tuberculous granulomas.** *Nat Immunol* 2004, **5**:828–835. <http://dx.doi.org/10.1038/ni1091>. Epub 2004/06/29. ni1091 [pii]. PubMed PMID: [15220915](#).
 119. Bodnar KA, Serbina NV, Flynn JL: **Fate of Mycobacterium tuberculosis within murine dendritic cells.** *Infect Immun* 2001, **69**:800–809. <http://dx.doi.org/10.1128/IAI.69.2.800-809.2001>. Epub 2001/02/13. PubMed PMID: [11159971](#); PubMed Central PMCID: [PMC97955](#).
 120. Remoli ME, Giacomini E, Petruccioli E, Gafa V, Severa M, Gagliardi MC, Iona E, Pine R, Nisini R, Coccia EM: **Bystander inhibition of dendritic cell differentiation by Mycobacterium tuberculosis-induced IL-10.** *Immunol Cell Biol* 2010, <http://dx.doi.org/10.1038/icb.2010.106>. Epub 2010/09/02. [pii] 10.1038/icb.2010.106. PubMed PMID: [20805841](#).
 121. Robinson RT, Khader SA, Martino CA, Fountain JJ, Teixeira-Coelho M, Pearl JE, Smiley ST, Winslow GM, Woodland DL, Walter MJ, Conejo-Garcia JR, Gubler U, Cooper AM: **Mycobacterium tuberculosis infection induces il12rb1 splicing to generate a novel IL-12Rbeta1 isoform that enhances DC migration.** *J Exp Med* 2010, **207**:591–605. <http://dx.doi.org/10.1084/jem.20091085>. Epub 2010/03/10. [pii] 10.1084/jem.20091085. PubMed PMID: [20212068](#); PubMed Central PMCID: [PMC2839154](#).
 122. Tian T, Woodworth J, Skold M, Behar SM: **In vivo depletion of CD11c+ cells delays the CD4+ T cell response to Mycobacterium tuberculosis and exacerbates the outcome of infection.** *J Immunol* 2005, **175**:3268–3272. PubMed PMID: [16116218](#).
 123. Scott-Browne JP, Shafiani S, Tucker-Heard G, Ishida-Tsubota K, Fontenot JD, Rudensky AY, Bevan MJ, Urdahl KB: **Expansion and function of Foxp3-expressing T regulatory cells during tuberculosis.** *J Exp Med* 2007, **204**:2159–2169. <http://dx.doi.org/10.1084/jem.20062105>. Epub 2007/08/22. [pii] 10.1084/jem.20062105. PubMed PMID: [17709423](#); PubMed Central PMCID: [PMC2118702](#).
 124. Windish HP, Lin PL, Mattila JT, Green AM, Onuoha EO, Kane LP, Flynn JL: **Aberrant TGF-beta signaling reduces T regulatory cells in ICAM-1-deficient mice, increasing the inflammatory response to Mycobacterium tuberculosis.** *J Leukoc Biol* 2009,

- 86:713–725, <http://dx.doi.org/10.1189/jlb.1208740>. Epub 2009/05/21, [pii] 10.1189/jlb.1208740. PubMed PMID: 19454651.
125. Kursar M, Koch M, Mittrucker HW, Nouailles G, Bonhagen K, Kamradt T, Kaufmann SH: **Cutting edge: regulatory T cells prevent efficient clearance of *Mycobacterium tuberculosis*.** *J Immunol* 2007, **178**:2661–2665. PubMed PMID: 17312107.
 126. Burl S, Hill PC, Jeffries DJ, Holland MJ, Fox A, Lugos MD, Adegbola RA, Rook GA, Zumla A, McAdam KP, Brookes RH: **FOXP3 gene expression in a tuberculosis case contact study.** *Clin Exp Immunol* 2007, **149**:117–122, <http://dx.doi.org/10.1111/j.1365-2249.2007.03399.x>. Epub 2007/05/01, [pii] 10.1111/j.1365-2249.2007.03399.x. PubMed PMID: 17465993; PubMed Central PMCID: PMC1942016.
 127. Green AM, Mattila JT, Bigbee CL, Bongers KS, Lin PL, Flynn JL: **CD4(+) regulatory T cells in a cynomolgus macaque model of *Mycobacterium tuberculosis* infection.** *J Infect Dis* 2010, **202**:533–541, <http://dx.doi.org/10.1086/654896>. Epub 2010/07/14, PubMed PMID: 20617900.
 128. Lazarevic V, Nolt D, Flynn JL: **Long-term control of *Mycobacterium tuberculosis* infection is mediated by dynamic immune responses.** *J Immunol* 2005, **175**:1107–1117. Epub 2005/07/09, [pii]. PubMed PMID: 16002712.
 129. Lin F, Butcher EC: **T cell chemotaxis in a simple microfluidic device.** *Lab Chip* 2006, **6**:1462–1469, <http://dx.doi.org/10.1039/b607071j>. Epub 2006/10/27, PubMed PMID: 17066171.
 130. Darrah PA, Patel DT, De Luca PM, Lindsay RW, Davey DF, Flynn BJ, Hoff ST, Andersen P, Reed SG, Morris SL, Roederer M, Seder RA: **Multifunctional TH1 cells define a correlate of vaccine-mediated protection against *Leishmania major*.** *Nat Med* 2007, **13**:843–850, <http://dx.doi.org/10.1038/nm1592>. Epub 2007/06/15, [pii] 10.1038/nm1592. PubMed PMID: 17558415.
 131. Cooper AM, Dalton DK, Stewart TA, Griffin JP, Russell DG, Orme IM: **Disseminated tuberculosis in interferon gamma gene-disrupted mice.** *J Exp Med* 1993, **178**:2243–2247. Epub 1993/12/01. PubMed PMID: 8245795; PubMed Central PMCID: PMC2191280.
 132. Al-Muhsen S, Casanova JL: **The genetic heterogeneity of mendelian susceptibility to mycobacterial diseases.** *J Allergy Clin Immunol* 2008, **122**:1043–1051, <http://dx.doi.org/10.1016/j.jaci.2008.10.037>. quiz 52-3. Epub 2008/12/17, [pii] S0091-6749(08)01919-2. PubMed PMID: 19084105.
 133. Peters W, Scott HM, Chambers HF, Flynn JL, Charo IF, Ernst JD: **Chemokine receptor 2 serves an early and essential role in resistance to *Mycobacterium tuberculosis*.** *Proc Natl Acad Sci U S A* 2001, **98**:7958–7963. PubMed PMID: 11438742; PubMed Central PMCID: PMC650000.
 134. Scott HM, Flynn JL: ***Mycobacterium tuberculosis* in chemokine receptor 2-deficient mice: influence of dose on disease progression.** *Infect Immun* 2002, **70**:5946–5954. PubMed PMID: 12379669.
 135. Algood HM, Flynn JL: **CCR5-deficient mice control *Mycobacterium tuberculosis* infection despite increased pulmonary lymphocytic infiltration.** *J Immunol* 2004, **173**:3287–3296. Epub 2004/08/24, [pii]. PubMed PMID: 15322191.
 136. Algood HM, Lin PL, Flynn JL: **Tumor necrosis factor and chemokine interactions in the formation and maintenance of granulomas in tuberculosis.** *Clin Infect Dis* 2005, **41**(Suppl. 3):S189–S193, <http://dx.doi.org/10.1086/429994>. Epub 2005/06/29, [pii] 10.1086/429994. PubMed PMID: 15983898.
 137. Chakravarty SD, Xu J, Lu B, Gerard C, Flynn J, Chan J: **The chemokine receptor CXCR3 attenuates the control of chronic *Mycobacterium tuberculosis* infection in BALB/c mice.** *J Immunol* 2007, **178**:1723–1735. Epub 2007/01/24, [pii]. PubMed PMID: 17237422.
 138. Sallusto F: **The role of chemokines and chemokine receptors in T cell priming and Th1/Th2-mediated responses.** *Haematologica* 1999, **84**(Suppl. EHA-4):28–31. Epub 2000/07/25. PubMed PMID: 10907461.
 139. Rhoades ER, Cooper AM, Orme IM: **Chemokine response in mice infected with *Mycobacterium tuberculosis*.** *Infect Immun* 1995, **63**:3871–3877. Epub 1995/10/01. PubMed PMID: 7558294; PubMed Central PMCID: PMC173545.
 140. Khader SA, Rangel-Moreno J, Fountain JJ, Martino CA, Reiley WW, Pearl JE, Winslow GM, Woodland DL, Randall TD, Cooper AM: **In a murine tuberculosis model, the absence of homeostatic chemokines delays granuloma formation and protective immunity.** *J Immunol* 2009, **183**:8004–8014, <http://dx.doi.org/10.4049/jimmunol.0901937>. Epub 2009/11/26, [pii] 10.4049/jimmunol.0901937. PubMed PMID: 19933855; PubMed Central PMCID: PMC2799945.
 141. Tsai MC, Chakravarty S, Zhu G, Xu J, Tanaka K, Koch C, Tufariello J, Flynn J, Chan J: **Characterization of the tuberculous granuloma in murine and human lungs: cellular composition and relative tissue oxygen tension.** *Cell Microbiol* 2006, **8**:218–232. PubMed PMID: 16441433.
 142. Capuano 3rd SV, Croix DA, Pawar S, Zinovic A, Myers A, Lin PL, Bissel S, Fuhrman C, Klein E, Flynn JL: **Experimental *Mycobacterium tuberculosis* infection of cynomolgus macaques closely resembles the various manifestations of human *M. tuberculosis* infection.** *Infect Immun* 2003, **71**:5831–5844. Epub 2003/09/23. PubMed PMID: 14500505; PubMed Central PMCID: PMC201048.
 143. Swaim LE, Connolly LE, Volkman HE, Humbert O, Born DE, Ramakrishnan L: ***Mycobacterium marinum* infection of adult zebrafish causes caseating granulomatous tuberculosis and is moderated by adaptive immunity.** *Infect Immun* 2006, **74**:6108–6117, <http://dx.doi.org/10.1128/IAI.00887-06>. Epub 2006/10/24, [pii] 10.1128/IAI.00887-06. PubMed PMID: 17057088; PubMed Central PMCID: PMC1695491.
 144. Volkman HE, Pozos TC, Zheng J, Davis JM, Rawls JF, Ramakrishnan L: **Tuberculous granuloma induction via interaction of a bacterial secreted protein with host epithelium.** *Science* 2010, **327**:466–469, <http://dx.doi.org/10.1126/science.1179663>. Epub 2009/12/17, [pii] 10.1126/science.1179663. PubMed PMID: 20007864.
 145. Davis JM, Ramakrishnan L: **The role of the granuloma in expansion and dissemination of early tuberculous infection.** *Cell* 2009, **136**:37–49, <http://dx.doi.org/10.1016/j.cell.2008.11.014>. Epub 2009/01/13, [pii] S0092-8674(08)01443-8. PubMed PMID: 19135887.
 146. Gammack D, Ganguli S, Marino S, Segovia-Juarez J, Kirschner DE: **Understanding the immune response in tuberculosis using different mathematical models and biological scales.** *Multiscale Model Sim* 2005, **3**:312–345. PubMed PMID: n/a; PubMed Central PMCID: PMC650000.
 147. Lauffenburger DA, Linderman JJ: **Cell surface receptor/ligand binding fundamentals. Receptors: models for binding, trafficking, and signaling.** New York: Oxford University Press; 1993:9–72.
 148. Cilfone NA, Kirschner DE, Linderman JJ: **Strategies for efficient numerical implementation of hybrid multi-scale agent-based models to describe biological systems.** *Cel Mol Bioeng* 2014; 1–18, <http://dx.doi.org/10.1007/s12195-014-0363-6>.
 149. Kirschner DE, Chang ST, Riggs TW, Perry N, Linderman JJ: **Toward a multiscale model of antigen presentation in immunity.** *Immunol Rev* 2007, **216**:93–118, <http://dx.doi.org/10.1111/j.1600-065X.2007.00490.x>. Epub 2007/03/21, PubMed PMID: 17367337; PubMed Central PMCID: PMCn/a.
 150. Agliari E, Asti L, Barra A, Scrivo R, Valesini G, Wallis RS: **Application of a stochastic modeling to assess the evolution of tuberculous and non-tuberculous mycobacterial infection in patients treated with tumor necrosis factor inhibitors.** *PLoS One* 2013, **8**:e55017, <http://dx.doi.org/10.1371/journal.pone.0055017>. PubMed PMID: 23383039; PubMed Central PMCID: PMC3557254.
 151. Mattila JT, Ojo OO, Kepka-Lenhart D, Marino S, Kim JH, Eum SY, Via LE, Barry 3rd CE, Klein E, Kirschner DE, Morris Jr SM, Lin PL, Flynn JL: **Microenvironments in tuberculous granulomas are delineated by distinct populations of macrophage subsets and expression of nitric oxide synthase and arginase isoforms.** *J Immunol* 2013, **191**:773–784, <http://dx.doi.org/10.4049/jimmunol.1300113>. Epub 2013/06/12, PubMed PMID: 23749634; PubMed Central PMCID: PMC3746594.
 152. Lin PL, Coleman T, Carney JP, Lopresti BJ, Tomko J, Fillmore D, Dartois V, Scanga C, Frye LJ, Janssen C, Klein E, Barry 3rd CE, Flynn JL: **Radiologic responses in cynomolgus macaques for assessing tuberculosis chemotherapy regimens.**

- Antimicrob Agents Chemother* 2013, <http://dx.doi.org/10.1128/AAC.00277-13>. Epub 2013/06/26, PubMed PMID: 23796926; PubMed Central PMCID: PMC3754323.
153. Linderman JJ, Kirschner DE: **In silico models of M. tuberculosis infection provide a route to new therapies.** *Drug Discov Today Dis Models* 2014, <http://dx.doi.org/10.1016/j.ddmod.2014.02.006>. ePub 2014/02/04, PubMed PMID: 24487820.
 154. Egelund EF, Barth AB, Peloquin CA: **Population pharmacokinetics and its role in anti-tuberculosis drug development and optimization of treatment.** *Curr Pharm Des* 2011, **17**: 2889–2899. PubMed PMID: 21834766.
 155. Wilkins JJ, Savic RM, Karlsson MO, Langdon G, McIlleron H, Pillai G, Smith PJ, Simonsson US: **Population pharmacokinetics of rifampin in pulmonary tuberculosis patients, including a semimechanistic model to describe variable absorption.** *Antimicrob Agents Chemother* 2008, **52**:2138–2148, <http://dx.doi.org/10.1128/AAC.00461-07>. Epub 2008/04/09, PubMed PMID: 18391026; PubMed Central PMCID: PMC2415769.
 156. Wilkins JJ, Langdon G, McIlleron H, Pillai G, Smith PJ, Simonsson US: **Variability in the population pharmacokinetics of isoniazid in South African tuberculosis patients.** *Br J Clin Pharmacol* 2011, **72**:51–62, <http://dx.doi.org/10.1111/j.1365-2125.2011.03940.x>. PubMed PMID: 21320152; PubMed Central PMCID: PMC3141186.
 157. Peloquin CA, Hadad DJ, Molino LP, Palaci M, Boom WH, Dietze R, Johnson JL: **Population pharmacokinetics of levofloxacin, gatifloxacin, and moxifloxacin in adults with pulmonary tuberculosis.** *Antimicrob Agents Chemother* 2008, **52**: 852–857, <http://dx.doi.org/10.1128/AAC.01036-07>. PubMed PMID: 18070980; PubMed Central PMCID: PMC2258503.
 158. Jonsson S, Davidsen A, Wilkins J, Van der Walt JS, Simonsson US, Karlsson MO, Smith P, McIlleron H: **Population pharmacokinetics of ethambutol in South African tuberculosis patients.** *Antimicrob Agents Chemother* 2011, **55**: 4230–4237, <http://dx.doi.org/10.1128/AAC.00274-11>. PubMed PMID: 21690284; PubMed Central PMCID: PMC3165318.
 159. Chigutsa E, Meredith S, Wiesner L, Padayatchi N, Harding J, Moodley P, Mac Kenzie WR, Weiner M, McIlleron H, Kirkpatrick CM: **Population pharmacokinetics and pharmacodynamics of ofloxacin in South African patients with multidrug-resistant tuberculosis.** *Antimicrob Agents Chemother* 2012, **56**:3857–3863, <http://dx.doi.org/10.1128/AAC.00048-12>. PubMed PMID: 22564839; PubMed Central PMCID: PMC3393408.
 160. Svensson EM, Murray S, Karlsson MO, Dooley KE: **Rifampicin and rifapentine significantly reduce concentrations of bedaquiline, a new anti-TB drug.** *J Antimicrob Chemother* 2015, **70**: 1106–1114, <http://dx.doi.org/10.1093/jac/dku504>. PubMed PMID: 25535219; PubMed Central PMCID: PMC4356204.
 161. Denti P, Jeremiah K, Chigutsa E, Faurholt-Jepsen D, PrayGod G, Range N, Castel S, Wiesner L, Hagen CM, Christiansen M, Chagalucha J, McIlleron H, Friis H, Andersen AB: **Pharmacokinetics of isoniazid, pyrazinamide, and ethambutol in newly diagnosed pulmonary TB patients in Tanzania.** *PLoS One* 2015, **10**:e0141002, <http://dx.doi.org/10.1371/journal.pone.0141002>. PubMed PMID: 26501782; PubMed Central PMCID: PMC4621059.
 162. Prideaux B, Via LE, Zimmerman MD, Eum S, Sarathy J, O'Brien P, Chen C, Kaya F, Weiner DM, Chen PY, Song T, Lee M, Shim TS, Cho JS, Kim W, Cho SN, Olivier KN, Barry 3rd CE, Dartois V: **The association between sterilizing activity and drug distribution into tuberculosis lesions.** *Nat Med* 2015, **21**:1223–1227, <http://dx.doi.org/10.1038/nm.3937>. PubMed PMID: 26343800; PubMed Central PMCID: PMC4598290.
 163. Lenaerts A, Barry 3rd CE, Dartois V: **Heterogeneity in tuberculosis pathology, microenvironments and therapeutic responses.** *Immunol Rev* 2015, **264**:288–307, <http://dx.doi.org/10.1111/immr.12252>. Epub 2015/02/24, PubMed PMID: 25703567.
 164. Prideaux B, ElNaggar MS, Zimmerman M, Wiseman JM, Li X, Dartois V: **Mass spectrometry imaging of levofloxacin distribution in TB-infected pulmonary lesions by MALDI-MSI and continuous liquid microjunction surface sampling.** *Int J Mass Spectrom* 2015, **377**:699–708, <http://dx.doi.org/10.1016/j.jms.2014.08.024>. PubMed PMID: 26185484; PubMed Central PMCID: PMC4501920.
 165. Dartois V: **The path of anti-tuberculosis drugs: from blood to lesions to mycobacterial cells.** *Nat Rev Microbiol* 2014, **12**: 159–167, <http://dx.doi.org/10.1038/nrmicro3200>. Epub 2014/02/04, PubMed PMID: 24487820.
 166. Pienaar E, Sarathy J, Prideaux B, Dartois V, Kirschner D, Linderman JJ: **Multi-scale systems pharmacology comparison of moxifloxacin, levofloxacin and gatifloxacin efficacy in tuberculosis granulomas.** 2016. Submitted.
 167. Dheda K, Barry 3rd CE, Maartens G: **Tuberculosis.** *Lancet* 2016, **387**:1211–1226, [http://dx.doi.org/10.1016/S0140-6736\(15\)00151-8](http://dx.doi.org/10.1016/S0140-6736(15)00151-8). PubMed PMID: 26377143.
 168. Johnson PL, Kochin BF, McAfee MS, Stromnes IM, Regoes RR, Ahmed R, Blattman JN, Antia R: **Vaccination alters the balance between protective immunity, exhaustion, escape, and death in chronic infections.** *J Virol* 2011, **85**:5565–5570, <http://dx.doi.org/10.1128/JVI.00166-11>. PubMed PMID: 21411537; PubMed Central PMCID: PMC3094965.
 169. Akondy RS, Johnson PL, Nakaya HI, Edupuganti S, Mulligan MJ, Lawson B, Miller JD, Pulendran B, Antia R, Ahmed R: **Initial viral load determines the magnitude of the human CD8 T cell response to yellow fever vaccination.** *Proc Natl Acad Sci U S A* 2015, **112**:3050–3055, <http://dx.doi.org/10.1073/pnas.1500475112>. PubMed PMID: 25713354; PubMed Central PMCID: PMC4364194.
 170. Wherry EJ, Teichgraber V, Becker TC, Masopust D, Kaech SM, Antia R, von Andrian UH, Ahmed R: **Lineage relationship and protective immunity of memory CD8 T cell subsets.** *Nat Immunol* 2003, **4**:225–234, <http://dx.doi.org/10.1038/ni889>. PubMed PMID: 12563257.
 171. Gong C, Linderman JJ, Kirschner D: **Harnessing the heterogeneity of T cell differentiation fate to fine-tune generation of effector and memory T cells.** *Front Immunol* 2014, **5**:57, <http://dx.doi.org/10.3389/fimmu.2014.00057>. Epub 2014/03/07, PubMed PMID: 24600448; PubMed Central PMCID: PMC3928592.
 172. Nunes-Alves C, Booty MG, Carpenter SM, Jayaraman P, Rothchild AC, Behar SM: **In search of a new paradigm for protective immunity to TB.** *Nat Rev Microbiol* 2014, **12**: 289–299, <http://dx.doi.org/10.1038/nrmicro3230>. Epub 2014/03/05, nrmicro3230 [pii]. PubMed PMID: 24590243.
 173. Wallis RS, Kim P, Cole S, Hanna D, Andrade BB, Maeurer M, Schito M, Zumla A: **Tuberculosis biomarkers discovery: developments, needs, and challenges.** *Lancet Infect Dis* 2013, **13**:362–372, [http://dx.doi.org/10.1016/S1473-3099\(13\)70034-3](http://dx.doi.org/10.1016/S1473-3099(13)70034-3). Epub 2013/03/28, S1473-3099(13)70034-3 [pii]. PubMed PMID: 23531389.
 174. Walzl G, Ronacher K, Hanekom W, Scriba TJ, Zumla A: **Immunological biomarkers of tuberculosis.** *Nat Rev Immunol* 2011, **11**:343–354, <http://dx.doi.org/10.1038/nri2960>. Epub 2011/04/09, nri2960 [pii]. PubMed PMID: 21475309.
 175. Whitworth HS, Aranday-Cortes E, Lavalani A: **Biomarkers of tuberculosis: a research roadmap.** *Biomark Med* 2013, **7**: 349–362, <http://dx.doi.org/10.2217/bmm.13.53>. Epub 2013/06/06, PubMed PMID: 23734796.
 176. Flynn JL, Chan J: **Immunology of tuberculosis.** *Annu Rev Immunol* 2001, **19**:93–129, <http://dx.doi.org/10.1146/annurev.immunol.19.1.93>. Epub 2001/03/13, [pii] 10.1146/annurev.immunol.19.1.93. PubMed PMID: 11244032; PubMed Central PMCID: PMC4598290.
 177. Saltini C: **Schedule or dosage? The need to perfect intermittent regimens for tuberculosis.** *Am J Respir Crit Care Med* 2006, **174**:1067–1068, <http://dx.doi.org/10.1164/rccm.200608-1194ED>. PubMed PMID: 17085748.
 178. Chang KC, Leung CC, Grosset J, Yew WW: **Treatment of tuberculosis and optimal dosing schedules.** *Thorax* 2011, **66**: 997–1007, <http://dx.doi.org/10.1136/thx.2010.148585>. PubMed PMID: 21169286.
 179. Zumla A, Gillespie SH, Hoelscher M, Phillips PP, Cole ST, Abubakar I, McHugh TD, Schito M, Maeurer M, Nunn AJ: **New**

- antituberculosis drugs, regimens, and adjunct therapies: needs, advances, and future prospects. *Lancet Infect Dis* 2014, **14**:327–340, [http://dx.doi.org/10.1016/S1473-3099\(13\)70328-1](http://dx.doi.org/10.1016/S1473-3099(13)70328-1). PubMed PMID: 24670627.
180. Ford CB, Shah RR, Maeda MK, Gagneux S, Murray MB, Cohen T, Johnston JC, Gardy J, Lipsitch M, Fortune SM: **Mycobacterium tuberculosis mutation rate estimates from different lineages predict substantial differences in the emergence of drug-resistant tuberculosis.** *Nat Genet* 2013, **45**:784–790, <http://dx.doi.org/10.1038/ng2656>. PubMed PMID: 23749189; PubMed Central PMCID: PMC3777616.
 181. Colijn C, Cohen T, Ganesh A, Murray M: **Spontaneous emergence of multiple drug resistance in tuberculosis before and during therapy.** *PLoS One* 2011, **6**:e18327, <http://dx.doi.org/10.1371/journal.pone.0018327>. Epub 2011/04/12, PubMed PMID: 21479171; PubMed Central PMCID: PMC3068161.
 182. Hansen E, Woods RJ, Read AF: **How to use a chemotherapeutic agent when resistance to it threatens the patient.** *PLoS Biol* 2017, **15**:e2001110, <http://dx.doi.org/10.1371/journal.pbio.2001110>. PubMed PMID: 28182734; PubMed Central PMCID: PMC5300106.
 183. Lin YJ, Liao CM: **Quantifying the impact of drug combination regimens on TB treatment efficacy and multidrug resistance probability.** *J Antimicrob Chemother* 2015, **70**:3273–3282, <http://dx.doi.org/10.1093/jac/dkv247>. PubMed PMID: 26311836.
 184. Luria SE, Delbruck M: **Mutations of bacteria from virus sensitivity to virus resistance.** *Genetics* 1943, **28**:491–511. PubMed PMID: 17247100; PubMed Central PMCID: PMC1209226.
 185. Kirschner DE, Hunt CA, Marino S, Fallahi-Sichani M, Linderman JJ: **Tuneable resolution as a systems biology approach for multi-scale, multi-compartment computational models.** *Wiley Interdiscip Rev Syst Biol Med* 2014, **6**:289–309, <http://dx.doi.org/10.1002/wsbm.1270>. PubMed PMID: 24810243; PubMed Central PMCID: PMC4102180.
 186. Conzelmann H, Saez-Rodriguez J, Sauter T, Bullinger E, Allgower F, Gilles ED: **Reduction of mathematical models of signal transduction networks: simulation-based approach applied to EGF receptor signalling.** *Syst Biol (Stevenage)* 2004, **1**:159–169. PubMed PMID: 17052126; PubMed Central PMCID: PMCexempt.
 187. Maurya MR, Bornheimer SJ, Venkatasubramanian V, Subramaniam S: **Reduced-order modelling of biochemical networks: application to the GTPase-cycle signalling module.** *Syst Biol (Stevenage)* 2005, **152**:229–242. Epub 2006/09/22. PubMed PMID: 16986265; PubMed Central PMCID: PMC3417759.
 188. Maurya MR, Katore S, Patkar PR, Rundell AE, Venkatasubramanian V: **A systematic framework for the design of reduced-order models for signal transduction pathways from a control theoretic perspective.** *Comput Chem Eng* 2006, **30**:437–452, <http://dx.doi.org/10.1016/j.compchemeng.2005.10.003>.
 189. Guzzetta G, Kirschner D: **The roles of immune memory and aging in protective immunity and endogenous reactivation of tuberculosis.** *PLoS One* 2013, **8**:e60425, <http://dx.doi.org/10.1371/journal.pone.0060425>. Epub 2013/04/13, PubMed PMID: 23580062; PubMed Central PMCID: PMC3620273.
 190. Bhunu CP, Garira W, Mukandavire Z, Magombedze G: **Modelling the effects of pre-exposure and post-exposure vaccines in tuberculosis control.** *J Theor Biol* 2008, **254**:633–649, <http://dx.doi.org/10.1016/j.jtbi.2008.06.023>. Epub 2008/07/23, [pii] S0022-5193(08)00309-3. PubMed PMID: 18644386.
 191. Holland DP, Sanders GD, Hamilton CD, Stout JE: **Strategies for treating latent multiple-drug resistant tuberculosis: a decision analysis.** *PLoS One* 2012, **7**:e30194, <http://dx.doi.org/10.1371/journal.pone.0030194>. PubMed PMID: 22272302; PubMed Central PMCID: PMC3260212.
 192. Rejniak KA, Anderson AR: **Hybrid models of tumor growth.** *Wiley Interdiscip Rev Syst Biol Med* 2011, **3**:115–125, <http://dx.doi.org/10.1002/wsbm.102>. PubMed PMID: 21064037; PubMed Central PMCID: PMC3057876.
 193. Mac Gabhann F, Popel AS: **Systems biology of vascular endothelial growth factors.** *Microcirculation* 2008, **15**:715–738, <http://dx.doi.org/10.1080/10739680802095964>. PubMed PMID: 18608994; PubMed Central PMCID: PMC2722038.
 194. Vodovotz Y, Clermont G, Chow C, An G: **Mathematical models of the acute inflammatory response.** *Curr Opin Crit Care* 2004, **10**:383–390. PubMed PMID: 15385756.
 195. Flamm MH, Colace TV, Chatterjee MS, Jing H, Zhou S, Jaeger D, Brass LF, Sinno T, Diamond SL: **Multiscale prediction of patient-specific platelet function under flow.** *Blood* 2012, **120**:190–198, <http://dx.doi.org/10.1182/blood-2011-10-388140>. PubMed PMID: 22517902; PubMed Central PMCID: PMC3390957.
 196. Peirce SM, Mac Gabhann F, Bautch VL: **Integration of experimental and computational approaches to sprouting angiogenesis.** *Curr Opin Hematol* 2012, **19**:184–191, <http://dx.doi.org/10.1097/MOH.0b013e3283523ea6>. PubMed PMID: 22406822; PubMed Central PMCID: PMC4132663.
 197. Siewe N, Yakubu AA, Satoskar AR, Friedman A: **Granuloma formation in leishmaniasis: a mathematical model.** *J Theor Biol* 2016, **412**:48–60, <http://dx.doi.org/10.1016/j.jtbi.2016.10.004>. PubMed PMID: 27769685.