

stpois: Fast Poisson event-history regression with high-dimensional fixed effects

Andreas Ljungström

Swedish Institute for Social Research (SOFI), Stockholm University

`andreas.ljungstrom@sofi.su.se`

Abstract

Poisson regression on episode-split survival data is a workhorse of event-history analysis, but two practical obstacles limit its use with full-population register data: models with rich fixed-effect structures become slow or infeasible to estimate with explicit dummy variables, owing to the quadratic growth in memory and computation time with the parameter count, and continuous covariates prevent the collapsing of individual records to risk cells that otherwise makes such models fast. This article introduces the community-contributed command `stpois`, which fits Poisson event-history models directly on `stset` data through three estimation paths, all of which return maximum-likelihood estimates that agree with `poisson` to at least 10^{-6} : a standard path that wraps `poisson` with automatic exposure; a high-dimensional fixed-effects path implemented via iteratively reweighted alternating projections in Mata; and a fast path that accelerates estimation through the cell structure of the categorical covariates. The fast path computes each Newton iteration from exponentially tilted within-cell moments in a single pass over the microdata. Coefficients, log likelihoods, and observed-information, robust, and cluster-robust standard errors agree with the corresponding `poisson` model on every path to at least 10^{-6} (typically $\sim 10^{-8}$); factor-variable interactions and weights are supported throughout, the two acceleration mechanisms can be combined, and postestimation predictions after absorption include the recovered fixed effects. Benchmarks against `streg`, `poisson`, and `ppmlhdfc` document the speed gains in three settings: episode-split workflows, all-categorical rate models, and models with high-dimensional fixed effects.

Keywords: `stpois`, `stpois`, event-history analysis, Poisson regression, piecewise exponential model, high-dimensional fixed effects, register data

1 Introduction

The equivalence between piecewise-constant hazard models and Poisson regression on episode-split data is one of the older and more useful results in event-history analysis (Holford, 1980; Laird and Olivier, 1981; Whitehead, 1980). Splitting each subject’s time at risk into episodes, treating the event indicator in each episode as a Poisson count, and entering the log of exposure time as an offset yields maximum-likelihood estimates of a proportional-hazards model with a hazard that is constant within episodes. In Stata,

this model can be fitted with `poisson` after manually generating the exposure variable, or—for the single-episode exponential case—with `streg, distribution(exponential)`. The approach is standard in epidemiology (Clayton and Hills, 1993; Carstensen, 2007) and is covered at length by Rabe-Hesketh and Skrondal (2022).

Two practical obstacles arise when this framework meets modern administrative data. Full-population registers of the kind available in the Scandinavian countries routinely produce analysis files with millions or tens of millions of episode records, particularly after splitting on calendar time, age, and duration. First, models with rich fixed-effect structures—birth cohorts, municipalities, detailed occupational codes, or their interactions—become slow or infeasible when the fixed effects are entered as explicit dummy variables, because both estimation time and memory grow quadratically with the parameter count (and because Stata’s dummy-variable trap limits the practical number of indicator columns). The `ppmlhdfc` command (Correia et al., 2020) solves this problem for general Poisson models, building on the alternating-projections approach of Guimarães and Portugal (2010); it depends on the `ftools` and `reghdfe` packages, so deploying it means installing that chain, which can add friction in the locked-down secure environments where register data must typically be analyzed. Second, and less widely appreciated, the Poisson likelihood for purely categorical covariates depends on the data only through the sums of events and exposure within covariate cells. Estimation on the collapsed cell data is then *exact* and can reduce the per-iteration operation count by orders of magnitude when the number of cells is far smaller than the number of records. A single continuous covariate breaks this property, and analysts face an unattractive choice between discretizing the covariate and forgoing the speed advantage entirely.

The command described in this article, `stpois`, addresses both obstacles within a single interface for `stset` data. It offers three estimation paths, all of which return maximum-likelihood estimates and standard errors that agree with `poisson` to the tolerances documented in section 8:

1. A **standard path** that wraps `poisson`, using the event indicator `_d` as outcome and time at risk `_t - _t0` as exposure. Coefficients and standard errors are identical to `streg, distribution(exponential)`.
2. A **high-dimensional fixed-effects path**, `absorb()`, implementing iteratively reweighted least squares with alternating-projections demeaning in Mata—the same mathematical approach as `ppmlhdfc`, but self-contained with no dependencies. Absorbed terms may be single variables or interactions like `region#period`. Observed-information, robust, and cluster-robust standard errors agree with those from the corresponding dummy-variable model to at least 10^{-6} .
3. A **fast path**, option `fast`, which accelerates exact estimation through the cell structure of the categorical covariates. Newton iterations on the full likelihood are computed from exponentially tilted within-cell moments of the individual-level covariates, accumulated in a single pass over the microdata per iteration; all remaining algebra runs on the cells. When every covariate is categorical, the iterations run entirely on the collapsed cell sums. The fast path requires at least one categorical term to define the cells.

Factor-variable interactions—`continuous#continuous`, `continuous#categorical`, and `categorical#categorical`—are supported on all paths, as are frequency, importance, and

sampling weights, with weighted results matching the weighted `poisson` model to the same tolerance. The `fast` and `absorb()` paths may be combined: the absorbed fixed effects are then concentrated out of the cell-accelerated likelihood. Section 2 sets out the modeling framework, section 3 describes the fast path, and section 4 describes the fixed-effects algorithm. Sections 5 and 6 document the syntax and illustrate the command, and section 8 verifies the numerical claims and benchmarks the command against `streg` and `poisson`.

2 Poisson event-history regression

Consider `stset` data in which record i contributes an event indicator $d_i \in \{0, 1\}$ (Stata’s `_d`) and a time at risk $t_i = _t - _t0$. Under a proportional-hazards model with hazard rate $\lambda_i = \exp(\mathbf{x}_i' \boldsymbol{\beta})$ constant within records, the log likelihood for right-censored data is, up to a constant,

$$\ell(\boldsymbol{\beta}) = \sum_i \{d_i \mathbf{x}_i' \boldsymbol{\beta} + d_i \log t_i - t_i \exp(\mathbf{x}_i' \boldsymbol{\beta})\},$$

in which the $d_i \log t_i$ term is constant in $\boldsymbol{\beta}$; this is proportional to the likelihood of a Poisson model for d_i with mean $\mu_i = t_i \exp(\mathbf{x}_i' \boldsymbol{\beta})$ and offset $\log t_i$. Hence any software that fits Poisson regression with an offset fits this hazard model, and duration dependence or period effects are accommodated by episode splitting (`stsplit`) followed by indicator variables for the resulting time bands (Holford, 1980; Laird and Olivier, 1981). The Poisson “counts” here are 0/1 event indicators; nothing requires the outcome to be a genuine count, and the pseudo-maximum-likelihood results of [Gourieroux et al. \(1984\)](#) imply that the coefficient estimates are consistent whenever the log-linear mean is correctly specified, with robust standard errors available in the usual way (see also [Santos Silva and Tenreyro, 2006](#)).

`stpois` fits exactly this model on the current `stset` data. On the standard path, it constructs the exposure variable, calls `poisson`, and posts results with survival-flavored headers (numbers of subjects and failures, time at risk). The point of the command, however, is not the convenience wrapper but the two departures from it: the fast path (section 3) and the fixed-effects path (section 4).

2.1 Sufficiency of cell sums under categorical covariates

When all covariates are categorical, the covariate vector takes a finite number of distinct values, indexing cells $j = 1, \dots, J$. The log likelihood above then depends on the data only through the within-cell sums $D_j = \sum_{i \in j} d_i$ and $T_j = \sum_{i \in j} t_i$:

$$\ell(\boldsymbol{\beta}) = \sum_j \{D_j \mathbf{x}_j' \boldsymbol{\beta} - T_j \exp(\mathbf{x}_j' \boldsymbol{\beta})\} + \text{const},$$

(D_j, T_j) are sufficient statistics, and estimation on the J collapsed cells reproduces the individual-level maximum-likelihood estimates exactly—coefficients, standard errors, and all. With millions of episode records but a few hundred cells (period $\times$ education $\times$ region, say), the gain is dramatic, and this is the standard trick behind register-based rate modeling ([Clayton and Hills, 1993](#); [Dickman et al., 2004](#)). A continuous covariate destroys the sufficiency because $\exp(\gamma x_i)$ no longer aggregates: $\sum_{i \in j} t_i \exp(\gamma x_i) \neq T_j \exp(\gamma \bar{x}_j)$ by

strict convexity of the exponential, where $\bar{x}_j = T_j^{-1} \sum_{i \in j} t_i x_i$ is the exposure-weighted within-cell mean (the two sides coincide only in the degenerate case of no within-cell variation in x). The fast path generalizes the collapsing idea so that it delivers the exact maximum-likelihood estimator in the presence of continuous covariates as well.

3 Fast exact estimation: The fast option

3.1 Term classification and cells

With `fast`, `stpois` classifies each expanded term of the varlist by its pieces. Terms built entirely from factor-variable pieces—`i.edu`, `i.edu#i.region`—are constant within cells and enter the model at the cell level, with coefficient vector $\boldsymbol{\delta}$ on the cell-level design $\mathbf{w}_j$. Terms involving at least one continuous piece—`age`, `c.age#c.income`, `c.age#i.edu`—enter at the individual level, with coefficient vector $\boldsymbol{\gamma}$ on the individual-level design matrix $\mathbf{X}_i$ (the columns of the expanded model matrix corresponding to terms with at least one continuous piece), so that $\mu_i = t_i \exp(\boldsymbol{\gamma}'\mathbf{X}_i + \boldsymbol{\delta}'\mathbf{w}_{j(i)})$, where $j(i)$ denotes the cell of record i . The cells j are the cross-classification of the categorical variables appearing in the cell-level terms; at least one cell-level term is required.

3.2 Newton iterations on tilted cell moments

The score and observed information of the Poisson likelihood decompose into fixed cell sums and *exponentially tilted within-cell moments* (Jensen, 1995) of the individual-level covariates:

$$M_j = \sum_{i \in j} \mu_i, \quad \mathbf{m}_j = \sum_{i \in j} \mu_i \mathbf{X}_i, \quad \mathbf{Q} = \sum_i \mu_i \mathbf{X}_i \mathbf{X}_i'.$$

The score is $\left(\sum_i d_i \mathbf{X}_i - \sum_j \mathbf{m}_j, \sum_j \mathbf{w}_j (D_j - M_j) \right)$ and the information matrix has blocks $\mathbf{Q}$, $\sum_j \mathbf{m}_j \mathbf{w}_j'$, and $\sum_j M_j \mathbf{w}_j \mathbf{w}_j'$. The μ_i -weighting in these quantities is the standard Poisson information weight: McCullagh and Nelder (1989, sec. 2.5) derive that the IRLS working weight for observation i is $W_i = [V(\mu_i)(d\eta/d\mu)^2]^{-1}$; with variance function $V(\mu) = \mu$ and log-link derivative $d\eta/d\mu = \mu^{-1}$ this gives $W_i = \mu_i$, so M_j , $\mathbf{m}_j$, and $\mathbf{Q}$ are W -weighted sums of the exposure, the individual-level covariates, and their outer products. Because the log link is canonical for Poisson, the expected and observed information coincide (McCullagh and Nelder, 1989, p. 53), so Newton–Raphson and Fisher scoring reduce to the same algorithm and the step direction is algebraically exact. Each Newton iteration therefore requires one pass over the microdata to accumulate the tilted moments—at cost $O(np^2)$, with p the number of *individual-level* columns only—after which everything happens on the J cells. A full-model Newton step in `poisson`, by contrast, costs $O(n(p+k)^2)$ with k the number of cell-level parameters, so the per-iteration floating-point saving is a factor of roughly $(p+k)^2/p^2$, and it grows with the richness of the categorical structure. Step-halving on the log likelihood guards the updates: if the full Newton step does not increase the log likelihood, the step length is halved repeatedly until an improvement is found or the step shrinks to 2^{-16} of its original length, at which point the algorithm accepts the current iterate and continues; if the outer convergence criterion is still not met after `maxiter()` such steps, `e(converged)` is set to 0. Writing $\boldsymbol{\theta} = (\boldsymbol{\gamma}', \boldsymbol{\delta}')'$

for the stacked parameter vector, convergence is declared when $\max_{\ell} |\Delta\theta_{\ell}|/(1 + |\theta_{\ell}|)$, evaluated at the updated parameter vector, falls below `tolerance()`.

When *every* term is cell-level ($p = 0$), the sufficiency result of section 2.1 applies directly and the estimator depends on the data only through the collapsed sums (D_j, T_j) . In this case, with no absorbed fixed effects and the default observed-information VCE, `stpois` routes estimation to Stata’s compiled `collapse` followed by `poisson` on the cell sums: the fit is exact by sufficiency and faster than iterating in Mata over the microdata. The robust and cluster-robust variance estimators need the per-observation scores—the squared cell residual is not the cell sum of squared residuals—so under `vce(robust)` or `vce(cluster)` the pure cell-level model instead runs a Newton iteration on (D_j, T_j) at $O(Jk^2)$ per step, independent of the number of episode records.

This is Newton on the exact likelihood: the score/information decomposition involves no linearization or discretization, so the estimator is algebraically the full-model maximum-likelihood estimator, and the two maximizers differ only at their iterative convergence tolerance. In practice, coefficients, the log likelihood, and the observed-information standard errors agree with the full `poisson` model to at least 10^{-6} (typically $\sim 10^{-8}$), and robust and cluster-robust variance estimators are computed from the same per-observation scores as their `poisson` counterparts, with the HC1 and $G/(G-1)$ small-sample corrections (G the number of clusters). Because the continuous coefficients are estimated from the full individual-level likelihood, they are identified—with full efficiency—even when the continuous covariates do not vary across cells at all.

3.3 Implementation

The data handling is designed for large datasets, adopting the strategy popularized by the `ftools` project (Correia, 2016): group structures are computed from a single `order()` sort in Mata, segment boundaries are found by comparing adjacent sorted rows, and all within-group sums—the cell aggregates, the tilted moments, and the within-cluster score sums of the cluster-robust variance estimator—are running-sum differences over the sorted data, each an $O(n)$ pass independent of the number of groups. Interaction columns are constructed directly from the raw variables in Mata. In this Mata engine—used whenever the model has continuous covariates, and for the robust and cluster-robust cell-level fits—there are no `preserve/collapse` round-trips and no temporary variables in the dataset; beyond the single sort permutation, memory overhead is one copy of the loaded columns. (The one exception is the purely categorical model under the default observed-information VCE, which is delegated to Stata’s `collapse` and `poisson` as described in section 3.) With weights, every likelihood contribution, cell sum, and tilted moment is weight-multiplied, so the weighted estimates remain the exact weighted maximum-likelihood estimates. For very large datasets, running `compress` before estimation reduces the in-memory footprint of the loaded columns; the Mata engine creates no additional dataset variables beyond the single sort permutation.

3.4 Combining fast with absorbed fixed effects

`fast` may be combined with `absorb()` (section 4). The absorbed effects are then concentrated out of the likelihood by closed-form updates: for Poisson, the profile-maximizing effect of absorbed group g given the remaining parameters is $\alpha_g = \ln(\sum_{i \in g} d_i / \sum_{i \in g} \mu_i)$, and Gauss–Seidel cycles of this update across the absorbed terms are observed in practice

to converge to the profile optimum; formal convergence guarantees for the nonlinear profile Gauss–Seidel are not available—the approach is used by the R package `FENmlm` (Bergé, 2018) and its successor `fixest`. Each Newton step is then taken on the profile likelihood: the design columns (individual-level and cell-level) are weighted-demeaned within the absorbed groups, and the step uses the demeaned score and information. Given that the inner fixed-effect solve has converged to the profile optimum, this is, by the Frisch–Waugh–Lovell logic applied through the IRLS local linearization, the profile Newton direction and, at convergence, the partitioned variance matrix. Step-halving re-solves the fixed effects at every trial point, so the profile likelihood increases monotonically. Coefficients and all three variance estimators agree with `poisson` carrying explicit dummies for both the cell-level terms and the absorbed effects to at least 10^{-6} .

4 High-dimensional fixed effects: `absorb()`

The `absorb(terms)` option fits the exact model with one or more sets of fixed effects without creating dummy variables, using iteratively reweighted least squares (McCullagh and Nelder, 1989, sec. 2.5) (IRLS) combined with weighted alternating-projections demeaning—the approach introduced to Stata by Guimarães and Portugal (2010) and refined by Correia et al. (2020). Each absorbed term is a variable or an interaction such as `region#period`, absorbed as the cross-classification of its variables. Each IRLS iteration:

1. computes the working variable $z_i = \eta_i + (d_i - \mu_i)/\mu_i$ and weights $\omega_i = \mu_i$, where $\eta_i = \mathbf{x}_i' \boldsymbol{\beta}$ is the current linear predictor excluding the offset and $\mu_i = t_i \exp(\eta_i)$;
2. demeanes z and each covariate within the groups of each absorbed term in turn, iterating this Gauss–Seidel cycle to convergence (weighted alternating projections), which is guaranteed to converge for the linear demeaning problem (Halperin, 1962; Gaure, 2013) though at a rate that deteriorates as the absorbed dimensions become more correlated;
3. regresses the demeaned $\tilde{z}$ on the demeaned $\tilde{\mathbf{X}}$ by weighted least squares to update $\boldsymbol{\beta}$; and
4. recovers the fixed-effect contribution as the complement of the projection and updates η .

Convergence is declared when $\max_\ell |\Delta \beta_\ell| / (1 + |\beta_\ell|)$, evaluated at the updated coefficient vector, falls below `tolerance()` (default 10^{-8}). Sort permutations and group boundaries for each absorbed term are precomputed once, so every demeaning pass is a single $O(n)$ sweep independent of the number of levels. The implementation is pure Mata with no dependencies, and the non-absorbed varlist may include factor variables and interactions.

Standard errors come from the partitioned information at convergence: $\mathbf{V} = (\tilde{\mathbf{X}}' \tilde{\boldsymbol{\Omega}} \tilde{\mathbf{X}})^{-1}$, with $\tilde{\boldsymbol{\Omega}} = \text{diag}(\omega_i)$, for `vce(oim)`, the HC1 sandwich with meat $\sum_i \mathbf{s}_i \mathbf{s}_i' \cdot n / (n - 1)$ and scores $\mathbf{s}_i = (d_i - \mu_i) \tilde{\mathbf{x}}_i$ for `vce(robust)`, and the $G/(G - 1)$ -corrected clustered sandwich with scores summed within clusters for `vce(cluster)`. By the Frisch–Waugh–Lovell logic applied through the IRLS local linearization, these are not linear approximations but the partitioned estimators; given that the fixed effects are solved to tolerance, all three match the corresponding dummy-variable `poisson` output to within 10^{-6} (typically $\sim 10^{-8}$), which the test suite verifies (section 8). The model test reported after `absorb()` is a

Wald test of the non-absorbed coefficients, $\hat{\beta}'\mathbf{V}^{-1}\hat{\beta}$, which matches the joint Wald test on the dummy-variable model; a naive likelihood-ratio test formed from the reported profile log likelihood against a null without the fixed effects would not be valid, since the test's degrees of freedom would ignore the absorbed effects.

Groups of an absorbed term in which no events occur are separated in the maximum-likelihood sense—their fixed effects diverge to $-\infty$ —and `stpois` drops such observations before estimation with a message, mirroring the practice of `ppmlhdfe` (Correia et al., 2020, which treats separation in far greater generality). The header statistics (subjects, failures, time at risk) are computed on the final estimation sample after any such drops, so they always agree with `e(N)` and `e(sample)`.

At convergence, the total fixed-effect contribution $\hat{\eta}_i - \mathbf{x}_i'\hat{\beta}$ is stored in a variable—`_stpois_fe` by default, or a name supplied in the `d()` option, mirroring `ppmlhdfe`'s `d()` and the `fixef()` accessor of `fixest`. `predict` adds this variable to the linear predictor, so the full set of postestimation statistics—linear predictor, expected events, hazard, and survival—is available after `absorb()` and reproduces the corresponding dummy-variable poisson predictions.

5 The stpois command

5.1 Syntax

```
stpois varlist [if] [in] [weight] [, irr nolog level(#) noconstant vce(vcetype) robust
      cluster(varname) absorb(terms) d(newvar) tolerance(#) maxiter(#) fast
      poisson_options]
```

The data must be `stset`; see `[st] stset`. `fweights`, `iweights`, and `pweights` are allowed on all paths; `pweights` imply robust standard errors (see option descriptions below), and weighted coefficients and standard errors agree with the corresponding weighted poisson model to at least 10^{-6} . `fast` and `absorb()` may be combined (section 3.4). The varlist may include factor variables and interactions on every path. Variables named in `absorb()` and, with `fast`, the cluster variable must be numeric.

`stpois` supports the `svy:` prefix for design-based variance estimation. `svy: stpois` produces Taylor-linearized standard errors that agree with `svy: poisson` on both point estimates and standard errors; see section 6.7. `svy bootstrap:` and `svy jackknife:` are also supported on all paths; `absorb()` and `fast` are restricted to these resampling-based variants, since Taylor linearization requires per-observation scores that are not available after absorption or cell acceleration.

5.2 Options

`irr` reports exponentiated coefficients (incidence-rate ratios, equal to hazard ratios in this model). Display only.

`nolog` suppresses iteration logs.

`level`(#) sets the confidence level; the default is `level(95)`.

`noconstant` suppresses the constant term.

`vce`(*vcetype*), `robust`, and `cluster`(*varname*) specify the variance estimator. `vce(oim)` (the default), `vce(robust)`, and `vce(cluster clustvar)` are supported on all paths and reproduce the corresponding full poisson model to at least 10^{-6} ; the standard

path additionally accepts anything `poisson` accepts. On the `fast` and `absorb()` paths, supplying `pweights` without an explicit `vce()`, `robust`, or `cluster()` option automatically sets `vce(robust)`; on the standard path, `poisson` applies the same convention. When comparing results with `ppmlhdfc`—which reports robust standard errors by default—ensure both commands use `vce(robust)` to avoid misinterpreting the difference between `vce(oim)` and `vce(robust)` as an algorithmic discrepancy.

`fast` selects cell-accelerated exact estimation; see section 3. With `fast`, terms built only from factor-variable pieces define the cells and enter at the cell level; at least one such term is required.

`absorb(terms)` absorbs the fixed effects of one or more terms, each a variable name or a `var1#var2` interaction; see section 4. May be combined with `fast`.

`d(newvar)` stores the total fixed-effect contribution of `absorb()` in *newvar*; the default is `_stpois_fe`, replaced on each run. The variable is used by `predict`.

`tolerance(#)` and `maxiter(#)` control the iteration convergence criterion and cap for `absorb()` and `fast`; defaults are `1e-8` and `100`.

Any remaining options on the standard path are passed to `poisson`.

5.3 Stored results

`stpois` stores the following in `e()`:

	<code>e(N)</code>	number of observations (sum of the weights with <code>fweights</code>)
	<code>e(N_sub)</code>	number of subjects
	<code>e(N_fail)</code>	number of failures (a weighted sum with <code>fweights</code>)
	<code>e(risk)</code>	total time at risk (a weighted sum with <code>fweights</code>)
	<code>e(ll)</code>	log likelihood (individual-level, comparable across paths)
	<code>e(ll_0)</code>	log likelihood, constant-only model (paths without <code>absorb()</code>)
<i>Scalars</i>	<code>e(chi2)</code>	model test statistic
	<code>e(df_m)</code>	model degrees of freedom
	<code>e(rank)</code>	rank of <code>e(V)</code>
	<code>e(k)</code>	number of parameters (standard path)
	<code>e(ic)</code>	number of iterations
	<code>e(converged)</code>	1 if converged, 0 otherwise
	<code>e(N_cells)</code>	number of cells (<code>fast</code>)
	<code>e(N_clust)</code>	number of clusters (with <code>vce(cluster)</code>)
	<code>e(cmd)</code>	<code>stpois</code>
	<code>e(cmdline)</code>	command as typed
	<code>e(depvar)</code>	<code>_d</code> (the failure indicator)
	<code>e(t0)</code>	<code>_t0</code> (the entry-time variable)
	<code>e(title)</code>	title in estimation output
	<code>e(chi2type)</code>	LR or Wald; type of model test
	<code>e(vce)</code>	<code>vcetype</code> specified in <code>vce()</code>
<i>Macros</i>	<code>e(predict)</code>	<code>stpois_p</code> (program used to implement <code>predict</code>)
	<code>e(properties)</code>	<code>b V</code>
	<code>e(fast)</code>	<code>fast</code> , if the fast path was used
	<code>e(cont_vars)</code>	individual-level (continuous-involving) terms (<code>fast</code>)
	<code>e(cat_vars)</code>	cell-defining categorical terms (<code>fast</code>)
	<code>e(absorb)</code>	absorbed terms (<code>absorb()</code>)
	<code>e(fes_var)</code>	name of the stored fixed-effect variable (<code>absorb()</code>)
	<code>e(clustvar)</code>	name of the cluster variable (with <code>vce(cluster)</code>)
<i>Matrices</i>	<code>e(b)</code>	coefficient vector
	<code>e(V)</code>	variance-covariance matrix of the estimators

Functions `e(sample)` marks the estimation sample

All quantities are computed on the final estimation sample (after any separation-induced drops on the `absorb()` path), so they agree with `e(N)` and `e(sample)`. Paths without absorbed effects store a likelihood-ratio model test, with `e(chi2type)` set to LR; with `absorb()` the Wald test of section 4 is stored, with `e(chi2type)` set to Wald. Because every path reports the individual-level log likelihood, likelihood-based statistics are comparable across paths.

Because `e(predict)` is set to `stpois_p`, `predict` offers the linear predictor (`xb`, the default, including the log-exposure offset unless `nooffset`), predicted numbers of events (`n`), hazard rates (`hazard` or `hr`), and survival probabilities (`survival`). After `absorb()`, `predict` adds the stored fixed-effect contribution to the linear predictor, so all four statistics include the absorbed effects and match the dummy-variable `poisson` predictions. Postestimation commands that operate on the coefficient vector and its variance matrix—`lincom`, `test`, `nlcom`, and `estimates`—work as usual on every path. `margins` is available for quantities that are functions of the reported coefficients; on the standard and `fast` paths it and `predict` behave exactly as after `poisson`, since no stored fixed-effect variable

is involved. Note, however, that after `absorb()` the absorbed fixed-effect contribution enters `predict` through the stored `_stpois_fe` variable, which `margins` holds fixed at its estimated values. Margins of the response therefore condition on the absorbed effects and their delta-method standard errors do not propagate the sampling uncertainty in those effects, exactly as for any fixed-effect contribution recovered post-absorption in `ppmlhdfc` or `reghdfe`. Predictions based solely on the non-absorbed coefficient vector—setting the absorbed effects to zero—are not directly available through `predict`; they can be obtained with `matrix score newvar = e(b)`, which scores only the columns of `e(b)` without adding the fixed-effect variable. This is useful, for example, to compute marginal effects at the mean of the non-absorbed covariates while holding the fixed effects at zero.

On the `fast` and `absorb()` paths, `stpois` stores the convergence flag `e(converged)` and the iteration count `e(ic)`. If the cap `maxiter()` is reached before the tolerance in `tolerance()` is met, the command still posts the last iterate, sets `e(converged)` to 0, and issues a warning, so a noninteractive pipeline can detect the failure by inspecting `e(converged)`. Cell-level and non-absorbed terms that are collinear—including collinearity with the absorbed effects—are dropped through a generalized inverse and reflected in `e(rank)`. Separation is detected only for the absorbed terms, whose all-zero-event groups are dropped before estimation (section 4); separation induced by the non-absorbed varlist is not given a targeted drop and instead surfaces as nonconvergence with `e(converged)` equal to 0. In practice, if `stpois` does not converge, the recommended diagnostic steps are:

1. Check `e(ic)` against `maxiter()`. If the two are equal, the iteration cap was hit rather than the tolerance, and increasing `maxiter()` or loosening `tolerance()` may help.
2. Inspect for quasi-complete separation in the non-absorbed terms: a binary covariate that perfectly predicts the event indicator within some stratum will cause the corresponding coefficient to diverge. Tabulating `_d` against the suspect covariate, or running `poisson _d varlist, exposure(exp)` on a small subsample, can isolate the issue.
3. Check for collinearity and poor scaling. Continuous covariates on very different scales can slow convergence or trigger numerical issues; standardizing them before estimation is a simple precaution.
4. If using `absorb()`, verify with `tabulate absorbvar _d` that every absorbed group contains at least one event; any all-zero group should have been dropped automatically, but interactions can create groups that pass the check individually while being jointly separated.
5. As a cross-check, fit the model with `ppmlhdfc` (Correia et al., 2020), which provides a more comprehensive separation-detection algorithm that can identify separated observations even in the non-absorbed terms.

6 Examples

6.1 Standard path

The default path reproduces `streg`, `distribution(exponential)` exactly (the log likelihoods differ by a constant because `streg` includes the $d_i \log t_i$ terms). We load the Stanford heart transplant data and `stset` them:

```
. webuse stan3
(Heart transplant data)
. stset t1, failure(died) id(id)
(output omitted)

. stpois age posttran, nolog
Poisson event-history regression
No. of subjects =          103          Number of obs   =          172
No. of failures =           75          LR chi2(2)      =          36.75
Time at risk   =        31938          Prob > chi2     =          0.0000
Log likelihood =   -233.83404
```

	_d	Coefficient	Std. err.	z	P> z	[95% conf. interval]	
	age	.0605817	.015235	3.98	0.000	.0307216	.0904417
	posttran	-1.285206	.2377048	-5.41	0.000	-1.751099	-.8193135
	_cons	-7.8376	.7094787	-11.05	0.000	-9.228153	-6.447047

6.2 Absorbing fixed effects

`absorb()` returns the same coefficients and standard errors as the explicit dummy-variable model, without estimating the absorbed effects:

```
. stpois age posttran, absorb(surgery) nolog
(HDFE: 6 IRLS iterations, converged=1)
Poisson EHA - HDFE (absorb: surgery)
No. of subjects =          103          Number of obs   =          172
No. of failures =           75          Wald chi2(2)    =          35.17
Time at risk   =        31938          Prob > chi2     =          0.0000
Log likelihood =   -227.61949
```

	_d	Coefficient	Std. err.	z	P> z	[95% conf. interval]	
	age	.0610952	.014454	4.23	0.000	.0327659	.0894244
	posttran	-1.141144	.2402034	-4.75	0.000	-1.611934	-.6703538

For comparison, `poisson _d age posttran i.surgery, exposure(exp)` on the same sample gives `age` = .0610952 (SE .014454) and `posttran` = -1.141144 (SE .2402034): identical to all printed digits. The Wald statistic equals the joint test of `age` and `posttran` on the dummy-variable model.

6.3 Fast estimation on a large dataset

The following simulation generates 500,000 subjects in 200 risk cells (4 education $\times$ 5 region $\times$ 10 period), with two continuous covariates (`age`, `income`):

```
. clear
. set obs 500000
. set seed 20260713
. gen edu    = ceil(4 * runiform())
. gen region = ceil(5 * runiform())
. gen period = ceil(10 * runiform())
. gen age    = 30 + 20 * runiform()
```

```

. gen income = rnormal(10, 2)
. gen lambda = exp(-6 + 0.03*age + 0.05*income
> + 0.3*(edu==2) + 0.5*(edu==3) - 0.2*(edu==4)
> + 0.1*(region==2) - 0.2*(region==3) + 0.2*(period >= 6))
. gen ptime = 0.5 + 0.5 * runiform()
. gen failed = (runiform() < 1 - exp(-lambda * ptime))
. gen id2 = _n
. stset ptime, failure(failed) id(id2)
(output omitted)

. stpois age income i.edu i.region i.period, fast nolog
(fast) 2 individual-level + 16 cell-level parameters...
(fast) 500000 obs, 200 cells, 5 Newton iterations, converged=1
Poisson EHA - fast (cell-accelerated exact MLE)
No. of subjects =      500,000      Number of obs   =   500,000
No. of failures =        6,759
Time at risk    =       375045      LR chi2(18)     =    850.69
Log likelihood   = -35297.23308     Prob > chi2    =    0.0000

```

_d	Coefficient	Std. err.	z	P> z	[95% conf. interval]	
age	.0265379	.002121	12.51	0.000	.0223808	.0306951
income	.0518263	.0060872	8.51	0.000	.0398956	.0637571
edu						
2	.3070999	.0353078	8.70	0.000	.237898	.3763019
3	.5094171	.0339083	15.02	0.000	.442958	.5758762
4	-.1907793	.0398372	-4.79	0.000	-.2688587	-.1126998
(output omitted)						
_cons	-5.885082	.1191483	-49.39	0.000	-6.118608	-5.651556

Running the same model on the standard path returns identical output—the same coefficients, standard errors, and log likelihood to all digits—in five Newton iterations versus a full poisson maximization; the timing comparison is in section 8.

6.4 Interactions

All interaction types are accepted with `fast`; terms involving a continuous piece, such as the `edu#c.age` terms produced by `c.age##i.edu` below, are estimated at the individual level, while the categorical main effects and any categorical#category interactions run at the cell level:

```

. stpois c.age##i.edu income i.region i.period, fast nolog
(fast) 5 individual-level + 16 cell-level parameters...
(fast) 500000 obs, 200 cells, 5 Newton iterations, converged=1
Poisson EHA - fast (cell-accelerated exact MLE)
No. of subjects =      500,000      Number of obs   =   500,000
No. of failures =        6,759
Time at risk    =       375045      LR chi2(21)     =    851.30
Log likelihood   = -35296.92816     Prob > chi2    =    0.0000

```

_d	Coefficient	Std. err.	z	P> z	[95% conf. interval]	
age	.0255056	.0046717	5.46	0.000	.0163492	.0346621
edu#c.age						
2	-.0009954	.0061531	-0.16	0.871	-.0130552	.0110645
3	.0021302	.0059106	0.36	0.719	-.0094545	.0137148
4	.0034035	.0069472	0.49	0.624	-.0102128	.0170198
(output omitted)						
_cons	-5.842836	.2075639	-28.15	0.000	-6.249653	-5.436018

Interacted fixed effects are absorbed with `absorb(edu#region)`, which matches `poisson` with explicit `i.edu#i.region` dummies on coefficients and standard errors.

6.5 Combining fast with absorb(), and predictions

Returning to the Stanford heart-transplant data, the two acceleration mechanisms combine, and both absorption engines store the fixed-effect contribution that `predict` uses:

```
. webuse stan3, clear
(Heart transplant data)

. stset t1, failure(died) id(id)
(output omitted)

. stpois age i.posttran, fast absorb(surgery) nolog
(fast) 1 individual-level + 1 cell-level parameters...
(fast) 172 obs, 2 cells, 5 Newton iterations, converged=1
Poisson EHA - fast + HDFE (absorb: surgery)
No. of subjects =          103          Number of obs   =          172
No. of failures =           75          Wald chi2(2)    =          35.17
Time at risk   =        31938          Prob > chi2     =          0.0000
Log likelihood =    -227.61949
```

	_d	Coefficient	Std. err.	z	P> z	[95% conf. interval]	
	age	.0610952	.014454	4.23	0.000	.0327659	.0894244
	1.posttran	-1.141144	.2402034	-4.75	0.000	-1.611934	-.6703538

The same coefficients and standard errors are returned by the IRLS engine (`absorb()` without `fast`) and by `poisson` with explicit `i.surgery` dummies. Predictions after `absorb()` include the absorbed effects:

```
. stpois age posttran, absorb(surgery) nolog
(HDFE: 6 IRLS iterations, converged=1)
Poisson EHA - HDFE (absorb: surgery)
No. of subjects =          103          Number of obs   =          172
No. of failures =           75          Wald chi2(2)    =          35.17
Time at risk   =        31938          Prob > chi2     =          0.0000
Log likelihood =    -227.61949
```

	_d	Coefficient	Std. err.	z	P> z	[95% conf. interval]	
	age	.0610952	.014454	4.23	0.000	.0327659	.0894244
	posttran	-1.141144	.2402034	-4.75	0.000	-1.611934	-.6703538

```
. predict double haz, hazard
. summarize haz
```

Variable	Obs	Mean	Std. dev.	Min	Max
haz	172	.0050208	.0039522	.0003224	.0212179

6.6 Episode-split workflow

The typical episode-split workflow starts with one record per subject, splits spells into shorter episodes, adds indicators for the resulting time bands to model the baseline hazard, and then estimates the piecewise-constant hazard model. The following self-contained example illustrates each step.

We simulate 1,000 subjects observed for up to ten years, with age, sex, and log-income as covariates:

```
. clear
. set obs 1000
. set seed 20260716
. gen id = _n
. gen female = (runiform() < 0.5)
. gen age = 18 + floor(42 * runiform()) // age 18--59
. gen income = rnormal(10, 2)
```

```
. gen years = 1 + floor(10 * runiform()) // 1--10 years at risk
. gen event = (runiform() < 1 - exp(-exp(-5 + 0.04*age - 0.3*female) * years))
. stset years, failure(event) id(id)
```

The dataset at this point has one record per subject, with `_t0 = 0`, `_t = years`, and `_d = event`. `stsplit` splits each spell into annual episodes:

```
. stsplit fu, at(1(1)10)
(output omitted)
. list id female age _t0 _t _d in 1/6, noobs sep(3)
```

id	female	age	_t0	_t	_d
1	0	52	0	1	0
1	0	52	1	2	0
1	0	52	2	3	1
2	1	34	0	1	0
2	1	34	1	2	0
2	1	34	2	3	0

After splitting, each original subject-year becomes a separate episode record, with `_t0` and `_t` marking the episode boundaries. The variable `fu` records the follow-up band (0 = first year, 1 = second year, and so on). The failure indicator `_d` is 1 only in the episode in which the event occurred and 0 in all preceding episodes. The exposure for each record, `_t - _t0`, is one year except possibly for the last episode, which may be shorter.

A piecewise-constant hazard model with ten follow-up bands, a linear age effect, and sex is then estimated on the episode file:

```
. stpois i.fu age female income, irr
. stpois i.fu age female income, fast
. stpois age female income, absorb(fu) vce(robust)
```

The first command uses the standard path and produces identical results to `streg, distribution(exponential)` on the same data. The second uses the fast path, which treats `i.fu` as the cell-defining categorical term and estimates `age`, `female`, and `income` at the individual level from exponentially tilted within-cell moments. The third absorbs the ten follow-up dummies, estimating only the covariate coefficients; because the follow-up baseline hazard is of secondary interest here, absorption avoids placing nine dummy columns in the output while still controlling for duration dependence. The fast and absorb approaches can be combined: `stpois age female income i.edu, fast absorb(fu region) vce(cluster region)` absorbs two high-dimensional nuisance effects simultaneously and clusters the standard errors, while the individual-level covariates and the education cells are estimated from the full episode file.

6.7 Survey data

`stpois` declares `properties(st svyr svyb svyj)`, so `svyset` data can be analyzed with the full `svy:` prefix family. For Taylor-linearized standard errors, `svy:` `stpois` calls `stpois` to obtain point estimates, then requests per-observation scores $d_i - \hat{\mu}_i$ from `stpois_p` and constructs the sandwich variance matrix using the survey design; this is the same linearization as `svy: poisson` and produces identical results:

```
. sysuse cancer, clear
. stset studytime, failure(died)
. svyset _n [pweight=age]
. svy: stpois age drug, irr
(output omitted)
. svy: poisson _d age drug, exposure(studytime) irr // identical output
```


`stpois` inherits sampling weights through the `stset` weight mechanism: when `svy:` is active, it injects the survey `pweight` into the dataset’s `stset` characteristics before calling the subcommand, and `stpois` reads those characteristics to apply the weights to the internal `poisson` call. This mirrors the behavior of `streg` under `svy:` and means that `svy: stpois` and `svy: poisson` agree on both point estimates and linearized standard errors to at least 10^{-6} . `svy, subpop()`: subpopulation estimation and stratified or multi-stage designs are handled automatically by Stata’s `svy` machinery.

Taylor linearization requires per-observation scores and is therefore restricted to the standard path. With `absorb()` or `fast`, use `svy bootstrap: stpois` or `svy jackknife: stpois`: these resampling-based variance estimators re-run the full estimation on each replicate and are therefore valid for any estimation path.

7 Relation to existing commands

Table 1 places `stpois` among the Stata commands that fit closely related models. For one-way fixed effects, the official `xtpoisson, fe` conditional estimator is dependency-free and consistent, but it conditions the fixed effects out of the likelihood rather than absorbing them, so it does not extend to multi-way fixed effects, does not recover or predict the fixed-effect contribution, and its clustered-variance conventions differ from those of `poisson` with explicit dummies; `stpois`, `absorb()` targets exactly the cases—multi-way absorption, recovered effects, and `poisson`-matching robust and clustered standard errors—in which the conditional estimator does not apply. Against `ppmlhdfe`, the closest command for the absorption path, `stpois` adds `stset` integration and the cell-accelerated fast path while requiring only Stata 14 or later and carrying no external dependencies, at the cost of a narrower separation-detection scheme (section 4).

Table 1: Feature coverage relative to related commands (● supported natively, ○ available only indirectly, e.g. through explicit dummies or with restrictions, blank not available)

Feature	<code>poisson</code>	<code>streg</code>	<code>stcox</code>	<code>xtpoisson, fe</code>	<code>ppmlhdfe</code>	<code>stpois</code>
<code>stset</code> -native interface		●	●			●
Survival headers/predict		●	●			●
One-way fixed effects	○			●	●	●
Multi-way fixed effects	○				●	●
Cell-accelerated continuous fit						●
Robust/cluster VCE	●	●	●	●	●	●
fw/iw/pw weights	●	●	●	○	●	●
Separation detection					●	○
Recover/predict FE contribution					●	●
<code>svy:</code> prefix (linearized)	●	●	●	○	○	●
No external dependencies	●	●	●	●		●

8 Validation and performance

8.1 Numerical agreement with `streg` and `poisson`

The standard and fast estimators are algebraically exact in that they solve the same likelihood as `poisson` with no mathematical approximation; the `absorb()` estimator converges to the same maximum-likelihood optimum to within its specified tolerance. “Exact” throughout this article therefore means agreement with the `streg/poisson` benchmarks to at least 10^{-6} on coefficients *and* all three variance estimators, with observed agreement typically on the order of 10^{-8} ; it does not assert bit-for-bit equality of two independent maximizers. The package ships with a test suite (`test_stpois.do` and `test_fast_exact.do` in the `tests` directory) that runs on the `stan3` heart-transplant data and on simulated data—not on register data, which cannot leave the secure environments that motivate the command. Each exactness claim is held to a numerical benchmark:

1. The **standard path** matches `streg, distribution(exponential)` on every coefficient and standard error to 10^{-6} . On the `stan3` data, both report $\hat{\beta}_{\text{age}} = 0.0605817$ (SE 0.0152350) and $\hat{\beta}_{\text{posttran}} = -1.2852060$ (SE 0.2377048); the log likelihoods differ only by the fixed $\sum_i d_i \log t_i$ term that `streg` includes and the Poisson formulation drops.
2. **absorb()** matches the explicit dummy-variable `poisson` model on coefficients *and* standard errors to 10^{-6} (observed agreement is typically on the order of 10^{-8}), for the observed-information, robust, and cluster-robust variance estimators. The check covers single fixed effects, interacted fixed effects (`edu#region` against explicit `i.edu#i.region` dummies), multiple simultaneous fixed-effect terms (`absorb(edu region period)`), and factor-variable interactions in the varlist. The Wald model statistic matches the joint test on the dummy-variable model, and a separation test verifies that the header statistics are computed on the post-drop estimation sample.
3. **fast** matches the full `poisson` model on coefficients, standard errors (observed-information, robust, and clustered), and log likelihood to 10^{-6} . The check covers continuous#continuous, continuous#categorical, and categorical#categorical interactions, all-categorical models, and a two-cell setting in which the continuous covariate is constructed to have exactly zero between-cell variation, so that its coefficient is identified purely from within-cell variation.
4. **Weights** (frequency, importance, and sampling) on the fast and absorbed paths match the corresponding weighted `poisson` model on coefficients and standard errors to 10^{-6} , with `e(N)` following Stata’s conventions and `pweights` implying robust standard errors.
5. **fast + absorb()** matches `poisson` with explicit dummies for both the cell-level terms and the absorbed effects—coefficients and observed-information, robust, and clustered standard errors to 10^{-6} —including multi-term absorbs and weighted estimation.
6. **predict** after `absorb()` reproduces the dummy-variable `poisson` predictions of expected events, and the hazard equals the exponentiated no-offset linear predictor, on both the IRLS and the fast+absorb engines.

7. **svy**: on the standard path, **svy: stpois** agrees with **svy: poisson** on point estimates and Taylor-linearized standard errors to 10^{-6} (observed agreement at the level of floating-point rounding, $\sim 10^{-15}$), including with stratified designs and subpopulation estimation.

All tests pass on Stata 19 (BE); the command requires version 14. The suite was run on a clean Stata installation with no other user-contributed packages present, confirming that **stpois** draws on no external dependencies.

8.2 Agreement with **ppmlhdfe** and with **collapse+poisson**

Because **absorb()** targets the same problem as **ppmlhdfe** and **fast** generalizes the classical collapse-and-fit workflow, the accompanying script **benchmark_ppmlhdfe.do** checks **stpois** directly against both on a 10^6 -record design with a 1,000-level fixed effect. The point estimates are identical: on the **absorb()** path **stpois** and **ppmlhdfe** agree on $\hat{\beta}_{\text{age}}$ to 2×10^{-13} , and on the combined **fast + absorb()** path to 1×10^{-14} . Compared like for like, the standard errors agree to about 3×10^{-9} : **ppmlhdfe** reports robust standard errors by default, as is standard for Poisson pseudo-maximum-likelihood, and these match **stpois** under **vce(robust)** and **vce(cluster)** to that tolerance. **stpois** additionally provides **vce(oim)** as its default, reproducing **poisson**'s classical observed-information standard errors—an option **ppmlhdfe** does not offer (as of version 2.4.1, the version benchmarked here). Against the folk baseline of collapsing to cells and fitting weighted **poisson** on the sums, the **fast** path reproduces the all-categorical estimates and their standard errors to machine precision ($< 10^{-15}$)—as it must, since for this case (no absorbed effects, default VCE) **fast** *is* that collapse-and-fit workflow, performed internally so the user neither writes the **collapse** nor leaves the **stset** data (section 3). What **fast** adds beyond the manual workflow is support for continuous covariates, for which no sufficient cell reduction exists (section 2.1): the command accelerates through the cell structure while retaining the continuous terms in the individual-level tilted moments. On timing (medians of ten runs, same machine as the tables), **stpois**, **absorb()** fits the 1,000-level design of Table 5 in 2.9 s against **ppmlhdfe**'s 4.6 s, and the combined **fast + absorb()** path in 5.0 s against 11.6 s; **stpois** is thus competitive with, and here faster than, the established command while adding the **stset** interface and survival postestimation. On the all-categorical design, the internal collapse route fits in 0.5 s versus 0.1 s for hand-collapsing and calling **poisson**—roughly $5\times$ slower—the gap arising from **stpois**'s fixed overhead of sample marking, term parsing, and result posting; what the command adds is the **stset** workflow and continuous-covariate support without requiring the user to write a **collapse**.

8.3 Timing benchmarks

All timings are medians over ten runs on Stata 19 (BE, Apple silicon, warm cache), produced by the benchmark scripts in the package repository, which write their results to comma-separated files versioned alongside the paper; the tables below reproduce those files. Reported speedup ratios are computed from the unrounded median timings, so they need not equal the ratio of the rounded cells shown. In every comparison the estimates agree across methods to the tolerances above—the scripts assert this inside the timing loops—so the tables compare identical output.

Table 2 times a mixed model with two continuous and 16 categorical parameters (education, region, period), comparing `streg`, `distribution(exponential)`, the `stpois` standard path (a `poisson` call), and `stpois, fast`.

Table 2: Mixed continuous + categorical model, seconds per fit. DGP: 200 cells (4 education $\times$ 5 region $\times$ 10 period), two continuous covariates (age, income), $\approx 1\%$ event rate. Medians of 10 runs, Stata 19 BE, Apple silicon, warm cache.

n	<code>streg</code>	<code>stpois</code> (standard)	<code>fast</code>	speedup vs. <code>streg</code>
100,000	0.37	0.34	0.11	$3.3\times$
1,000,000	2.73	2.52	1.10	$2.5\times$
5,000,000	13.26	12.41	6.72	$2.0\times$

Table 3 times the canonical episode-split workflow: 100,000 subjects observed for up to 30 years are split into one-year episodes with `stsplit` (1,888,654 episode records), and a piecewise-constant hazard model with 30 follow-up bands, education, region, and two continuous covariates is estimated on the episode file.

Table 3: Episode-split workflow (1,888,654 episode records). DGP: 100,000 subjects split into annual episodes over up to 30 years; piecewise-constant hazard with 30 follow-up bands, education, region, and two continuous covariates. Medians of 10 runs, Stata 19 BE, Apple silicon, warm cache.

Method	Time	speedup vs. <code>streg</code>
<code>streg, distribution(exponential)</code>	5.84 s	—
<code>stpois</code> (standard)	5.12 s	$1.1\times$
<code>stpois, fast</code>	2.19 s	$2.7\times$

Table 4 times an all-categorical rate model (education $\times$ region $\times$ period, 200 cells), the case where the sufficiency result of section 2.1 lets `fast` fit on the collapsed cell sums alone through the internal `collapse + poisson` route.

Table 4: All-categorical rate model, seconds per fit. DGP: 200 cells (4 education $\times$ 5 region $\times$ 10 period), no continuous covariates, $\approx 1\%$ event rate. Medians of 10 runs, Stata 19 BE, Apple silicon, warm cache. For this case `fast` internally routes through `collapse+poisson` on the 200 cell sums; hand-collapsing and calling `poisson` directly is $\approx 5\times$ faster (section 8.2).

n	<code>streg</code>	<code>stpois</code> (standard)	<code>fast</code>	speedup vs. <code>streg</code>
1,000,000	3.60	3.43	0.60	$6.0\times$
5,000,000	13.05	12.06	2.86	$4.6\times$

In tables 2–4 the complexity analysis implies that the remaining **fast** cost is dominated by the single aggregating pass over the microdata: the per-iteration work is $O(np^2)$ in the few individual-level columns only (Table 2), and for the all-categorical model (Table 4) it reduces to one **collapse** plus a **poisson** fit on a few hundred cells. The gains are therefore largest relative to estimators that must pass over the microdata repeatedly, which is what the fixed-effects benchmark isolates. Table 5 estimates a model with two continuous covariates and a single fixed effect with 1,000 levels on one million observations, comparing **poisson** with explicit dummies, **stpois**, **absorb()**, and **ppmlhdfe** on identical output.

Table 5: High-dimensional fixed effect: 1,000 levels, $n = 10^6$, two continuous covariates. **ppmlhdfe** reports robust standard errors by default and offers no observed-information option (section 8.2). Rows 1–3: medians of 10 runs from **benchmark_hdfe.do**; row 4: medians of 10 runs from **benchmark_ppmlhdfe.do** on the same DGP (robust SEs; **stpois** SE omitted for brevity). Stata 19 BE, Apple silicon, warm cache.

Method	Time	$\hat{\beta}_{\text{age}}$	SE
poisson with 1,000 dummies	34.31 s	0.0266067	0.0017000
stpois , absorb(g)	2.90 s	0.0266067	0.0017000
ppmlhdfe , absorb(g)	4.59 s	0.0266067	—
stpois , fast absorb(g)	5.03 s	(see section 8.2)	
ppmlhdfe , absorb(g) [fast comp.]	11.64 s		

Nearly a $12\times$ saving at identical output; the dummy-variable model’s cost grows roughly quadratically in the number of levels while the absorption cost grows only linearly, so the complexity analysis implies that the gap should widen further in higher-dimensional settings. For register applications with 10^7 or more episodes, detailed period $\times$ education $\times$ region structures, and high-dimensional nuisance effects—a regime beyond the scale of the benchmarks reported here—cell acceleration and fixed-effect absorption together may reduce runtimes from minutes to seconds, which would change what is practical to do interactively: specification searches, bootstrap loops, or estimation inside secure environments with tight session limits.

9 Conclusion

stpois packages four things that belong together but have required separate workflows: the Poisson formulation of event-history models on **stset** data, absorption of high-dimensional fixed effects without external dependencies, cell-accelerated estimation that retains continuous covariates, and design-based variance estimation via the **svy:** prefix. The standard and fast paths are algebraically exact—they solve the same likelihood as **poisson** with no mathematical approximation—and the **absorb()** path converges to the same maximum-likelihood optimum to within its specified tolerance; the test suite holds coefficients *and* variance estimators, not merely point estimates, to the **streg** and **poisson** benchmarks at 10^{-6} (typically $\sim 10^{-8}$), and factor-variable interactions are supported throughout, including interacted fixed effects. Weights are accepted on every path, the fast and absorbed paths can be combined, predictions after **absorb()** include the stored

fixed-effect contributions, and `svy: stpois` produces Taylor-linearized standard errors that agree with `svy: poisson` to at least 10^{-6} . The command, its help file, the test suite, and the benchmark scripts are available from the author’s repository and are installable with `net install`.

10 Acknowledgments

TBW

References

- Laurent Bergé. Efficient estimation of maximum likelihood models with multiple fixed-effects: The R package FENmlm. *CREA Discussion Papers*, (13), 2018.
- Bendix Carstensen. Age-period-cohort models for the Lexis diagram. *Statistics in Medicine*, 26(15):3018–3045, 2007.
- David Clayton and Michael Hills. *Statistical Models in Epidemiology*. Oxford University Press, Oxford, 1993.
- Sergio Correia. Ftools: Stata module to provide alternatives to common stata commands optimized for large datasets. Statistical Software Components S458213, Boston College Department of Economics, 2016.
- Sergio Correia, Paulo Guimarães, and Tom Zylkin. Fast Poisson estimation with high-dimensional fixed effects. *Stata Journal*, 20(1):95–115, 2020.
- Paul W. Dickman, Andy Sloggett, Michael Hills, and Timo Hakulinen. Regression models for relative survival. *Statistics in Medicine*, 23(1):51–64, 2004. doi: 10.1002/sim.1597.
- Simen Gaure. OLS with multiple high dimensional category variables. *Computational Statistics & Data Analysis*, 66:8–18, 2013.
- Christian Gourieroux, Alain Monfort, and Alain Trognon. Pseudo maximum likelihood methods: Applications to Poisson models. *Econometrica*, 52(3):701–720, 1984.
- Paulo Guimarães and Pedro Portugal. A simple feasible procedure to fit models with high-dimensional fixed effects. *Stata Journal*, 10(4):628–649, 2010.
- Israel Halperin. The product of projection operators. *Acta Scientiarum Mathematicarum*, 23:96–99, 1962.
- Theodore R. Holford. The analysis of rates and of survivorship using log-linear models. *Biometrics*, 36(2):299–305, 1980.
- Jørgen Ledet Jensen. *Saddlepoint Approximations*. Oxford University Press, Oxford, 1995.
- Nan Laird and Donald Olivier. Covariance analysis of censored survival data using log-linear analysis techniques. *Journal of the American Statistical Association*, 76(374):231–240, 1981.

Peter McCullagh and John A. Nelder. *Generalized Linear Models*. Chapman & Hall, London, 2 edition, 1989.

Sophia Rabe-Hesketh and Anders Skrondal. *Multilevel and Longitudinal Modeling Using Stata*. Stata Press, College Station, TX, 4 edition, 2022.

J. M. C. Santos Silva and Silvana Tenreiro. The log of gravity. *Review of Economics and Statistics*, 88(4):641–658, 2006.

John Whitehead. Fitting Cox’s regression model to survival data using GLIM. *Journal of the Royal Statistical Society. Series C (Applied Statistics)*, 29(3):268–275, 1980.

About the author. Andreas Ljungström is a PhD student at the Swedish Institute for Social Research (SOFI), Stockholm University. His work concerns social stratification broadly, but with a special focus on integration in education and early labor market outcomes using full-population Swedish register data.