Dynamical Localization of Quantum Walks on the hexagonal lattice in the regime of strong disorder

Mathematical Challenges in Quantum Mechanics

Andreas Schäfer

Université Grenoble Alpes

11/02/2025









Quantum Walks on the hexagonal lattice

Intuition: Quantum particle moving in discrete time on discrete grid Γ.
State of quantum particle with internal degree of freedom s is given by $\psi \in l^2(\Gamma) \otimes \mathbb{C}^s$.

Definition

One time-step of a QW is given by $U\psi$, where U is a unitary, local operator on $l^2(\Gamma) \otimes \mathbb{C}^s$.

Quantum Walks on the hexagonal lattice


Intuition: Quantum particle moving in discrete time on discrete grid Γ.
State of quantum particle with internal degree of freedom s is given by $\psi \in l^2(\Gamma) \otimes \mathbb{C}^s$.

Definition

One time-step of a QW is given by $U\psi$, where U is a unitary, local operator on $l^2(\Gamma) \otimes \mathbb{C}^s$.

Equip every $x \in \Gamma$ with coin matrix $C_x \in U(3)$

$$\mathcal{C}(|x\rangle \otimes e_i) = |x\rangle \otimes C_x\, e_i$$



Quantum Walks on the hexagonal lattice

Intuition: Quantum particle moving in discrete time on discrete grid Γ.
State of quantum particle with internal degree of freedom s is given by $\psi \in l^2(\Gamma) \otimes \mathbb{C}^s$.


Definition

One time-step of a QW is given by $U\psi$, where U is a unitary, local operator on $l^2(\Gamma) \otimes \mathbb{C}^s$.

Equip every $x \in \Gamma$ with coin matrix $C_x \in U(3)$

$$\mathcal{C}(|x\rangle \otimes e_i) = |x\rangle \otimes C_x\, e_i$$

S_j act on $l^2(\Gamma)$, set $\mathcal{S} = \sum_{i=1}^{3} S_i \otimes |e_i\rangle\langle e_i|$



Quantum Walks on the hexagonal lattice

Intuition: Quantum particle moving in discrete time on discrete grid Γ.
State of quantum particle with internal degree of freedom s is given by $\psi \in l^2(\Gamma) \otimes \mathbb{C}^s$.

Definition


One time-step of a QW is given by $U\psi$, where U is a unitary, local operator on $l^2(\Gamma) \otimes \mathbb{C}^s$.

Equip every $x \in \Gamma$ with coin matrix $C_x \in U(3)$

$$\mathcal{C}(|x\rangle \otimes e_i) = |x\rangle \otimes C_x\, e_i$$

S_j act on $l^2(\Gamma)$, set $\mathcal{S} = \sum_{i=1}^{3} S_i \otimes |e_i\rangle\langle e_i|$

$\Longrightarrow$ Quantum Walk: $U = \mathcal{SC}$



Transport Properties

Translation-invariant case:

Fourier Transform: $\widehat{U} = \mathcal{F} U \mathcal{F}^{-1}$ is multiplication operator on $L^2(\mathbb{T}^2, \mathbb{C}^6)$

$\rightsquigarrow \sigma_{\text{cont}}(U) = \varnothing \iff k \in \mathbb{T}^2 \mapsto \operatorname{tr}\Big(\widehat{U}(k)\Big) = \text{const.}$

$\rightsquigarrow$ a.a. coin matrices $C \in U(3)$ lead to continuous spectrum, i.e. transport

Transport Properties

Translation-invariant case:

Fourier Transform: $\widehat{U} = \mathcal{F} U \mathcal{F}^{-1}$ is multiplication operator on $L^2(\mathbb{T}^2, \mathbb{C}^6)$

$\rightsquigarrow \sigma_{\text{cont}}(U) = \varnothing \iff k \in \mathbb{T}^2 \mapsto \operatorname{tr}\left(\widehat{U}(k)\right) = \text{const.}$

$\rightsquigarrow$ a.a. coin matrices $C \in U(3)$ lead to continuous spectrum, i.e. transport

General case:

Topological index proves transport depending on $\lim_{x \to \infty} C_x$:

Transport Properties

Translation-invariant case:

Fourier Transform: $\widehat{U} = \mathcal{F} U \mathcal{F}^{-1}$ is multiplication operator on $L^2(\mathbb{T}^2, \mathbb{C}^6)$

$\rightsquigarrow \sigma_{\text{cont}}(U) = \varnothing \iff k \in \mathbb{T}^2 \mapsto \operatorname{tr}\left(\widehat{U}(k)\right) = \text{const.}$


$\rightsquigarrow$ a.a. coin matrices $C \in U(3)$ lead to continuous spectrum, i.e. transport

General case:

Topological index proves transport depending on $\lim_{x\to\infty} C_x$:

Projection P onto "half-space" of edges, consider

$$\operatorname{ind}(U) = \operatorname{ind}(U^* P U, P) \in \mathbb{Z}$$



Transport Properties

Translation-invariant case:

Fourier Transform: $\widehat{U} = \mathcal{F} U \mathcal{F}^{-1}$ is multiplication operator on $L^2(\mathbb{T}^2, \mathbb{C}^6)$

$\rightsquigarrow \sigma_{\text{cont}}(U) = \varnothing \iff k \in \mathbb{T}^2 \mapsto \text{tr}\left(\widehat{U}(k)\right) = \text{const.}$

$\rightsquigarrow$ a.a. coin matrices $C \in U(3)$ lead to continuous spectrum, i.e. transport


General case:

Topological index proves transport depending on $\lim_{x\to\infty} C_x$:

Projection P onto "half-space" of edges, consider

$$\text{ind}(U) = \text{ind}(U^* P U, P) \in \mathbb{Z}$$

$\text{ind}(U) \neq 0 \Rightarrow$ criteria on $\sigma(U), \sigma_{\text{ac}}(U) = \mathbb{S}^1$



Transport Properties

Translation-invariant case:

Fourier Transform: $\widehat{U} = \mathcal{F} U \mathcal{F}^{-1}$ is multiplication operator on $L^2(\mathbb{T}^2, \mathbb{C}^6)$

$\rightsquigarrow \sigma_{\text{cont}}(U) = \varnothing \iff k \in \mathbb{T}^2 \mapsto \text{tr}\left(\widehat{U}(k)\right) = \text{const.}$

$\rightsquigarrow$ a.a. coin matrices $C \in U(3)$ lead to continuous spectrum, i.e. transport


General case:

Topological index proves transport depending on $\lim_{x\to\infty} C_x$:

Projection P onto "half-space" of edges, consider

$$\text{ind}(U) = \text{ind}(U^* P U, P) \in \mathbb{Z}$$

$\text{ind}(U) \neq 0 \Rightarrow$ criteria on $\sigma(U), \sigma_{\text{ac}}(U) = \mathbb{S}^1$, e.g. $|(C_x)_{2,1}|, |(C_x)_{1,2}|, |(C_x)_{3,3}| \to 1$ ($|x| \to \infty$ on boundary), then $\sigma(U) = \mathbb{S}^1$



Dynamical Localization

Equip every site $x \in \Gamma$ with i.i.d. RV ω_x uniformly distributed on $[0, 2\pi]$
$\Longrightarrow$ define $D_\omega(|x\rangle \otimes e_i) = e^{i\omega_x} |x\rangle \otimes e_i$ and $U_\omega = D_\omega U$

Definition

Dynamical localization: $\exists\, g > 0$ such that $\forall\, x, y \in \Gamma$:

$$\mathbb{E}\Big(\sup_{n\in\mathbb{Z}} |\langle x|\, U_\omega^n\, |y\rangle| \Big) \leqslant c\, e^{-g|x-y|}$$

i.e. probability to move from site y to x decreases exponentially in distance $|x-y|$, independently of how many steps QW may take

Dynamical Localization

Equip every site $x \in \Gamma$ with i.i.d. RV ω_x uniformly distributed on $[0, 2\pi]$
$\implies$ define $D_\omega(|x\rangle \otimes e_i) = e^{i\omega_x} |x\rangle \otimes e_i$ and $U_\omega = D_\omega U$

Definition

Dynamical localization: $\exists\, g > 0$ such that $\forall\, x, y \in \Gamma$:

$$\mathbb{E}\Big(\sup_{n\in\mathbb{Z}} |\langle x| U_\omega^n |y\rangle| \Big) \leqslant c\, e^{-g|x-y|}$$

i.e. probability to move from site y to x decreases exponentially in distance $|x-y|$, independently of how many steps QW may take


Dyn. Loc. implies spectral localization, i.e. U_ω has pure point spectrum a.s.
Fractional moments criterion: for some $s \in (0,1)$ and $1 - \epsilon < |z| < 1$

$$\mathbb{E}\Big(|\langle x| (U_\omega - z)^{-1} |y\rangle|^s \Big) \leqslant c\, e^{-g|x-y|}$$

The result

E. Hamza, A. Joye, G. Stolz (2009): Frac. mom. crit $\Longrightarrow$ dyn. loc.


$C_0 = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}$ induces the dynamics



The result

E. Hamza, A. Joye, G. Stolz (2009): Frac. mom. crit $\Longrightarrow$ dyn. loc.

$C_0 = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}$ induces the dynamics



Theorem 1

For any $s \in (0, \frac{1}{3})$ there exists $\varphi, c, g > 0$ such that for any $\{C_x\}_{x\in\Gamma} \subset U(3)$ with $\sup_x \|C_x - C_0\| \leqslant \varphi$ and all $x, y \in \Gamma$, $z \in \mathbb{C}$ with $|z| < 1$:

$$\mathbb{E}\left(|\langle x \otimes e_i| \, (U_\omega - z)^{-1} \, |y \otimes e_j\rangle|^s\right) \leqslant c \, e^{-g|x-y|}.$$

Proof: Finite volume criterion

Finite volume criterion

Restrict QW U_ω to box Λ_L of size $L \in 2$

Iterative step:

$\exists\, q < 1$ s.t. for $s < \frac{1}{3}$, L big enough $\exists\, \varphi > 0$:

$$\mathbb{E}\left(\left|\langle x^{(i)}|\,(U_\omega - z)^{-1}\,|y^{(j)}\rangle\right|^s\right) \leqslant q \max_{v^{(k)} \in \partial(\Lambda_L + x)} \mathbb{E}\left(\left|\langle v^{(k)}|\,(U_\omega - z)^{-1}\,|y^{(j)}\rangle\right|^s\right)$$

for all $\|C - C_0\| \leqslant \varphi$ and y outside of box $\Lambda_L + x$

Finite volume criterion


Restrict QW U_ω to box Λ_L of size $L \in^2$

Iterative step:

$\exists\, q < 1$ s.t. for $s < \frac{1}{3}$, L big enough $\exists\, \varphi > 0$:

$$\mathbb{E}\left(\left|\langle x^{(i)}|\,(U_\omega - z)^{-1}\,|y^{(j)}\rangle\right|^s\right) \leqslant q \max_{v^{(k)}\in\partial(\Lambda_L + x)} \mathbb{E}\left(\left|\langle v^{(k)}|\,(U_\omega - z)^{-1}\,|y^{(j)}\rangle\right|^s\right)$$

for all $\|C - C_0\| \leqslant \varphi$ and y outside of box $\Lambda_L + x$



Finite volume criterion


Restrict QW U_ω to box Λ_L of size $L \in^2$

Iterative step:

$\exists\, q < 1$ s.t. for $s < \frac{1}{3}$, L big enough $\exists\, \varphi > 0$:

$$\mathbb{E}\left(\left|\langle x^{(i)}|\,(U_\omega - z)^{-1}\,|y^{(j)}\rangle\right|^s\right) \leqslant q \max_{v^{(k)} \in \partial(\Lambda_L + x)} \mathbb{E}\left(\left|\langle v^{(k)}|\,(U_\omega - z)^{-1}\,|y^{(j)}\rangle\right|^s\right)$$

for all $\|C - C_0\| \leqslant \varphi$ and y outside of box $\Lambda_L + x$



Finite volume criterion


Restrict QW U_ω to box Λ_L of size $L \in^2$

Iterative step:

$\exists\, q < 1$ s.t. for $s < \frac{1}{3}$, L big enough $\exists\, \varphi > 0$:

$$\mathbb{E}\left(\left|\langle x^{(i)}|\,(U_\omega - z)^{-1}\,|y^{(j)}\rangle\right|^s\right) \leqslant q \max_{v^{(k)} \in \partial(\Lambda_L + x)} \mathbb{E}\left(\left|\langle v^{(k)}|\,(U_\omega - z)^{-1}\,|y^{(j)}\rangle\right|^s\right)$$

for all $\|C - C_0\| \leqslant \varphi$ and y outside of box $\Lambda_L + x$



Thank you for your attention!