

Assignment 1: Poisson–Gamma Posterior

CSDS 611 — Advanced Topics and Methods of Data Science I

Andrei Cozma

Model and parameterization

Consider the Bayesian model

$$w_n \mid \lambda \sim \text{Poisson}(\lambda), \quad n = 1, \dots, N, \quad \lambda \sim \text{Gamma}(\phi, \psi).$$

Assume that the observations are conditionally independent given λ , with $w_n \in \{0, 1, 2, \dots\}$ and $\phi, \psi > 0$. We use the Gamma shape–scale parameterization. Let $\mathbf{w} = (w_1, \dots, w_N)$ and $S = \sum_{n=1}^N w_n$. We seek the posterior density $p(\lambda \mid \mathbf{w})$.

1. Likelihood and prior

Conditional independence gives the joint likelihood

$$\begin{aligned} p(\mathbf{w} \mid \lambda) &= \prod_{n=1}^N \frac{\lambda^{w_n} e^{-\lambda}}{w_n!} \\ &= \frac{\lambda^S e^{-N\lambda}}{\prod_{n=1}^N w_n!}. \end{aligned}$$

The Gamma prior has density

$$p(\lambda) = \frac{1}{\Gamma(\phi)\psi^\phi} \lambda^{\phi-1} e^{-\lambda/\psi}, \quad \lambda > 0.$$

2. Apply Bayes' rule

By Bayes' rule,

$$\begin{aligned} p(\lambda \mid \mathbf{w}) &\propto p(\mathbf{w} \mid \lambda) p(\lambda) \\ &\propto \lambda^S e^{-N\lambda} \lambda^{\phi-1} e^{-\lambda/\psi} \\ &= \lambda^{\phi+S-1} \exp\left[-\left(N + \frac{1}{\psi}\right)\lambda\right]. \end{aligned}$$

The final proportionality omits only factors that do not depend on λ .

3. Normalize and identify the posterior

Introduce a normalizing constant C and write

$$p(\lambda \mid \mathbf{w}) = C \lambda^{\phi+S-1} \exp \left[- \left(N + \frac{1}{\psi} \right) \lambda \right], \quad \lambda > 0.$$

The posterior must integrate to one. Using the substitution $t = (N + 1/\psi)\lambda$, we obtain

$$\begin{aligned} 1 &= \int_0^\infty p(\lambda \mid \mathbf{w}) d\lambda \\ &= C \int_0^\infty \lambda^{\phi+S-1} e^{-(N+1/\psi)\lambda} d\lambda \\ &= C \left(N + \frac{1}{\psi} \right)^{-(\phi+S)} \int_0^\infty t^{\phi+S-1} e^{-t} dt \\ &= C \frac{\Gamma(\phi+S)}{\left(N + \frac{1}{\psi} \right)^{\phi+S}}. \end{aligned}$$

The integral in t equals $\Gamma(\phi+S)$, by normalization of the Gamma density with shape $\phi+S$ and scale 1. Therefore,

$$C = \frac{\left(N + \frac{1}{\psi} \right)^{\phi+S}}{\Gamma(\phi+S)}.$$

Substituting C gives the normalized posterior density:

$$p(\lambda \mid \mathbf{w}) = \frac{\left(N + \frac{1}{\psi} \right)^{\phi+S}}{\Gamma(\phi+S)} \lambda^{\phi+S-1} \exp \left[- \left(N + \frac{1}{\psi} \right) \lambda \right], \quad \lambda > 0.$$

Recognizing the Gamma density, with scale $1/(N + 1/\psi) = \psi/(1 + N\psi)$, we conclude that

$$\lambda \mid w_1, \dots, w_N \sim \text{Gamma} \left(\phi + \sum_{n=1}^N w_n, \frac{\psi}{1 + N\psi} \right) \quad (\text{shape--scale}).$$

The posterior remains in the Gamma family, so the Gamma prior is conjugate to the Poisson likelihood.