

## COURSE 4

### 2.3. Hermite interpolation

**Example 1** *In the following table there are some data regarding a moving car. We may estimate the position (and the speed) of the car when the time is  $t = 10$  using Hermite interpolation.*

<i>Time</i>	0	3	5	8	13
<i>Distance</i>	0	225	383	623	993
<i>Speed</i>	75	77	80	74	72

Let  $x_k \in [a, b]$ ,  $k = 0, 1, \dots, m$  be such that  $x_i \neq x_j$ , for  $i \neq j$  and let  $r_k \in \mathbb{N}$ ,  $k = 0, 1, \dots, m$ . Consider  $f : [a, b] \rightarrow \mathbb{R}$  such that there exist  $f^{(j)}(x_k)$ ,  $k = 0, 1, \dots, m$ ;  $j = 0, 1, \dots, r_k$  and  $n = m + r_0 + \dots + r_m$ .

**The Hermite interpolation problem (HIP)** consists in determining the polynomial  $P$  of the smallest degree for which

$$P^{(j)}(x_k) = f^{(j)}(x_k), \quad k = 0, \dots, m; \quad j = 0, \dots, r_k.$$

**Definition 2** A solution of (HIP) is called **Hermite interpolation polynomial**, denoted by  $H_n f$ .

**Hermite interpolation polynomial**,  $H_n f$ , satisfies the interpolation conditions:

$$(H_n f)^{(j)}(x_k) = f^{(j)}(x_k), \quad k = 0, \dots, m; \quad j = 0, \dots, r_k.$$

Hermite interpolation polynomial is given by

$$(H_n f)(x) = \sum_{k=0}^m \sum_{j=0}^{r_k} h_{kj}(x) f^{(j)}(x_k) \in \mathbb{P}_n, \quad (1)$$

where  $h_{kj}(x)$  denote **the Hermite fundamental interpolation polynomials**. They fulfill the relations:

$$h_{kj}^{(p)}(x_\nu) = 0, \quad \nu \neq k, \quad p = 0, 1, \dots, r_\nu$$

$$h_{kj}^{(p)}(x_k) = \delta_{jp}, \quad p = 0, 1, \dots, r_k, \quad \text{for } j = 0, 1, \dots, r_k \text{ and } \nu, k = 0, 1, \dots, m,$$

$$\text{with } \delta_{jp} = \begin{cases} 1, & j = p \\ 0, & j \neq p. \end{cases}$$

We denote by

$$u(x) = \prod_{k=0}^m (x - x_k)^{r_k+1} \quad \text{and} \quad u_k(x) = \frac{u(x)}{(x - x_k)^{r_k+1}}.$$

We have

$$h_{kj}(x) = \frac{(x - x_k)^j}{j!} u_k(x) \sum_{\nu=0}^{r_k-j} \frac{(x - x_k)^\nu}{\nu!} \left[ \frac{1}{u_k(x)} \right]_{x=x_k}^{(\nu)}. \quad (2)$$

**Example 3** Find the Hermite interpolation polynomial for a function  $f$  for which we know  $f(0) = 1$ ,  $f'(0) = 2$  and  $f(1) = -3$  (equivalent with  $x_0 = 0$  multiple node of order 2 or double node,  $x_1 = 1$  simple node).

**Sol.** We have  $x_0 = 0, x_1 = 1, m = 1, r_0 = 1, r_1 = 0, n = m + r_0 + r_1 = 2$

$$\begin{aligned}(H_2 f)(x) &= \sum_{k=0}^1 \sum_{j=0}^{r_k} h_{kj}(x) f^{(j)}(x_k) \\ &= h_{00}(x)f(0) + h_{01}(x)f'(0) + h_{10}(x)f(1).\end{aligned}$$

We have  $h_{00}, h_{01}, h_{10}$ . These fulfill relations:

$$h_{kj}^{(p)}(x_\nu) = 0, \quad \nu \neq k, \quad p = 0, 1, \dots, r_\nu$$

$$h_{kj}^{(p)}(x_k) = \delta_{jp}, \quad p = 0, 1, \dots, r_k, \quad \text{for } j = 0, 1, \dots, r_k \text{ and } \nu, k = 0, 1, \dots, m.$$

We have  $h_{00}(x) = a_1 x^2 + b_1 x + c_1 \in \mathbb{P}_2$ , with  $a_1, b_1, c_1 \in \mathbb{R}$ , and the system

$$\begin{cases} h_{00}(x_0) = 1 \\ h'_{00}(x_0) = 0 \\ h_{00}(x_1) = 0 \end{cases} \Leftrightarrow \begin{cases} h_{00}(0) = 1 \\ h'_{00}(0) = 0 \\ h_{00}(1) = 0 \end{cases}$$

that becomes

$$\begin{cases} c_1 = 1 \\ b_1 = 0 \\ a_1 + b_1 + c_1 = 0. \end{cases}$$

Solution is:  $a_1 = -1, b_1 = 0, c_1 = 1$  so  $h_{00}(x) = -x^2 + 1$ .

We have  $h_{01}(x) = a_2x^2 + b_2x + c_2 \in \mathbb{P}_2$ , with  $a_2, b_2, c_2 \in \mathbb{R}$ . The system is

$$\begin{cases} h_{01}(x_0) = 0 \\ h'_{01}(x_0) = 1 \\ h_{01}(x_1) = 0 \end{cases} \Leftrightarrow \begin{cases} h_{01}(0) = 0 \\ h'_{01}(0) = 1 \\ h_{01}(1) = 0 \end{cases}$$

and we get  $h_{01}(x) = -x^2 + x$ .

We have  $h_{10}(x) = a_3x^2 + b_3x + c_3 \in \mathbb{P}_2$ , with  $a_3, b_3, c_3 \in \mathbb{R}$ . The system is

$$\begin{cases} h_{10}(x_0) = 0 \\ h'_{10}(x_0) = 0 \\ h_{10}(x_1) = 1 \end{cases} \Leftrightarrow \begin{cases} h_{10}(0) = 0 \\ h'_{10}(0) = 0 \\ h_{10}(1) = 1 \end{cases}$$

and we get  $h_{10}(x) = x^2$ .

The Hermite polynomial is

$$(H_2f)(x) = -x^2 + 1 - 2x^2 + 2x - 3x^2 = -6x^2 + 2x + 1.$$

**The Hermite interpolation formula is**

$$f = H_n f + R_n f,$$

where  $R_n f$  denotes the remainder term (the error).

**Theorem 4** *If  $f \in C^n[\alpha, \beta]$  and  $f^{(n)}$  is derivable on  $(\alpha, \beta)$ , with  $\alpha = \min\{x, x_0, \dots, x_m\}$  and  $\beta = \max\{x, x_0, \dots, x_m\}$ , then there exists  $\xi \in (\alpha, \beta)$  such that*

$$(R_n f)(x) = \frac{u(x)}{(n+1)!} f^{(n+1)}(\xi). \quad (3)$$

**Proof.** Consider

$$F(z) = \begin{vmatrix} u(z) & (R_n f)(z) \\ u(x) & (R_n f)(x) \end{vmatrix}.$$

$F \in C^n[\alpha, \beta]$  and there exists  $F^{(n+1)}$  on  $(\alpha, \beta)$ .

We have

$$F(x) = 0, \quad F^{(j)}(x_k) = 0, \quad k = 0, \dots, m; \quad j = 0, \dots, r_k;$$

because

$$u(x) = \prod_{k=0}^m (x - x_k)^{r_k+1} \Rightarrow u^{(j)}(x_k) = 0, \quad j = 0, \dots, r_k$$

and

$$(R_m f)^{(j)}(x_k) = f^{(j)}(x_k) - (H_n f)^{(j)}(x_k) = f^{(j)}(x_k) - f^{(j)}(x_k) = 0.$$

So,  $F$  and its derivatives have  $n + 2$  distinct zeros in  $(\alpha, \beta)$ . Applying successively Rolle's theorem it follows that  $F'$  has at least  $n + 1$  zeros in  $(\alpha, \beta) \Rightarrow \dots \Rightarrow F^{(n+1)}$  has at least one zero  $\xi \in (\alpha, \beta)$ ,  $F^{(n+1)}(\xi) = 0$ .

We have

$$F^{(n+1)}(z) = \begin{vmatrix} u^{(n+1)}(z) & (R_n f)^{(n+1)}(z) \\ u(x) & (R_n f)(x) \end{vmatrix},$$

with  $u(z) = \prod_{k=0}^m (z - z_k)^{r_k+1} \in \mathbb{P}_{n+1} \Rightarrow u^{(n+1)}(z) = (n + 1)!$ , and  $(R_n f)^{(n+1)}(z) = f^{(n+1)}(z) - (H_n f)^{(n+1)}(z) = f^{(n+1)}(z)$  (as,  $H_n f \in$

$\mathbb{P}_n$ ). We get

$$F^{(n+1)}(\xi) = \begin{vmatrix} (n+1)! & f^{(n+1)}(\xi) \\ u(x) & (R_n f)(x) \end{vmatrix} = 0,$$

whence it follows (3). ■

**Corollary 5** *If  $f \in C^{n+1}[a, b]$  then*

$$|(R_n f)(x)| \leq \frac{|u(x)|}{(n+1)!} \|f^{(n+1)}\|_{\infty}, \quad x \in [a, b]$$

where  $\|\cdot\|_{\infty}$  denotes the uniform norm ( $\|f\|_{\infty} = \max_{x \in [a, b]} |f(x)|$ ).

**Remark 6** *In case of  $m = 0$ , i.e.,  $n = r_0$ , (HIP) becomes **Taylor interpolation problem**. Taylor interpolation polynomial is*

$$(T_n f)(x) = \sum_{j=0}^n \frac{(x - x_0)^j}{j!} f^{(j)}(x_0).$$

**Example 7** *Find the Hermite interpolation formula for the function  $f(x) = xe^x$  for which we know  $f(-1) = -0.3679$ ,  $f(0) = 0$ ,  $f'(0) = 1$ ,*



$f(1) = 2.7183$ , (equivalent with  $x_0 = -1$  simple,  $x_1 = 0$  multiple of order 2 and  $x_2 = 1$  simple). Which is the limit of the error for approximating  $f(\frac{1}{2})$ ?