(3)	@	@ @	0 .	Mak	
Water = 1	pyp:	4.65	2.50	· Making Viscon	
0.765	NATER: (5050) (PVP Glycero 1	ip solute prP		
1	6 6 640 5)			suspensions.	
1.611 ml	4.018 m.2	32.2.67.8	Solutionaler (neight)	(hoider ve	
8 3		00	64+)	(besides water - ICP)	
				2	

40, 32,	pox objective, committee along drawter -	20x stjecture , 10 mm: # of pixels in a single particle & along a Odiantic: 80/13 18, 20, 21, 23, 21, 23 ang: 21	Calibration and Testing - made water # 10	2016/1/22
2, 24	white law	reds of	for testing.	BMC
AD, 32, 27, AB, 39, 3	Dans curted	x objective , 10 mm: of pixels in a single particle	mude water # 101 nm / palystyren	()
11	six article	3 . 23 . 23 . 23 . 23 . 23 . 23 . 23 .	1017	
32 47 47 64	Sixii counted	tricle	2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	
3.57×106 Piers 3.57×106 Piers 40.767 10.767	direct	alon	Lopen & polys	
1 1 2 8 8 6 1 8 8 6 1 8 8 8 6 1 8 8 8 8 8 8 8	# 210/4"	ang a od		
3/5 × ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~ ~	10.48 m	diavete	E S	28
3/3 2	7.5	1		

4 10	· v		4	1				4	
7.30	3 4 2	2, 7.30		to sin	4 10	W	2 -	made.	
7.3024e-13	7-30-24 ×10 7-2188 3-2950 3-9260 ×10-13 2-13 1-680 12-15 3-9266-13 3-9266-13	7.3024 e -13 7.29 88 3.2730e-15	£ 336-12 (simulated	2,30	4.65	H20	haly	
	3 3 4	-13 17.	12 12 12 12 12 12 12 12 12 12 12 12 12 1	0	2.50 p glywrol to,47 AM	1,65 d by 40 47 m	420 + 0,47 mm (diameter 2,50 ppp + 0,47 m	Polystynere	
7.32910-13	2188	2988	400		derived +	+	0,47 P+	20 11	
	, is g	W. 2	7.7701 e-15		4 + 1,01/4/1/4	0,43	67,0	inspensions for:	
3.2063 e-15	680	7300	016.		14/4	My 1	3 (2)	Inas	
3.2063 e-15	le-l	1 4	4 5		3		anet	for	
15	7 01						er		
5.0	4.0	1.18	1 10	B	To X			Sid	
5,0073 # e-13	4.0377 € -13	1.1515e-12	1.7864e	dexperimental.	X	F7.45	5440	Stides wood:	
0	1 -13	-12	र प्राचीत	inna					30