


## Lesson 5-Unit 2 – Derivatives (3)

### Derivatives of exponential and trigonometric functions

#### Review of Exponential Functions

The exponential function is defined as:  $y = f(x) = b^x$ ;  $b > 0, b \neq 1$ . The graph of the exponential function is represented below:



The x-axis ( $y = 0$ ) is a horizontal asymptote.

#### Number e

The number e is defined by:

$$e = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n \quad (1)$$

which can be written also as:

$$e = \lim_{u \rightarrow 0} (1 + u)^{\frac{1}{u}} \quad (2)$$

#### Exponential Function

The exponential function  $e^x$  may be evaluate using the limit:

$$e^x = \lim_{n \rightarrow \infty} \left(1 + \frac{x}{n}\right)^n \quad (3)$$

#### Derivative of $e^x$

$$(e^x)' = e^x \quad (4)$$

#### Proof

$$\lim_{h \rightarrow 0} \frac{e^{x+h} - e^x}{h} = \lim_{h \rightarrow 0} \frac{e^x(e^h - 1)}{h} = \left(\lim_{h \rightarrow 0} e^x\right) \left(\lim_{h \rightarrow 0} \frac{e^h - 1}{h}\right) = e^x$$

We used that  $\lim_{h \rightarrow 0} \frac{e^h - 1}{h} = 1$  ( substitute h by  $\pm 0.1, \pm 0.01, \pm 0.001$  to estimate  $\frac{e^h - 1}{h}$  ).

#### Derivative of $e^{f(x)}$

$$(e^{f(x)})' = e^{f(x)} f'(x).$$

Ex. Differentiate and simplify.

a.  $x^3 e^x$

$$(x^3 e^x)' = 3x^2 e^x + x^3 e^x = (3x^2 + x^3) e^x$$

b.  $e^{x^2+x+2}$

$$(e^{x^2+x+2})' = (2x+1) e^{x^2+x+2}$$

Derivative of  $b^x$ ,  $b > 0, b \neq 1$

$$(b^x)' = b^x \ln b$$


Proof

$$(b^x)' = (e^{x \ln b})' = e^{x \ln b} \ln b = b^x \ln b$$

## Review of Logarithmic Function

$$y = b^x \Leftrightarrow x = \log_b y$$

$$y = f(x) = \log_b y, \quad b > 0, b \neq 1, x > 0$$



Derivative of  $\ln x$

$$(\ln x)' = \frac{1}{x}$$

Proof

$$y = \ln x \Leftrightarrow x = e^y$$

$$x' = (e^y)' \Leftrightarrow 1 = e^y y' \Leftrightarrow y' = \frac{1}{e^y} \Leftrightarrow y' = \frac{1}{x}$$

Derivative of  $\log_b x$

$$(\log_b x)' = \frac{1}{x \ln b}$$

(Hint:  $\log_b x = \frac{\ln x}{\ln b}$ )

## Derivative of Trigonometric Functions

### Review of Trigonometric Functions

$$\begin{aligned} \sin x : \mathbb{R} &\rightarrow [-1, 1], & \sin(x + 2\pi) &= \sin x \\ \cos x : \mathbb{R} &\rightarrow [-1, 1], & \cos(x + 2\pi) &= \cos x \end{aligned}$$



Derivative of  $\sin x$  and  $\cos x$

$$(\sin x)' = \cos x$$

$$(\cos x)' = -\sin x$$

Proof

$$\begin{aligned} \lim_{h \rightarrow 0} \frac{\sin(x+h) - \sin x}{h} &= \lim_{h \rightarrow 0} \frac{\sin x \cos h + \sin h \cos x - \sin x}{h} = \left( \lim_{h \rightarrow 0} \sin x \right) \left( \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} \right) + \left( \lim_{h \rightarrow 0} \cos x \right) \left( \lim_{h \rightarrow 0} \frac{\sin h}{h} \right) \\ &= \cos x \end{aligned}$$

Ex. Differentiate.

a.  $x^2 \tan x$

b.  $\cot x$

c.  $\tan \sqrt{x^2 + 1}$

*Solution*

$$\begin{aligned} \text{a. } (x^2 \tan x)' &= 2x \tan x + x^2 (\tan x)' \\ &= 2x \tan x + x^2 \left( \frac{\sin x}{\cos x} \right)' \\ &= 2x \tan x + x^2 \left( \frac{\sin^2 x + \cos^2 x}{\cos^2 x} \right)' \\ &= 2x \tan x + \frac{x^2}{\cos^2 x} \end{aligned}$$

b.  $(\cot x)' = -\frac{1}{\sin^2 x}$

$$\begin{aligned} \text{c. } (\tan \sqrt{x^2 + 1})' &= \frac{1}{\cos^2 \sqrt{x^2 + 1}} (\sqrt{x^2 + 1})' \\ &= \frac{1}{\sqrt{x^2 + 1} \cos^2 \sqrt{x^2 + 1}} \end{aligned}$$